CASE REPORT

UPDATE Sirenomelia in a Cameroonian woman: a case report and review of the literature [version 2; peer review: 2 approved]

Frederick LI Morfaw, Philip N Nana

Department of Obstetrics and Gynaecology, Faculty of Medicines and Biomedical Sciences, University of Yaoundé, Yaoundé, Cameroon

v2 First published: 26 Jul 2012, 1:6
<https://doi.org/10.12688/f1000research.1-6.v1>
 Latest published: 06 Sep 2012, 1:6
<https://doi.org/10.12688/f1000research.1-6.v2>

Abstract

Sirenomelia is a rare congenital malformative disorder characterized by fusion of the lower limbs giving a characteristic mermaid-like appearance to the affected foetus. We report a case of sirenomelia occurring in a 19 year old Cameroonian woman following premature rupture of membranes and associated cord prolapse. This is the first documented case in this country. We highlight some of the cultural myths associated with this disorder and discuss our findings relative to the present literature and related controversies on its etiopathogenesis.

Open Peer Review

Approval Status

	1	2
version 2 (update) 06 Sep 2012		view
version 1 26 Jul 2012	view	view

1. **Laxmi Baxi**, Columbia University Medical Center, New York, NY, USA
2. **John Svigos**, University of Adelaide, Adelaide, Australia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Frederick LI Morfaw (ikomi_fred@yahoo.com)

Competing interests: The authors do not declare any competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2012 Morfaw FL and Nana PN. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Morfaw FL and Nana PN. **Sirenomelia in a Cameroonian woman: a case report and review of the literature [version 2; peer review: 2 approved]** F1000Research 2012, 1:6 <https://doi.org/10.12688/f1000research.1-6.v2>

First published: 26 Jul 2012, 1:6 <https://doi.org/10.12688/f1000research.1-6.v1>

UPDATED Changes from Version 1

This article has been revised for Version 2 to take into consideration the pertinent comments from the referees. In particular, statements presumed to be judgmental by referee Laxmi Baxi were either duly referenced or eliminated. The section under risk factors that confused genes with chromosomes has been reviewed and clarified. More emphasis has also been placed on scientific data in this version by including the reports from [Garrido *et al.* \(2011\)](#) on the possible role of genetic factors in sirenomelia from experimental work done in mice. Historical data has been removed from the report as requested by Dr Baxi, and identified repetitions were curtailed.

In response to the comments made by John Svigos, we have clarified the actual incidence of the condition (1 in 100,000 pregnancies), as well as the number of cases occurring in diabetic mothers (0.5–3.7%), and in twin pregnancies (15–20%). Finally, the Etiopathogenesis section has been shortened and made more precise to better fit a case report.

See referee reports

Introduction

Sirenomelia is an extremely rare congenital malformative disorder, which is often fatal¹. Its incidence is estimated at about 1 in 100,000 pregnancies², and cases have been reported from all ethnic groups worldwide³. This anomaly predominantly affects males (sex ratio 2.7:1)⁴, and is frequent among one of two monozygotic twins⁵. The most prominent yet inconstant feature of this malformative disorder is the complete or partial fusion of the lower limbs into a single lower limb³. The resultant infant bears a resemblance to the mermaid of ancient Greek mythology⁶. The disorder has equally been referred to as symmelia, sympodia monopodia, sympus, but most commonly as the ‘mermaid syndrome’ since the fusion of the lower limbs gives a characteristic mermaid-like appearance⁷.

In the African context, such mermaid-like babies are referred to as ‘mammy-water babies’, and bear an evil connotation associated with witchcraft and sorcery.

The underlying visceral anomalies are usually such that the syndrome is incompatible with life³, yet there are a number of reported cases of surviving infants with this condition in the English literature^{8–12}.

We report here the first documented case of sirenomelia in Cameroon, and discuss our findings in relation to the present literature and related controversies of its etiopathogenesis.

Case report

Miss N.N, 19 years old G2P0010, a single student at 38 weeks gestation according to her last menstrual period was referred to the Yaoundé Central Maternity for the management of cord prolapse at term. Her history revealed premature rupture of membranes 04 days prior to her consultation, with continuous per vaginal flow of clear liquor. She declares that foetal movements had been present prior to membrane rupture. She eventually developed uterine contractions 4 days later with an associated cord prolapse visible at the introitus.

This pushed her to consult at a health centre from where she was referred to our hospital.

In her past medical history, she was not diabetic nor did she bear any known chronic pathologies. There was no family history of diabetes nor malformations. Her first pregnancy two years earlier ended up in a clandestine voluntary termination of pregnancy by endo-uterine aspiration in the first trimester without any recorded complications. This pregnancy resulted from a non-consanguineous union with a 23 year old student. This pregnancy was very poorly followed up in a local health centre with two antenatal consultations. The few tests that were carried out did not reveal any anomalies, but no ultrasound was done. The evolution of the pregnancy so far had been uneventful without any history of teratogenic drug intake or traditional concoctions.

On admission the patient was hemodynamically stable, afebrile, with a symphysis-fundal height of 26 cm, absent foetal heart tones, a foetus in cephalic presentation, and a non-pulsating third degree cord prolapse. The working diagnosis of a prolonged rupture of membranes complicated by third degree cord prolapse and intrauterine foetal death was made. The patient was induced using oxytocin infusions, and was placed on prophylactic parenteral antibiotics. Labour evolved normally and within 5 hours she expelled a dead first degree macerated foetus with the following morphological abnormalities: (see [Figure 1–Figure 6](#)).

- Distended abdomen
- umbilical cord with a single artery
- Undetermined sex (absent external genitalia with only a 2x3 cm tag marking the position of the genitalia).
- Imperforate anus
- absence of a urinary meatus

Figure 1. Complete morphological view showing distended abdomen, fused lower limbs and maceration around the neck.

Figure 2. Complete morphological view of sirenomelic foetus.

Figure 4. Imperforate anus.

Figure 3. Posterior view showing fused lower limbs, imperforate anus.

Figure 5. Photograph of the groin area showing the undetermined external genitalia, and absent urinary meatus.

Figure 6. Photograph showing the fused feet with 11 toes.

- fused lower segment of the body below the pelvis into a single lower limb, with two feet fused posteriorly giving a single flipper-like foot with eleven toes spread out in a fan-like pattern (Mermaid-like or ‘mammy-water’). The foot was oriented anteriorly relative to the trunk, and external palpation gave the impression of probably two femurs and two tibias.

There was complete placental delivery and uterine revision was done removing membranous debris.

The birth weight of 2.3Kg at 38 weeks gestation reflected intra-uterine growth restriction.

Traditional and cultural beliefs precluded autopsy, and the corpse was handed over to the family for burial. The patient was maintained on parenteral antibiotics for 48 hours then oral relay was done on day 2 post-partum. She received adequate post-partum counselling and was discharged on that day.

Discussion

Sirenomelia or mermaid syndrome is an abnormal development of the caudal region of the body involving varying degrees of fusion of the lower limbs with or without bony defects¹³. It is usually associated with other visceral defects such as hypoplastic lungs, cardiac agenesis, absent genitalia, digestive defects, absent kidney and bladder, vertebral and central nervous system defects^{14,15}. Death usually results from obstructive renal failure due to renal agenesis or dysgenesis, with survival depending on adequate kidney functioning and renal outflow¹⁶.

In our patient however we could not ascertain the nature of the full foetal anomalies given that an autopsy was never performed, and no pre-birth imaging was available. Furthermore, the 3rd degree

cord prolapse was an obvious cause of fetal death. The history was suggestive of foetal movements in the few days prior to membrane rupture, hence this foetus could have been born alive.

Etiology and risk factors

The etiology of this multisystemic human malformation is unknown³ and no teratogens have been found in humans¹⁷. Recent evidence from mice models suggests it can have a genetic basis, yet much still needs to be done to define the role of candidate genetic factors in humans³. Certain risk factors however exist.

Maternal diabetes has been described as an important risk factor for caudal malformations in general³. However, with only about 0.5–3.7% of sirenomelia cases occurring in diabetic mothers^{13,18,19}, the association between maternal diabetes and sirenomelia has been described as weak²⁰. Our patient was not known to be diabetic.

The syndrome is also reported to be associated with twins, with about 15% to 20% of cases being derived from products of twin pregnancies^{5,13}, most of them monozygotic⁵. Reports actually indicate a 100–150 times higher incidence in monozygotic twins relative to dizygotic twins or singletons²⁰. In our patient, this was not a twin gestation, and there was no family history of twinning.

Also, exposure to heavy metals has been shown to be associated sirenomelia in humans^{22,23}.

It is worth noting that although Lynch *et al.*²⁴ recognised an autosomal form of caudal dysgenesis, no chromosomal abnormalities are found in sirenomelia and it does not recur in families¹⁴. This was a reassuring feature for our patient and should serve as a counseling feature for mothers bearing babies with this distressing anomaly.

Etiopathogenesis

The etiopathogenesis of this syndrome has been subject to a lot of debate over time. Numerous theories have been proposed to explain its origin.

Stevenson *et al.*²¹ proposed the vascular steal theory. This theory suggests that there is shunting of blood via an abnormal abdominal artery arising from high up in the aorta towards the placenta. This leaves the caudal part of the embryo poorly perfused. Consequently there is hypoplasia of the vasculature distal to the artery leading to nutritional deficiency of the caudal half of the body²³. Hence there may be complete/incomplete agenesis of the caudal structures (kidneys, sacrum, and lower portions of the digestive tract) except the gonads which are intra-abdominal. There could also be vertebral dysgenesis, lower limb atrophy and inconstant lower limb fusion¹⁴. The single umbilical artery in our patient favours this theory. However, Jaiyessimi *et al.*²⁰ reported a case of sirenomelia without this vitelline artery steal, indicating that factors other than vitelline artery steal could be responsible for sirenomelia in humans.

The defective blastogenesis theory regards sirenomelia as part of the caudal regression syndrome (CRS)³, more recently referred to as caudal dysgenesis^{25,26}. This is a rare congenital defect characterized

by a broad spectrum of lumbosacral dysgenesis. According to the this theory, during the ultimate stages of gastrulation occurring by the third week of intra-uterine life, there is a defect in blastogenesis leading to a wide range of phenotypic manifestations on the caudal extremity^{25,27}. Even though the syndrome was initially described by Duhamel²⁸ to include genitourinary and vertebral anomalies, phenotypic expression depends on the intensity, duration and initiation time of the underlying event³. Some authors consider sirenomelia to be the most extreme form of this relentless condition³.

However for authors such as Pinette *et al.*¹⁶, the distinction between caudal regression syndrome and sirenomelia remains speculative.

A further theory described in the literature regards sirenomelia as part of the VACTERL syndrome. VACTERL syndrome involves vertebral, anal, cardiovascular, tracheal esophageal, renal and limb dysgenesis. There is a major overlap in the phenotypic manifestations of sirenomelia and VACTERL²⁰. In most cases, the distinction between sirenomelia sequence and VACTERL lies within the severity of the component defects, and the single lower limb in sirenomelia can be regarded as an indicator of other severe malformations, especially in the gastrointestinal and genito-urinary systems²⁰. In our patient the lack of an autopsy did not permit any assertions to be made as to the relationship with VACTERL.

Other theories exist, but given their controversies in the literature, they are not considered here. Yet the overlap in these syndromes/theories waters the debate in the scientific world as to the uniqueness or diversities of these syndromes.

Classification

Stocker and Heifetz¹³ classified the sirenomelia sequence into 7 types as shown be in Table 1.

We did not have any radiographs and could therefore not classify our patient into any of these categories with certainty even though external palpation was in favour of a type I.

Table 1. Classification of the Sirenomelic sequence adapted from Stocker and Heifetz¹³.

Type	Characteristic
I	All thigh and leg bones are present
II	Single fibula
III	Absent fibula
IV	Partially fused femurs, fused fibulae
V	Partially fused femurs
VI	Single femur, single tibia
VII	Single femur, absent tibia

Diagnosis

The past medical history of the patient could identify patients at risk. Antenatal diagnosis is possible on ultrasound and x-ray. Ultrasound is however the predominant diagnostic tool. Moreover, there

is no open defect which could cause abnormal increases in alpha foeto-protein levels²⁹.

Ultrasound features permitting confirmation of the diagnosis include lack of tibia/fibula, a single femur, convergent femoral bones, bilateral renal agenesis, polycystic kidneys/renal agenesis, obstructive uropathy and intra-uterine growth retardation³⁰. The use of ultrasound in diagnosis is however not without difficulty. In sirenomelic fetuses, bilateral renal agenesis causes severe oligohydramnios thus limiting ultrasound evaluation of the limbs in the second and third trimesters^{31,32}. However, in earlier gestational ages, the amniotic fluid volume may be sufficient to detect abnormal lower limbs. In such cases, we may notice in addition to abnormal lower limbs, bilateral renal dysgenesis, absent bladder, undetermined external genitalia, anorectal agenesis and lumbosacral agenesis⁴. Other abnormalities may touch the cardiovascular system and abdominal walls.

Antenatal confirmation of the diagnosis justifies a therapeutic termination of the pregnancy.

At delivery, clinical evaluation is usually sufficient to confirm the diagnosis. In our case, the diagnosis was obvious given the fused lower limbs, single umbilical artery, and imperforate anus. However radiographic images are important if we are to be able to classify the condition according to the Stocker and Heifer classification (Table 1)¹³. An autopsy permits determination of the extent of the associated anomalies, but is usually of limited use in the African context where cultural norms and beliefs largely precludes its practice.

Management and prognosis

Sirenomelia carries with it a very poor prognosis. Survival is largely dependent on the extent of visceral anomalies, especially obstructive renal failure due to renal agenesis/dysgenesis¹⁶. In the case of antenatal diagnosis, a voluntary termination of pregnancy is advisable in order to avoid the physical and psychological stress to parents and the family. This decision however depends on the gestational age of the pregnancy, the severity of the malformations and of course the desires of the parents¹⁷.

Recent reports indicate that about 50% of these infants are born alive after 8–9 months gestation¹⁷. However most of them die within 5 days of life¹³. The management of sirenomelia is difficult and expensive, and the outcome is unpredictable¹⁷. The main therapeutic modality involves surgical and medical compensation aimed mainly at maintaining adequate renal function. Surgery to correct the anomaly and separate the fused limbs is usually not a priority as there is no guarantee of its success, and it carries with it an increased risk of compromising the life of an already delicate infant.

There have been reports of surviving sirenomelic fetuses^{8–12}. Pertinent amongst these is the case of the surviving infant with sirenomelia associated with absent bladder reported by Stanton *et al.*¹². This infant underwent 5 surgeries before the age of 4 years and continues to be bedridden and dependent. In the case described by Pinette *et al.*¹⁶, the infant received a renal transplant using a cadaveric donor

kidney. By publication time, this I infant was 5 years old with normal renal and cognitive development for her age. However, separation of the lower limbs was indefinitely delayed due to concerns regarding disruption of blood supply to abdominal organs and the transplanted kidney.

These reflect the constraints, both financial and physical to conservatively manage sirenomelia and reiterate the importance of antenatal diagnosis and voluntary termination of pregnancy especially in our resource limited settings like ours.

Conclusion

Sirenomelia remains a rare but peculiar syndrome. Controversies on its etiopathogenesis persist. Its antenatal diagnosis is possible albeit difficult by ultrasound. The associated visceral anomalies are usually incompatible with life. However surviving sirenomelic foetuses have been described with costly conservative management and mediocre results. In the African context, these mermaid-like foetuses are described as 'mammy-water babies'. This carries with it the connotation of sorcery and witchcraft with which no family wishes to be identified. Therefore antenatal diagnosis and termination of pregnancy is advisable. Knowledge of this rare syndrome is

important to dissipate cultural myths whenever it occurs, and free the family from stigmatization.

Consent

Written informed consent for publication of their clinical details and/or clinical images was obtained from the patient/parent/guardian/relative of the patient.

Author contributions

F.M received the patient and discussed the case history and management with P.N. F.M wrote the first draft of the article. F.M and P.N read and revised several versions of the manuscript. Both authors read and approved the final manuscript.

Competing interests

The authors do not declare any competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. Taori KB, Mitra K, Ghonga NP, *et al.*: **Sirenomelia sequence (mermaid): report of three cases.** *Indian J Radiol Imaging.* 2002; **12**: 399–401.
[Reference Source](#)
2. Martinez-Frias ML, Garcia A, Bermejo E, *et al.*: **Cyclopia and sirenomelia in a liveborn infant.** *J Med Genet.* 1998; **35**(3): 263–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Garrido-Allepuz C, Haro E, Gonzalez-Lamuno D, *et al.*: **A clinical and experimental overview of sirenomelia: insight into the mechanisms of congenital limb malformations.** *Dis Model Mech.* 2011; **4**(3): 289–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Valenzano M, Paoletti R, Rossi A, *et al.*: **Sirenomelia. Pathological features, antenatal ultrasonographic clues, and a review of current embryogenic theories.** *Hum Reprod Update.* 1999; **5**(1): 82–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Di LM, Brandt ML, Veilleux A, *et al.*: **Sirenomelia in an identical twin: a case report.** *J Pediatr Surg.* 1991; **26**(11): 1334–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Messineo A, Innocenti M, Gelli R, *et al.*: **Multidisciplinary surgical approach to a surviving infant with sirenomelia.** *Pediatrics.* 2006; **118**(1): e220–e223.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Kulkarni ML, Sureshkumar C, Sindhur PS, *et al.*: **Sirenomelia with spina bifida.** *Indian Pediatr.* 1994; **31**(1): 51–5.
[PubMed Abstract](#)
8. Savader SJ, Savader BL, Clark RA, *et al.*: **Sirenomelia without Potter syndrome: MR characteristics.** *J Comput Assist Tomogr.* 1989; **13**(4): 689–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Murphy JJ, Fraser GC, Blair GK, *et al.*: **Sirenomelia: case of the surviving mermaid.** *J Pediatr Surg.* 1992; **27**(10): 1265–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Clarke LA, Stringer DA, Fraser GC, *et al.*: **Long term survival of an infant with sirenomelia.** *Am J Med Genet.* 1993; **45**(3): 292–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. McCoy MC, Chescheir NC, Kuller JA, *et al.*: **A fetus with sirenomelia, omphalocele, and meningomyelocele, but normal kidneys.** *Teratology.* 1994; **50**(2): 168–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Stanton MP, Penington EC, Hutson JM, *et al.*: **A surviving infant with sirenomelia (Mermaid syndrome) associated with absent bladder.** *J Pediatr Surg.* 2003; **38**(8): 1266–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Stocker JT, Heifetz SA: **Sirenomelia. A morphological study of 33 cases and review of the literature.** *Perspect Pediatr Pathol.* 1987; **10**: 7–50.
[PubMed Abstract](#)
14. Tang TT, Oechler HW, Hinkle DH, *et al.*: **Limb body-wall complex in association with sirenomelia sequence.** *Am J Med Genet.* 1991; **41**(1): 21–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Rodriguez JL, Palacios J, Razquin S, *et al.*: **Sirenomelia and anencephaly.** *Am J Med Genet.* 1991; **39**(1): 25–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Pinette MG, Hand M, Hunt RC, *et al.*: **Surviving sirenomelia.** *J Ultrasound Med.* 2005; **24**(11): 1555–9.
[PubMed Abstract](#)
17. Fadhlou A, Khrouf M, Gaigi S, *et al.*: **The sirenomelia sequence: a case history.** *Clin Med Insights Case Rep.* 2010; **3**: 41–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Duncan PA, Shapiro LR, Klein RM, *et al.*: **Sacroccygeal dysgenesis association.** *Am J Med Genet.* 1991; **41**(2): 153–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Lynch SA, Wright C: **Sirenomelia, limb reduction defects, cardiovascular malformation, renal agenesis in an infant born to a diabetic mother.** *Clin Dysmorphol.* 1997; **6**(1): 75–80.
[PubMed Abstract](#)
20. Jaayesimi F, Gomathinayagam T, Dixit A, *et al.*: **Sirenomelia without vitelline artery steal.** *Ann Saudi Med.* 1998; **18**(6): 542–4.
[PubMed Abstract](#)
21. Stevenson RE, Jones KL, Phelan MC, *et al.*: **Vascular steal: the pathogenic mechanism producing sirenomelia and associated defects of the viscera and soft tissues.** *Pediatrics.* 1986; **78**(3): 451–7.
[PubMed Abstract](#)
22. Castilla EE, Mastroiacovo P, Lopez-Camelo JS, *et al.*: **Sirenomelia and cyclopia cluster in Cali, Colombia.** *Am J Med Genet A.* 2008; **146A**(20): 2626–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Orioli IM, Mastroiacovo P, Lopez-Camelo JS, *et al.*: **Clusters of sirenomelia in South America.** *Birth Defects Res A Clin Mol Teratol.* 2009; **85**(2): 112–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Lynch SA, Bond PM, Copp AJ, *et al.*: **A gene for autosomal dominant sacral agenesis maps to the holoprosencephaly region at 7q36.** *Nat Genet.* 1995; **11**(1): 93–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Opitz JM, Zanni G, Reynolds JF Jr, *et al.*: **Defects of blastogenesis.** *Am J Med Genet.* 2002; **115**(4): 269–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Thottungal AD, Charles AK, Dickinson JE, *et al.*: **Caudal dysgenesis and sirenomelia-single centre experience suggests common pathogenic basis.** *Am J Med Genet A.* 2010; **152A**(10): 2578–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Duesterhoeft SM, Ernst LM, Siebert JR, *et al.*: **Five cases of caudal regression**

- with an aberrant abdominal umbilical artery: Further support for a caudal regression-sirenomelia spectrum. *Am J Med Genet A*. 2007; **143A**(24): 3175–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Duhamel B: **From the Mermaid to Anal Imperforation: The Syndrome of Caudal Regression.** *Arch Dis Child*. 1961; **36**(186): 152–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Davari F, Googol N, Kaveh M, *et al.*: **Sirenomelia (mermaid syndrome) in an infant of a diabetic mother.** *Acta Medica Iranica*. 2003; **41**(1): 69–72.
[Reference Source](#)
 30. Sirtori M, Ghidini A, Romero R, *et al.*: **Prenatal diagnosis of sirenomelia.** *J Ultrasound Med*. 1989; **8**(2): 83–8.
[PubMed Abstract](#)
 31. Chenoweth CK, Kellogg SJ, Abu-Yousef MM, *et al.*: **Antenatal sonographic diagnosis of sirenomelia.** *J Clin Ultrasound*. 1991; **19**(3): 167–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. van Zalen-Sprock MM, van Vugt JM, van der Harten JJ, *et al.*: **Early second-trimester diagnosis of sirenomelia.** *Prenat Diagn*. 1995; **15**(2): 171–7.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 24 September 2012

<https://doi.org/10.5256/f1000research.550.r259>

© 2012 Svigos J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

John Svigos

Discipline of Obstetrics & Gynaecology, School of Paediatrics & Reproductive Health, Faculty of Health Sciences, University of Adelaide, Adelaide, Australia

The article reads much better now, and it represents the most comprehensive current review of sirenomelia in the available literature.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 27 July 2012

<https://doi.org/10.5256/f1000research.102.r258>

© 2012 Svigos J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

John Svigos

Discipline of Obstetrics & Gynaecology, School of Paediatrics & Reproductive Health, Faculty of Health Sciences, University of Adelaide, Adelaide, Australia

This is an interesting article with some quaint comments indicative of local community attitudes to bizarre abnormalities, which increase the appeal of the article.

The author(s) should consider being more specific about the actual incidence of the abnormality itself and the actual incidence associated with twins and diabetes. The author(s) should also review the level of detail under the 'Ethiopathogenesis' and 'Classification' subheadings, as I believe that these sections should be shortened/abbreviated if it is to fulfill the criteria of a 'Case Report' presentation.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 12 Sep 2012

Frederick Morfaw, Department of Obstetrics and Gynaecology, Faculty of Medicines and Biomedical Sciences, University of Yaoundé, Cameroon

Thank you for your very pertinent comments. With these taken on board we have now reviewed the work and made changes accordingly in a new version of the article.

Competing Interests: No competing interests were disclosed.

Reviewer Report 26 July 2012

<https://doi.org/10.5256/f1000research.102.r257>

© 2012 Baxi L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Laxmi Baxi

Department of Obstetrics and Gynecology, Columbia University Medical Center, New York, NY, USA

This is an interesting article. However there are a few things that need addressing:

1. There should be no judgmental statements.
2. The author(s) should not confuse between gene and chromosomes.
3. There should be more emphasis on scientific data and information.
4. If the author(s) do need to address historical data, then it should be in full.

It is too bad there isn't any pre- or post-birth imaging, however the photographs presented are interesting. There also seems to be a repetition of some facts, which need to be curtailed.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard.

Author Response 12 Sep 2012

Frederick Morfaw, Department of Obstetrics and Gynaecology, Faculty of Medicines and Biomedical Sciences, University of Yaoundé, Cameroon

Thank you for your very pertinent comments. With these taken on board we have now reviewed the work and made changes accordingly in a new version of the article.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research