Check for updates

RESEARCH ARTICLE

Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study [version 1; peer review: 1 approved, 1 approved with reservations, 1 not approved]

Bikal Shrestha¹, Stuti Yadav^{id}¹, Subodh Dhakal^{id}¹, Pooja Ghimire¹, Yubika Shrestha¹, Ela Singh Rathaure^{id}²

¹Nepalese Army Institute of Health Sciences (NAIHS), Kathmandu, 44600, Nepal

²Epidemiology and Disease control division, USAID, Kathmandu, 44600, Nepal

V1 First published: 11 Oct 2021, 10:1031
<https://doi.org/10.12688/f1000research.72824.1>
 Latest published: 11 Apr 2022, 10:1031
<https://doi.org/10.12688/f1000research.72824.2>

Abstract

Background: Self-esteem is vital to living a happy, confident and content life. Medical students experience various forms of stress due to academic, financial and social pressures which could affect their levels of self-esteem. This study aims to study the status of self-esteem among undergraduates of a medical college at Tribhuvan University, Nepal.

Methods: After receiving the ethical approval from the Institutional Review Committee (IRC) of NAIHS; we conducted a descriptive cross-sectional study among the first- to fifth-year medical students from December 2020 to April 2021. 190 were selected for the study using a stratified random sampling technique. This study used the Rosenberg self-esteem scale to measure self-esteem of the participants. A Google Forms questionnaire was sent to the participants via email. Then, the data obtained were entered in the Google sheet and later analyzed using SPSS 27. A Chi-square test was used to identify potential differences in self-esteem scores among different variables. A p-value of < 0.05 was considered statistically significant.

Results: This study included a total of 180 participants, among which, 18.9% (13.19% to 24.61%; at 95% CI) students showed low self-esteem. 74.4% (68.02% to 80.78%; at 95% CI) students had normal self-esteem and 6.7% (3.05% to 10.35%; at 95% CI) students had high self-esteem. The mean self-esteem score was 19.19 (15.01 to 23.37; at 95% CI). Female participants suffered more than males from low self-esteem, and third-year students had the highest percentage of low self-esteem (30.77%).

Conclusion: The majority (74.4%) of medical students had normal self-

Open Peer Review

Approval Status

	1	2	3
version 2
(revision)
11 Apr 2022
version 1
11 Oct 2021

1. **Dr. Richard Mottershead** ^{id}, RAK Medical and Health Sciences University, Ras Al Khaimah, United Arab Emirates
2. **Christian Sunday Ugwuanyi** ^{id}, University of Nigeria, Nsukka, Nigeria
3. **Sunil Kumar Joshi** ^{id}, Kathmandu Medical College, Kathmandu, Nepal

Any reports and responses or comments on the article can be found at the end of the article.

esteem. However, 18.9% students had low self-esteem, among which, third-year students suffered the most (30.77%). Likewise, females exhibited higher prevalence of low self-esteem compared to males. Interventions to boost the level of self-esteem should be carried out to help medical students become confident and efficient doctors.

Keywords

self-esteem, college, medical students, Kathmandu, Nepal, undergraduates, doctors, self-perception

Corresponding author: Stuti Yadav (stuti4542@gmail.com)

Author roles: **Shrestha B:** Conceptualization, Software, Supervision, Visualization, Writing – Review & Editing; **Yadav S:** Conceptualization, Investigation, Methodology, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Dhakal S:** Conceptualization, Investigation, Methodology, Software, Writing – Original Draft Preparation, Writing – Review & Editing; **Ghimire P:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Shrestha Y:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Singh Rathaur E:** Conceptualization, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Shrestha B *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Shrestha B, Yadav S, Dhakal S *et al.* **Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study [version 1; peer review: 1 approved, 1 approved with reservations, 1 not approved]** F1000Research 2021, 10:1031 <https://doi.org/10.12688/f1000research.72824.1>

First published: 11 Oct 2021, 10:1031 <https://doi.org/10.12688/f1000research.72824.1>

Introduction

Self-esteem is one's sense of self-worth, acceptance, and confidence that individuals feel toward themselves.¹ Mentally and physically active people are the foundation of a healthy society and serve the community. Doctors and medical students in particular must have a healthy mental state. A doctor suffering from anxiety, dissatisfaction, and low self-esteem will not be able to fulfil his/her duty properly,² and a medical student with low self-esteem will never grow to become a healthy, efficient doctor.

Having balanced self-esteem is highly important for healthcare professionals, including medical students.³ Medical students undergo major stress in the form of academic workload,⁴ family expectations, peer pressure, financial issues^{5,6} and excessive working hours.⁵⁻⁷ Studies show that a large proportion of medical students are unhappy and have low self-esteem,⁸ and that students with low self-esteem are likely to suffer from anxiety and mental breakdowns, doubt themselves and feel a sense of inadequacy.⁹ These students tend to appear less competent among their peers¹⁰ which contributes to depression and unhappiness. Self-development and well-being is also negatively affected by inadequate self-esteem.⁸

Self-esteem influences a student's performance, as well as their physical and psychological health and therefore is important to be maintained. A person who has high self-esteem can cope more effectively with stress and anxiety-inducing events in life without any negativity or impact on their mental health.¹¹ It helps one to foster confidence, appear competent among their colleagues, tackle challenges with optimism and take on criticism without it affecting their self-image.⁸

In Nepal, studies evaluating the self-esteem of medical students are lacking, but existing literature studying nursing students found that nearly 78% of the students had low self-esteem.¹² Producing healthy doctors is a crucial part of human resource planning.¹³ Hence, assessing their self-esteem and designing necessary interventions for students with inadequate self-esteem can help us produce confident and optimistic doctors. Since confidence is regarded as a subjective marker of competence,¹⁴ we can expect to produce highly competent doctors from the timely intervention. Having this competency in student life will help them acquire good knowledge and later, as a doctor, enable them to effectively manage patients. This study aims to report the status of self-esteem among medical students at a college in Kathmandu.

Methods

Study design and setting

A descriptive cross-sectional study was conducted from December 2020 to April 2021. Data collection took place from 30th of December, 2020 to 25th of January, 2021 among medical students at the Nepalese Army Institute of Health Sciences-College of Medicine (NAIHS-COM) situated in Kathmandu. Medical undergraduates from all years of study were included in the study. The study was conducted after receiving ethical approval from the Institutional Review Committee (IRC), NAIHS; on December 2020 (Reg. No. 373).

Sample size

We calculated the adequate sample size using the formula for infinite population and then adjusted it according to our population size. Based on Syed *et al's* study, we estimated the proportion of students with low self-esteem to be 18%.¹⁵ Using this estimation, we calculated the minimum sample size that could represent our population.

$$N = z^2 \times p \times (1 - p) / d^2$$

[where, N = sample size for infinite population, z = factor to achieve (1 - a) % level of confidence (z = 1.96 at 95% C.I) p = estimated rate of population proportion i.e. 18% here.¹⁵ d = margin of error to be tolerated (5% here)]

$$N = (1.96)^2 \times 0.18 \times (1 - 0.18) / 0.05^2$$

$$N = 226.80 \quad \text{i.e. 227}$$

For our study population, adjusted sample size is:

$$n = N / (1 + (N - 1) / P)$$

[where, n = adjusted sample size for our finite population, P = our study population i.e. 550 students.]

$$n = 227 / (1 + 226 / 550) = 160.89 \quad \text{i.e. 161}$$

Table 1. Socio-demographic characteristics and RSES of the students along with p-value.

Variables		RSES SCORE			Total	p-value
		< 16	16 – 25	>25		
Sex	Male	22 (17.89%)	91 (73.98%)	10 (8.13%)	123	0.037
	Female	12 (21.05%)	43 (75.44%)	2 (3.51%)	57	
Type of primary schooling	Government	2 (10%)	16 (80%)	2 (10%)	20	0.508
	Private	32 (20%)	118 (73.75%)	10 (6.25%)	160	
Year of medical college	First year	6 (17.65%)	26 (76.47%)	2 (5.88%)	34	0.660
	Second year	5 (16.13%)	23 (74.19%)	3 (9.68%)	31	
	Third year	12 (30.77%)	25 (64.10%)	2 (5.13%)	39	
	Fourth year	5 (14.71%)	28 (82.35%)	1 (2.94%)	34	
	Fifth year	6 (14.29%)	32 (76.19%)	4 (9.52%)	42	
Level of education	Pre-clinical	11 (16.92%)	49 (75.38%)	5 (7.69%)	65	0.972
	Clinical	23 (20%)	85 (73.91%)	7 (6.09%)	115	
Province	Province No. 1	1 (9.09%)	9 (81.82%)	1 (9.09%)	11	0.939
	Province No. 2	9 (26.47%)	23 (67.65%)	2 (5.88%)	34	
	Bagmati Province	13 (16.25%)	61 (76.25%)	6 (7.5%)	80	
	Gandaki Province	5 (20.83%)	18 (75.00%)	1 (4.17%)	24	
	Lumbini Province	5 (21.74%)	16 (69.57%)	2 (8.7)	23	
	Karnali Province	0 (0%)	3 (100%)	0 (0%)	3	
	Sudurpashchim Province	1 (20.00%)	4 (80.00%)	0 (0%)	5	

Hence, our sample size (n) is 161.

Adjusting the sample size by taking non-response rate at 15%, the final sample size becomes 190.

Sampling strategy and eligibility criteria

Students 18 years or older who gave consent were eligible to participate in the study. For the purpose of sampling, students were divided into two strata: pre-clinical (first, second) and clinical year (third, fourth, fifth). This was done because students of pre-clinical years shared some common characteristics – such as studying basic medical subjects without clinical exposure, while students of clinical years had clinical exposure. Students were selected through a Proportionate Stratified Random sampling technique so that every student from the first to fifth year had an equal probability of being selected in the study. This would ensure that the sample within each stratum properly represented our study population. The sampling frame was prepared by collecting the list of students from administrative section of the institute, who were numbered from 1 to 550. Within that limit, 190 random numbers were generated with the help of Google Random Number Generator as per the proportion of students from each year and the proportion of males and females in each year. The random numbers that would represent the student researchers involved in this study were excluded.

Survey instrument

To measure self-esteem, we used a questionnaire based on the Rosenberg Self Esteem Scale (RSES).^{16,17} The questionnaire had two parts; demographic details and the Rosenberg Self Esteem Scale. Demographic details covered general information including: age, sex, province of permanent address, year of study and type of institution they had attended for primary level education. A copy of the questionnaire can be found in the extended data.¹⁸

The RSES was developed by sociologist Morris Rosenberg.¹⁶ It is a 10-item Likert-type scale with items answered on a four-point scale from strongly agree to strongly disagree. For items 1, 3, 4, 7, 10: Strongly Agree = 3, Agree = 2, Disagree = 1, and Strongly Disagree = 0. For items 2, 5, 6, 8, 9 (which are negative statements; hence reversed in score): Strongly Agree = 0, Agree = 1, Disagree = 2, and Strongly Disagree = 3.^{16,18} A total is obtained by adding these markings which ranges from 0–30. A score less than 16 indicates low self-esteem, more than 25 indicated high self-esteem and scores from 16 to 25 shows normal self-esteem in the respondents.¹⁷

The RSES has high reliability: test-retest correlations are in the range of .82 to .88 and Cronbach's alpha for various samples are in the range of .77 to .88.^{19,20} We also calculated Cronbach's alpha for our study sample and the value for it was 0.811 which showed good internal consistency and reliability of the instrument for our population as well.

Data collection and statistical analysis

The questionnaire was sent to 190 students through Google forms via email. Out of 190 students, 182 responded among which 2 refused to give consent. Hence, we obtained complete data from 180 students which was then entered into a Google spreadsheet¹⁸ and later analyzed using Microsoft Excel and SPSS version 27. Chi Square Test of Association was used to compare between and among different categorical variables by taking the median values of RSES score. A p-value of < 0.05 was considered statistically significant.

Potential biases

The participant responses could have been subjected to response bias and subjective bias. Response bias includes responding to questionnaire items without understanding them and also giving socially desirable answers. Subjective bias could have occurred as the questionnaire is based on the subjective feelings of the respondents. We could not control subjective bias as it is inevitable when measuring abstract parameters such as self-esteem. But for response bias, we employed certain techniques to minimize it as far as possible. The questionnaire was self-administered, without individual interpretation by the researcher, but with explanation for some possible ambiguities in it. We checked the clarity of the items by translating the items to Nepali and retranslating it back to English with the help of two individuals who were not involved in the study. Moreover, we assured participants regarding concealed identity and confidentiality issues at the beginning and the Likert scale used was a 4-point Likert without the neutral responding.

Ethical statement

Ethical approval was obtained from the Institutional Review Committee (IRC), NAIHS in December 2020 (Reg. No. 373). There was no patient or public involvement in the design, conduct or reporting of our research. The first section of our Google forms survey had an information sheet with information about our study and an informed consent section where the participants could choose to agree or disagree to participate in the study via a tick-box. Participants who chose to agree could proceed to the questionnaire section. They were informed that their identity would be concealed and their de-identified data would be accessed only by the researchers. In this way, only after obtaining informed consent, the randomly selected medical students were involved as participants for data collection.

Results

180 responses were received from 190 study participants (94.73% response rate). Out of the 180 participants, 123 (68.3%) were male and 57 (31.7%) were female. Majority of the respondents were from clinical years (third, fourth and fifth year) making up 63.5%, whereas only 36.1% respondents were from Pre-Clinical years (first and second year). The average age of the students was 22.17 ± 1.78 years. The mean self-esteem score of the respondents was 19.19 ± 4.18 (15.01 to 23.37 at 95% CI) which lies in the normal range of RSES score.

Among 180 participants, 18.9% (13.19 to 24.61 at 95% CI) were found to have low self-esteem. 74.4% (68.02% to 80.78% at 95% CI) students had normal self-esteem and 6.7% (3.05% to 10.35% at 95% CI) students had high self-esteem.

Female participants were found to have suffered lower self-esteem than males. The mean self-esteem score of males was 19.63 ± 4.29 and that of females was 18.26 ± 3.81 . There is an association between sex and median values of RSES scores (i.e., whether students had higher or lower self-esteem than the median value). Significant difference in self-esteem was found between males and females at 5% level of significance ($\chi^2 = 4.33$, $p = 0.037$). Not enough evidence was found to suggest a significant difference in RSES scores between and among any other variables. Out of 65 pre-clinical year students, 11 (16.92%) and out of 115 clinical year students 23 (20%) were found to have low self-esteem. Third year medical students had the highest percentage of low self-esteem i.e., 12 (30.77%) as shown in Table 1.

It was found that 20% (14.16% to 25.84% at 95% CI) of the students who had attended private school for their primary level of schooling had low self-esteem whereas its prevalence among government school students was reported to be 10% (5.62% to 14.38% at 95% CI). Further details of the findings have been presented in Table 1.

Discussion

Self-esteem refers to an individual's overall sense of value or worth.¹⁹ Various factors including age,²¹ sex,^{22,23} socio-economic status,^{23,24} level of study,²⁵ academic stress¹² and family environment²⁶ influence self-esteem. A reasonably high degree of self-esteem is considered an attribute of good mental health,²⁷ and students with high self-esteem are found

to have better academic achievements compared to those with low self-esteem.²⁸ So, in order to improve the academic performance of medical students, healthy self-esteem is imperative. Moreover, healthcare professionals with high self-esteem show higher tendency to influence and induce positive well-being not only in patients but also in the healthcare team.²⁹ Empathy is an essential element of the physician-patient relationship and is also an essential quality for medical students.³⁰ A medical student with low self-esteem is likely to grow less empathetic since self-esteem and empathy are positively correlated.³⁰ Low self-esteem has also been allied with unhappiness,³¹ suicidal ideation,³² isolation and avoidance of social settings³³ and psychological disorders.³⁴

Numerous studies have been conducted on self-esteem among the general population but there have been limited studies conducted among medical students. According to studies carried out by Syed, *et al.* and Nair, *et al.* among medical students, the prevalence of low self-esteem was found to be 18%¹⁵ and 19.4%³⁵ respectively. According to our study, the prevalence of low self-esteem among the medical students of NAIHS-COM was found to be 18.9% which was similar to the above mentioned studies. This prevalence of low self-esteem may be due to high academic stress,¹² low perceived family support¹² and learning environment.²⁴ Since not many studies have been conducted on self-esteem among medical students in Nepal, other contributing factors are yet to be explored. 74.4% of the students had normal levels of self-esteem. Having high but realistic self-esteem is important for sound mental health. But considering the potential for subjective bias in this study, it cannot be implied that the high scores (>25) attained by the respondents are an accurate reflection of their actual self-esteem status, and not a subjective, narcissistic trait. The prevalence of high self-esteem was reported to be 6.7% in our study. Studies among Arab participants suggest that self-esteem positively and significantly linked with attitude towards life, mental health, and general satisfaction.³⁶⁻⁴¹ People with high self-esteem are expected to persist in the face of challenges than those with low self-esteem.⁴² But it is also suggested that people with high self-esteem are more likely to be conceited, arrogant and sometimes narcissistic.⁴³

The mean RSES score of males was found to be significantly higher at 19.63 ± 4.29 , whereas females had a score of 18.26 ± 3.8 . Studies conducted among Malaysian and Indian students showed no significant difference between males and females.^{44,45} A study done among four Arab counties showed Kuwaiti men had a significantly higher mean score than their female peers.⁴⁰ A study conducted in Loni, India also showed higher prevalence of low self-esteem in males than females,¹⁵ whereas our study indicated that low self-esteem was more prevalent among females.

In our study, significant differences between self-esteem were not found among clinical and pre-clinical year students. However, third year medical students had the highest prevalence of low self-esteem. This could be due to lack of adaptation to academic burden while transitioning from pre-clinical phase to clinical phase. A study conducted in India by Syed *et al* had a dissimilar finding that the cohort of first year males had the highest prevalence of low self-esteem.¹⁵ In a study conducted by Naz, *et al*, the highest prevalence of low self-esteem was found in fourth year medical students.⁴⁶ As the third year medical students had the highest prevalence of low self-esteem, proper orientation and pre-exposure sessions should be conducted before entering the clinical phase. Psychological counselling should be frequently provided to all medical students, focusing more on the students with low self-esteem. Medical students should also be encouraged to participate in extracurricular activities which may help in boosting their self-esteem. This is because involvement in extracurricular activities can help students achieve personal identity development and a sense of self-worth⁴⁷ which will in turn, add to their self-esteem.

A Nigerian study showed that self-esteem is positively and significantly correlated with the primary school attended.²⁴ The self-esteem of medical students who attended private schools was significantly higher than those of public school students.^{24,48} In our study, we tried to figure out whether there was any difference in self-esteem between students who had attended government schools for primary level of schooling and those who had attended private schools. Although we found that a proportionately higher number of students who had acquired primary levels of education through private schools had low self-esteem, the difference was not significant. We propose that this could be because the majority of the students (88.9%) were from private schools and a comparatively higher percentage of low self-esteem was seen among them. But the statistical significance of this difference was not obtained which could have been due to the very low number of students from government schools for the comparison.

We were also interested to know if the permanent address of students created an impact on their self-esteem since not all parts of Nepal are equally developed and students from some areas may be deprived of opportunities. But we did not find any significant difference in the self-esteem of students from different provinces.

Self-esteem can be classified as explicit and implicit self-esteem.⁴⁹ Explicit self-esteem is presented in the reflective and deliberative evaluations that one makes about oneself, while implicit self-esteem results from responses to self-relevant stimuli and is the function of automatic process.⁵⁰ It is found that people with high explicit self-esteem and low implicit

self-esteem showed higher levels of narcissism⁴⁹ and the self-esteem level measured with RSES is usually the explicit one.⁵¹ So, a self-esteem score of more than 25 cannot always be guaranteed to be healthy since high self-esteem can be relatively secure or defensive.⁴⁹ Our study showed that 6.7% students had high self-esteem. The normal range of self-esteem falls within the 16-25 score range which 74.4% of the students come under.

Healthy self-esteem in healthcare professionals is a personal quality which, when combined with professionalism and accountability, can greatly reinforce patient satisfaction.⁵² This is the reason why strengthening medical students' self-esteem warrants special consideration from the beginning of their medical studies as early intervention could yield better results.

Strengths and limitations

The findings of this study contribute evidence to the limited literature on the status of self-esteem in medical undergraduates at a college in Nepal. Our findings provide estimations of our future doctors' levels of self-esteem. This would help the policy makers and medical institution to develop interventions that could support those with low self-esteem. The study used well-known standard screening tools i.e. RSES to assess self-esteem and measured its difference among several variables. The RSES has high reliability.^{19,20} We also calculated Cronbach's alpha for our study sample and the value for it was 0.811 which showed good internal consistency of the instrument for our population as well.

As the study was conducted among medical students at a single medical college, the results cannot be generalized to the larger populations of the students of other medical colleges of Nepal as well as the general population. Therefore, multi-centric studies on self-esteem should be conducted in order to expand the generalizability of the findings. Likewise, Rosenberg's self-esteem scale only measures general self-esteem neglecting self-esteem in relation to peers, parents and schooling.⁵³ This leads to partial assessment of levels of self-esteem. The participants may have felt pressure to give answers that were socially desirable which could have contributed to response bias. Response bias includes responding to questionnaire items without understanding them and also, providing what they consider to be socially desirable answers. There might also be subjective bias associated with our study as the questionnaire is based on the subjective feelings of the respondents.

Conclusion

This study found that the majority of medical students had normal self-esteem. Several factors including age, gender, schooling, and year of study could have an impact upon self-esteem. Further studies on self-esteem should be conducted on larger populations to determine the contributing factors of low self-esteem in medical students. In our study, low self-esteem was most prevalent in third year students which may be due to the transition from pre-clinical to clinical study. Proper counseling and clinical exposure in pre-clinical years may help students to manage this transition more easily. Considering the conducted study and the related studies, we can conclude that self-esteem along with medical knowledge and skills play a great role in shaping future doctors. Hence, it is recommended that essential measures such as clinical exposure in pre-clinical years and proper counseling should be used to enhance the levels of self-esteem and mental health among medical students.

Conflict of interest

No competing interests were disclosed.

Funding

The author(s) declared that no grants were involved in supporting this work.

Data availability

Underlying data

Figshare: 'Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study'.
<https://doi.org/10.6084/m9.figshare.15149589>.¹⁸

Extended data

Figshare: 'Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study'.
<https://doi.org/10.6084/m9.figshare.15149589>.¹⁸

This project contains the following extended data:

- Questionnaire (includes consent form)

Reporting guidelines

STROBE checklist for 'Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study', <https://doi.org/10.6084/m9.figshare.15149589>.¹⁸

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

We would like to thank the students of NAIHS for their participation and the Community Medicine Department, NAIHS for allowing and helping us to conduct this study.

References

- Van Vugt M, Howard C, Moss S: **Being better than some but not better than average: Self-enhancing comparisons in aerobics.** *Br. J. Soc. Psychol.* 1998 Jun; **37**(2): 185–201.
[Publisher Full Text](#)
- Hadinezhad P, Masoudzadeh A: **A study of the relationship between self-esteem and academic achievement in medical students of Sari Medical College.** *Life.* 2018 Apr 1; **50**: 1.
- Hudd SS, Dumlao J, Erdmann-Sager D, et al.: **Stress at college: effects on health habits, health status and self-esteem.** *College Student Journal.* 2000 Jun 1; **34**(2).
- Stewart SM, Lam TH, Betson CL, et al.: **A prospective analysis of stress and academic performance in the first two years of medical school.** *Med. Educ.* 1999 Apr; **33**: 243–250.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wolf TM, Faucett JM, Randall HM, et al.: **Graduating medical students' ratings of stresses, pleasures, and coping strategies.** *J. Med. Educ.* 1988 Aug.; **63**: 636–642.
[Publisher Full Text](#)
- Dahlin ME, Runeson B: **Burnout and psychiatric morbidity among medical students entering clinical training: a three year prospective questionnaire and interview-based study.** *BMC Med. Educ.* 2007 Dec 1; **7**(1): 6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guthrie EA, Black D, Shaw CM, et al.: **Embarking upon a medical career: psychological morbidity in first year medical students.** *Med. Educ.* 1995 Sep; **29**(5): 337–341.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Iqbal A, Nawaz H, Sikandar MZ, et al.: **A Cross-Sectional Study of Happiness, Self-Esteem and Optimism in Medical Students.** *Ann. King Edward Med. Univ.* 2019 Oct 12; **25**(3).
- McAuley E, Mihalko SL, Bane SM: **Exercise and self-esteem in middle-aged adults: multidimensional relationships and physical fitness and self-efficacy influences.** *J. Behav. Med.* 1997 Feb 1; **20**(1): 67–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu M, Wu L, Ming Q: **How does physical activity intervention improve self-esteem and self-concept in children and adolescents? Evidence from a meta-analysis.** *PLoS one.* 2015 Aug 4; **10**(8): e0134804.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Motl RW, Konopack JF, McAuley E, et al.: **Depressive symptoms among older adults: Long-term reduction after a physical activity intervention.** *J. Behav. Med.* 2005 Aug 28 [cited 2021 May 30]; **28**(4): 385–394.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pandey RA, Chalise HN: **Self-esteem and academic stress among nursing students.** *Kathmandu Univ. Med. J.* 2015; **13**(4): 298–302.
- Amirkhani M, Habibi F, Hakimi A, et al.: **A cross-sectional study of relationship between self-esteem and self-concept of students of medical sciences.** *Inter. J. Res. Med. Sci.* 2018 Mar; **6**(3): 747.
[Publisher Full Text](#)
- Cohen A, Cohen R: **Development of competence in clerkship.** *Med. Teach.* 1990 [cited 2021 May 30]; **12**(1): 47–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Aarif SM, Mishra B: **Are the future doctors low on mental health and self esteem: a cross sectional study from a rural health university.** *Indian J Prev Soc Med.* 2009; **40**(3,4): 189–193.
- Society and the Adolescent Self-Image [Morris Rosenberg, 1965]. pdf | DocDroid: [cited 2021 May 22].
[Reference Source](#)
- Rosenberg self-esteem scale: callhelpline.org.uk, by Betsy Cadwaladr Univ Heal Board. [cited 2021 Apr 20].
[Reference Source](#)
- Status of self-esteem in medical students of a college in Kathmandu: A descriptive cross sectional study.** [cited 2021 Sep 14].
[Reference Source](#)
- Rosenberg M: *Conceiving the self* Malabar. FL: Krieger; 1986.
- Blascovich J, Tomaka J: **Measures of "Self-esteem"**. Robinson JP, Shaver PR, Wrightsman LS, editors. *Measures of Personality and Social Psychological Attitudes*. 3rd Edition. Ann Arbor: Institute for Social Research; 1993.
- Tsai I, Ying Y-W, Lee PA: **Cultural predictors of self-esteem: A study of Chinese American female and male young adults.** *Cult. Divers. Ethn. Minor. Psychol.* 2001; **7**: 284–297.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Robins RW, Tracy JL, Trzesniewski K, et al.: **Personality correlates of self-esteem.** *J. Res. Pers.* 2001; **35**(4): 463–482.
[Publisher Full Text](#)
- McMullin JA, Cairney J: **Self-esteem and the intersection of age, class, and gender.** *J. Aging Stud.* 2004 Feb 1; **18**(1): 75–90.
[Publisher Full Text](#)
- Housley K, Martin S, Mc Coy H, et al.: **Self-esteem of adolescent females as related to race, economic status, and area of residence.** *Percept. Mot. Skills.* 1987 Apr; **64**(2): 559–566.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Egwurugwu J, Ohamaeme M, Dike E, et al.: **Assessment of Socio-demographic Factors and Self-esteem among Imo State University Medical Students, Nigeria.** *J Adv Med Med Res.* 2017 Jan 10; **24**(4): 1–12.
[Publisher Full Text](#)
- Johari Talib ZM, Mamat M: **Effects of parenting style on children development.** *World J. Soc Sci.* 2011 May; **1**(2): 14–35.
- self-esteem – APA Dictionary of Psychology: [cited 2021 Apr 16].
[Reference Source](#)
- Sepahi V, Niroumand E, Keshavarzi F, et al.: **The Relationship between Self-esteem and Academic Achievement in Pre-clinical and Clinical Medical Students.** *J. Babol Uni. Med. Sci.* 2015; **3**(1): 32–38.
- Pais M: **Perceived self-esteem amongst first-year nursing students-a cross-sectional survey.** *Int. J. Heal. Rehab. Sci.* 2012; **1**(2): 74–80.
- Huang L, Thai J, Zhong Y, et al.: **The Positive Association Between Empathy and Self-Esteem in Chinese Medical Students: A Multi-Institutional Study.** *Front. Psychol.* 2019 August; **10**: 1921.
[Publisher Full Text](#)
- Roberts JE, Gotlib IH, Kassel JD: **Adult attachment security and symptoms of depression: the mediating roles of dysfunctional attitudes and low self-esteem.** *J. Pers. Soc. Psychol.* 1996 Feb; **70**(2): 310–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mruck CJ: *Self-esteem research, theory, and practice: Toward a positive psychology of self-esteem*. Springer Publishing Company; 2006 May 9.

33. Mahon NE, Yarcheski A, Yarcheski TJ, *et al.*: **A meta-analytic study of predictors for loneliness during adolescence.** *Nurs. Res.* 2006 Sep 1; **55**(5): 308–315.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Burgoon JM, Meece JL, Granger NA: **Self-efficacy's influence on student academic achievement in the medical anatomy curriculum.** *Anat. Sci. Educ.* 2012 Sep; **5**(5): 249–255.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Nguyen DT, Wright EP, Dedding C, *et al.*: **Low self-esteem and its association with anxiety, depression, and suicidal ideation in Vietnamese secondary school students: a cross-sectional study.** *Front. Psych.* 2019 Sep 27; **10**: 698.
[Publisher Full Text](#)
36. Abdel-Khalek AM: **Love of life as a new construct in the well-being domain.** *Soc. Behav. Personal. Int. J.* 2007 Jan 1; **35**(1): 125–134.
[Publisher Full Text](#)
37. Abdel-Khalek AM: **The development and validation of the Arabic Scale of Mental Health (ASMh).** *Psychol. Rep.* 2011 Dec; **109**(3): 949–964.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Abdel-Khalek AM: **Associations between religiosity, mental health, and subjective well-being among Arabic samples from Egypt and Kuwait.** *Ment. Health Relig. Cult.* 2012 Oct 1; **15**(8): 741–758.
[Publisher Full Text](#)
39. Abdel-Khalek AM: **The Arabic Scale of Happiness (ASH): psychometric characteristics.** *Comprehensive Psychology.* 2013 Jan 1; **2**: Article 5–Article 9.
[Publisher Full Text](#)
40. Abdel-Khalek AM, Korayem AS, El-Nayal MA: **Self-esteem among college students from four Arab countries.** *Psychol. Rep.* 2012 Feb; **110**(1): 297–303.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Abdel-Khalek A, Snyder CR: **Correlates and predictors of an Arabic translation of the Snyder Hope Scale.** *J. Posit. Psychol.* 2007 Oct 1; **2**(4): 228–235.
[Publisher Full Text](#)
42. Baumeister RF, Campbell JD, Krueger JI, *et al.*: **Does high self-esteem cause better performance, interpersonal success, happiness, or healthier lifestyles?.** *Psychol. Sci. Public Interest.* 2003 May; **4**(1): 1–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Baumeister RF, Smart L, Boden JM: **Relation of threatened egotism to violence and aggression: The dark side of high self-esteem.** *Psychol. Rev.* 1996 Jan; **103**(1): 5–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Ahmat SN, Muda MR, Neoh CF: **Self-esteem level and its relationship to academic performance among undergraduate pharmacy students in a Malaysian public university.** *Indian J Pharm Educ Res.* 2018 Apr 1; **52**(2): 197–201.
45. Chaudhari B: **The Relationship of Eating Disorders Risk with Body Mass Index, Body Image and Self-Esteem among Medical Students.** *Ann. Med. Health Sci. Res.* 2017 [cited 2021 Apr 15]; Vol. 7. *Annals of Medical and Health Sciences Research.*
[Reference Source](#)
46. Naz F, Aijaz S, Ullah Khan S: **Level of self esteem in medical students according to their educational year, gender and socio economic status.** [cited 2021 Apr 15].
[Reference Source](#)
47. Extracurricular Activity Involvement and Adolescent Self-Esteem: [cited 2021 Sep 14].
[Reference Source](#)
48. Eremie MD, Chikweru AE: **Self Esteem among Private and Public Secondary Schools Students in Rivers State: Implications for Counselling.** *Kuwait Chapter of the Arabian J. Busi. Manage. Rev.* 2015 Jul 1; **4**(11): 1–6.
[Publisher Full Text](#)
49. Zeigler-Hill V, Jordan CH: **Two faces of self-esteem: implicit and explicit forms of self-esteem.** Gawronski B, Payne BK, editors. *Handbook of Implicit Social Cognition: Measurement, Theory, and Applications.* New York: Guilford Press; 2010; p. 392–407.
50. Jordan CH, Whitfield M, Zeigler-Hill V: **Intuition and the Correspondence Between Implicit and Explicit Self-Esteem.** *J. Pers. Soc. Psychol.* 2007 Dec; **93**(6): 1067–1079.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Di Pierro R, Mattavelli S, Gallucci M: **Narcissistic Traits and Explicit Self-Esteem: The Moderating Role of Implicit Self-View.** *Front. Psychol.* 2016 November 22; **7**: 1815.
[Publisher Full Text](#)
52. Decker PJ: **The hidden competencies of healthcare: Why self-esteem, accountability, and professionalism may affect hospital customer satisfaction scores.** *Hosp. Top.* 1999; **77**(1): 14–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Potard C: **Self-Esteem Inventory (Coopersmith).** *Encyclopedia of Personality and Individual Differences.* Springer International Publishing; 2017; p. 1–3.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 09 March 2022

<https://doi.org/10.5256/f1000research.76429.r123050>

© 2022 Joshi S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sunil Kumar Joshi

Nepal Injury Research Centre, Department of Community Medicine, Kathmandu Medical College, Kathmandu, Nepal

I appreciate the efforts put forward by a group of young students to participate in such a novel study in Nepal. This study on self-esteem has been conducted in a single medical college in Kathmandu.

Here are a few comments from me:

There is no explanation on the classification of self-esteem and its relation to the RSES scoring in the Introduction section, it comes for the first time in the Methods section.

Few sentences need to be rephrased or restructured as they are not very clear to me. The manuscript needs to be checked by an English language expert. The authors should elaborate on the technical words mentioned in the text, e.g., narcissism.

In the pdf format of the manuscript "Table 1. Socio-demographic characteristics and RSES of the students along with p-value" comes in the Methods section, it should go to the Results section.

I have some reservations about the sample size calculation. An estimated rate of population proportion from a study in a rural Indian medical college i.e. 18% has been cited to calculate the sample size (ref 15). NAIHS is not a rural medical college, and as it is the first of its kind study in Nepal, I suggest using the population proportion of 50%. It would bring a sample size of approximately 384. I do not support the adjusted sample size calculation for the finite population in this study. As the total number of students in this medical college is 550, the researchers would be easily able to get the required study respondents. So, I suggest the students do recalculate the sample size as I suggested, extend the data collection and reanalyse data and resubmit for publication. This will definitely make the findings more robust. I suggest being considerate in citing the published literature. I think a total of 53 references is a little high for this article.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public health, Occupational health and safety, injury and violence prevention.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 05 Apr 2022

Stuti Yadav, Nepalese Army Institute of Health Sciences (NAIHS), Kathmandu, Nepal

Thank you for the comments, sir. The manuscript has been changed as per the comments and the table has also been placed in the results section.

Competing Interests: none

Reviewer Report 03 February 2022

<https://doi.org/10.5256/f1000research.76429.r98205>

© 2022 Ugwuanyi C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Christian Sunday Ugwuanyi

Department of Science Education, Faculty of Education, University of Nigeria, Nsukka, Nigeria

The study explored the state of self-esteem among undergraduates at Tribhuvan University in Nepal's medical college. The majority of medical students reported normal self-esteem, according to the findings of a descriptive cross-sectional survey conducted among first- to fifth-year medical students from December 2020 to April 2021. The findings, however, revealed that 18.9% of

students showed low self-esteem, with third-year students being the most affected (30.77 percent). The researchers recommended that to help medical students become confident and effective doctors, interventions to improve self-esteem should be implemented. This study was properly conceptualised and documented as an academic paper, but more efforts should be made by the authors to use recent empirical studies to support their line of argument. The methodology section of the manuscript was properly handled to allow for its replication. In other words, sufficient details of methods and analysis were provided to allow replication by others. The results were also adequately presented.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Educational research, measurement and evaluation

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 22 November 2021

<https://doi.org/10.5256/f1000research.76429.r100126>

© 2021 Mottershead D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dr. Richard Mottershead

Ras Al Khaimah College of Nursing, RAK Medical and Health Sciences University, Ras Al Khaimah,

United Arab Emirates

I would like to say thank you for asking me to review this research. I have read the article and have the following comments to make:

This is a generally well-composed paper, which will be of interest to healthcare educators who wish to be aware of issues which may hamper student performance and well-being. The authors appear to have a creditable background and research experience which gives the article credibility and allows for an interesting insight into student mental well-being within Nepal. The paper proposes that female students are at greater risk of reduced self-esteem and the causality needs now to be ascertained in order to reduce further stigma and shame. Comparisons have been made with relevant contemporary research and this allows for a global overview of the topic and how this article fits within the body of evidence.

I think that the article is correct not to generalize and to highlight the subjective nature of assessing self-esteem but no doubt the data will highlight the need to support and the participants may have developed a collective consciousness on the necessity to self-care in order to be effective healthcare practitioners.

I wish the authors well with their future efforts to support this cohort and in progressing future research and support networks which I am sure will provide long-term benefits to the health and well-being of medical students with results that can be referred to the wider student healthcare field.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: I am a qualified researcher and mental health specialist with experience as an educator within higher education in undergraduate and postgraduate programmes across nursing, medical and health and social care provision.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research