RESEARCH ARTICLE

Cookies rich in iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants: a novel functional food potential for adolescent with anemia [version 1; peer review: 1 approved, 1 approved with reservations]

Alexander Sam Leonard Bolang¹, Mochammad Rizal², Fahrul Nurkolis³, Nelly Mayulu⁴, Narpudji Astuti Taslim⁵, Son Radu⁶, Mrinal Samtiya⁶, Youla Annatje Assa⁷, Hendra Agung Herlambang⁸, Alpinia Shinta Pondagitan¹, Christopherous Diva Vivo⁸

¹Food and Nutrition, Sam Ratulangi University, Manado, North Sulawesi, 95115, Indonesia

²Division of Nutritional Sciences, Cornell University, Ithaca, New York, 14850, USA

³Biological Sciences, State Islamic University of Sunan Kalijaga (UIN Sunan Kalijaga Yogyakarta), Yogyakarta, Yogyakarta, 55281, Indonesia

⁴Nutritional Sciences, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia

⁵Food Sciences, Universiti Putra Malaysia, Selangor Darul Ehsan, 43400, Malaysia

⁶Nutrition Biology, Central University of Haryana, Haryana, 123031, India

⁷Biochemistry Department, Sam Ratulangi University, Manado, North Sulawesi, 95115, Indonesia

⁸Dentistry, University of Indonesia, Depok, 16424, Indonesia

V1 First published: 25 Oct 2021, 10:1075
<https://doi.org/10.12688/f1000research.74045.1>

Latest published: 04 Nov 2022, 10:1075
<https://doi.org/10.12688/f1000research.74045.2>

Abstract

Background: This study aims to process porang flour, moringa leaves, and tempe into cookies that are high in iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants as potential functional food snacks for adolescents with anemia.

Methods: This study is experimental with a completely randomized trial design (CRD) with three treatments and triplicates. There are three comparison formulations of a combination of porang, moringa leaves, and black soybean-based tempe; respectively F1 (3:3:3)%, F2 (3:6:6)%, and F3 (3:10:10)%. The cookie sample was then analyzed for water content, ash, iron, folic acid, cobalt, and antioxidant activity. The differences between the samples were analyzed based on the activity of antioxidants, iron, folic acid, and vitamin B12 (cobalamin) in data obtained from triplicates using multivariate ANOVA analysis.

Results: The results of the ash and water content tests of all three cookie formulations showed values in accordance with the Indonesian National Standard (SNI) for cookie products. F3 has significantly higher iron, folic acid, and cobalt levels than F1 and F2 ($p < 0.05$).

Open Peer Review

Approval Status ? ✓

	1	2
version 2 (revision) 04 Nov 2022		
version 1 25 Oct 2021	? view	✓ view
1. Emmanuel Oladeji Alamu , International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria 2. Yaktiworo Indriani , Universitas Lampung, Bandar Lampung, Indonesia Any reports and responses or comments on the		

Antioxidant activity is highest in F3, but there is no meaningful difference between F2 and F3 ($p>0.05$).

Conclusions: The combination of porang's tubers, moringa leaves, and tempe made from black soybeans, especially F3, has the potential to be used as a functional cookie processed product source of iron (Fe), folic acid, cobalamin, and antioxidants for adolescent anemia.

Keywords

Iron, Folic Acid, Cobalamin, Antioxidant, Anemia, Functional Food, Cookies

article can be found at the end of the article.

This article is included in the **Agriculture, Food and Nutrition** gateway.

Corresponding authors: Alexander Sam Leonard Bolang (bolangasl@gmail.com), Mochammad Rizal (mr855@cornell.edu), Fahrul Nurkolis (20106040012@student.uin-suka.ac.id)

Author roles: **Bolang ASL:** Conceptualization, Data Curation, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Rizal M:** Data Curation, Formal Analysis, Funding Acquisition, Writing – Original Draft Preparation, Writing – Review & Editing; **Nurkolis F:** Conceptualization, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; **Mayulu N:** Conceptualization, Supervision, Validation, Writing – Review & Editing; **Taslim NA:** Supervision, Validation, Writing – Review & Editing; **Radu S:** Formal Analysis, Writing – Review & Editing; **Samtiya M:** Writing – Review & Editing; **Assa YA:** Writing – Review & Editing; **Herlambang HA:** Writing – Review & Editing; **Pondagitan AS:** Writing – Review & Editing; **Vivo CD:** Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Lembaga Pengelola Dana Pendidikan, Indonesia Endowment Fund for Education (LPDP: Lembaga Pengelola Dana Pendidikan) Ministry of Finance, Republic of Indonesia.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Bolang ASL *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bolang ASL, Rizal M, Nurkolis F *et al.* **Cookies rich in iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants: a novel functional food potential for adolescent with anemia [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:1075 <https://doi.org/10.12688/f1000research.74045.1>

First published: 25 Oct 2021, 10:1075 <https://doi.org/10.12688/f1000research.74045.1>

Introduction

Anemia is a serious public health problem faced by all countries in the world. According to the World Health Organization (WHO), 42% of children <5 years, 33% of women of productive age, and 40% of pregnant women in the world have anemia. In Indonesia, Basic Health Research (Riskesdas) shows an increase in the prevalence of anemia in adolescents aged 15-24 years. In 2007, the prevalence of adolescent anemia in Indonesia was only 6.90% (Riskesdas, 2007), which then increased to 18.40% in 2013 (Riskesdas, 2013) and 32% in 2018 (Riskesdas, 2018). The impact of anemia is complex, ranging from immunity (Hasan *et al.*, 2016) to work productivity (Blakstad *et al.*, 2020), growth (Soliman *et al.*, 2009) to intelligence (Bahrami *et al.*, 2020), and physical fitness (Tsai *et al.*, 2019) to maternal, fetal and child health (Georgieff, 2020).

Various interventions need to be done to speed up the treatment of anemia. One of the Government of Indonesia's programs is the provision of iron and folic acid supplementation in the form of Blood Added Tablets (TTD) in school-aged teenagers. Although the program's scope is quite high, which is 76.2%, its effectiveness is relatively low because only 2.13% consume TTD as recommended ≥ 52 items per year (Riskesdas, 2018). Therefore, alternative efforts such as food fortification become important to do, especially in snacks that are mostly preferred by teenagers. In addition to promising results in terms of decreasing the problem of micronutrient deficiency (Dewi *et al.*, 2021), fortification in snacks is also potentially effective because adolescents aged 18-25 years tend to consume snacks ≥ 3 times per day which contributes to 25% recommended dietary allowance (RDA) or average daily dietary intake level (USDA, 2012).

Many local Indonesian foodstuffs are of high nutritional value (Permatasari *et al.*, 2021), for example, Porang tubers. Porang's tubers (*Amorphophallus oncophyllus*) can be used as potential glucomannan source flour in Indonesia (especially the original regency, Madiun, East Java, Indonesia) because of the high glucomannan content (Harmayani *et al.*, 2014). The bulbs are cultivated in Indonesia as secondary plants under teak, mahogany, and sonokeling plantations (Yanuriati *et al.*, 2017). As a source of glucomannan, people have many benefits as functional foodstuffs (Harmayani *et al.*, 2014; Yanuriati *et al.*, 2017). Glucomannan properties can bind to other components such as proteins, minerals, and active compounds such as antioxidants that form a matrix of tissues in the food developed from it (Herlina *et al.*, 2015). Glucomannan has been widely used in the food industry as a thickener, gel-forming, texture, water binder, and emulsion stabilizer, including in cookie making (Imeson, 2009; Mikkonen *et al.*, 2009; Zhang *et al.*, 2005).

In addition to porang's tubers, another local foodstuff that is also known to have many health benefits is moringa leaves. *Moringa oleifera* is known as an excellent food source that is easy to digest and rich in protein (Fahey, 2005). According to Sultana and Anwar (2007), moringa leaves have many valuable compounds such as protein, vitamins, calcium, iron, folic acid, and antioxidants (carotenoids, flavonoids, and phenols) (Sultana & Anwar 2007). Various developing countries in the world feed their children with moringa (Kasolo *et al.*, 2010). Busani *et al.* (2011) reported that the presence of various minerals and vitamins helps boost immunity against various diseases, especially moringa leaves contain various amino acids (Busani *et al.*, 2011). In addition, the levels of iron, folic acid, and antioxidants contained in it also have the potential to become an anti-anemia agent (Mo *et al.*, 2013; Kurtoglu *et al.*, 2003).

Indonesia's third potential foodstuff, which has been known globally, is tempe. Tempe is made from soybeans fermented by *Rhizopus* spp mushrooms such as *R. oligosporus*, *R. stolonifer*, and *R. oryzae* with several characteristics such as white, compact texture, and mixed distinctive flavors of mushrooms and soy flavors (Astuti *et al.*, 2013). Black soybean tempe has potential functional characteristics, as black soybeans contain phenolics, tannins, anthocyanins, cobalamin (B12), and isoflavones, as well as a higher antioxidant activity than yellow soybeans (Xu dan Chang, 2007). According to Nurrahman *et al.* (2011) black soybean tempe has the same level of fondness as yellow soybean tempe, and its antioxidant activity is higher (Nurrahman *et al.*, 2011). In addition, Vitamin B12 (Cobalamin) in the tempe has the potential in the formation of red blood cells and its antioxidants to overcome red blood cell damage (Mo *et al.*, 2013; Kurtoglu *et al.*, 2003).

Therefore, the combination of porang's tubers, moringa leaves, and tempe made from black soybeans can be used as processed products such as functional cookies as a snack for anemia adolescents. This study utilizes and formulates all three ingredients into functional cookies and determines levels of iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants.

Methods

This study is experimental with a completely randomized trial design (CRD) with three treatments and triplicates.

Cookies formulation

Cookie processing

Cookie making refers to Nugraha (2009), which consists of the preparation of materials, dough formation (cream formation and flour mixing), dough printing, roasting, refrigeration, and packaging. The preparation of materials starts from weighing the material according to treatment. The formation of the dough begins by mixing margarine, egg yolk, stevia sugar powder, and baking powder using a mixer so that cream is formed. Furthermore, we added flour, moringa leaf flour (freeze-drying methods), and tempe flour (made from black soybean), corresponding to the treatment. Then the dough is formed into sheets and printed using a molding tool. Cookies are then put into the oven at 140°C for 15-20 minutes. After the cookies are done cooking, the cookies are then cooled so that the cookies are perfectly served.

Ash levels and water content analysis

The procedure for determining ash levels was carried out using the Official Methods of Analysis of Association of Official Agricultural Chemists (AOAC) International 2005 (AOAC 2005). First, the cup was dried in the oven at a temperature of 105°C for one hour. Then, the cup was cooled for 15 minutes in a desiccator and weighed. A sample of 2 grams was put into a furnace with a temperature of 550 °C for three hours. It was then cooled outside the furnace until the temperature was $\pm 120^{\circ}\text{C}$ and put into a desiccator to remove purified water and crystals. Plates and ash were weighed so that constant weight was obtained.

The determination of the moisture content used was the AOAC method of drying (thermogravimetry). The principle of this method was based on the evaporation of water in materials by heating, then weighing to a constant weight. The weight loss that occurred is the water content contained in the material. The empty plate was heated in the oven at 105°C for 30 minutes, cooled in a desiccator for 15 minutes, then weighed (W0). A sample of 2 grams was then put in a cup that had a known weight, weighed (W1), then dried in the oven at a temperature of 105°C for three hours, cooled in a desiccator for 15-30 minutes, then the cup and its contents were weighed, re-dried for an hour, cooled in an extractor, and re-weighed (W2).

Iron (Fe) ortho-phenantroline analysis

The principle of determining total iron content by the method o-phenanthroline was the same as the method of $\alpha.\alpha'$ – the principle of iron (III) was reduced by hydroxyl amine to form iron (II). Iron (II), both present in foodstuffs and iron reduction (III), was then reacted with o-phenanthroline to form a red Fe-phenanthroline complex that can be measured with visible spectrophotometers at a wavelength of 510 nm.

Folic acid and B12 (cobalamin) analysis

This analysis used HPLC-UV with a gradient elevation system in which the composition of the phase of motion changes during the measurement process. The phase of motion used was dapi phosphate 10 mM pH 4, and the dapi phosphate mixture: acetonitrile, 7:3. As a stationary phase, the Atlantis column T3 C18 (4.6 mm $\times$ 150 mm), 5 μm was used. Oven temperature were 35 °C, with a flow rate of 0.5 mL/min and injection volume of 20 μL . UV detectors were set at wavelengths of 280 for folic acid and 275 nm for cobalt (vitamin B12). A raw solution of cobalt (vitamin B12) with a concentration of 2 $\mu\text{g/mL}$ and folic acid of 0.4 $\mu\text{g/mL}$ was injected into HPLC of 20 μL under appropriate analytical conditions.

Antioxidant activity assay

Antioxidant activity was determined using DPPH (2,2-diphenyl-1-picrylhydrazil). Each sample consisting of 100 μL was placed in a microplate of 96 and added 100 μL DPPH 0.3 mM, then incubated for 30 minutes in a dark room. The absorbance of the sample was measured using an ELISA reader (# Allsheng) at a wavelength of 517 nm.

Data management and analysis

Statistical analysis was done using SPSS 26.0 for apple's version of MacBook. The differences between the samples were analyzed based on the activity of antioxidants, iron, folic acid, and vitamin B12 (cobalamin). Data obtained from three repeats (triplicates) was analyzed with multivariate ANOVA at 95% CI ($p < 0.05$). The result was declared significant if the p -value < 0.05 .

Results

Test of normality

Ash levels and water content analysis

The results of the ash and water content tests of the three cookie formulations above showed a value in accordance with the Indonesian National Standard for cookie products (SNI 01-2973-1992), which is a maximum ash content of 1.5% and water content of no more than 5%. Raw details of ash level and ash level test results are found in (Underlying Data) (Nurkolis & Bolang, 2021).

The results of the ANOVA multivariate test above showed that there was a significant difference ($p < 0.05$) between all treatments in each test result or variable (Fe, folic acid, and vitamin B12). However, in antioxidant activity, that has a significant difference only F2 & F3. While between F1 & F2 equal F1 & F3 there is no significant difference ($p > 0.05$). Details of raw data from laboratory tests on levels of antioxidant activity (%), iron (Fe), folic acid, and cobalamin (vitamin B12) can be seen in Underlying Data (Nurkolis & Bolang, 2021).

Table 1. Cookies formulation.

Ingredients	F1 (%)	F2 (%)	F3 (%)
Porang Flour	3	3	3
Moringa Leaf Flour	3	6	10
Tempe Flour (Black-Soybean)	3	6	10
Stevia	10	10	10
Skimmed Milk	5	5	5
Egg Yolk	10	10	10
Baking Powder	1	1	1
Margarine	20	20	20

Table 2. Tests of normality.

	Formulasi	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Fe (mg/100 g)	F1	.322	3	.	.880	3	.324
	F2	.212	3	.	.990	3	.811
	F3	.382	3	.	.757	3	.015
Folate (mcg/100 g)	F1	.270	3	.	.949	3	.565
	F2	.249	3	.	.967	3	.652
	F3	.358	3	.	.813	3	.146
B12 (mcg/100 g)	F1	.212	3	.	.990	3	.813
	F2	.297	3	.	.917	3	.441
	F3	.227	3	.	.983	3	.749
Antioxidant activity (%)	F1	.216	3	.	.988	3	.793
	F2	.283	3	.	.935	3	.506
	F3	.253	3	.	.964	3	.637

Based on the normality test results above, the data obtained is homogeneous ($p > 0.05$), followed by the Anova multivariate test.

^aLilliefors significance correction.

Discussion

Anemia is a severe public health problem faced by all countries globally, and various interventions need to be done to speed up the treatment of anemia. One of them is utilizing local natural ingredients as functional food sources of iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants for adolescent anemia. In this study, researchers utilized and formulated all three local natural ingredients (a combination of people's tubers, moringa leaves, and black soy-based tempe) into functional cookies and determining levels of iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants. The selection of processing into cookie products is wrong as an alternative effort instant snack food that is majority preferred by teenagers without having to process it again (e.g., iron source chicken liver but must go through the cooking process, etc.). This research is a basic study, whose certainty of clinical benefits needs to be done further through pre-clinical studies and clinical trials in humans. However, the results of this study (cookie formulation) can be used as a

Table 3. Results of ash levels and water content analysis.

Formulation	Ash	Moisture content
F1	1.30 ± 0.35	3.42 ± 1.35
F2	1.37 ± 0.07	3.76 ± 1.26
F3	1.46 ± 0.16	3.73 ± 1.77

Table 4. Multiple comparisons.

Tukey HSD							
Dependent variable	(I) Formulasi	(J) Formulasi	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
						Lower bound	Upper bound
Fe (mg/100 g)	F1	F2	1.81667*	.47005	.019	.3744	3.2589
		F3	-4.16667*	.47005	.000	-5.6089	-2.7244
	F2	F1	-1.81667*	.47005	.019	-3.2589	-.3744
		F3	-5.98333*	.47005	.000	-7.4256	-4.5411
	F3	F1	4.16667*	.47005	.000	2.7244	5.6089
		F2	5.98333*	.47005	.000	4.5411	7.4256
Folate (mcg/100 g)	F1	F2	4.65333*	.71293	.002	2.4659	6.8408
		F3	-4.79000*	.71293	.001	-6.9775	-2.6025
	F2	F1	-4.65333*	.71293	.002	-6.8408	-2.4659
		F3	-9.44333*	.71293	.000	-11.6308	-7.2559
	F3	F1	4.79000*	.71293	.001	2.6025	6.9775
		F2	9.44333*	.71293	.000	7.2559	11.6308
B12 (mcg/100 g)	F1	F2	5.39000*	1.02781	.005	2.2364	8.5436
		F3	-6.31667*	1.02781	.002	-9.4703	-3.1631
	F2	F1	-5.39000*	1.02781	.005	-8.5436	-2.2364
		F3	-11.70667*	1.02781	.000	-14.8603	-8.5531
	F3	F1	6.31667*	1.02781	.002	3.1631	9.4703
		F2	11.70667*	1.02781	.000	8.5531	14.8603
Antioxidant activity (%)	F1	F2	3.46667	1.42911	.112	-.9182	7.8516
		F3	-2.14667	1.42911	.355	-6.5316	2.2382
	F2	F1	-3.46667	1.42911	.112	-7.8516	.9182
		F3	-5.61333*	1.42911	.018	-9.9982	-1.2284
	F3	F1	2.14667	1.42911	.355	-2.2382	6.5316
		F2	5.61333*	1.42911	.018	1.2284	9.9982

reference in the dosing of preclinical trials and future clinical trials. The results of the ash and water content tests of these three cookie formulations showed a value in accordance with the Indonesian National Standard for cookie products (SNI 01-2973-1992), which is a maximum ash content of 1.5% and water content of no more than 5% (Table 3).

Formulation 3 (F3) is a high iron cookie, amounting to 29.41 ± 0.65 mg / 100g, higher than F1 and F2 significantly ($p < 0.05$) (Table 4; Figure 1). In the scheme (Figure 5), iron is needed by the body for erythropoiesis or the process of red blood cell synthesis, precisely as the constituent material of hemoglobin (Singh, 2018). F3 cookies in this study have higher iron levels than the functional cookie of previous research 2020 which only has a level of 2.51 mg/100g (Rahmat *et al.*, 2020). In line with AKG (Angka Kecukupan Gizi) Indonesia in 2019, iron adequacy rates for ages 10-18 years in men range from 8-11 mg, in women 8-15 mg per person per day. Seeing this, F3 cookies are potentially a food source of iron, especially for adolescents with anemia.

In addition to iron height, F3 cookies have a fairly high antioxidant content of $24.41 \pm 1.31\%$, although there is no significant difference with F2 cookies (Table 4; Figure 4). In addition to needing iron food sources to avoid anemia is also

Figure 1. F3 cookie has the highest Fe content. In formulation 3 (F3) based on multivariate results, iron (Fe) levels were 29.41 ± 0.65 mg/100 g, significantly higher than F1 and F2 ($p < 0.05$).

Figure 2. F3 cookie has the highest folic acid content. In formulation 3 (F3) based on multivariate results, folic acid (vitamin B9) or folate levels were 308.24 ± 1.30 mcg/100 g, significantly higher than F1 and F2 ($p < 0.05$).

Figure 3. F3 cookie has the highest cobalamin content. In formulation 3 (F3) based on multivariate results, the levels of vitamin B12 (cobalamin) owned amounted to 109.76 ± 1.74 mcg/100 g, higher than F1 and F2 significantly ($p < 0.05$).

Figure 4. F3 Cookie has the highest antioxidant activity. Antioxidant activity is highest in F3 with a percentage of 24.41 ± 1.31 . However, there is no meaningful difference ($p > 0.05$) between F2 and F3 ($p = 0.355$).

needed foods that have antioxidant abilities to protect red blood cell damage due to the occurrence of oxidative stress (Reactive Oxygen Species) (Lobo *et al.*, 2010; Gwozdinski *et al.*, 2021).

Folic acid or vitamin B9 and cobalt (vitamin B12) also have an important role in minimizing the incidence of anemia. Folic acid collaborates with vitamin B12 in making red blood cells and helps iron function properly in the body (Figure 5) (Mahmood, 2014). Vitamin B9 works with vitamins B6 and B12 and other nutrients in controlling the amino acid homocysteine levels in increasing the effectiveness of red blood cell formation (Figure 5) (Mahmood, 2014). Formulation 3 (F3) is the best formulation when viewed from iron (Fe), folic acid (vitamin B9), cobalamin (vitamin B12), and antioxidant activity. F3 cookies have the highest levels of folic acid and cobalamine (308.24 ± 1.30 mcg/100 g and 109.76 ± 1.74 mcg/100 g) compared to F1 and F2 (Table 4, Figures 2 and 3). Cookie F3 has folic acid that is closest to AKG (Angka Kecukupan Gizi), which recommends 400 mcg per person per day (for men and women aged 10-18 years).

Cookie innovation made from a combination of porang's tubers, moringa leaves, and tempe made from black soybeans has potential to be made a functional anti-anemia food or to minimize anemia comprehensively (see Figure 5). In addition,

Figure 5. Potential F3 cookie scheme as anti-anaemic.

the sugar or sweetener used in this formulation is stevia which is clinically safer (not hyperglycemia) (Ashwell, 2015). However, further studies are needed in both experimental animals and humans to find out more about the clinical benefits.

Conclusion

The combination of porang's tubers, moringa leaves and tempe made from black soybeans, especially F3 has the potential to be used as a functional cookie processed product source of iron (Fe), folic acid, cobalamin, and antioxidants for adolescent anemia.

Data availability

Underlying data

Figshare: Cookies Rich in Iron (Fe), Folic Acid, Cobalamin, and Antioxidants: A Novel Functional Food for Adolescent with Anemia. <https://doi.org/10.6084/m9.figshare.15173451.v1> (Nurkolis & Bolang, 2021).

The project contains the following underlying data:

- Raw Data: Water and Ash Content, Iron (Fe), Folic Acid, Cobalamin, and Antioxidants.

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgment

We thank all contributors for their outstanding help in formatting the paper. I would also like to thank Prof. Ir. Hardinsyah, MS., Ph.D. (as President of the Federations of Asian Nutrition Societies; President of Food and Nutrition Society of Indonesia; and Chair of Southeast Asia Probiotics Scientific and Regulatory Experts Network), and Prof. Dr. Nurpudji A Taslim, M.D., MPH., Sp.GK(K) (Chairman of the Indonesian Association of Clinical Nutritionists), who has provided commentary, advice, and input in the research and writing of this paper, as well as the motivation that has been given to the author to keep the spirit of research during the pandemic. This research was supported by the Indonesia Endowment Fund for Education (LPDP: Lembaga Pengelola Dana Pendidikan) Ministry of Finance, Republic of Indonesia.

References

- Ashwell M: **Stevia, Nature's Zero-Calorie Sustainable Sweetener: A New Player in the Fight Against Obesity.** *Nutr. Today.* 2015; **50**(3): 129–134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Astuti MN, Suparmo M, Soesatyo HNE: **The role of black soybean tempe in increasing antioxidant enzyme activity and human lymphocyte proliferation in vivo.** *Int. J. Curr. Microbiol. App. Sci.* 2013; **2**: 316–327.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bahrami A, Khorasanchi Z, Tayefi M, et al.: **Anemia is associated with cognitive impairment in adolescent girls: A cross-sectional survey.** *Appl. Neuropsychol. Child.* 2020; **9**: 165–171.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blakstad MM, Nevins JEH, Venkatramanan S, et al.: **Iron status is associated with worker productivity, independent of physical effort in Indian tea estate workers.** *Appl. Physiol. Nutr. Metab.* 2020; **45**: 1360–1367.
[Publisher Full Text](#)
- Busani M, Patrick JM, Arnold H, et al.: **Nutritional characterization of Moringa (Moringa oleifera Lam) leaves.** *Afr. J. Biotechnol.* 2011; **10**(60): 12925–12933.
[Publisher Full Text](#)
- Dewi NU, Mahmudiono T: **Effectiveness of Food Fortification in Improving Nutritional Status of Mothers and Children in Indonesia.** *Int. J. Environ. Res. Public Health.* 2021; **18**: 2133.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fahey JW: **Moringa oleifera: a review of the medical evidence for its nutritional, therapeutic and prophylactic properties. Part 1.** *Trees for Life Journal.* 2005; **1**(5): 1–15.
- Georgieff MK: **Iron deficiency in pregnancy.** *Am. J. Obstet. Gynecol.* 2020; **223**: 516–524.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gwozdziński K, Pieniazek A, Gwozdziński L: **Reactive Oxygen Species and Their Involvement in Red Blood Cell Damage in Chronic Kidney Disease.** *Oxidative Med. Cell. Longev.* 2021; vol.2021: 19 pages. Article ID 6639199.
[Publisher Full Text](#)
- Harmayani E, Aprilia V, Marsono Y: **Characterization of glucomannan from Amorphophallus oncophyllus and its prebiotic activity in vivo.** *Carbohydr. Polym.* 2014; **112**: 475–479.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hassan TH, Badr MA, Karam NA, et al.: **Impact of iron deficiency anemia on the function of the immune system in children.** *Medicine (Baltimore).* 2016; **95**: e5395.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Herlina H, Darmawan I, Rusdianto AS: **Penggunaan tepung glukomanan umbi gembili (Dioscorea esculenta L.) sebagai bahan tambahan makanan pada pengolahan sosis daging ayam.** *Jurnal Agroteknologi.* 2015; **9**(02): 134–144.
- Imeson A: *Food Stabilisers, Thickeners and Gelling Agents.* Oxford, UK: Wiley-Blackwell; 1st ed. 2009.
[Publisher Full Text](#)
- Kasolo JN, Bimenya GS, Ojok L, et al.: **Phytochemicals and uses of Moringa oleifera leaves in Ugandan rural communities.** *J. Med. Plant Res.* 2010; **4**(9): 753–757.
- Kurtoglu E, Ugur A, Baltaci AK, et al.: **Effect of iron supplementation on oxidative stress and antioxidant status in iron-deficiency anemia.** *Biol. Trace Elem. Res.* 2003; **96**(1-3), 117–124.
[PubMed Abstract](#) | [Publisher Full Text](#)
- RISKESDAS: *Kementerian Kesehatan Republik Indonesia. 2007. Riset Kesehatan Dasar Tahun.* Jakarta: Kementerian Kesehatan; 2007.
- RISKESDAS: *Kementerian Kesehatan Republik Indonesia. 2013. Riset Kesehatan Dasar Tahun.* Jakarta: Kementerian Kesehatan; 2013.
- RISKESDAS: *Kementerian Kesehatan Republik Indonesia. 2018. Riset Kesehatan Dasar Tahun.* Jakarta: Kementerian Kesehatan; 2018.
- Lobo V, Patil A, Phatak A, et al.: **Free radicals, antioxidants and functional foods: Impact on human health.** *Pharmacogn. Rev.* 2010; **4**(8): 118–126.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mahmood L: **The metabolic processes of folic acid and Vitamin B12 deficiency.** *J. Health Res. Rev.* 2014; **1**(1): 5.
[Publisher Full Text](#)
- Mikkonen KS, Tenkanen M, Cooke P, et al.: **Mannans as stabilizers of oil-in-water beverage emulsions.** *LWT-Food Sci. Technol.* 2009; **42**: 849–855.
[Publisher Full Text](#)
- Mo H, Kariluoto S, Piironen V, et al.: **Effect of soybean processing on content and bioaccessibility of folate, vitamin B12 and isoflavones in tofu and tempe.** *Food Chem.* 2013; **141**(3): 2418–2425.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nugraha A: *Evaluasi mutu kukis dengan substitusi minyak sawit merah, tepung tempe dan tepung udang rebon (Acetes erythraeus).* Pekanbaru: Skripsi Fakultas Pertanian Universitas Riau; 2009.
- Nurrahman, Astuti M, Soesatyo SdMHNE: **The effect of black soybeans tempe and its ethanol extract on lymphocyte proliferation and IgA secretion in Salmonella typhimurium induced rat.** *Afr. J. Food Sci.* 2011; **5**(14): 775–779.
[Publisher Full Text](#)
- Permatasari HK, Nurkolis F, Augusta PS, et al.: **Kombucha tea from seagrapes (Caulerpa racemosa) potential as a functional anti-ageing food: in vitro and in vivo study.** *Heliyon.* 2021; **7**(9): e07944.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rahmat M, Priawantiputri W, Pusparini P: **Cookies Bayam Dan Tepung Sorgum Kaya Akan Zat Besi Sebagai Makanan Tambahan Untuk Ibu Hamil Dengan Anemia.** *Jurnal Riset Kesehatan Poltekkes Depkes Bandung.* 2020; **12**(2): 245–254.
- Singh AK: **Erythropoiesis: The Roles of Erythropoietin and Iron.** *Textbook of Nephro-Endocrinology.* Academic Press; 2018 (pp. 207–215).
[Publisher Full Text](#)
- Soliman AT, Al Dabbagh MM, Habboub AH, et al.: **Linear growth in children with iron deficiency anemia before and after treatment.** *J. Trop. Pediatr.* 2009; **55**: 324–327.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sultana B, Anwar F: **Flavonols (kaempferol, quercetin, myricetin) contents of selected fruits, vegetables and medicinal plants.** *Food Chem.* 2008; **108**(3): 879–884.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tsai KZ, Lai SW, Hsieh CJ, et al.: **Association between mild anemia and physical fitness in a military male cohort: The CHIEF study.** *Sci. Rep.* 2019; **9**: 11165.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- US Department of Agriculture, Agricultural Research Service: **Snacks: percentages of selected nutrients contributed by food and beverages consumed at snack occasions, by gender and age, What We Eat in America, NHANES 2011–2012.** [cited 2021 Aug 31].
[Reference Source](#)
- Xu BJ, Chang SKS: **A Comparative study on phenolic profiles and antioxidant of legums as affected by extraction solvents.** *J. Food Sci.* 2007; **72**(2): S159–S166.
[Publisher Full Text](#)
- Yanuriati A, Marseno DW, Rochmadi HE: **Characteristics of glucomannan isolated from fresh tuber of Porang (Amorphophallus muelleri Blume) Carbohydr. Polym. 2017; **156**: 56–63.
[PubMed Abstract](#) | [Publisher Full Text](#)**
- Zhang YQ, Xie BJ, Gan X: **Advance in the applications of konjac glucomannan and its derivatives.** *Carbohydr. Polym.* 2005; **60**: 27–31.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 23 June 2022

<https://doi.org/10.5256/f1000research.77755.r140091>

© 2022 Indriani Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yaktiworo Indriani

Department of Agribusiness, Faculty of Agriculture, Universitas Lampung, Bandar Lampung, Indonesia

I think this article deserves to be indexed and has benefits for improving the health of adolescent women in Indonesia, especially reducing anemia through the consumption of cookies that are high in Fe, folic acid, Vitamin B12 and antioxidants. In general, I consider that the background and benefits of the research are clearly written. However, it is still found Indonesian, namely AKG (Angka Kecukupan Gizi), it must be translated into English as the Recommended Dietary Allowance (RDA). The data and processing are presented quite well through pictures but the discussion is too short and there is no comparison with the results of other studies. I hope that the author can provide an explanation why the content of Fe, Folate, vitamin B12 and antioxidants in F2 are lower than F1?

As a suggestion for further research, it is necessary to carry out a sensory evaluation of the cookies produced, considering that the addition of porang and tempeh can change the color, taste and aroma of the product.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Nutrition and Food

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 16 February 2022

<https://doi.org/10.5256/f1000research.77755.r101869>

© 2022 Alamu E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Emmanuel Oladeji Alamu

Food and Nutrition Sciences Laboratory, International Institute of Tropical Agriculture (IITA),
Ibadan, Nigeria

Dear Editor,

This research article contains information on using porang tubers, moringa leaves, and tempe to develop cookies with high iron (Fe), folic acid, cobalamin (vitamin B12), and antioxidants for adolescents with anemia. The paper is not well written, and I have issues with the methodology, results, discussion, and conclusion sections. The methods were not well described and lacked basic information and references. These sections needed additional information. The authors should work on the results and discussion sections because they are not well presented in their current forms. The authors need to discuss their results better; currently, it lacks comparisons with the previous work on cookies.

I have the full comments directly inserted in the reviewed version of the manuscript that has been attached and can be accessed through this link:

<https://f1000researchdata.s3.amazonaws.com/supplementary/74045/5e3c9f70-90f1-4240-ac31-37489c75df0e.pdf>

I recommend that the authors do thorough work on the manuscript by responding to all my comments before publication.

Thank you.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Food Science and Technology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research