Check for updates

CASE REPORT

Case Report: The importance of examining colon and rectum in patients with appendiceal cancer [version 1; peer review: 3 approved with reservations]

Hugin Reistrup ¹, Siv Fonnes ¹, Jacob Rosenberg¹, Kristoffer Andresen¹

V1 First published: 26 Feb 2021, 10:152
<https://doi.org/10.12688/f1000research.50909.1>
 Latest published: 03 Mar 2022, 10:152
<https://doi.org/10.12688/f1000research.50909.2>

Abstract

Appendiceal cancer is rare and is often diagnosed incidentally in patients undergoing appendectomy for acute appendicitis. However, patients with appendiceal cancer are at increased risk of synchronous malignancy. In this case report, we present a 58-year-old man initially diagnosed with acute appendicitis after presenting to the emergency department with abdominal pain. He had an appendectomy and was discharged the following day. Unexpectedly, the postoperative histopathologic examination showed a primary adenocarcinoma in the appendix. A computed tomography scan showed rectal wall thickening and the patient was referred to colonoscopy where an experienced endoscopist found a rectal tumor during the digital rectal examination prior to the colonoscopy. The tumor was initially missed by the newly qualified doctor who examined the patient during his first admittance to hospital. The patient's two primary cancers were treated with a laparoscopic right hemicolectomy for the appendiceal cancer and a low anterior resection for the rectal cancer. This case supports the importance of a full colorectal workup in patients with appendiceal cancer. It also emphasizes the value of a thorough digital rectal examination and the need for improved focus on teaching and practice of the procedure.

Keywords

Appendicitis, appendix cancer, appendiceal cancer, synchronous, rectum cancer, rectal cancer, digital rectal examination, case report

This article is included in the **Oncology** gateway.

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)
03 Mar 2022
version 1
26 Feb 2021

1. Susanne Merkel ,
2. Cemil Yüksel ,
3. Wilhelm Graf,

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Hugin Reistrup (hugin.reistrup@gmail.com)

Author roles: **Reistrup H:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Fonnes S:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Rosenberg J:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Andresen K:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Reistrup H *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Reistrup H, Fonnes S, Rosenberg J and Andresen K. **Case Report: The importance of examining colon and rectum in patients with appendiceal cancer [version 1; peer review: 3 approved with reservations]** F1000Research 2021, 10:152 <https://doi.org/10.12688/f1000research.50909.1>

First published: 26 Feb 2021, 10:152 <https://doi.org/10.12688/f1000research.50909.1>

Introduction

Primary appendiceal neoplasms are rare and represent 1% of all gastrointestinal cancers, and the incidence is increasing^{1,2}. Typically, symptoms are vague and the patient often presents with acute appendicitis with the tumor being diagnosed incidentally during histopathological examination after appendectomy. In other cases, tumors can present as bowel obstruction or as a palpable pelvic mass³. Several studies have found that appendiceal neoplasms are associated with an increased risk of synchronous colorectal lesions^{1,3-6}. Multiple primary tumors can be divided into synchronous or metachronous tumors. Synchronous tumors are defined as tumors diagnosed less than six months apart while metachronous tumors are diagnosed more than six months apart⁷. It has been suggested, that the association between synchronous tumors in the appendix and the colon and rectum could be due to similar histological pattern and that the appendix derives embryologically from the cecum⁴. Guidelines recommend that patients with appendiceal neoplasms should undergo colonoscopy³. Also, guidelines call for abdominal examination and a digital rectal examination (DRE)³. DRE is a simple, quick, and inexpensive clinical procedure. One retrospective study found a sensitivity of 76% and specificity of 92% of the DRE for finding palpable rectal tumors⁸. Although it is a useful clinical tool for diagnostics and screening, DRE has a learning curve and the sensitivity is highly examiner-dependent⁹.

We present a patient with synchronous tumors in the appendix and rectum. This case underlines the importance of a full colorectal examination in patients diagnosed with primary appendiceal neoplasms and highlights the value of a thorough DRE. The case is presented in accordance with the CARE guideline¹⁰.

Case presentation

A 58-year-old Caucasian man with no prior medical or surgical history was admitted to the emergency department after referral from his general practitioner. The patient complained of constant diffuse abdominal pain for two days with exacerbation upon movement. He had had fever for one day. His stool was normal without blood and there was no nausea or vomiting. The patient had no family history of colorectal cancer. The abdominal examination revealed direct tenderness in both lower quadrants. The DRE that was performed by the newly qualified doctor on call was without palpable tumors. Laboratory blood tests showed elevated C-reactive protein (110 mg/L [normal value less than 3 mg/L]). All other blood tests were normal. An acute computed tomography (CT) scan of the abdomen showed signs of acute appendicitis. A diagnostic laparoscopy confirmed the diagnosis macroscopically and an uncomplicated laparoscopic appendectomy was performed. The patient was discharged the following day. The postoperative standard histopathologic examination of the appendix showed acute non-perforated appendicitis and, surprisingly, a 13 mm T1 goblet cell adenocarcinoma in the apex of the appendix with tumor-free resection margins. The patient was informed and underwent a new CT scan of the thorax and abdomen. The only new finding on the repeated CT scan was rectal wall thickening. The patient was discussed at a multidisciplinary team conference where it was decided to perform a colonoscopy due to

the rectal wall thickening and thereafter a laparoscopic right hemicolectomy due to the tumor in the appendix. Prior to the colonoscopy, the endoscopist, who is an experienced colorectal surgeon, performed a DRE. The endoscopist palpated the distal part of a tumor in the anterior wall of the rectum seven centimeters from the anal opening. The colonoscopy showed a tumor highly suspicious for malignancy with a central depression of 25 mm (Figure 1). It was classified as National Institute for Clinical Excellence (NICE) type 3. A magnetic resonance imaging (MRI) scan staged the tumor as a T2 without metastasis to lymph nodes or distant metastasis (Figure 2). The histopathologic examination of the rectal lesion showed a primary signet ring cell carcinoma that was histologically distinct from the appendiceal tumor. Hence, the patient had two primary synchronous

Figure 1. Colonoscopy showing the rectal tumor.

Figure 2. Magnetic resonance imaging scan showing the rectal tumor.

tumors. Eventually, the patient was treated with simultaneous laparoscopic right hemicolectomy and low anterior resection. **Figure 3** presents a timeline of the events. The postoperative course was complicated by an anastomotic leakage of the colorectal anastomosis. The leakage was treated with endoscopic vacuum-assisted closure. After surgery, the patient received adjuvant chemotherapy.

Discussion

This case illustrates the importance of a full colorectal workup in patients with a primary tumor in the appendix. This is supported by current clinical guidelines. The American Society of Colon and Rectal Surgeons' clinical guideline strongly recommends that a colonoscopy should be performed to exclude synchronous colorectal lesions in patients with appendiceal neoplasms, and also strongly recommends a complete physical examination including a DRE³.

In this case, the patient's rectal tumor was missed during the DRE performed by a newly qualified doctor in the emergency department. Studies have shown that newly qualified doctors lack confidence in performing a DRE, and the DRE is rarely repeated or supervised by a senior doctor^{11,12}. Also, clinicians often avoid performing a DRE^{12,13}. The DRE is a cost-efficient and quick procedure to assess and identify numerous conditions in both sexes, e.g. anal fissures, skin tags, pilonidal sinuses, anal fistulas, rectal prolapse, anal warts, skin diseases (e.g. dermatitis), anorectal tumors, gastrointestinal tract bleeding, abscesses, hemorrhoids, sphincter function, constipation, fecal impaction, prostatic hypertrophy, prostate tumors, prostatitis, pelvic inflammatory disease, pelvic floor prolapse (e.g. rectocele),

and pelvic floor dyssynergia^{12,14–16}. Hence, the importance of a sufficient DRE should not be underestimated.

Medical school is the foundation where students learn the necessary basic skills enabling them to practice medicine when becoming qualified doctors. Worryingly, senior medical students lack training in performing a DRE and up to 44% have never performed the examination on a human subject when graduating^{17,18}. A randomized controlled study found that medical students that had practiced DRE on phantoms and human volunteers had an increased confidence in performing and trusting the results of the examination compared with students who only practiced on phantoms¹⁹. Like many other practical clinical skills, the DRE has a learning curve. A questionnaire survey concluded that the more adequate the training in performing a DRE, the more confident the examiner was in making a diagnosis using the DRE¹². One study comparing novice to more experienced examiners on specially designed simulators found that the latter had a significantly higher detection rate for both prostate and rectal anomalies⁹. This highlights the importance of sufficient training and experience in DRE. When assessing a patient with abdominal or urogenital complaints, the abdominal examination is a core clinical skill. Many clinicians will experience that the abdominal examination is frequently repeated by a fellow and more experienced clinician, but this is not the case with the DRE^{11,18,20}. As both the abdominal examination and the DRE are standard and important clinical tools of screening in patients admitted to the emergency department, one could question why the abdominal examination is often supervised or repeated while the DRE is not. Nonetheless, the case presented here highlights the need for improved focus on the DRE in medical training.

Figure 3. Timeline of events. CT; computed tomography.

A strength of this case is that we have a very detailed report of the patient's history from his first contact to his general practitioner to several months after surgery. Also, we have full and detailed reports of the histopathologic examinations as well as both images and detailed reports of the CT scans, the MRI scan, and the colonoscopy. A limitation to this case is that we do not have information on the approximate total number of DREs performed by the newly qualified doctor initially admitting the patient to the hospital making it difficult to evaluate the doctor's experience in performing the procedure.

Conclusion

In this case report, we present a patient with primary synchronous cancers in the appendix and rectum. The case underlines

the importance of the DRE as a quick, inexpensive, and useful skill in the everyday clinic. Patients with primary appendiceal neoplasms should have a full colorectal workup and there should be an improved focus on teaching and training the DRE as it improves confidence and diagnostic accuracy of the procedure.

Data availability

All data underlying the results are available as part of the article and no additional source data are available.

Consent

Written informed consent for publication of clinical details and clinical images was obtained from the patient.

References

- Connor SJ, Hanna GB, Frizelle FA: **Appendiceal tumors: retrospective clinicopathologic analysis of appendiceal tumors from 7,970 appendectomies.** *Dis Colon Rectum.* 1998; **41**(1): 75–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Marmor S, Portschy PR, Tuttle TM, et al.: **The rise in appendiceal cancer incidence: 2000-2009.** *J Gastrointest Surg.* 2015; **19**(4): 743–750.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Glasgow SC, Gaertner W, Stewart D, et al.: **The American Society of Colon and Rectal Surgeons, clinical practice guidelines for the management of appendiceal neoplasms.** *Dis Colon Rectum.* 2019; **62**(12): 1425–1438.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Khan MN, Moran BJ: **Four percent of patients undergoing colorectal cancer surgery may have synchronous appendiceal neoplasia.** *Dis Colon Rectum.* 2007; **50**(11): 1856–1859.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Smeenk RM, van Velthuisen MLF, Verwaal VJ, et al.: **Appendiceal neoplasms and pseudomyxoma peritonei: a population based study.** *Eur J Surg Oncol.* 2008; **34**(2): 196–201.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lohsiriwat V, Vongjirad A, Lohsiriwat D: **Incidence of synchronous appendiceal neoplasm in patients with colorectal cancer and its clinical significance.** *World J Surg Oncol.* 2009; **7**: 51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vogt A, Schmid S, Heinimann K, et al.: **Multiple primary tumours: challenges and approaches, a review.** *ESMO Open.* 2017; **2**(2): e000172.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ang CW, Dawson R, Hall C, et al.: **The diagnostic value of digital rectal examination in primary care for palpable rectal tumour.** *Colorectal Dis.* 2008; **10**(8): 789–792.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Balkissoon R, Blossfield K, Salud L, et al.: **Lost in translation: unfolding medical students' misconceptions of how to perform a clinical digital rectal examination.** *Am J Surg.* 2009; **197**(4): 525–532.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gagnier JJ, Kienle G, Altman DG, et al.: **The CARE guidelines: consensus-based clinical case reporting guideline development.** *BMJ Case Rep.* 2013; **2013**: bcr2013201554.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yeung JMC, Yeeles H, Tang SW, et al.: **How good are newly qualified doctors at digital rectal examination?** *Colorectal Dis.* 2011; **13**(3): 337–340.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wong RK, Drossman DA, Bharucha AE, et al.: **The digital rectal examination: a multicenter survey of physicians' and students' perceptions and practice patterns.** *Am J Gastroenterol.* 2012; **107**(8): 1157–1163.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shrestha MP, Borgstrom M, Trowers E: **Digital rectal examination reduces hospital admissions, endoscopies, and medical therapy in patients with acute gastrointestinal bleeding.** *Am J Med.* 2017; **130**(7): 819–825.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Khadem TH, Rosenberg J, Achiam MP: **[Digital rectal examination is a useful clinical procedure in the work-up of patients with colorectal or urogenital symptoms].** *Ugeskr Laeger.* 2013; **175**(44): 2636–2639.
[PubMed Abstract](#)
- Talley NJ: **How to do and interpret a rectal examination in gastroenterology.** *Am J Gastroenterol.* 2008; **103**(4): 820–822.
[PubMed Abstract](#)
- Borkenhagen JF, Eastwood D, Kilari D, et al.: **Digital rectal examination remains a key prognostic tool for prostate cancer: a national cancer database review.** *J Natl Compr Canc Netw.* 2019; **17**(7): 829–837.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fitzgerald D, Connolly SS, Kerin MJ: **Digital rectal examination: national survey of undergraduate medical training in Ireland.** *Postgrad Med J.* 2007; **83**(983): 599–601.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lawrentschuk N, Bolton DM: **Experience and attitudes of final-year medical students to digital rectal examination.** *Med J Aust.* 2004; **181**(6): 323–325.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Patel MI, Kakala B, Beattie K: **Teaching medical students digital rectal examination: a randomized study of simulated model vs rectal examination volunteers.** *BJU Int.* 2019; **124** Suppl 1: 14–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Turner KJ, Brewster SF: **Rectal examination and urethral catheterization by medical students and house officers: taught but not used.** *BJU Int.* 2000; **86**(4): 422–426.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ? ? ?

Version 1

Reviewer Report 09 September 2021

<https://doi.org/10.5256/f1000research.54002.r91612>

© 2021 Graf W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Wilhelm Graf**

Thanks for letting me review this case report. My comments are as follows:

1. In the introduction it is stated that these tumors might present as bowel obstruction but in reality this is not true since the appendix is not in the GI passage.
2. The MTD conference should be added to the timeline figure.
3. Please state the body temperature, pulse and blood pressure.
4. Give more details of the MRI exam.
5. The listing of possible findings on DRE in the discussion could be shortened and focused just on malignant findings.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Surgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 Feb 2022

Hugin Reistrup

Dear Wilhelm Graf,

Thank you for your valuable comments, which we have read with great interest. We have answered your comments in a point-to-point format.

1) In the introduction it is stated that these tumors might present as bowel obstruction but in reality this is not true since the appendix is not in the GI passage.

Reply: We agree that this is an unlikely event, but although rare, bowel obstruction due to appendiceal neoplasm has been well described in the literature. We have also referred to a reference in the manuscript to back up this statement.

2) The MTD conference should be added to the timeline figure.

Reply: This is a very good point. We have now elaborated the decisions taken at the MDT conferences to the timeline figure.

3) Please state the body temperature, pulse and blood pressure.

Reply: We have now added this information to the "Case presentation" section.

4) Give more details of the MRI exam.

Reply: Unfortunately, there is no further information available on the MRI exam.

5) The listing of possible findings on DRE in the discussion could be shortened and focused just on malignant findings.

Reply: We fully understand your point regarding this. An important message in this study is to highlight the importance of a sufficient DRE. Also, we want to illustrate that the DRE can help diagnose a lot of different conditions, both benign and malign. Hence, we have chosen to make a comprehensive list of these conditions to underline this important point to the reader. As this case also has an educational aim, we are convinced that it is important to show the vast diagnostic benefits of this simple exam.

We hope that you find the comments and additions above to the manuscript sufficient.

Competing Interests: No competing interests were disclosed.

Reviewer Report 24 August 2021

<https://doi.org/10.5256/f1000research.54002.r91604>

© 2021 Yüksel C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cemil Yüksel

First of all, I would like to thank the authors for tackling a good topic. One of the most important topics in surgical training is rectal examination and it can sometimes be overlooked.

Some suggestions for the authors:

1. Are the tumor markers of the patient high? I think some blood values can be added to the article, only CRP is given.
2. Was there any peritoneal spread in the surgical findings? It is more likely to be seen in appendiceal tumors.
3. Colonoscopic view could have been better.
4. The tumoral lesion in colonoscopy could have been shown better.
5. Why was neoadjuvant treatment not planned for the patient?
6. Why was a protective ileostomy not planned for the patient? Because ileostomy should be performed for anastomosis safety in lower rectal tumors.
7. Has thoracic tomography been applied?
8. The postoperative condition of the patient has not been mentioned much. When did oral intake start?
9. The patient's family history was not mentioned.
10. Pathology results are not fully mentioned. Surgical margins? Number of lymph nodes removed? Metastasis?

As a result, the article has shortcomings in terms of surgery. However, the subject mentioned is important.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Partly

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Surgical Oncology, Gastric cancer, Colorectal cancer, Cytoreductive surgery, HIPEC

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 Feb 2022

Hugin Reistrup

Dear Cemil Yüksel,

Thank you for your valuable comments, which we have read with great interest. We have answered your comments in a point-to-point format.

1) Are the tumor markers of the patient high? I think some blood values can be added to the article, only CRP is given.

Reply: Unfortunately, we do not have tumor markers for this patient. In Denmark, tumor markers are not taken routinely when performing an uncomplicated appendectomy, and as soon as the appendiceal adenocarcinoma was diagnosed, a diagnostic colonoscopy was performed shortly afterwards. We agree that further blood samples would add value to the case and we therefore have added the leukocyte count to the "Case presentation" section.

2) Was there any peritoneal spread in the surgical findings? It is more likely to be seen in appendiceal tumors.

Reply: This is an interesting point, but no, there was no peritoneal spread. As stated in the manuscript, the laparoscopic appendectomy was uncomplicated and there were no other

abnormalities described during the laparoscopy. To further elaborate the laparoscopic findings, we have now added that the patient had a nonperforated appendicitis in the "Case presentation" section.

3) Colonoscopic view could have been better.

Reply: We completely agree that the photo (figure 1) from the colonoscopy could have been of a better quality. Unfortunately, this was the best colonoscopic view available. Still, we are convinced that it does add value to the case.

4) The tumoral lesion in colonoscopy could have been shown better.

Reply: Here, we presume that you are referring to the same photo as mentioned in comment no. 3. Therefore, we would like to refer to our response to that comment.

5) Why was neoadjuvant treatment not planned for the patient?

Reply: Unfortunately, this information is not supplied in the patient's journal. But according to Danish guidelines on neoadjuvant treatment of resectable rectal cancers, rectal tumors classified as T1 tumors are not treated with preoperative chemotherapy.

6) Why was a protective ileostomy not planned for the patient? Because ileostomy should be performed for anastomosis safety in lower rectal tumors.

Reply: This is a very good point. A protective loop ileostomy was actually performed. We have added this information to the "Case presentation" section.

7) Has thoracic tomography been applied?

Reply: Yes, as mentioned in the "Case presentation" section, a CT thorax and abdomen was performed. *"...The patient was informed and underwent a new CT scan of the thorax and abdomen."*

8) The postoperative condition of the patient has not been mentioned much. When did oral intake start?

Reply: The patient started oral intake the day after surgery, which is standard in Denmark following this type of surgery. We have deliberately chosen to omit details such as these from the manuscript to keep the manuscript short and concise.

9) The patient's family history was not mentioned.

Reply: As mentioned in the "Case presentation" section, there was no family history of colorectal cancer. *"...The patient had no family history of colorectal cancer"*.

10) Pathology results are not fully mentioned. Surgical margins? Number of lymph nodes removed? Metastasis?

Reply: We fully agree that this has not been elaborated sufficiently in the case, hence, we have added the following to the "Case presentation" section.

- "...Following surgery, the pTNM (pathological tumor-node-metastasis) classification for the appendiceal adenocarcinoma was T1N0M0 and for the rectal carcinoma T2N1M0"

- "...After the postoperative infection had been successfully managed, the patient received adjuvant chemotherapy for the rectal carcinoma. The decision to treat the patient with adjuvant chemotherapy was multifactorial: N1 staging, presence of tumor satellites, anastomotic leakage, and signet ring cell carcinoma with tumor tissue near the resection margin with a tumor deposit located 1.5 mm from the mesorectal fascia."

We hope that you find the comments and additions above to the manuscript sufficient.

Competing Interests: No competing interests were disclosed.

Reviewer Report 28 June 2021

<https://doi.org/10.5256/f1000research.54002.r87541>

© 2021 Merkel S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Susanne Merkel

The authors present an interesting case report of a common condition (acute appendicitis) with an incidental finding of increasing incidence (primary adenocarcinoma of the appendix). They emphasize the importance of rectal digital examination.

The case report is well written according to the CARE guidelines. However, I miss the pTNM classification of appendiceal carcinoma after appendectomy plus hemicolectomy and the pTNM classification of rectal carcinoma. I also miss the rationale for the indication for adjuvant chemotherapy: Synchronous multiple carcinomas? Signet ring cell carcinoma? Anastomotic leak? Lymph node metastases?

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Surgical oncology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 Feb 2022

Hugin Reistrup

Dear Susanne Merkel,

Thank you for your valuable comments, which we have read with great interest. We have answered your comments in a point-to-point format.

1) The case report is well written according to the CARE guidelines. However, I miss the pTNM classification of appendiceal carcinoma after appendectomy plus hemicolectomy and the pTNM classification of rectal carcinoma.

Reply: We agree that the pTNM classification would be of value to the present case. Therefore, we have added this to both the description of the appendiceal adenocarcinoma after appendectomy plus hemicolectomy (T1N0M0) and rectal carcinoma (T2N1M0) in the "Case presentation" section.

2) I also miss the rationale for the indication for adjuvant chemotherapy: Synchronous multiple carcinomas? Signet ring cell carcinoma? Anastomotic leak? Lymph node metastases?

Reply: We agree that the reason for choosing to treat the patient with adjuvant chemotherapy should be elaborated. The patient received neoadjuvant chemotherapy for the rectal carcinoma (signet cell ring carcinoma). The appendiceal carcinoma (goblet cell adenocarcinoma) did not require further treatment postoperatively. The oncologists based their decision on adjuvant chemotherapy on the following: N+, tumor satellite, anastomotic leakage, and signet ring cell carcinoma with tumor tissue near the resection margin with a tumor deposit located 1.5 mm from the mesorectal fascia. We have added these details to the "Case presentation" section.

We hope that you find the additions above to the manuscript sufficient.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research