RESEARCH ARTICLE

Regional disparities in postnatal care among mothers aged 15-49 years old in Indonesia [version 1; peer review: 2 approved]

Mochammad Nur Cahyono¹, Ferry Efendi ¹, Harmayetty Harmayetty¹, Qorinah Estiningtyas Sakilah Adnani^{2,3}, Hsiao Ying Hung⁴

¹Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

²Department of Midwifery, Karya Husada Institute of Health Science, Kediri, Indonesia

³Quality Maternal & Newborn Care Research Alliance, Yale University, Connecticut, USA

⁴Department of Nursing, College of Medicine, National Cheng Kung University, Tainan, Taiwan

V1 First published: 26 Feb 2021, 10:153
<https://doi.org/10.12688/f1000research.50938.1>
 Latest published: 16 Aug 2021, 10:153
<https://doi.org/10.12688/f1000research.50938.2>

Abstract

Background: In Indonesia, maternal mortality remains high, significantly 61.59% occur in the postnatal period. Postnatal care (PNC) provision is a critical intervention between six hours and 42 days after childbirth and is the primary strategy to reduce maternal mortality rates. However, underutilization of PNC in Indonesia still remains high, and limited studies have shown the regional disparities of PNC in Indonesia.

Methods: This study aims to explore the gaps between regions in PNC service for mothers who have had live births during the last five years in Indonesia. This study was a secondary data analysis study using the Indonesian Demographic and Health Survey (IDHS) in 2017. A total of 13,901 mothers aged 15-49 years having had live births within five years were included. Chi-squared test and binary logistic regression were performed to determine regional disparities in PNC.

Results: Results indicated that the prevalence of PNC service utilization among mothers aged 15-49 years was 70.94%. However, regional gaps in the utilization of PNC service were indicated. Mothers in the Middle of Indonesia have used PNC services 2.54 times compared to mothers in the East of Indonesia (OR = 2.54; 95% CI = 1.77-3.65, $p < 0.001$). Apart from the region, other variables have a positive relationship with PNC service, including wealth quintile, accessibility health facilities, age of children, childbirth order, mother's education, maternal occupation, husband's age, and husband's education.

Conclusion: Structured policies are needed to reduce gaps in areas with low service utilization. Developing innovative strategies to address PNC inequality in maternal services to improve maternal health is expected.

Open Peer Review

Approval Status

	1	2
version 2 (revision) 16 Aug 2021		
version 1 26 Feb 2021	view	view
1. Asmaa Salah Eldin Mohamed Saleh , Beni Suef University, Cairo, Egypt		
2. Kusrini S. Kadar , Universitas Hasanuddin, Makassar, Indonesia La Trobe University, Melbourne, Australia		

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

postnatal care, regional disparities, reduced inequalities

Corresponding author: Ferry Efendi (ferry-e@fkip.unair.ac.id)

Author roles: **Cahyono MN:** Conceptualization, Formal Analysis, Methodology, Writing – Original Draft Preparation; **Efendi F:** Conceptualization, Methodology, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Harmayetty H:** Conceptualization, Methodology, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Adnani QES:** Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Hung HY:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Cahyono MN *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Cahyono MN, Efendi F, Harmayetty H *et al.* **Regional disparities in postnatal care among mothers aged 15-49 years old in Indonesia [version 1; peer review: 2 approved]** F1000Research 2021, 10:153
<https://doi.org/10.12688/f1000research.50938.1>

First published: 26 Feb 2021, 10:153 <https://doi.org/10.12688/f1000research.50938.1>

Introduction

Maternal morbidity and mortality are a serious global health challenge. In 2019, the World Health Organization (WHO) revealed that 94% of maternal mortality occurred in low and middle-income countries, of which Indonesia is one (World Health Organization, 2019; World Health Organization *et al.*, 2015). During 2000 and 2017, the maternal mortality ratio plunged by about 38% worldwide. Even with this decline, Sub-Saharan Africa and Southern Asia accounted for approximately 86% of maternal deaths worldwide. Southern Asia alone accounted for nearly one-fifth (58,000), which demonstrated the struggle to improve maternal health (World Health Organization, 2019; World Health Organization *et al.*, 2015). In the Indonesia context, the government determined a target to reach an important goal of reducing maternal mortality rate to 102 per 100,000 live births in 2015 (Bappenas, 2015; Ministry of Health Republic of Indonesia, 2018). Despite the significant efforts to expand maternal health programmes, recent evidence showed that Indonesia was off-track to reach the Millennium Development Goals (MDGs) target by 2015. In 2015, maternal mortality deaths in Indonesia were three times higher than the MDGs target (Ministry of Health Republic of Indonesia, 2018; UNICEF & World Health Organization, 2015; World Health Organization *et al.*, 2015). Hence, reducing maternal mortality ratio to less than 70 per 100,000 by 2030 as one of the Sustainable Development Goals (SDGs) target could be a critical challenge for Indonesia (UNICEF & World Health Organization, 2015; United Nations, 2017; World Health Organization, 2012).

Most maternal mortality deaths are preventable or treatable if skilled healthcare, such as midwives, is provided during the postnatal period. Critical interventions during the postnatal period must be delivered to prevent maternal mortality deaths (Lawn *et al.*, 2016; World Health Organization *et al.*, 2015). In the Indonesian setting, the cause of maternal mortality is predominantly due to postpartum haemorrhage, followed by indirect causes, such as heart disease, severe anaemia, malaria, HIV/ AIDS and hepatitis. Many factors have been linked to interventions of postpartum haemorrhage which cause emergency cases, and skilled health care required to respond effectively to emergencies (Adisasmita *et al.*, 2015; Mahmood *et al.*, 2018). However, PNC remains a critical intervention to reduce maternal deaths in Indonesia. A new health programme called EMAS (Expanding Maternal and Neonatal Survival), was implemented by the Indonesian government in 2012, focused on improving maternal health care. The programme aimed at ensuring that every woman has access to quality maternal healthcare, including childbirth assistance by skilled health personnel in healthcare facilities (a target of 2018 strategic plan: 82%), four visits of Antenatal care (ANC) (78%), PNC, and providing ANC (87%). However, the national data revealed that utilization of PNC among 34 provinces in Indonesia remain varied and was considered lower compared to the childbirth assistance by skilled health personnel coverage. Socioeconomic, geographical, and demographic factors influence the underutilization of PNC. Current systematic reviews show that levels of

education, poverty, and limitations of access to PNC services are common issues in low-and middle-income countries linked to inequities in the use of PNC services (Langlois *et al.*, 2015).

In an Indonesia context, the national data in 2018 revealed that the average percentage of PNC visits for the first time in Indonesia was 93.3%. The highest percentage of visits occurred in Yogyakarta (99.6%) and the lowest percentage of visits was in Papua (56.3%). However, there were regional gaps in PNC visits across the provinces in Indonesia. Also, PNC utilization in rural areas was lower than urban areas in Indonesia (Kementerian Kesehatan Republik Indonesia, 2018; Ministry of Health Republic of Indonesia, 2018; Probandari *et al.*, 2017). It is worth noting that 61.59 % of maternal mortality rates occurred during the postnatal period in Indonesia. Additionally, evidence shows that the quality of PNC is lower in most districts and cities among the Eastern Region in Indonesia. PNC must be performed a minimum of three times: within the first six hours to the third day after the delivery, from the fourth day to the 28th after the delivery and the 29th day to the 42nd day after childbirth. The standards of PNC including examination for vital signs; the apex of the uterus; lochia and other per vagina fluids; breasts and counselling for exclusive breastfeeding; provision of communication, information about and education of PNC, and family planning (Ministry of Health Republic of Indonesia, 2018; Probandari *et al.*, 2017).

Several studies show that regional disparities in PNC occur in several countries. In Ethiopia, there were differences in each region and variations at regional levels at the utilization of PNC among women (Sisay *et al.*, 2019). PNC service in Zambia was also reported to experience regional disparities (Jacobs *et al.*, 2017). In the same vein, in Bangladesh, disparities in the utilization of maternal health services were also reported (Raheem *et al.*, 2019). However, research focused on regional disparities on PNC among mothers aged 15–49 years old in Indonesia are not well investigated. This study was conducted to analyze the gap between regions in PNC service utilization among mothers aged 15 – 49 years old who have had live births during the last five years in Indonesia. This study is significant because it can be a source of information and a reference regarding regional disparities in the utilization of PNC services in Indonesia. This research could complete a bigger picture for consideration in resolving disparities maternal services in Indonesia.

Methods

This study analyzed and reported unit data from the 2017 Indonesian Demographic Data Survey (IDHS) collected by the Inner City Fund (ICF). This study is part of the International Demographic and Health Survey (DHS) program. In this study, unit data consist of women aged 15–49 years old having had live births in the last five years preceding the survey.

The purpose of the cross-sectional study of the IDHS conducted by the ICF from July 24 to September 30, 2017 was to offer up-to-date projections of basic demographic and health indicators. The IDHS study demonstrates a broad overview of population problems in Indonesia.

We utilized the data conducted by the national and provincial representatives. This cross-sectional study represents 1,970 census blocks urban and rural areas in Indonesia. The census block obtained 59,100 female respondents aged 15–49 years old. The survey employed a two-stage stratified cluster sampling method. The first stage was the selection of several census blocks by systematic sampling proportional size. In the second stage, 25 ordinary households were selected with systematic sampling from the listing. In this study, a sample of 13,901 women aged 15–49 years from 34 provinces in Indonesia was analyzed. The inclusion criteria were taken from IDHS that included all women aged 15–49 years who had given birth in the last five years. The exclusion criteria were whether the variables incomplete or not available.

Ethical consideration

Ethical review boards approved ethical clearance for the Inner City Fund OCR Macro (number 45 CFR 46) and the national board review from the Ministry of Health of Republic Indonesia.

Before the survey, an informed consent was obtained from the respondents based on voluntary participation.

Variables

The dependent variable of this study was PNC visits. According to the recommendation of the Ministry of Health Republic of Indonesia, PNC must be performed at minimum three times: at the first six hours to the third day after the delivery, on the fourth day to the 28th after the delivery and the 29th day to the 42nd day after childbirth (Ministry of Health Republic of Indonesia, 2018). This data was based on the mother's perception of PNC utilization during the postnatal period. Independent variables analyzed in this study were the related geographic and socioeconomic factors, including a region of residence, the place of residence, wealth quintile, health insurance, access to health facility, age, gender, birth rank, education, and occupation.

The residence region was grouped as six regions, namely Sumatera, Jawa, Bali & Nusa Tenggara, Kalimantan, Sulawesi, Maluku & Papua, and of which were also categorized as West Indonesia, Middle Indonesia and East Indonesia. The place of residence was determined as rural and urban areas. The wealth quintile of households was set into five categories: poorest, poorer, middle, richer, and richest. The wealth quintile of households was scored based on wealth criteria (DHSProgram, 2016). Health insurance was divided into two categories, namely yes and no. Access to the health facility was categorized into two, namely difficult and not. Children's age was divided into five categories: less than one month, one month, two months, three months, four months. The gender of the child, namely female and male. Birth rank was categorized as first child, second, third, fourth and so on. Mother's age divided into six categories: 15–24, 25–29, 30–34, 35–39, 40–49, and 45–49, while husband's age divided into seven categories: 11–20, 21–30, 31–40, 41–50, 51–60, 61–70, 71–80. Mother's and husbands' education levels were grouped into no education, primary, secondary, and

higher education. Mother's and husband's occupation was divided into two categories: not working and working.

Statistical analysis

Data were analyzed using STATA version 16.0. The descriptive statistics method was also utilized for showing what kinds of data. Due to variables, the chi-square test was performed to determine variables correlated to the PNC utilization. Binary logistic regression was utilized to determine disparity in this study. Measurement of associations among variables was expressed as Odds Ratio (OR) and 95% Confidence Interval (CI). Significant variables were tested with a p-value of 0.05 and 95% CI, which are considered the disparity in PNC among mothers aged 15–49 years in Indonesia.

Results

A total of 13,901 women aged 15–49 years old with live births in the last five years preceding the survey were interviewed. Table 1 shows the bivariate analysis that there were ten categories among some variables associated with the utilization of PNC visits (p-value < 0.05). These variables include geographic factors, region, socio-economy (wealth quintile and access to the health facility), children (age of child and birth rank), mother factors (age, education and occupation), husband factors (age, education, and occupation). Residence, socioeconomic (health insurance ownership), child gender, mother's age and husband's occupation did not show associations with the utilization of PNC visits among mothers aged 15–49 years old in Indonesia (Table 1). More detail results can be found in Table 1.

Table 2 shows that the least distribution of respondents is in the East of Indonesia. Nearly half of participants who live in the East of Indonesia did not use the PNC services. More than half of participants who live in the East of Indonesia was classified as lowest. Interestingly, most participants who live in the East of Indonesia stated that there was no problem with access to the health facility. More detail characteristics of participants can be found in Table 2.

Table 3 reveals that nearly half of participants who live in Sumatera, Kalimantan, Maluku and Papua did not use the PNC service. The highest number of participants classified as lowest was found in Maluku & Papua. For health insurance ownership, more participants who live in Kalimantan did not have health insurance. The highest number participants having four and more children per household were found in Maluku & Papua (Table 3).

In multivariate analysis, the participants who live in the Middle of Indonesia utilized PNC services 2.54 times more than the participants who live in the West of Indonesia (OR = 2.54; 95% CI = 1.77–3.65). The participants who live in the East of Indonesia had 0.71 fewer odds than participants in Indonesia's West (OR = 0.71; 95% CI = 0.52–0.96). The participants who live in Jawa were 1.46 times more likely to use PNC services (OR = 1.46; 95% CI = 1.26–1.69) compared to participants living in Sulawesi (OR = 0.53; 95% CI = 0.35–0.80).

Table 1. Socio-demographic characteristic of participants by utilization of PNC in Indonesia (n=13,901).

Characteristic	Utilization of PNC				X ²	p-value
	No		Yes			
	n	%	n	%		
The geographic factors						
Indonesia						
West Indonesia	3265	29.1	7946	70.9	41.45***	0.00
Middle Indonesia	611	26.5	1693	73.5		
East Indonesia	164	42.4	222	57.6		
Region						
Sumatera	1056	34.3	2027	65.7	116.36***	0.00
Jawa	2028	26.3	5691	73.7		
Bali & Nusa Tenggara	232	26.5	642	73.5		
Kalimantan	291	33.5	578	66.5		
Sulawesi	270	27.8	701	72.2		
Maluku & Papua	164	42.4	222	57.6		
Place of residence						
Rural	2051	28.6	5116	71.4	1.45	0.49
Urban	1989	29.5	4745	70.5		
Socioeconomic factors						
Wealth quintile						
Poorest	928	34.6	1757	65.4	58.56***	0.00
Poorer	827	29.5	1980	70.5		
Middle	781	26.9	2127	73.1		
Richer	753	26.3	2109	73.7		
Richest	751	28.4	1888	71.6		
Health insurance						
No	1713	29.8	4026	70.2	3.00	0.19
Yes	2326	28.5	5835	71.5		
Access to the health facility						
Difficult	555	36.0	985	64.0	41.46***	0.00
Not	3485	28.2	8877	71.8		
Child's factors						
Age of child (in month)						
Less than one month	1007	33.0	2045	67.0	58.13***	0.00
One month	938	29.9	2198	70.1		
Two months	840	30.0	1956	70.0		
Three months	631	24.7	1929	75.3		
Four months	623	26.4	1733	73.6		
Sex of child						
Male	2118	29.9	4963	70.1	5.11	0.06
Female	1922	28.2	4898	71.8		

Characteristic	Utilization of PNC				X ²	p-value
	No		Yes			
	n	%	n	%		
Birth rank						
First child	1251	27.1	3368	72.9	41.25***	0.00
Second	1400	28.7	3469	71.3		
Third	761	29.0	1864	71.0		
Fourth and more	628	35.1	1160	64.9		
Mother's factors						
Age of mother in year						
15–24	796	30.6	1808	69.4	6.26	0.52
25–29	1032	29.2	2502	70.8		
30–34	995	27.7	2594	72.3		
35–39	808	29.0	1975	71.0		
40–44	336	29.3	809	70.7		
45–49	73	29.5	174	70.5		
Mother's level of education						
No education	63	50.0	63	50.0	55.19***	0.00
Primary	1117	31.2	2458	68.8		
Secondary	2357	28.9	5803	71.1		
Higher	503	24.6	1537	75.4		
Mother's occupation						
Working	1917	27.2	5136	72.8	24.97***	0.0001
Not working	2123	31.0	4725	69.0		
Husband's factor						
Age of husband in year						
11–20	44	45.8	52	54.2	23.83**	0.0068
21–30	1164	29.6	2772	70.4		
31–40	1901	28.2	4843	71.8		
41–50	817	29.2	1985	70.8		
51–60	106	35.7	192	64.3		
61–70	8	33.8	15	66.2		
71–80	0	0.0	2	100.0		
Husband's level of education						
No education	51	37.0	86	63.0	39.21***	0.0001
Primary	1254	32.5	2610	67.5		
Secondary	2255	28.0	5803	72.0		
Higher	479	26.0	1362	74.0		
Husband's occupation						
Working	4016	29.1	9799	70.9	0.15	0.72
Not working	23	27.2	63	72.8		

*p<0.05; **p<0.01; ***p<0.001

Table 2. Socio-demographic characteristic of study participants PNC in Indonesia based on region (n=13,901).

The use of PNC	Indonesia							
	West Indonesia		Middle Indonesia		East Indonesia		Total	
	n	%	N	%	n	%	n	%
No	3265	29.12	611	26.51	164	42.45	4039	29.06
Yes	7946	70.88	1693	73.49	222	57.55	9861	70.94
Wealth quintile								
Poorest	1686	15.04	772	33.51	227	58.75	2685	19.32
Poorer	2224	19.84	516	22.40	67	17.30	2807	20.19
Middle	2465	21.99	403	17.48	40	10.42	2908	20.92
Richer	2516	22.44	314	13.63	32	8.28	2862	20.59
Richest	2320	20.69	299	12.98	20	5.25	2639	18.99
Health insurance								
No	4731	42.20	870	37.77	138	35.77	5739	41.29
Yes	6480	57.80	1434	62.23	248	64.23	8162	58.71
Access to the health facility								
Difficult	1176	10.49	294	12.75	70	18.03	1539	11.07
Not	10035	89.51	2010	87.25	317	81.97	12361	88.93
Age of child (in month)								
Less than one month	2461	21.95	495	21.48	97	25.02	3052	21.96
One month	2533	22.59	509	22.11	93	24.21	3136	22.56
Two months	2230	19.89	485	21.07	82	21.17	2797	20.12
Three months	2084	18.59	411	17.84	64	16.64	2560	18.42
Four months	1903	16.98	403	17.50	50	12.96	2356	16.95
Sex of child								
Male	5671	50.58	1208	52.42	203	52.56	7081	50.94
Female	5540	49.42	1096	47.58	183	47.44	6819	49.06
Birth rank								
First child	3843	34.28	686	29.77	90	23.28	4619	33.23
Second	4015	35.81	752	32.65	101	26.27	4869	35.03
Third	2100	18.74	453	19.68	71	18.45	2625	18.88
Fourth and more	1252	11.17	412	17.89	124	32.00	1788	12.86
Age of mother in year								
15–24	2080	18.55	445	19.33	79	20.46	2604	18.73
25–29	2853	25.44	579	25.14	102	26.28	3533	25.42
30–34	2916	26.01	577	25.05	95	24.69	3589	25.82

The use of PNC	Indonesia							
	West Indonesia		Middle Indonesia		East Indonesia		Total	
	n	%	N	%	n	%	n	%
35–39	2277	20.31	435	18.87	72	18.54	2783	20.02
40–44	902	8.04	214	9.29	29	7.45	1145	8.23
45–49	183	1.63	53	2.32	10	2.57	247	1.77
Mother's level of education								
No education	63	0.56	43	1.87	21	5.35	127	0.91
Primary	2855	25.47	633	27.47	87	22.52	3575	25.72
Secondary	6739	60.11	1213	52.66	208	53.84	8160	58.70
Higher	1554	13.86	415	18.00	71	18.29	2039	14.67
Mother's occupation								
Working	5473	48.82	1352	58.68	228	59.12	7053	50.74
Not working	5738	51.18	952	41.32	158	40.88	6847	49.26
Age of husband in year								
11–20	70	0.63	20	0.85	6	1.48	96	0.69
21–30	3135	27.96	672	29.15	129	33.46	3936	28.31
31–40	5474	48.83	1093	47.43	177	45.91	6744	48.52
41–50	2295	20.47	445	19.31	62	16.06	2802	20.16
51–60	220	1.96	67	2.91	11	2.92	298	2.14
61–70	15	0.14	7	0.31	1	0.17	23	0.17
71–80	1	0.01	1	0.03	0	0.00	2	0.01
Husband's level of education								
No education	73	0.65	49	2.14	15	3.77	137	0.99
Primary	3066	27.35	716	31.10	81	20.98	3864	27.80
Secondary	6653	59.34	1175	51.01	230	59.65	8058	57.97
Higher	1418	12.65	363	15.75	60	15.60	1841	13.25
Husband's occupation								
Working	11154	99.49	2285	99.20	376	97.32	13815	99.38
Not working	57	0.51	18	0.80	10	2.68	86	0.62
Total	11211	100.00	2304	100.00	386	100.00	13901	100.00

(Table 4). Details of the results of a multivariate analysis shown in Table 4.

Table 5 reveals the middle category women based on wealth index had PNC service's utilization increased by 1.25 greater odds (OR = 1.25; 95% CI = 1.07-1.47) more than the richer category mothers (OR = 1.23; 95% CI = 1.04-1.45). Participants who thought that access to the health facility was not a problem

had an odds ratio of 1.29 greater than participants who considered access to the health facility to be a major problem in using PNC service (OR = 1.29; 95% CI = 1.10-1.51). Mothers having children aged three months had used PNC service 1.43 times (OR = 1.43; 95% CI = 1.23-1.66) more than mothers with children aged four months (OR = 1.30; 95% CI = 1.12-1.51). Mothers who had a second child had utilized PNC service 0.88 times (OR = 0.88; 95% CI = 0.78-0.99)

Table 3. Socio-demographic characteristic of participants PNC in Indonesia based on region (n=13,901).

	Region												
The use of PNC	Sumatera		Jawa		Bali & Nusa Tenggara		Kalimantan		Sulawesi		Maluku & Papua		Total
	n	%	n	%	n	%	n	%	n	%	n	%	n
No	1056	34.25	2028	26.27	232	26.52	291	33.46	270	27.79	164	42.45	4039
Yes	2027	65.75	5691	73.73	642	73.48	578	66.54	701	72.21	222	57.55	9861
Wealth quintile													
Poorest	711	23.05	838	10.85	357	40.83	211	24.34	342	35.21	227	58.75	2685
Poorer	725	23.52	1401	18.16	172	19.67	203	23.43	238	24.54	67	17.30	2807
Middle	657	21.32	1722	22.31	121	13.88	205	23.60	162	16.72	40	10.42	2908
Richer	546	17.72	1914	24.80	115	13.10	143	16.41	112	11.57	32	8.28	2862
Richest	444	14.39	1844	23.89	109	12.52	106	12.22	116	11.96	20	5.25	2639
Health insurance													
No	1259	40.83	3238	41.95	356	40.74	437	50.33	311	32.04	138	35.77	5739
Yes	1824	59.17	4481	58.05	518	59.26	431	49.67	659	67.96	248	64.23	8162
Access to the health facility													
Difficult	373	12.10	744	9.64	104	11.88	104	11.98	144	14.87	70	18.03	1539
Not	2710	87.90	6974	90.36	770	88.12	764	88.02	826	85.13	317	81.97	12361
Age of child (in month)													
Less than one month	731	23.70	1638	21.23	174	19.88	178	20.50	235	24.17	97	25.02	3052
One month	679	22.03	1770	22.93	193	22.05	175	20.18	225	23.22	93	24.21	3136
Two months	628	20.37	1515	19.62	184	21.01	201	23.15	188	19.34	82	21.17	2797
Three months	578	18.76	1438	18.63	165	18.91	152	17.54	162	16.65	64	16.64	2560
Four months	467	15.13	1358	17.59	159	18.15	162	18.63	161	16.62	50	12.96	2356
Sex of child													
Male	1589	51.55	3886	50.35	467	53.40	431	49.65	505	52.02	203	52.56	7081
Female	1494	48.45	3832	49.65	407	46.60	437	50.35	466	47.98	183	47.44	6819
Birth rank													
First child	882	28.60	2838	36.77	251	28.67	260	29.97	298	30.74	90	23.28	4619
Second	1025	33.24	2845	36.87	308	35.26	313	36.05	276	28.43	101	26.27	4869
Third	655	21.25	1357	17.58	160	18.26	178	20.44	205	21.11	71	18.45	2625
Fourth and more	521	16.91	678	8.79	156	17.81	118	13.54	191	19.72	124	32.00	1788

The use of PNC	Region												
	Sumatera		Jawa		Bali & Nusa Tenggara		Kalimantan		Sulawesi		Maluku & Papua		Total
	n	%	n	%	n	%	n	%	n	%	n	%	n
Age of mother in year													
15–24	512	16.60	1474	19.10	149	17.07	180	20.76	210	21.64	79	20.46	2604
25–29	783	25.41	1953	25.30	220	25.16	248	28.60	227	23.39	102	26.28	3533
30–34	886	28.73	1931	25.02	227	26.00	221	25.47	228	23.51	95	24.69	3589
35–39	604	19.59	1610	20.86	173	19.83	136	15.70	188	19.39	72	18.54	2783
40–44	249	8.09	625	8.09	84	9.57	62	7.12	96	9.91	29	7.45	1145
45–49	49	1.59	126	1.63	21	2.36	20	2.35	21	2.17	10	2.57	247
Mother's level of education													
No education	31	1.01	24	0.31	32	3.70	9	1.01	10	1.00	21	5.35	127
Primary	721	23.40	1988	25.76	250	28.58	269	30.98	260	26.75	87	22.52	3575
Secondary	1782	57.81	4739	61.40	445	50.94	479	55.17	506	52.14	208	53.84	8160
Higher	548	17.78	967	12.53	147	16.78	112	12.85	195	20.10	71	18.29	2039
Mother's occupation													
Working	1684	54.63	3590	46.51	559	64.00	448	51.63	543	55.96	228	59.12	7053
Not working	1399	45.37	4129	53.49	315	36.00	420	48.37	427	44.04	158	40.88	6847
Age of husband in year													
11–20	23	0.74	42	0.54	8	0.89	9	1.04	9	0.90	6	1.48	96
21–30	829	26.90	2169	28.10	244	27.87	258	29.67	307	31.61	129	33.46	3936
31–40	1557	50.50	3725	48.26	435	49.80	412	47.42	438	45.15	177	45.91	6744
41–50	606	19.66	1627	21.08	164	18.78	159	18.30	184	18.96	62	16.06	2802
51–60	63	2.04	146	1.89	22	2.53	30	3.42	26	2.72	11	2.92	298
61–70	4	0.12	11	0.14	1	0.13	1	0.15	6	0.57	1	0.17	23
71–80	1	0.04	0	0.00	0	0.00	0	0.00	1	0.08	0	0.00	2
Husband's level of education													
No education	21	0.67	46	0.60	30	3.45	11	1.29	14	1.49	15	3.77	137
Primary	786	25.50	2140	27.72	266	30.38	267	30.74	324	33.43	81	20.98	3864
Secondary	1865	60.51	4564	59.14	430	49.20	490	56.43	478	49.23	230	59.65	8058
Higher	410	13.31	968	12.55	148	16.97	100	11.55	154	15.85	60	15.60	1841
Husband's occupation													
Working	3072	99.65	7673	99.41	865	98.93	865	99.55	964	99.33	376	97.32	13815
Not working	11	0.35	45	0.59	9	1.07	4	0.45	6	0.67	10	2.68	86
Total	3083	100.00	7718	100.00	874	100.00	869	100.00	970	100.00	386	100.00	13901

Table 4. Binary logistic regression of PNC utilization among mothers aged 15–49 years old in Indonesia based on geographical location.

Variable	The use of PNC			
	P	OR	95% CI	
			lower	upper
Indonesia				
West Indonesia	Ref.	1.0		
Middle Indonesia	0.00***	2.54	1.77	3.65
East Indonesia	0.03*	0.71	0.52	0.96
Region				
Sumatera	Ref.	1.0		
Jawa	0.00***	1.46	1.26	1.69
Bali & Nusa Tenggara	0.007**	0.57	0.38	0.86
Kalimantan	0.007**	0.66	0.48	0.89
Sulawesi	0.002**	0.53	0.35	0.80
Maluku & Papua	Omitted			

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.**Table 5.** Binary logistic regression of PNC utilization among mothers aged 15–49 years old in Indonesia.

Variable	The use of PNC			
	P	OR	95% CI	
			lower	upper
Wealth quintile				
Poorest	Ref.	1.0		
Poorer	0.07	1.15	0.99	1.35
Middle	0.007**	1.25	1.07	1.47
Richer	0.017*	1.23	1.04	1.45
Richest	0.86	1.02	0.85	1.22
Access to the health facility				
Difficult	Ref.	1.0		
Not	0.001**	1.29	1.10	1.51
Age of child (in month)				
Less than one month	Ref.	1.0		
One month	0.06	1.13	0.99	1.28
Two months	0.11	1.11	0.98	1.26
Three months	0.00***	1.43	1.23	1.66
Four months	0.001**	1.30	1.12	1.51

Variable	The use of PNC			
	P	OR	95% CI	
			lower	upper
Birth rank				
First child	Ref.	1.0		
Second	0.04*	0.88	0.78	0.99
Third	0.04*	0.86	0.74	0.99
Fourth and more	0.00***	0.68	0.58	0.81
Mother's level of education				
No education	Ref.	1.0		
Primary	0.02*	1.80	1.09	2.98
Secondary	0.03*	1.77	1.06	2.96
Higher	0.006**	2.11	1.24	3.59
Mother's occupation				
Working	Ref.	1.0		
Not working	0.007**	0.87	0.79	0.96
Age of husband in year				
11–20	Ref.	1.0		
21–30	0.016*	1.83	1.12	3.00
31–40	0.004**	2.07	1.26	3.40
41–50	0.003**	2.18	1.31	3.63
51–60	0.06	1.73	0.97	3.07
61–70	0.17	2.19	0.72	6.68
71–80	Omitted			
Husband's level of education				
No education	Ref.	1.0		
Primary	0.98	1.01	0.67	1.51
Secondary	0.50	1.15	0.76	1.74
Higher	0.48	1.17	0.76	1.83

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

more than mothers who had a third child (OR = 0.86; 95% CI = 0.74–0.99). Mothers who had higher education had 2.11 times the chance to utilize PNC visits (OR = 2.11; 95% CI = 1.24–3.59) compared to those with lower level education (OR = 1.80; 95% CI = 1.09–2.98). Mothers with husband's aged 41–50 had a higher chance of utilization PNC visits (OR = 2.18; 95% CI = 1.31–3.63) compared to mothers with husband's aged 31–40 (OR = 2.07; 95% CI = 1.26–3.40). Details of the results of a multivariate analysis shown in Table 5.

Discussion

This study aimed to examine the gap across the region in Indonesia for PNC utilization among mothers aged 15–49 years old using the 2017 IDHS data sets. PNC has been the primary strategy to improve maternal health outcomes to reduce the high maternal mortality deaths in Indonesia. Therefore, assessing the regional disparities in PNC may provide evidence for the government to resolve discrepancies in maternal services in Indonesia. This study demonstrated that the prevalence of PNC service utilization among mothers aged 15–49 years was 70.94%. This finding was higher than other research in Sub-Saharan Africa, and Ethiopia with Abebo & Tesfaye (2018) and Tessema *et al.* (2020), respectively finding that 47.9% and 52.48% of women had used PNC service. An intertwined complex factor, such as the health system, maternal health policies, and socio-cultural variations across countries, may hinder women's use of the PNC service.

The findings in this study revealed the geographic influence on mothers for utilization of PNC service. Mothers who settled in the Middle of Indonesia had increased odds of using PNC service, while those which lived in the East of Indonesia had decreased odds. This result is congruent with previous research carried out in Ethiopia (Sisay *et al.*, 2019). In Ethiopia, the geographic factor is correlated to the utilization of PNC service due to the region's level of development and location. Evidence in Indonesia shows that the socio-economic development, such as industrial, housing, public transportation, road facilities and health facilities in the East of Indonesia, lagged compared to the West of Indonesia (Ministry of Health Republic of Indonesia, 2018; Soewondo *et al.*, 2019; Suparmi *et al.*, 2018). Therefore, it is a call of action to provide equality in developing human resources and infrastructure to reduce the possibility of gaps.

Additionally, mothers who live in Java are 1.46 times more likely to use PNC services than mothers who live in Sulawesi where they are 0.53 less likely to utilize PNC services. Consistent with previous studies performed in Indonesia, Java has dominance development compared to other islands because this island is the centre of the Indonesian government (Laksono *et al.*, 2020). The Java island oriented and centred development model has harmed maternal health outcomes in Indonesia (Bappenas, 2018). Natural resources, human resources and facilities must be equal throughout Indonesia, so the gap between islands could be minimized.

This study revealed that the wealth index was significantly linked to PNC service utilization among mothers aged 15–49 years old in Indonesia. Mothers from the middle households based on the wealth index had PNC service's utilization increased by 1.25 greater odds (OR = 1.25; 95% CI = 1.07–1.47) more than the richer mothers (OR = 1.23; 95% CI = 1.04–1.45). Interestingly, this result was not consistent with that of research done in Pakistan, Ethiopia, and Tanzania, where mothers from the richer wealth quintile were significantly associated with the utilization of PNC services (Berhe *et al.*, 2019; Mohan *et al.*, 2017; Yunus *et al.*, 2013). However, in this

study we have found that, statistically, the odds ratio was quite similar between middle and richer households. Additionally, the previous study in other parts of Ethiopia demonstrated that sociodemographic factors, such as income, did not correlate with the use of PNC services (Angore *et al.*, 2018). Mothers from richer households were more likely to access the PNC services. Ownerships of consumer goods at home, such as motorcycles and cars, may increase the risk at ease transportation, making them have no strain to access the health facility.

The present study showed that access to the health facility was correlated to PNC service in Indonesia. With mothers who thought that access to the health facility was not a problem, they had an odds ratio of 1.29 greater in using PNC service than mothers who considered access to the health facility difficult. Previous research conducted in Malawi has found a significant association between the health facility and the utilization of PNC services. Other Ethiopia studies have demonstrated that physical accessibility plays an essential variable in health service utilization (Kim *et al.*, 2019; Tarekegn *et al.*, 2014). Access to the health facility is related to the costs incurred, which is influenced by having transportation to the health facility, which is considered expensive. Long and shorter distances, better roads, and better public transportation may increase access to the health facility, primarily in Indonesia, with its massive gaps in development across the country (Bappenas, 2018).

In this study, mothers having children aged three months increased the likelihood to use PNC services about 1.43 times (OR = 1.43; 95% CI = 1.23–1.66) more than mothers with children aged four months (OR = 1.30; 95% CI = 1.12–1.51). A similar study in Nepal showed that PNC service utilization in the early postnatal period was most likely due to the motherhood transition period (Sanjel *et al.*, 2019). The possible reason could be that mothers with fewer children, and younger children, may want information and petrified of complications during the postnatal period. In the Indonesia setting, the first neonatal examination (KN1) is carried out at 6–48 hours after baby is born, which is at the same time for the first PNC visit (KF1). The second neonatal examination (KN2) is carried out between 3–7 days with the second PNC visit (KF2). The third neonatal examination (KN3) occurs alongside the third PNC visit (KF3), which is between 8–28 days after birth (Ministry of Health Republic of Indonesia, 2018).

Also, mothers who have second child had a 0.88 times likelihood to utilize PNC service (OR = 0.88; 95% CI = 0.78–0.99) than mothers having a third child (OR = 0.86; 95% CI = 0.74–0.99). This is congruent with previous studies conducted in Ethiopia and India, which showed that the higher the child's birth order, the lower the utilization of PNC services (Ali & Chauhan, 2020; Sisay *et al.*, 2019). A possible reason could be that mothers who had more children were more likely experienced in motherhood and had appropriate knowledge from previous maternal experiences and childcare, hence restraining PNC service.

This study revealed that mothers who hold higher education qualifications have 2.11 times the chance of utilizing PNC visits (OR = 2.11; 95% CI = 1.24-3.59) compared to those with lower level education (OR = 1.80; 95% CI = 1.09-2.98). The finding is similar to a study conducted in Ethiopia, which revealed that a mother's education was significant to PNC utilization (Tarekegn *et al.*, 2014). The possible reason could be that educated women have a greater opportunity to be informed and are more aware of seeking advice and treatment from skilled healthcare personnel than uneducated women.

This study demonstrated that the husband's age was significantly linked to the utilization of PNC among mothers aged 15–49 years in Indonesia. Mothers who have husbands aged 41–50 have a higher chance of utilizing PNC visits than mothers who have husbands aged 31–40. These findings were in line with another study in India that showed that the husband's age was associated with PNC's wives' service (Jungari & Paswan, 2019). Husband's autonomy and power in decision-making regarding wife's needs, including their health care needs, remained persistent. Husband's age may be linked to the level of maturity and primary controller, which exacerbates their wives' access to the health service (Jungari & Paswan, 2019; Sekine & Carter, 2019).

To the best of our knowledge, this is the first study to examine the gaps across the region in utilization PNC service among mothers aged 15–49 years old in Indonesia, which is one of a country in Southeast Asia that contribute to the global

burden of maternal mortality rates in the world. The strengths of our study that we utilized the national and provincial data representatives, which internationally standardized. However, we also note some limitations. The IDHS data analyzed in this study was collected using the cross-sectional method and mother's recall preceding survey prone to the possibility bias information.

Conclusion

This study reveals the gap across the region for PNC utilization among mothers aged 15–49 years old in Indonesia. This study's findings provide evidence to complete the bigger picture for the government to resolve discrepancies in maternal services in Indonesia. Structured policies are needed to reduce gaps in areas with low service utilization. Developing innovative strategies to address inequality in maternal services to improve maternal health is inescapable. Future research should explore the interregional gaps and factors that cause maternal health service utilization by using a different platform.

Data availability

Data used in this study is available online from the Indonesian 2017 Demographic and Health Survey (DHS) website under the DHS VII recode column. Access to the dataset requires registration and is granted only for legitimate research purposes. A guide for how to apply for dataset access is available at: <https://dhsprogram.com/data/Access-Instructions.cfm>.

References

- Abebo TA, Tesfaye DJ: **Postnatal care utilization and associated factors among women of reproductive age Group in Halaba Kulito Town, Southern Ethiopia.** *Arch Public Health.* 2018; **76**: 9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Adisasmita A, Smith CV, El-Mohandes AAE, *et al.*: **Maternal characteristics and clinical diagnoses influence obstetrical outcomes in Indonesia.** *Matern Child Health J.* 2015; **19**(7): 1624–1633.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ali B, Chauhan S: **Inequalities in the utilisation of maternal health care in rural India: Evidences from national family health survey III & IV.** *BMC Public Health.* 2020; **20**(1): 369.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Angore BN, Tufa EA, Bisetegen FS: **Determinants of postnatal care utilization in urban community among women in Debre Birhan Town, Northern Shewa, Ethiopia.** *J Health Popul Nutr.* 2018; **37**(1): 10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bappenas: **Laporan pencapaian tujuan pembangunan milenium di Indonesia 2014.** Jakarta, Indonesia: Badan Perencanaan Pembangunan Nasional. 2015.
- Bappenas: **Prakarsa pemerintah daerah dalam upaya pengurangan kesenjangan wilayah dan pembangunan daerah.** Jakarta, Indonesia: Bappenas. 2018.
[Reference Source](#)
- Berhe A, Bayray A, Berhe Y, *et al.*: **Determinants of postnatal care utilization in Tigray, Northern Ethiopia: A community based cross-sectional study.** *PLoS One.* 2019; **14**(8): e0221161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DHSProgram: **Wealth Index.** 2016.
[Reference Source](#)
- Jacobs C, Moshabela M, Maswenyeho S, *et al.*: **Predictors of antenatal care, skilled birth attendance, and postnatal care utilization among the remote and poorest rural communities of Zambia: a multilevel analysis.** *Front Public Health.* 2017; **5**: 11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jungari S, Paswan B: **What he knows about her and how it affects her? Husband's knowledge of pregnancy complications and maternal health care utilization among tribal population in Maharashtra, India.** *BMC Pregnancy Childbirth.* 2019; **19**(1): 70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kementerian Kesehatan Republik Indonesia: **Laporan nasional riset kesehatan dasar: Jakarta.** 2018.
- Kim ET, Singh K, Speizer IS, *et al.*: **Availability of health facilities and utilization of maternal and newborn postnatal care in rural Malawi.** *BMC Pregnancy Childbirth.* 2019; **19**(1): 503.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Laksono AD, Rukmini R, Wulandari RD: **Regional disparities in antenatal care utilization in Indonesia.** *PLoS One.* 2020; **15**(2): e0224006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Langlois EV, Miszkurka M, Zunzunegui MV, *et al.*: **Inequities in postnatal care in low-and middle-income countries: a systematic review and meta-analysis.** *Bull World Health Organ.* 2015; **93**(4): 259–270G.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lawn JE, Blencowe H, Waiswa P, *et al.*: **Stillbirths: rates, risk factors, and acceleration towards 2030.** *Lancet.* 2016; **387**(10018): 587–603.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mahmood MA, Mufidah I, Scroggs S, *et al.*: **Root-cause analysis of persistently high maternal mortality in a rural district of Indonesia: Role of clinical care quality and health services organizational factors.** *BioMed research international.* 2018; **2018**(3673265): 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ministry of Health Republic of Indonesia: **Indonesia health profile Jakarta**. Indonesia: Ministry of Health Republic of Indonesia. 2018.

Mohan D, LeFevre AE, George A, *et al.*: **Analysis of dropout across the continuum of maternal health care in Tanzania: findings from a cross-sectional household survey**. *Health Policy Plan*. 2017; **32**(6): 791–799. [PubMed Abstract](#) | [Publisher Full Text](#)

Probandari A, Arcita A, Kothijah K, *et al.*: **Barriers to utilization of postnatal care at village level in Klaten district, central Java Province, Indonesia**. *BMC Health Serv Res*. 2017; **17**(1): 541. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Raheem E, Khan JR, Hossain MS: **Regional disparities in maternal and child health indicators: Cluster analysis of districts in Bangladesh**. *PLoS One*. 2019; **14**(2): e0210697. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sanjel K, Onta SR, Amatya A, *et al.*: **Patterns and determinants of essential neonatal care utilization among underprivileged ethnic groups in Midwest Nepal: a mixed method study**. *BMC Pregnancy Childbirth*. 2019; **19**(1): 310. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sekine K, Carter DJ: **The effect of child marriage on the utilization of maternal health care in Nepal: A cross-sectional analysis of Demographic and Health Survey 2016**. *PLoS One*. 2019; **14**(9): e0222643. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sisay MM, Geremew TT, Demlie YW, *et al.*: **Spatial patterns and determinants of postnatal care use in Ethiopia: findings from the 2016 demographic and health survey**. *BMJ Open*. 2019; **9**(6): e025066. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Soewondo P, Johar M, Pujisubekti R, *et al.*: **Inspecting primary healthcare centers in remote areas: Facilities, activities, and finances**. *Jurnal Administrasi Kesehatan Indonesia*. 2019; **7**(1): 89–98. [Publisher Full Text](#)

Suparmi, Kusumawardani N, Nambiar D, *et al.*: **Subnational regional**

inequality in the public health development index in Indonesia. *Glob Health Action*. 2018; **11**(sup1): 1500133. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tarekegn SM, Lieberman LS, Giedraitis V: **Determinants of maternal health service utilization in Ethiopia: analysis of the 2011 Ethiopian Demographic and Health Survey**. *BMC Pregnancy Childbirth*. 2014; **14**(1): 161. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tesema ZT, Yazachew L, Tesema GA, *et al.*: **Determinants of postnatal care utilization in sub-Saharan Africa: a meta and multilevel analysis of data from 36 sub-Saharan countries**. *Ital J Pediatr*. 2020; **46**(1): 175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

UNICEF and World Health Organization: **A decade of tracking progress for maternal, newborn and child survival the 2015 report**. Geneva, Switzerland. 2015.

[Reference Source](#)

United Nations: **Sustainable Development Goals 17 goals to transform our world**. 2017.

[Reference Source](#)

World Health Organization: **Countdown to 2015 Maternal, Newborn & Child Survival**. 2012.

[Reference Source](#)

World Health Organization: **Maternal mortality**. 2019.

[Reference Source](#)

World Health Organization, UNICEF, UNFPA, *et al.*: **Trends in maternal mortality: 1990 to 2015 Estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division**. 2015.

[Reference Source](#)

Yunus A, Iqbal S, Munawar R, *et al.*: **Determinants of postnatal care services utilization in Pakistan-insights from Pakistan demographic and health survey (PDHS) 2006-07**. *Middle East J Sci Res*. 2013; **18**(10): 1440–1447.

[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 29 July 2021

<https://doi.org/10.5256/f1000research.54033.r89395>

© 2021 Kadar K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Kusrini S. Kadar

¹ Faculty of Nursing, Universitas Hasanuddin, Makassar, Indonesia

² School of Nursing and Midwifery, La Trobe University, Melbourne, Vic, Australia

This is a very interesting study providing information of the utilization of PNC services in Indonesia. The authors have described the gaps across the region clearly. We can see which regions in Indonesia utilized more PNC service as well as other factors contributing to this utilization.

As this study using secondary data from Indonesian DHS in 2017, this situation might have changed now, perhaps it is good to add in the title the year of the survey.

In the method section, it is not very clear whether the authors explain the method used by the IDHS or the method for this study. The information provided in this section hinders the possibility of others to replicate this study. Need more detailed information, step-by-step of what the authors did in this method section¹.

In the variable, for example, the authors categorized the mothers' age into six groups however there is no reference of this categorizing whether using WHO definition or else. Need to provide more information regarding this section.

Statistical analysis is already clear and can be replicated. The description of the result also has been clear.

In the discussion part, the authors' explanation is very repetitive. Every paragraph has a similar pattern where it started from the finding, comparing with previous study and additional explanation. Although this pattern can be accepted, however somehow make this part very boring. The authors tend to start the statement with: *this study inline with.... the result congruent with...consistent with previous study...* If it's possible to change the pattern in every paragraph not only the word that is used.

Lastly, there are some words that can be changed such as **husband** with **spouse**.

Term for explaining regions in Indonesia can be changed (follow the time zone of Indonesia):

1. "West of Indonesia" with Western part/regions of Indonesia/Western Indonesia.
2. "Middle of Indonesia" with Central part/regions of Indonesia/Central Indonesia.
3. "East of Indonesia" with Eastern part/regions of Indonesia/Eastern Indonesia.

For language clarity, I encourage the authors to use an English language editing service to produce more clear sentences and paragraphs.

References

1. Thamrin SA, Arsyad DS, Kuswanto H, Lawi A, et al.: Predicting Obesity in Adults Using Machine Learning Techniques: An Analysis of Indonesian Basic Health Research 2018. *Front Nutr.* 2021; **8**: 669155 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Community health; health education; health promotion; nursing workforce

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Aug 2021

Ferry Efendi, Universitas Airlangga, Surabaya, Indonesia

1. We appreciate the reviewer's positive feedback and recommendations. We have added in the title the year of the survey as suggested. It would now read "Regional disparities in postnatal care among mothers aged 15-49 years old: An analysis of the Indonesian Demographic and Health Survey 2017".

2. Thank you for your feedback. The sentence have been revised as follow: This study was a secondary data analysis using the most recent data from the 2017 Indonesian Demographic Data Survey (IDHS) collected by the Inner City Fund (ICF). This study is part of the International Demographic and Health Survey (DHS) program. We clarify that in the data availability section the information required for other researchers to replicate this study. Data used in this study is available online from the Indonesian 2017 Demographic and Health Survey (DHS) website under the DHS VII recode column. Access to the dataset requires registration and is granted only for legitimate research purposes. A guide for how to apply for dataset access is available at: <https://dhsprogram.com/data/Access-Instructions.cfm>. While the guide for using datasets for analysis is available at <https://dhsprogram.com/data/Using-DataSets-for-Analysis.cfm>. Other researchers will be able to access the data set in the same way as the authors and the authors do not have special access rights that others do not have. Further, we clarify that unlike other secondary data, DHS secondary data is structured and well organized to produce high-quality data. We have added more information on methods section for the purpose of replicability.

Authors categorized the mothers' age into six groups based on the reference from DHS report, we have added the information on the methods section.

3. We appreciate the reviewer's positive feedback and recommendations. The discussion has been revised according to the reviewer suggestion. The sentence have been revised as follows:

for example:

- Among the geographic groups analyzed in this study, mothers who settled in the Middle of Indonesia had increased odds of using PNC service, while those who lived in the East of Indonesia had decreased odds. Similarly, in Ethiopia.

Other paragraphs:

- However, earlier research is done in Pakistan, Ethiopia, and Tanzania, revealed where mothers from the richer wealth quintile were significantly associated with the utilization of PNC services...
- Other Ethiopia studies have demonstrated that physical accessibility plays an essential variable in health service utilization...
- In this study, mothers having children aged three months increased the likelihood to use PNC services about 1.43 times (OR = 1.43; 95% CI = 1.23-1.66) more than mothers with children aged four months (OR = 1.30; 95% CI = 1.12-1.51). A similar study in Nepal showed that PNC service utilization in the early postnatal period was most likely due to the motherhood transition period.

4. We appreciate the reviewer drawing our attention to check words, which has been corrected in our resubmission with "spouse" and term for explaining regions in Indonesia as suggested (Western Indonesia, Central Indonesia and Eastern Indonesia).

5. The professional English language editing service has been utilized to clarify sentences.

Competing Interests: No competing interests were disclosed.

Reviewer Report 16 March 2021

<https://doi.org/10.5256/f1000research.54033.r80365>

© 2021 Saleh A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Asmaa Salah Eldin Mohamed Saleh

Community Health Nursing, Beni Suef University, Cairo, Egypt

Hello, best greeting - It is my pleasure to review this article, thanks to the editors and authors.

I found this article was written in good condition by following the scientific rules for writing and I found nothing to be critic on it except just a few notes that do not affect the article structure; like authors mention Java and Jawa in entire the article, I suggest unifying the term mentioned in the article.

In the conclusion section, the authors mentioned that *"Structured policies are needed to reduce gaps in areas with low service utilization. Developing innovative strategies to address PNC inequality in maternal services to improve maternal health is expected"*. It would be preferred to have clearly applicable recommendations which serve as the solution for the article hypothesis.

The references in the article must be written in chronologically way in the same pattern for all.

Table 3: Socio-demographic characteristic of participants PNC in Indonesia based on region (n=13,901). The total sum of (No) use of PNC is not right (4039) it is 4040, you need to review it.

Many thanks, with my best wishes.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Community health nursing, maternal and children health nursing, geriatric health nursing, health promotion, health education, occupational health nursing, public health nursing, primary health nursing, rehabilitation nursing.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Aug 2021

Ferry Efendi, Universitas Airlangga, Surabaya, Indonesia

1. We thank the reviewer for the thoughtful review of our manuscript. The term mentioned in the article, which is Java, has been complete and corrected.
2. Thank you for your suggestions. We have revised as follows: The results suggest the need for national policy focuses on service equality, accessible and reliable implementation to improve postnatal care utilization among mothers to achieve the maximum results for the Indonesian Universal Health Coverage plan.
3. The reference list has been completed and corrected.
4. We appreciate the reviewer drawing our attention to this error, which has been corrected in our resubmission (4040).

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research