RESEARCH ARTICLE

REVISED Pan-cancer chromatin analysis of the human vtRNA genes uncovers their association with cancer biology [version 2; peer review: 2 approved, 1 approved with reservations]

Rafael Sebastián Fort ^{1,2}, María Ana Duhagon ^{1,3}

¹Laboratorio de Interacciones Moleculares, Facultad de Ciencias, Universidad de la República, Montevideo, Montevideo, 11400, Uruguay

²Depto. de Genómica, Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Montevideo, 11600, Uruguay

³Depto. de Genética, Facultad de Medicina, Universidad de la República, Montevideo, Montevideo, 11400, Uruguay

v2 First published: 05 Mar 2021, 10:182
<https://doi.org/10.12688/f1000research.28510.1>
 Latest published: 09 Jun 2021, 10:182
<https://doi.org/10.12688/f1000research.28510.2>

Abstract

Background: The vault RNAs (vtRNAs) are a class of 84-141-nt eukaryotic non-coding RNAs transcribed by RNA polymerase III, associated to the ribonucleoprotein complex known as vault particle. Of the four human vtRNA genes, vtRNA1-1, vtRNA1-2 and vtRNA1-3, clustered at locus 1, are integral components of the vault particle, while vtRNA2-1 is a more divergent homologue located in a second locus. Gene expression studies of vtRNAs in large cohorts have been hindered by their unsuccessful sequencing using conventional transcriptomic approaches.

Methods: VtRNA expression in The Cancer Genome Atlas (TCGA) Pan-Cancer cohort was estimated using the genome-wide DNA methylation and chromatin accessibility data (ATAC-seq) of their genes as surrogate variables. The association between vtRNA expression and patient clinical outcome, immune subtypes and transcriptionally co-regulated gene programs was analyzed in the dataset.

Results: VtRNAs promoters are enriched in transcription factors related to viral infection. VtRNA2-1 is likely the most independently regulated homologue. VtRNA1-1 has the most accessible chromatin, followed by vtRNA1-2, vtRNA2-1 and vtRNA1-3. VtRNA1-1 and vtRNA1-3 chromatin status does not significantly change in cancer tissues. Meanwhile, vtRNA2-1 and vtRNA1-2 expression is widely deregulated in neoplastic tissues and its alteration is compatible with a broad oncogenic role for vtRNA1-2, and both tumor suppressor and oncogenic functions for vtRNA2-1. Yet, vtRNA1-1, vtRNA1-2 and vtRNA2-1 promoter DNA methylation predicts a shorter patient overall survival cancer-wide. In addition, gene ontology analyses of vtRNAs co-regulated genes identify a chromosome regulatory domain, epithelial differentiation, immune and thyroid cancer gene sets for specific vtRNAs. Furthermore, vtRNA expression patterns are

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)	view	view	view
09 Jun 2021			
version 1			
05 Mar 2021	view		

1. **Daniele Hasler** , University of Regensburg, Regensburg, Germany
2. **Deepanwita Sengupta**, Stanford University, Stanford, USA
3. **Yong Sun Lee**, National Cancer Center, Goyang, South Korea

Any reports and responses or comments on the article can be found at the end of the article.

associated with cancer immune subtypes and vtRNA1-2 expression is positively associated with cell proliferation and wound healing.

Conclusions: Our study presents the landscape of vtRNA chromatin status cancer-wide, identifying co-regulated gene networks and ontological pathways associated with the different vtRNA genes that may account for their diverse roles in cancer.

Keywords

vault RNA, vtRNA1-1, vtRNA1-2, vtRNA1-3, vtRNA2-1, nc886, mir-886, cancer, TCGA, DNA methylation, chromatin accessibility

This article is included in the **Cell & Molecular Biology** gateway.

This article is included in the **Glioblastoma Heterogeneity** collection.

Corresponding author: María Ana Duhagon (mduhagon@fcien.edu.uy)

Author roles: **Fort RS:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Duhagon MA:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Comisión Académica de Posgrado, UDELAR (Student Fellowship RF 2020); Comisión Sectorial de Investigación Científica (CSIC, UDELAR) (Research Grant MAD I+D 2016 #487); Agencia Nacional de Investigación e Innovación (ANII) (Student Fellowship RF 2016-2019); and Programa para el Desarrollo de las Ciencias Básicas (PEDECIBA, MEC). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2021 Fort RS and Duhagon MA. This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Fort RS and Duhagon MA. **Pan-cancer chromatin analysis of the human vtRNA genes uncovers their association with cancer biology [version 2; peer review: 2 approved, 1 approved with reservations]** F1000Research 2021, 10:182 <https://doi.org/10.12688/f1000research.28510.2>

First published: 05 Mar 2021, 10:182 <https://doi.org/10.12688/f1000research.28510.1>

REVISED Amendments from Version 1

In Abstract and Conclusion we reduced the conclusion of the transcription factor analysis, eliminating the “co-regulation”, since it was too speculative.

We added vtRNA2-2P pseudogen to the vtRNA3-1P analysis, to include the two vtRNA pseudogenes currently annotated (introduction, results, materials and methods, figure legends were modified accordingly).

We included a study of Y-RNA genes ATAC-seq as an external family of RNAPol III transcripts (results, materials and methods, figure legends were modified accordingly).

The addition of vtRNA2-2P and Y-RNA genes required an new version of SF3, and an updated ST1.

“MicroRNA precursor” was substituted to “generate small RNA” since the evidence of vtRNA processing and function in the microRNA pathway is not demonstrated for all of them.

“Expression” of vtRNA was substituted to “promoter DNA methylation, chromatin status or inferred expression” to make clear that the chromatin status just a surrogate of vtRNA expression but not their expression itself.

“Chromatin polymorphism” was substituted for “epigenetic variation” to describe the chromatin variability of vtRNA2-1.

“High Chromatin accessibility” was substituted for “Low DNA methylation” on the results section title to be accurate.

A comment about antecedents of RNAPol III-RNAPol II interaction was added to transcription factor result section to further justify the pertinency of the analysis. The involvement of RNAPol III in innate immune response was also commented.

TGFB1 was substituted by TGFBI.

Zenodo link was updated to include the new supplementary figure. <http://doi.org/10.5281/zenodo.4784353>

Any further responses from the reviewers can be found at the end of the article

Abbreviations

vtRNA	vault RNA
TCGA	The Cancer Genome Atlas
ATAC-seq	Assay for Transposase Accessible Chromatin with high-throughput sequencing
NoMe-seq	Nucleosome Occupancy and Methylome sequencing
OG	Oncogene
TSG	Tumor Suppressor Gene
TCGA	The Cancer Genome Atlas
TSS	Transcription Start Site
Ave	Average
SD	Standard Deviation
ACC	Adrenocortical carcinoma
BLCA	Bladder Urothelial Carcinoma
BRCA	Breast invasive carcinoma

CESC	Cervical squamous cell carcinoma and endocervical adenocarcinoma
CHOL	Cholangiocarcinoma
COAD	Colon adenocarcinoma
DLBC	Lymphoid Neoplasm Diffuse Large B-cell Lymphoma
ESCA	Esophageal carcinoma
GBM	Glioblastoma multiforme
HNSC	Head and Neck squamous cell carcinoma
KICH	Kidney Chromophobe
KIRC	Kidney renal clear cell carcinoma
KIRP	Kidney renal papillary cell carcinoma
LGG	Brain Lower Grade Glioma
LIHC	Liver hepatocellular carcinoma
LUAD	Lung adenocarcinoma
LUSC	Lung squamous cell carcinoma
MESO	Mesothelioma
OV	Ovarian serous cystadenocarcinoma
PAAD	Pancreatic adenocarcinoma
PCPG	Pheochromocytoma and Paraganglioma
PRAD	Prostate adenocarcinoma
READ	Rectum adenocarcinoma
SARC	Sarcoma
SKCM	Skin Cutaneous Melanoma
STAD	Stomach adenocarcinoma
TGCT	Testicular Germ Cell Tumors
THCA	Thyroid carcinoma
THYM	Thymoma
UCEC	Uterine Corpus Endometrial Carcinoma
UCS	Uterine Carcinosarcoma
UVM	Uveal Melanoma

Introduction

The vault RNAs (vtRNAs) are a class of eukaryotic middle size non-coding RNAs (84-141 nt) transcribed by the RNA polymerase III, which associates to the vault particle (<5 % of the total mass of the vault particle) (Boivin *et al.*, 2019; Kedersha & Rome, 1986; Kedersha *et al.*, 1991). Although this particle is the largest cellular ribonucleoprotein complex, its function remains scarcely understood. Most mammals have a single vtRNA gene (i.e., mouse and rat), but the human genome

has three copies, annotated as vtRNA1-1 (98bp), vtRNA1-2 and vtRNA1-3 (88bp) genes, which are located in the chromosome region 5q31.3 (locus vtRNA1) and code for the RNAs that are integral components of the vault particle. Lately, a transcript initially annotated as the human microRNA precursor hsa-mir-886 (Landgraf *et al.*, 2007), was re-classified as another human vtRNA homologue and consequently renamed as vtRNA2-1 (Stadler *et al.*, 2009). VtRNA2-1 is also located in chromosome 5 (5q31.1, locus vtRNA2) at 0.5Mb distance from the vtRNA1 cluster, where is placed between SMAD5 and TGFBI genes in an antisense direction. Existing evidence indicates that despite being a duplication of the vtRNAs of locus 1, the vtRNA2-1 is neither associated with the vault particle nor co-regulated with the vtRNA1 locus (Lee *et al.*, 2011; Stadler *et al.*, 2009). Nevertheless, the realization that only 5–20% of all the cellular vtRNA transcripts are associated to the vault particle suggests additional roles for the majority of vtRNA transcripts, independent of the vault particle (Kickhoefer *et al.*, 1998; Nandy *et al.*, 2009; van Zon *et al.*, 2001). The vtRNA3-1P and vtRNA2-2P are two additional vault RNA genes situated in chromosome X and 2 respectively, but were classified as a pseudogenes due to the absence of expression in cell lines and the presence of silencing mutations at the promoter (Büscher *et al.*, 2020; van Zon *et al.*, 2001).

Most of the current knowledge of vtRNA function focuses on vtRNA1-1 and vtRNA2-1 roles in viral infection and cancer. Functional studies found that vtRNA1-1 and vtRNA1-2 (but not vtRNA1-3) can interact directly with drugs as mitoxantrone, doxorubicin and etoposide (Gopinath *et al.*, 2005; Gopinath *et al.*, 2010; Mashima *et al.*, 2008). In addition, p62 protein interacts with vtRNA1-1 and inhibits the p62 dependent autophagy *in vitro* and *in vivo* (Horos *et al.*, 2019). The vault RNAs have also been linked to native immune response, because of their strong upregulation during the infection of Influenza A virus (IAV) and Epstein-Barr virus (EBV) (Amort *et al.*, 2015; Li *et al.*, 2015; Nandy *et al.*, 2009). These studies proposed a viral activation of host vtRNAs as a mechanism to maintain the inhibition of PKR signaling pathway, representing a viral strategy to circumvent host innate immunity.

Initial reports about vtRNA2-1 (locus vtRNA2) revealed that it is abundant in normal tissues while lowly expressed in cancer cell lines from different tissue origin (breast, melanoma, cervix, lung, oral and prostate), which is consistent with a tumor suppressive role in cancer (Lee *et al.*, 2011; Treppendahl *et al.*, 2012). Moreover, vtRNA2-1 was proposed as a new type of ncRNA functioning as a tumor suppressor gene (TSG) that inhibits PKR, and was consequently renamed as “nc886” (Golec *et al.*, 2019; Jeon *et al.*, 2012a; Jeon *et al.*, 2012b; Lee *et al.*, 2011). Then, its anti-proliferative and TSG function was described in prostate (Aakula *et al.*, 2016; Fort *et al.*, 2018; Ma *et al.*, 2020), skin (Lee *et al.*, 2019a), gastric (Lee *et al.*, 2014b) esophageal (Im *et al.*, 2020; Lee *et al.*, 2014a) and cholangiocarcinoma cells (Kunkeaw *et al.*, 2012). Conversely, a pro-proliferative and anti-apoptotic oncogenic role was proposed

for vtRNA2-1 in renal (Lei *et al.*, 2017), ovarian (Ahn *et al.*, 2018), thyroid (Lee *et al.*, 2016), cervical (Li *et al.*, 2017) and endometrial (Hu *et al.*, 2017) tissues. A recent finding proposing a vtRNA2-1/PKR loss mediated doxorubicin cytotoxicity introduced a novel view about its contribution to chemotherapy response (Kunkeaw *et al.*, 2019). In addition, a potential TSG role has been put forward for vtRNA1-3 in Myelodysplastic syndrome (MDS) (Helbo *et al.*, 2015).

Apart from their role as full length RNAs, vtRNAs are precursors of small RNAs. Indeed, small RNAs with microRNA like function were demonstrated to derive from vtRNA1-1 (svRNAs) and to repress the expression CYP3A4, a key enzyme of drug metabolism (Meng *et al.*, 2016; Persson *et al.*, 2009). VtRNA2-1 has also been shown to generate small RNAs that could act like microRNAs in different tissues, serving both as TSG in prostate (Aakula *et al.*, 2016; Fendler *et al.*, 2011; Fort *et al.*, 2020), bladder (Nordentoft *et al.*, 2012), breast (Tahiri *et al.*, 2014), colon (Yu *et al.*, 2011), lung (Bi *et al.*, 2014; Cao *et al.*, 2013; Gao *et al.*, 2011; Shen *et al.*, 2018) and thyroid cancers (Dettmer *et al.*, 2014; Xiong *et al.*, 2011) and as oncogene (OG) in renal (Yu *et al.*, 2014), colorectal (Schou *et al.*, 2014) and esophagus cancer (Okumura *et al.*, 2016) through the repression of specific mRNA transcripts in human cancer.

Different lines of evidence revealed that the vtRNA transcription can be controlled by promoter DNA methylation. The epigenetic control of vtRNA2-1 is complex and owns clinical relevance in several solid tumors including breast, lung, colon, bladder, prostate, esophagus, hepatic and stomach cancer (Cao *et al.*, 2013; Fort *et al.*, 2018; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Romanelli *et al.*, 2014; Treppendahl *et al.*, 2012; Yu *et al.*, 2020). Intriguing aspects of the epigenetic regulation of vtRNA2-1 locus comprise its dependence on the parental origin of the allele (Joo *et al.*, 2018; Paliwal *et al.*, 2013), and its sensitivity to the periconceptional environment ((Silver *et al.*, 2015) and subsequent independent studies (Richmond *et al.*, 2018; van Dijk *et al.*, 2018)). In addition, vtRNA1-3 promoter methylation was associated with significantly poor outcome in lower risk myelodysplastic syndrome patients (Helbo *et al.*, 2015).

Currently, transcriptomic sequencing is the benchmark technique to study global RNA expression (Stark *et al.*, 2019). Nonetheless, some classes of RNAs are elusive to the standard transcriptomics due to their stable RNA structure and the presence of modified bases or ends, which impair the cDNA synthesis and/or adapter ligation during the sequencing library preparation (Sendler *et al.*, 2011; Zheng *et al.*, 2015). The vtRNAs belong to this group due to their conserved stable stem/hairpin loop secondary structure. The latter, together with the lack of sequencing of 40–200bp-long RNAs in the conventional transcriptomic studies, which are mostly intended for small and long RNAs, probably delayed their study in comparison with other regulatory RNAs. In fact, the 50–300 nucleotides long RNAs are considered the black hole of RNA biology (Boivin *et al.*, 2018; Steitz, 2015). Nonetheless, since chromatin

accessibility plays a critical role in the regulation of gene expression, at some extent the transcription of an RNA can be inferred from the chromatin status of its promoter (Corces *et al.*, 2018; Duren *et al.*, 2017; Liu *et al.*, 2018; Park *et al.*, 2017b). Since mounting evidence has shown that vtRNAs expression is tightly controlled by chromatin accessibility, dependent on nucleosome positioning and promoter DNA methylation (Ahn *et al.*, 2018; Fort *et al.*, 2018; Helbo *et al.*, 2015; Helbo *et al.*, 2017; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Park *et al.*, 2017a; Park *et al.*, 2017b; Sallustio *et al.*, 2016; Treppendahl *et al.*, 2012), the chromatin structure is a suitable surrogate marker of vtRNA transcription and could be used as a proxy of their expression.

The expression, regulation, and role of the four human vtRNAs in normal and disease conditions are still poorly understood, and available knowledge indicates that they hold diverse tissue specific activities. The availability of genomic data of large sets of human tissues provided by The Cancer Genome Atlas (TCGA), allows the study of the human vtRNAs across 16 tissue types and normal/cancer conditions. Here, we analyze vtRNA genes chromatin of the TCGA patient cohort withdrawn from two approaches: the assay for transposase-accessible chromatin followed by NGS sequencing (ATAC-seq) to analyze chromatin accessibility (Buenrostro *et al.*, 2013; Buenrostro *et al.*, 2015; Corces *et al.*, 2018; Thurman *et al.*, 2012) and the Illumina Infinium Human Methylation 450K BeadChip to analyze the CpG methylation of DNA (Moran *et al.*, 2016). This led us to determine the patterns of transcriptional regulation of the four human vtRNAs throughout cancer tissues. We also evaluate the association between vtRNA promoter DNA methylation and the patient clinical outcome in all the available cancer types. Finally, looking for functional discoveries, we analyze vtRNA relations with immune subtypes and transcriptionally co-regulated gene programs. Our study reveals specific patterns of inferred expression for each vtRNA that support previous evidence and poses potential new roles and molecular programs in which they may participate across and intra cancer types, increasing the comprehension of the role of the human vtRNAs in health and disease.

Methods

ATAC-seq data

The ATAC-seq data obtained by The Cancer Genome Atlas (TCGA) consortium were retrieved from UCSC Xena Browser (Goldman *et al.*, 2020) on 03/10/2018. It comprises the genomic matrix TCGA_ATAC_peak_Log2Counts_dedup_promoter with normalized count values for 404 samples (385 primary solid tumor samples across 23 tissues). As is described in UCSC Xena Browser to calculate the average ATAC-seq values a prior count of 5 was added to the raw counts, then put into a “counts per million”, then log₂ transformed, then quantile normalized; the result is the average value in the file (log₂(count+5)-qn values) across all technical replicates and all biospecimens belonging to the same TCGA sample group. The gene promoter for ATAC-seq is defined as a region within -1000 to +100bp from the TSS site. Assignment of promoter peak to gene

mapping information derives from the peak summit within the promoter region. Peak location information of the ATAC-seq values for 500 bp gene promoter regions was retrieved from the file TCGA_ATAC_Log2Counts_Matrix.180608.txt.gz. All the analyses were performed for tissues with at least five primary tumor samples (21 tissues), which resulted in the exclusion of CESC (2 primary tumor samples) and SKCM (4 primary tumor samples). Correlations between ATAC-seq and DNA promoter methylation were performed for the 329 primary tumors that are studied by both strategies (Extended Data: Tables S1 and S2).

DNA methylation data

The DNA methylation data obtained from The Cancer Genome Atlas (TCGA) consortium was retrieved from UCSC Xena Browser (Goldman *et al.*, 2020) on date 03/10/2018. It comprises the normalized beta-value of DNA methylation obtained using Illumina Infinium Human Methylation 450 Bead-Chip arrays of the total (9149 samples) normal adjacent tissues (746 samples across 23 tissues) and primary solid tumors (8403 samples across 32 tissues) of the Pan-Cancer TCGA cohort. The current study is limited to solid tumors since only AML DNA methylation is available at the TCGA. All the analyses were performed for tissues with at least five samples. Normal adjacent tissues comprise 16 tissues, excluding CESC (two samples), GBM (two samples), PCPG (three samples), SARC (four samples), SKCM (two samples), STAD (two samples) and THYM (two samples). Primary tumors comprise 32 tissues. The normalized promoter average beta-values (500 bp bin) comprise the following CpG sites for vtRNA1-1: cg12532653, cg05913451, cg14633504, cg16615348, cg25602765, cg13323902, cg18296956; for vtRNA1-2: cg21161173, cg13303313, cg00500100, cg05174942, cg25984996, cg11807153, cg15697852; for vtRNA1-3: cg23910413, cg19065177, cg02053188, cg01063759, cg07379832, cg07741016; and for vtRNA2-1: cg16615357, cg08745965, cg00124993, cg26896946, cg25340688, cg26328633, cg06536614, cg18678645, cg04481923 (Extended Data: Tables S2–S5).

Analysis of transcription factors associated to vtRNA genes

The transcription factors (TFs) occupancy analysis of vtRNAs was performed at a ± 3000 bp region centered at each vtRNA gene. The CHIP-seq data was retrieved from the UCSC genome browser (GRCh37/hg19) (Kent *et al.*, 2002) and the K562 chronic myelogenous leukemia cell line was chosen since it has the richest TFs data of ENCODE dataset (ENCODE Transcription Factor Binding tracks - ENCODE 3 TFBS) (Euskirchen *et al.*, 2007). The KEGG pathways enrichment analysis of the core 23 TFs common to all vtRNAs was done using STRING software (Szklarczyk *et al.*, 2017) (Extended Data: Table S6).

Overall survival analysis

The Kaplan Meier curve analysis (Bland & Altman, 1998) was performed with software GraphPad Prism 6 using Log-ranked Mantel-Cox test. We analyzed the ATAC-seq normalized

values and DNA methylation average normalized beta-values for vtRNA promoters and survival data until 4000 days (99% of data). The log-rank statistics was calculated for quartiles of vtRNAs promoter chromatin accessibility comparison groups (25th and 75th) (*Extended Data: Tables S7 and S8*).

Pathway enrichment and cluster chromosome localization analyses

The genome wide correlation analysis of promoter chromatin accessibility (ATAC-seq normalized values) for each vtRNA in comparison with all the annotated genes was performed using the genomic matrix TCGA_ATAC_peak_Log2Counts_dedup_promoter retrieved from UCSC Xena Browser (385 primary tumor samples across 23 tissues) (Goldman *et al.*, 2020) on date 03/10/2018. The protein coding genes showing a Spearman correlation $r \geq 0.4$ were selected for pathway and cluster localization analysis. The pathway enrichment analysis was done using STRING software for the Gene Ontology Biological Process and KEGG pathways categories (Szkarczyk *et al.*, 2017). The cluster localization analysis was performed with two approaches, the Cluster Locator algorithm (Pazos Obregón *et al.*, 2018) and the Enrichr software (Kuleshov *et al.*, 2016) (*Extended Data: Tables S9 and S10*).

Immune subtypes data

The data of Immune Subtypes defined by Thorsson *et al.* (Thorsson *et al.*, 2018) is available at the Pan-Cancer TCGA and was retrieved from UCSC Xena Browser (Goldman *et al.*, 2020) on date 03/10/2018 and from the supplementary material of Thorsson *et al.* article (Thorsson *et al.*, 2018). ATAC-seq data of vtRNA promoters for the six Immune Subtype comprised a total of 364 samples: wound healing (105 samples), IFN- γ dominant (93 samples), inflammatory (110 samples), lymphocyte depleted (43 samples), immunologically quiet (nine samples), TGF- β dominant (four samples). DNA methylation average normalized beta-values of vtRNA promoters for the six Immune Subtype comprised 7693 samples including wound healing (1963 samples), IFN- γ dominant (2137 samples), inflammatory (2126 samples), lymphocyte depleted (933 samples), immunologically quiet (383 samples), TGF- β dominant (151 samples) (*Extended Data: Tables S11–S14*).

Statistical analysis

Unless specified all the variables are expressed as average value $\pm$ standard deviation (SD). Statistical analyses were performed with one-way ANOVA for multiple comparison tests, including Sidak's Honest Significant Difference test as a post-hoc test, as referred in the legend of the Figures. Spearman equations were applied to determine the correlations. All the analyses were done in R software (version 3.6) using libraries ggcorrplot, corrrplot, xlsx, heatmap.2 (clustering distance measured by Euclidean and Ward clustering algorithms) and software GraphPad Prism 6. The statistical significance of the observed differences was described using the p-value (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** p -value < 0.0001). Differences with a p-value of < 0.05 were considered significant.

Results

VtRNA genes have different chromatin accessibility at the promoter region

The Cancer Genome Atlas consortium incorporated ATAC-seq data for tumor samples of the GDC Pan-Cancer dataset (385 primary tumors samples across 23 cancer types (Corces *et al.*, 2018)). The ATAC-seq strategy is a genome-wide sequencing profiling approximation to chromatin accessibility using the hyperactive Tn5 transposase that simultaneously cut and ligate adapters preferentially to the accessible DNA regions. The counts of sequencing reads in a particular genomic region provides a direct measurement of chromatin accessibility (Buenrostro *et al.*, 2015; Tsompana & Buck, 2014). For those gene promoters regulated by DNA methylation, ATAC-seq is a more direct approximation to chromatin accessibility than promoter CpG DNA methylation measurements because the latter is well correlated but not an exact measurement of chromatin accessibility. Indeed, the effect of DNA methylation on chromatin structure depends on the gene and the distribution of the methylation sites along it. It is also important to mention that gene expression interpretations based solely on chromatin accessibility are blind to post-transcriptional regulatory events such as processing, localization, and stability of the RNA transcript. Nevertheless, since the chromatin status of the different vtRNA promoters has been positively correlated with the abundance of their transcripts (Ahn *et al.*, 2018; Fort *et al.*, 2018; Helbo *et al.*, 2015; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Park *et al.*, 2017a; Park *et al.*, 2017b; Sallustio *et al.*, 2016; Treppendahl *et al.*, 2012), ATAC-seq is likely a good surrogate of vtRNA expression. DNA CpG methylation data is available for 746 normal adjacent (across 23 tissues) and 8403 primary tumor (across 32 tissues) tissue samples of the Pan-Cancer TCGA cohort (*Extended Data: Tables S1–S5 and Figure S1*). The full data is listed in Tables S1–S5 and the tissue composition of the three datasets of the cohort, comprising tumor ATAC-seq, normal and tumor DNA methylation, is shown in Figure S1. The DNA methylation data has nine more tissue types and a higher number of samples than the others, while the abundance of some tissues is skewed to twice enrichment in 1-4 specific tissue types.

Although several reports demonstrated that the DNA methylation of vtRNA promoters is well correlated to vtRNA transcript expression in various tissues (Ahn *et al.*, 2018; Fort *et al.*, 2018; Helbo *et al.*, 2015; Helbo *et al.*, 2017; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Park *et al.*, 2017b; Sallustio *et al.*, 2016; Treppendahl *et al.*, 2012), the association between the DNA methylation and chromatin accessibility at their promoters was not previously investigated.

To assess the expression of vtRNAs in cancer tissues, we analyzed a 500 bp region, containing the full vtRNA genes, the proximal promoter region and the RNA polymerase III A and B box elements located downstream of the transcription start site (TSS) (Figure 1). Indeed, Vilalta *et al.*, 1994 demonstrated that the deletion of the 300bp 5'-flanking region of the rat

Figure 1. Genomic view of human vtRNA genes epigenetic features. Genomic view of 1.5 kb region of human vtRNA genes in UCSC Genome browser (GRCh37/hg19) centered at the 500bp bin highlighted in yellow, which was used for ATAC-seq and CpG methylation analyses. VtRNAs of locus 1 cluster (vtRNA1-1, vtRNA1-2, vtRNA1-3) and of locus 2 (vtRNA2-1) are in the sense and antisense orientation respectively. Several Gene annotation and ENCODE Project tracks for seven cell lines (GM12878, H1-hESC, HSMM, HUVEC, K562, NHEK, NHL) are displayed: DNA accessibility (DNaseI hypersensitivity clusters (color intensity is proportional to the maximum signal strength)), DNA methylation (for CpG islands length greater than 200 bp), histone modification (H3K27Ac, H3K4me1, H3K4me3 marks), conservation of the region in 100 Vertebrates (log-odds PhyloP scores). The vertical viewing range of the tracks displays the default settings of the browser for each variable.

vtRNA transcript largely reduced its transcription greater than 30-fold (Vilalta *et al.*, 1994). As depicted in Figure 1, this region bears epigenetic marks of transcriptionally active chromatin (Li *et al.*, 2007), including histone H3K27ac, H3K4me1, H3K4me3 at the promoters and DNaseI hypersensitivity clusters around the TSS in the cell types compiled by UCSC browser (Goldman *et al.*, 2013). Interestingly, vtRNA2-1 is the only vtRNA immersed in a CpG island and vtRNA1-1 displays the strongest epigenetic marks of transcriptional activity (Figure 1). The analysis of the ATAC-seq values of this 500 bp region of all the TCGA primary tumors reveals that the four human vtRNAs have different levels of chromatin accessibility (Figure 2A). Moreover, the average DNA methylation of the promoters (Figure 2B) in primary tumor samples supports this finding (32 tissues, *Extended Data: Figure S1* and *Tables S2* and *S4*). VtRNA1-1 promoter has the highest chromatin accessibility and the lowest DNA methylation, showing the smallest dispersion of values among the samples (Ave. 4.1, 0.6 SD) relative to the other three vtRNAs. Meanwhile, vtRNA1-2 (Ave. 2.9, 1.6 SD), vtRNA1-3 (Ave. 1.7, 1.2 SD) and vtRNA2-1 (Ave. 2.5, 1.4 SD) have broader ranges of ATAC-seq and DNA methylation values (Figure 2, Table 1, *Extended Data: Tables S1* and *S2* and *Figure S1*). A comparison among the vtRNAs shows that the average chromatin accessibility of their promoters is inversely proportional to their average DNA methylation (Table 1 and Figure 2) (relative ATAC-seq values are

vtRNA1-1 (4.1) > vtRNA1-2 (2.9) > vtRNA2-1 (2.5) and accordingly, relative DNA methylation are vtRNA1-1 (0.09) < vtRNA1-2 (0.4) < vtRNA2-1 (0.5)). Yet vtRNA1-3 average low chromatin accessibility is not accompanied by a comparatively denser promoter methylation (values 1.7 and 0.22 respectively). Nevertheless, the intra-sample correlation between the ATAC-seq and DNA methylation for the vtRNAs is between -0.24 and -0.78 (see rs-values in Table 1 and *Extended Data: Figure S2*), suggesting that regardless of the relatively smaller average effect of DNA methylation on vtRNA1-3 promoter accessibility, all the vtRNAs are regulated by DNA methylation to some extent (Table 1). In addition, the CpG DNA methylation negatively correlates with ATAC-seq values in the 500bp of the promoters of the four vtRNAs (Table 1 and *Extended Data: Figure S2*). The strength of the correlation for each vtRNA is higher for the two vtRNAs that have a broader spectrum of DNA methylation, i.e. vtRNA1-2 and vtRNA2-1 ($r_s = -0.74$ and $r_s = -0.78$ respectively) (Table 1 and *Extended Data: Figure S2B* and *S2D*). Meanwhile, vtRNA1-1 and vtRNA1-3 promoters, which have narrower range of DNA methylation (vtRNA1-1 has also a small range of chromatin accessibility) present a smaller association between both values ($r_s = -0.24$ and $r_s = -0.54$ respectively) (Table 1 and *Extended Data: Figure S2A* and *S2C*). Technical differences in ATAC-seq and DNA methylation array approaches, as well as naturally occurring non-linear correlation between average

Figure 2. Chromatin accessibility and DNA methylation of the vtRNA promoters. Analysis of vtRNA 500bp promoter region. **A.** Average chromatin accessibility ATAC-seq values for the 385 primary tumors available at the Pan-Cancer TCGA dataset (23 tissues types). **B.** Average DNA methylation beta-values for the 8403 primary tumors (32 tissues types). Dashed horizontal lines denote unmethylated (bottom gray, average beta-value ≤ 0.2), 50% methylated (middle red, average beta-value = 0.5) and highly methylated (top gray, average beta-value ≥ 0.8) promoters. The box plots show the median and the lower and upper quartile, and the whiskers the 2.5 and 97.5 percentile of the distribution.

Table 1. Correlation between vtRNA promoters ATAC-seq and DNA methylation values in the Pan-Cancer TCGA samples.

vtRNA	Average $\pm$ SD		Spearman correlation (ATAC-seq vs DNA methylation)	
	Promoter ATAC-seq	Promoter DNA methylation	R_s	p -value
vtRNA1-1	4.1 $\pm$ 0.6	0.09 $\pm$ 0.07	-0.24	<0.0001
vtRNA1-2	2.9 $\pm$ 1.6	0.42 $\pm$ 0.20	-0.74	<0.0001
vtRNA1-3	1.7 $\pm$ 1.2	0.22 $\pm$ 0.15	-0.54	<0.0001
vtRNA2-1	2.5 $\pm$ 1.4	0.47 $\pm$ 0.21	-0.78	<0.0001

The Spearman correlation was calculated for the 329 primary tumors studied by ATAC-seq and DNA methylation.

DNA methylation and chromatin structure may account for these differences (Corces *et al.*, 2018; Liu *et al.*, 2018). The low ATAC-seq value of the pseudogenes vtRNA3-1P (Ave. -1.2, 0.6 SD) and vtRNA2-2P (Ave. 0.1, 0.6 SD), represents a proof of concept of the analyses (Extended Data: Figure S3 and Tables S1-S5). The same approach was performed for the Y-RNAs genes RNY1 (Ave. 3.7, 0.4 SD), RNY3 (Ave. 2.1, 0.6 SD), RNY4 (Ave. 1.9, 0.6 SD) and RNY5 (Ave. 3.1, 0.4 SD) (Canella *et al.*, 2010), which code for ubiquitously expressed RNA polymerase III transcripts (Extended Data: Figure S3 and Tables S1).

Chromatin accessibility, DNA methylation and transcription factor binding at the four vtRNA promoters are correlated

Aiming to investigate a possible transcriptional co-regulation of vtRNA genes, we determined the pairwise correlations in chromatin accessibility and DNA methylation among the four vtRNA promoters. We found that all pairs of vtRNAs, except vtRNA1-2 and vtRNA2-1, show positive correlations in both datasets (Figure 3A and 3B) (r_s -values 0.23-0.41 for ATAC-seq and r_s -values 0.20-0.50 for DNA methylation). The unsupervised clustering of these epigenetic marks indicates that the three vtRNAs clustered at vtRNA1 locus (vtRNA1-1, vtRNA1-2 and vtRNA1-3), which are the vault particle associated vtRNAs, are more similar than vtRNA2-1 (Figure 3A and 3B).

Given that the chromatin accessibility of a DNA region is partially controlled by transcription and chromatin remodeler's factors (Corces *et al.*, 2018; Klemm *et al.*, 2019), we analyzed transcription factors (TFs) occupancy at a $\pm$ 3000 bp region centered at each vtRNA gene. Since there is no such study of the TCGA samples, we investigated the CHIP-seq data of the K562 chronic myelogenous leukemia cell line, which has the richest TF data at ENCODE dataset (Euskirchen *et al.*, 2007; Goldman *et al.*, 2013). We observed that the four vtRNAs share a common core of 23 TFs (26%). The three members of the vtRNA1 cluster share 16 additional TFs (18%), reaching

39 common TFs (44%), while only 2-4 TFs with vtRNA2-1 (Figure 3C and 3D and Extended Data: Table S6). We finally asked whether this common core of TFs is linked to a specific biological process. The STRING enrichment analysis of the core of 23 TFs common to all vtRNAs identified HTLV-I infection (ATF1, ATF3, E2F1, EP300, NFATC3, TBP), Hepatitis B (E2F1, EP300, NFATC3) and Transcriptional misregulation in cancer (AFF1, ATF1, MAX) as the top three enriched KEGG pathway terms (FDR < 0.05) (Figure 3D and Extended Data: Table S6).

Taken together these findings indicate that the transcriptional activity of the vtRNAs is gene specific, being vtRNA1-1 the most accessible and possibly the most expressed. In addition, the data suggest that the regulatory status of the vtRNA promoters could be coordinately controlled, and the three vtRNAs are more co-regulated among themselves. Yet, the core of TFs shared by the four vtRNAs is associated with viral infection and cancer related terms in the myeloid cell line K562.

The vtRNA promoters present gene and tissue specific patterns of chromatin accessibility in primary tumors

In order to investigate the expression of vtRNAs in different tissues we analyzed the Pan-Cancer TCGA ATAC-seq data discriminating the tissue of origin. As expected, vtRNA1-1 has the highest promoter chromatin accessibility among tissues and the smallest variation (Figure 4A, Extended Data: Tables S1-S5 and Figure S1). Remarkably, low-grade glioma tumor samples (LGG) show a global reduction in chromatin accessibility at the four vtRNA gene promoters (Figure 4A). An opposite pattern is seen in adrenocortical carcinoma tissue (ACC), where the vtRNAs have the highest concerted chromatin accessibility (Figure 4A). Individually, vtRNAs promoter accessibility is maximum and minimum in THCA (4.7) and LGG (3.3) for vtRNA1-1, ACC (4.2) and LGG (0.29) for vtRNA1-2, ACC (3.7) and LGG (0.24) for vtRNA1-3 and KIRC (4.2) and LGG (1.0) for vtRNA2-1 (Figure 4A). VtRNA3-1P reaches a maximum of promoter chromatin accessibility in Testicular Germ Cell Tumors (TGCT) (0.65) with an extremely low average ATAC-seq value and its minimum in PRAD (-1.5) (Extended Data: Figure S3C). VtRNA2-2P reaches a maximum and minimum promoter chromatin accessibility in Lung Adenocarcinoma (LUAD) (1.0) and ACC (-0.34) respectively (Extended Data: Figure S3C). Remarkably, although vtRNA1-1 and vtRNA1-3 have the highest and lowest promoter accessibility in the majority of tissues respectively, vtRNA1-2 and vtRNA2-1 are the most variable (vtRNA1-1 SD = 0.34, vtRNA1-2 SD = 1.4, vtRNA1-3 SD = 0.75 and vtRNA2-1 SD = 0.78) (Figure 4A). In addition, in 12 tissues the chromatin accessibility of vtRNA1-2 is higher than vtRNA2-1 (57%) whereas in nine tissues the chromatin accessibility of vtRNA1-2 is lower than vtRNA2-1 (43%) (Figure 4A).

Using the vtRNAs promoter DNA methylation data, we extended the analysis to 32 primary tumor tissues, incorporating nine more tissues than ATAC-seq samples (DLBC, KICH, OV, PAAD, READ, SARC, THYM, UCS, UVM) (Figure 4B, Extended Data: Figure S1) and 16 normal adjacent tissues (Figure 4C,

Figure 3. Comparative chromatin accessibility and transcription factor occupancy at the vtRNA loci. **A-B** Matrix of pairwise Spearman correlations (two-way hierarchical clustering distance measured by Euclidean and Ward clustering algorithms) of vtRNA 500 bp promoter region for ATAC-seq data (385 primary tumors samples across 23 tissues) (**A**) and DNA methylation average beta-values data (8403 primary tumors samples across 32 tissues) (**B**). **C**. Venn diagram of transcription factors identified as ChIP-seq Peaks by ENCODE 3 project in the cell line K562 (Venny 2.1; <https://bioinfogp.cnb.csic.es/tools/venny/index.html>). The region for TFs assignment was defined as ± 3000 bp from the vtRNA transcript sequence boundaries. **D**. Interaction cluster of the core 23 TFs common to all vtRNAs performed with STRING (Szklarczyk *et al.*, 2017). The colored labels of the top 3 enriched KEGG pathway terms (FDR < 0.05) are indicated.

Extended Data: Figure S1). Again, the DNA methylation profile mirrors the chromatin accessibility in the individual tissue types (*Figure 4*). Remarkably, the association between the average vtRNA promoter chromatin accessibility and DNA methylation of the vtRNA1 cluster in each tissue type is higher than that observed for the average of all tissues (vtRNA1-1 $r_s = -0.28$, vtRNA1-2 $r_s = -0.90$ vtRNA1-3 $r_s = -0.74$ and vtRNA2-1 $r_s = -0.58$) (*Table 1* and *Extended Data: Figure S4*). The opposite finding for vtRNA2-1 may be explained by the impact of the previously recognized epigenetic variation in the locus

(Silver *et al.*, 2015), which may prevail over the tissue specific variation for this gene.

In primary tumor tissues, promoter DNA methylation of vtRNA2-1 is higher than vtRNA1-2 in 17 tissues (53%) and the opposite is observed in the remaining 15 tissues (47%) (*Figure 4B*). Meanwhile in normal tissues, promoter DNA methylation of vtRNA2-1 is higher than vtRNA1-2 in 4 tissues (25%) and lower in 12 tissues (75%) (*Figure 4C*), indicating that although vtRNA1-2 is more methylated in

Figure 4. Chromatin accessibility and DNA methylation of the vtRNA promoters in tumor and normal tissues. A. ATAC-seq values for vtRNA promoters in 21 primary tumor tissues. **B.** Average beta-values of promoter DNA methylation for vtRNAs in 32 primary tumors tissues. **C.** Average beta-values of promoter DNA methylation for vtRNAs in 16 normal adjacent tissues. The charts show the average value and standard deviation of each vtRNAs for the tissues with at least five samples available at Pan-Cancer TCGA dataset.

normal samples, vtRNA2-1 gains methylation in neoplastic tissue, becoming more methylated than vtRNA1-2 (in 7 tissue types and vtRNA1-2 loose methylation: BLCA, BRCA, CHOL, HNSC, LUAD, LUSC, UCEC). Indeed, the Fisher's exact test identifies a mirrored profile of chromatin accessibility between normal (4 tissues with high vtRNA1-2 and 12 tissues with high vtRNA2-1 from 16 total tissues) and tumor tissues (17 tissues with high vtRNA1-2 and 15 tissues with high vtRNA2-1 from 32 total tissues) for vtRNA1-2 and vtRNA2-1 (p-value = 0.07). Despite this deregulation in transformed tissues, we wondered if the tissue specific differences in vtRNA promoters were explained by their pre-existing status in the normal tissues. The correlation among average vtRNA promoter DNA methylation in normal and tumor tissues suggest that the variation in chromatin accessibility among tissue types is already established in the normal tissue counterparts (vtRNA1-1 $r_s = 0.53$, vtRNA1-2 $r_s = 0.84$, vtRNA1-3 $r_s = 0.47$ and vtRNA2-1 $r_s = 0.75$, *Extended Data: Figure S5*).

VtRNA promoter's DNA methylation is associated with tumor stage and tissue of origin

The assessment of differential vtRNA expression from normal to tumor condition at the TCGA cohort can only be performed using DNA methylation dataset since no ATAC-seq analyses were performed in the normal tissues. The average beta-value of CpG sites in the 500bp vtRNAs promoter of normal and tumor tissue evidenced gene specific patterns of deregulation in neoplastic tissues (*Figure 5*). As depicted in *Figure 4*, vtRNA1-1 and vtRNA1-3 promoter DNA methylation and chromatin accessibility is low and high tissue-wide respectively, in both normal and tumor tissues (yet, there are highly methylated tumor outliers as LGG, that lack normal counterpart) (*Figure 4* and *Figures 5A and 5C*). Conversely, vtRNA1-2 and vtRNA2-1 revealed significant

differences in average promoter DNA methylation among the tissue categories (*Figures 5B and 5D*). In agreement with previous results, the methylation of vtRNA1-2 promoter decreases from normal to tumor samples, while vtRNA2-1's increases (*Figures 5B and 5D* and *Figure 4*).

We next asked whether the promoter status of the vtRNAs is deregulated during the normal and tumor transformation of different tissue types. This comparison was restricted to the 16 tissues with normal samples (with data of at least five samples, *Figure 6* and *Extended Data: Figure S1, Table S5*). The results revealed that vtRNA1-1 is globally unmethylated in normal and tumor from different tissue origins, but presents small but statistically significant changes in 4 tissue types (OG profile in BLCA, THCA UCEC and TSG in LUSC), whose biological impact remains to be determined (*Figure 6A*). Similarly, vtRNA1-3 is globally unmethylated, showing larger variability and significant upregulation in BRCA and PRAD, compatible with a TSG function (TSG trends for LIHC p-value = 0.08 and LUSC p-value = 0.09, *Figure 6C*). In agreement with the literature, vtRNA2-1 presents a TSG pattern of deregulation in PRAD, LUSC and BRCA (*Cao et al., 2013; Fort et al., 2018; Romanelli et al., 2014*), and an OG pattern in KIRP that has not been previously described (*Figure 6D*). Finally, a statistically significant dysregulation of vtRNA1-2 promoter DNA methylation across almost all cancer types (13 out of 16 tissues analyzed) poses it as a candidate OG in cancer (*Figure 6B*).

Overall, the data shows different DNA methylation changes in the vtRNAs promoters upon malignant transformation in several tissues. In addition, the deregulation of each vtRNA occurs mostly in a gene specific direction, where vtRNA1-2 has

Figure 5. Promoter DNA methylation of vtRNAs in Normal and Tumor tissues of Pan-Cancer TCGA dataset. Average beta-values of promoter DNA methylation of vtRNA1-1 (A), vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D), assessed in 746 normal and 8403 tumor tissues respectively. The box plots show the median line and lower and upper quartile and the whiskers the 2.5 and 97.5 percentile. Horizontal gray striped and red dotted lines denote unmethylated (average beta-value ≤ 0.2), 50% methylated (average beta-value = 0.5) and highly methylated (average beta-value ≥ 0.8) promoters. One-way ANOVA multiple test analysis with Sidak as posthoc test was performed. **** p-value < 0.0001.

Figure 6. VtRNA promoters DNA methylation in Normal vs. Tumor samples. Average beta-values of promoter DNA methylation for vtRNA1-1 (A) vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D). Acronyms indicate the tissue condition (normal (N) and tumor (T)). Blue and red colors indicate an increase or reduction in promoter methylation in tumor vs their normal tissues counterparts. The box plots show the median and the lower and upper quartile, and the whiskers the 2.5 and 97.5 percentile of the distribution. Horizontal, lines denote the methylation level of the promoters: grey striped bottom and top for unmethylated (average beta-value ≤ 0.2) or highly methylated (average beta-value ≥ 0.8) respectively, and red dotted for 50% methylated (average beta-value = 0.5) promoters. One-way ANOVA multiple test analysis with Sidak as posthoc test was performed. * p-value < 0.05 ; ** p-value < 0.01 ; *** p-value < 0.0001 .

an OG like epigenetic de-repression while vtRNA2-1 (and to a lesser extent vtRNA1-1, vtRNA1-3) has a TSG like repressive methylation in tumor tissues.

Low DNA methylation at the promoter of vtRNA1-1, vtRNA1-2 and vtRNA2-1 is associated with low patient overall survival

Seeking for a possible clinical significance of the chromatin accessibility change of the vtRNA promoters, we studied its association with the overall survival of patients (Figure 7 and

Extended Data: Tables S7 and S8). The analysis of patients stratified by vtRNA promoter accessibility quartiles did not show differences in overall survival (data available at Extended Data: Table S8). However, a significantly lower patient survival probability when patient tumors have lower promoter DNA methylation is observed for vtRNA1-1 (p-value = 0.003), vtRNA1-2 (p-value = 0.004) and vtRNA2-1 (p-value = 0.002) (patient stratified in quartiles of the promoter DNA methylation cohort values, Figure 7). Therefore, a lower promoter DNA methylation of vtRNA1-1, vtRNA1-2 and vtRNA2-1, a

Figure 7. Association between patient Overall survival and vtRNA promoters DNA methylation status in Pan-Cancer TCGA dataset. Kaplan-Meier curves of overall patient survival probability over the time (4000 days) discriminating the primary tumors into two cohorts with relatively higher (75th) and lower (25th) DNA methylation at each vtRNA promoter. Promoter DNA methylation data of the 8060 primary tumors was used to stratify the patients in two quartiles, based on the expression of vtRNA1-1 (A) vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D). Patient survival probability of the two groups was analyzed using Log-rank (Mantel-Cox) test. The dotted lines represent the 95% confidence interval for each curve.

surrogate of their high expression, might be associated with poor patient overall survival cancer-wide.

Genome-wide chromatin accessibility correlations with vtRNA promoters reveal their link to specific cancer related functions

Seeking to get insight into the function of the vtRNAs from their transcription regulatory marks, we searched for the genes co-regulated at chromatin level. We calculated the correlation of the ATAC-seq promoter values of each vtRNA and all the individual genes in the 385 primary tumor cohort and selected those showing a Spearman correlation $r_s \geq 0.4$. Using STRING software analysis, we then investigated their connection with Biological Process categories and KEGG pathways (Szklarczyk *et al.*, 2017). A few enriched terms for the genes correlated with vtRNA1-2, vtRNA1-3 and vtRNA2-1 is identified ($FDR < 1 \times 10^{-3}$) (Extended Data: Tables S9 and S10). The 196 protein coding genes correlated with vtRNA1-2 are enriched in the Biological Process “Epithelial cell differentiation” ($FDR < 1 \times 10^{-4}$) (Extended Data: Table S10). Meanwhile, the 358 protein coding genes correlated with vtRNA2-1 are enriched in the Biological Processes “Immune system process”, “Inflammatory response” and “Immune response” ($FDR < 1 \times 10^{-5}$) and the KEGG pathways term “Cytokine-cytokine receptor interaction” ($FDR < 1 \times 10^{-4}$) (Extended Data: Table S10). The only six protein coding genes correlated with vtRNA1-3 are enriched in the “Thyroid cancer” KEGG pathway term ($FDR < 1 \times 10^{-4}$). Remarkably, although STRING analysis of the vtRNA1-1 associated 37 protein coding genes did not find enriched process or pathways, 36 of the 37 correlated genes are located at chromosome 5q region. Using the Cluster Locator algorithm (Pazos Obregón *et al.*, 2018), we found a statistically significant non-random clustering behavior for the chromosomal location of the genes co-regulated with vtRNA1-1 ($p\text{-value} < 1 \times 10^{-10}$). Indeed, when the analysis was extended to 173 protein coding genes with Spearman correlation $r_s \geq 0.3$, 139 genes of 173 were situated at chromosome 5q region ($p\text{-value} < 1 \times 10^{-10}$). These genes are positioned in particular regions at chr5q31, chr5q35, chr5q23, chr5q14, chr5q32, chr5q34 ($FDR < 1 \times 10^{-4}$) (Kuleshov *et al.*, 2016) (Extended Data: Table S10). We did not find a similar phenomenon for the other three vtRNAs. In summary, vtRNA1-3, vtRNA1-2 and vtRNA2-1 transcription may be co-regulated with genes belonging to specific biological processes located elsewhere, while vtRNA1-1 transcription may be controlled by a locally specialized chromatin domain at chromosome 5q region lacking apparent functional relatedness.

Immune Subtypes associated with the vtRNAs profile

Taking into account that vtRNAs have been previously related to the immune response (Golec *et al.*, 2019; Li *et al.*, 2015; Nandy *et al.*, 2009), and that we found an association of vtRNA2-1 with immune related terms, we sought to investigate a putative relation between the vtRNAs expression and the six Immune Subtypes defined by Thorsson *et al.* (Thorsson *et al.*, 2018), compiled at Pan-Cancer TCGA (Extended Data: Tables S11-S14). These six Immune Subtypes are named because of the foremost immune characteristic, comprising wound

healing, IFN-g dominant, lymphocyte depleted, inflammatory, immunologically quiet and TGF-b dominant types (Thorsson *et al.*, 2018). As was expected, mirrored patterns were observed for promoter ATAC-seq and DNA methylation data of the six groups (Figure 8). The immunologically quiet subtype shows a concerted shutdown of all vtRNA promoters (Figure 8), which is largely composed by LGG samples showing low vtRNA ATAC-seq and high promoter methylation average values in LGG tumors (Figures 4A and 4B). The immunological quiet subtype exhibits the lowest leukocyte fraction (very low Th17 and low Th2) and the highest macrophage:lymphocyte ratio (high M2 macrophages) (Thorsson *et al.*, 2018). The remaining subtypes present a similar vtRNA profile except for the inversion of vtRNA1-2/vtRNA2-1 accessibility ratio, which is higher than 1 for the wound healing and IFN-g dominant subtypes while is lower than 1 for the lymphocyte depleted and inflammatory subtypes (Figure 8A). As expected, a mirrored pattern was observed for vtRNA1-2 and vtRNA2-1 at DNA methylation data (Figure 8B). Since the wound healing and IFN-g dominant subtype have high proliferation rate opposite to the lymphocyte depleted and inflammatory subtypes (Thorsson *et al.*, 2018), we evaluated the correlation between vtRNA1-2 and vtRNA2-1 chromatin status and proliferation rate or wound healing status defined by Thorsson *et al.* (Extended Data: Tables S12 and S14) (Thorsson *et al.*, 2018). A negative correlation between vtRNA1-2 promoter DNA methylation and the proliferation rate ($r_s = -0.44$) and wound-healing features of the tumors is revealed ($r_s = -0.48$) (Extended Data: Tables S12 and S14). These findings suggest that vtRNA transcriptional regulation may be associated with tumor immunity.

Discussion

Although important insight into the vtRNA function has been recently gained, the landscape of vtRNA expression in human tissues and cancer was unknown, perhaps for the absence of vtRNA data in transcriptomic studies. Here we used ATAC-seq and methylation arrays data from the Pan-Cancer TCGA consortium (Weinstein *et al.*, 2013), as surrogate variables to uncover the expression of vtRNAs across multiple cancer types. A limitation of our study is its blindness to post-transcriptional controls of vtRNA expression, such as RNA processing, transport, localization and/or stability. Indeed, there is evidence of vtRNA transcript regulation via RNA methylation by NSUN2, or via its association to other proteins like DUSP11, DIS3L2, SSB, SRSF2 and DICER (Burke *et al.*, 2016; Hussain *et al.*, 2013; Kickhoefer *et al.*, 2002; Labno *et al.*, 2016; Persson *et al.*, 2009; Sajini *et al.*, 2019). Nonetheless, the chromatin status of the vtRNA promoters has been correlated with the expression of their transcripts in several reports (Ahn *et al.*, 2018; Fort *et al.*, 2018; Helbo *et al.*, 2015; Helbo *et al.*, 2017; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Sallustio *et al.*, 2016; Treppendahl *et al.*, 2012). These evidences unequivocally stand for the use of chromatin structure as a suitable proxy for vtRNA expression.

An integrated genomic view of the regulatory features of the vtRNA genes compiled in UCSC genome browser indicates that the four vtRNAs have chromatin signs of active transcription in different cell lines and defines a proximal

Figure 8. Chromatin accessibility and DNA methylation of vtRNA promoters in the Six Immune Subtypes analyzed in the PANCAN TCGA dataset. **A.** ATAC-seq values expressed as log2 normalized values for vtRNA promoters grouped in six Immune Subtype: wound healing (105 samples), IFN- γ dominant (93 samples), inflammatory (110 samples), lymphocyte depleted (43 samples), immunologically quiet (9 samples), TGF-b dominant (4 samples). **B.** DNA methylation average beta-values for vtRNA promoters grouped in six Immune Subtype: wound healing (1963 samples), IFN- γ dominant (2137 samples), inflammatory (2126 samples), lymphocyte depleted (933 samples), immunologically quiet (383 samples), TGF-b dominant (151 samples) (Thorsson *et al.*, 2018). The charts show the average and standard deviation for each vtRNA promoter in each Immune Subtype group.

promoter regulatory region of approximately 500 bp (Nikitina *et al.*, 2011; Vilalta *et al.*, 1994). High resolution chromatin accessibility mapping studies by NOMe-Seq of the vtRNA promoters in cell lines validate this assumption (Helbo *et al.*, 2017). Particularly, vtRNA2-1 is singular for being immersed in a CpG island, for lacking the TATA box, DSE or PSE elements characteristic of the others vtRNA promoters, and for bearing a cAMP-response (CRE) element (Canella *et al.*, 2010; Helbo *et al.*, 2017; Nandy *et al.*, 2009). Since ATAC-seq approach is currently the best high throughput method to determine the

chromatin accessibility of a DNA region directly (Chen *et al.*, 2016; Sun *et al.*, 2019; Tsompana & Buck, 2014), we interrogated the ATAC-seq data of TCGA samples as a surrogate marker of vtRNA expression (Corces *et al.*, 2018). Although DNA methylation is a less direct and less accurate measurement of chromatin accessibility compared to ATAC-seq, 25 times more TCGA samples have been analyzed by DNA Methylation Arrays, including matched normal tissues samples, thus this dataset is of great worth for the current study. Our analyses revealed that vtRNA1-1 has accessible chromatin and small

variation in all the evaluated tissues, which is consistent with its identity as the ancestor vtRNA gene and its significance as the major RNA component of the ubiquitous vault particle (Stadler *et al.*, 2009; van Zon *et al.*, 2001). In addition, vtRNA1-2, vtRNA1-3 and vtRNA2-1 have lower chromatin accessibility at their promoters and a larger variation among the samples. As expected, the average DNA methylation assessed by microarrays is negatively correlated with the chromatin accessibility assessed by ATAC-seq. Therefore, the landscape of vtRNA chromatin status was corroborated by both approaches. Yet, the low chromatin accessibility of vtRNA1-3 promoter is not reflected by a concomitant high DNA methylation, suggesting that additional regulatory features of this gene, acting in trans or cis, play a major influence in its chromatin structure. As was reported by Helbo *et al.* using NOME-seq assay (Nucleosome Positioning), the chromatin at vtRNA1-3 may be poised for transcription, and still require TFs to achieve physiological levels of transcription (Helbo *et al.*, 2017).

The relative abundance of the four vtRNAs determined in the current study has been previously observed in various cell lines, including H157, H1299, CRL2741, WI-38 and NTERA-2 (Lee *et al.*, 2011), HSC CD34+, HL60 (Helbo *et al.*, 2015), T24, colorectal RKO, human diploid fibroblasts IMR90 (Helbo *et al.*, 2017) and MCF7 (Chen *et al.*, 2018). Yet, others found different comparative levels of the vtRNAs in MCF7, SW1573, GLC4, 8226, KB, SW620, MCF-7, HeLa, HEK-293 and L88/5 (van Zon *et al.*, 2001), CBL, BL2 and BL41 (Nandy *et al.*, 2009), EBV-negative Burkitt's lymphoma cell lines and A549 (Li *et al.*, 2015) and in LNCaP, PC3, DU145, RWPE-1, HEK, HeLa and MCF-7 (Stadler *et al.*, 2009) cell lines, suggesting that the expression of vtRNAs may differ in laboratory cell lines compared to tissues. Alternatively, the methods used to quantify the vtRNAs are not comparable among the studies.

Seeking to investigate regulatory connections between the vtRNAs as a valuable tool to study their function, we determine the pairwise correlations between the promoters of the vtRNAs. The positive correlations found are compatible with a coordinated transcriptional control of the vtRNAs, that is greater for the three members of the vtRNA1 cluster (specially vtRNA1-1 and vtRNA1-3). The exception is the pair vtRNA1-2 and vtRNA2-1, whose chromatin accessibility and promoter methylation are not linked in the cohort and can be indeed negatively associated.

As an alternative approach to investigate the vtRNA transcriptional co-regulation, we analyzed transcription and chromatin remodeling factors associated with their proximal promoter region. It is worth to mention that RNA polymerase II TFs clustered near TSS of RNA polymerase III transcripts could modulate occupancy and transcription rate of RNA polymerase III through RNA pol II activity (Campbell & White, 2014; Gerber *et al.*, 2020; Raha *et al.*, 2010; Zhang *et al.*, 2019). Specifically, RNA polymerase II could assist chromatin opening, thus allowing accessibility of RNA polymerase III or its associated factors in the human genome (Raha *et al.*, 2010). Interestingly, the 23 TFs common to all vtRNAs are related to viral infections and

cancer. In agreement with their higher chromatin correlation, the vtRNA1 clustered genes have a larger core of common TFs (39) in comparison with vtRNA2-1 (23 TFs). The divergence of the transcriptional control of vtRNA2-1 was previously recognized given its unique regulatory elements and the complex developmental methylation it undergoes (Canella *et al.*, 2010; Carpenter *et al.*, 2018; Helbo *et al.*, 2017; Nandy *et al.*, 2009). The upregulation of the four vtRNAs upon viral infection has been reported for several viruses, so it is possible that this core of TFs participate in the coordinated expression of vtRNA during viral infection and related responses (Amort *et al.*, 2015; Li *et al.*, 2015; Nandy *et al.*, 2009). These TFs regulate cell cycle and translational arrest as well as the inhibition of host innate immune response (Attar & Kurdastani, 2017; Ertesun *et al.*, 2016; Horsley & Pavlath, 2002; Jean *et al.*, 2000; Persengiev & Green, 2003; Rohini *et al.*, 2018). Since these processes are also hallmarks of cancer, the dysregulation of the normal function of vtRNAs in viral response may provide a selective advantage for cancer cell to overcome the many sources of stress faced during neoplastic evolution. If that holds true, the core vtRNA TFs may be co-opted in the malignant context to tune vtRNA expression favoring tumor progression. In agreement with that hypothesis, ATF1 and ATF3 were associated with CREB response and increased cell viability (Persengiev & Green, 2003). Likewise, Golec *et al.* showed that altered levels of vtRNA2-1 modulate PKR activation and consequently altered the levels of CREB phosphorylation during T cell activation, which is a prerequisite for IFN- γ activity (Golec *et al.*, 2019). Furthermore, E2F1, a cell cycle regulator, was described as a direct transcriptional regulator of vtRNA2-1 in cervical cancer cells (Li *et al.*, 2017). Moreover, it was shown that MYC binding to vtRNA2-1 promoter raises its expression and could be the explanation for the increased levels of vtRNA2-1 in some tumors (Park *et al.*, 2017a). Similarly, TGFB1 provokes the demethylation of vtRNA2-1 promoter and consequently increases its expression in ovarian cancer (Ahn *et al.*, 2018).

Alternatively, since RNA polymerase III transcripts are involved in virus replication and immune response (Graczyk *et al.*, 2015), it is possible that the TF bound to vtRNA vicinity are part of a global and not a vtRNA specific viral response.

Due to its importance for cancer biology, vtRNA2-1 transcriptional regulation was recently more investigated (Yeganeh & Hernandez, 2020). Various studies reported tissue specific roles of vtRNA2-1 in normal (Golec *et al.*, 2019; Lee *et al.*, 2019; Miñones-Moyano *et al.*, 2013; Sallustio *et al.*, 2016; Suojalehto *et al.*, 2014) and cancer tissues (Ahn *et al.*, 2018; Fort *et al.*, 2018; Hu *et al.*, 2017; Im *et al.*, 2020; Jeon *et al.*, 2012a; Kunkeaw *et al.*, 2012; Lee *et al.*, 2011; Lee, 2015; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Lee *et al.*, 2016; Lei *et al.*, 2017; Li *et al.*, 2017; Ma *et al.*, 2020; Treppendahl *et al.*, 2012) and epigenetic alterations of the locus have been described in different malignancies (Ahn *et al.*, 2018; Cao *et al.*, 2013; Fort *et al.*, 2018; Helbo *et al.*, 2015; Helbo *et al.*, 2017; Joo *et al.*, 2018; Lee *et al.*, 2014a; Lee *et al.*, 2014b; Park *et al.*, 2017a; Romanelli *et al.*, 2014; Treppendahl *et al.*, 2012). Meanwhile, except for Kirsten Grønbaek group's contributions on the chromatin

characterization of vtRNA1-3 and vtRNA1-2 promoters (Helbo *et al.*, 2015; Helbo *et al.*, 2017), little is known about the transcriptional regulation and expression of the other vtRNAs. Likewise, to our knowledge, the global landscape of vtRNA expression across different cancer types has not been investigated. Our study shows that, except for vtRNA1-1, which has a high promoter chromatin accessibility and low DNA methylation in all the tissues, the other vtRNA promoters exhibit different levels of chromatin accessibility and DNA methylation among the tissue types. The relative status of the chromatin in different tissues is accurately mirrored by both approaches in the tissues examined by ATAC-seq and Illumina DNA methylation arrays. Interestingly, the same analysis performed in the normal tissue samples reveals that the tissue specific variation in DNA methylation of vtRNA promoters in the neoplastic tissues is already established in their normal tissue counterparts, indicating that the regulation of vtRNA expression is tuned during normal development and cell differentiation.

While there is no apparent concerted modulation of the chromatin accessibility of the vtRNA promoters in most tissues, ACC and LGG tumors seem to be the exception. Concordantly, a global hypomethylation of malignant ACC tumors (Legendre *et al.*, 2016; Rechache *et al.*, 2012), and an aberrantly methylation processes associated to altered DNMTs activity in LGG tumors were reported (Nomura *et al.*, 2019). However, more research is necessary to understand the molecular basis of these observations.

The differential expression of the vtRNAs from normal to primary tumor tissues revealed different patterns depending on tissue origin. The average DNA methylation of the vtRNA1-1 and vtRNA1-3 promoters is not significantly deregulated, whereas vtRNA1-2 and vtRNA2-1 are decreased and increased respectively in tumor tissues. The latter finding favors a candidate OG and TSG function of vtRNA1-2 and vtRNA2-1 in cancer, respectively. Indeed, the epigenetic repression of vtRNA2-1 was previously reported by Romanelli *et al.* in BLCA, BRCA, COAD and LUSC analyzing the same data from TCGA (Romanelli *et al.*, 2014). Furthermore, functional studies in LUSC, PRAD, ESCA and AML provided experimental support to that hypothesis (Cao *et al.*, 2013; Fort *et al.*, 2018; Lee *et al.*, 2014a; Treppendahl *et al.*, 2012). Yet, vtRNA2-1 has been also proposed as an OG in ovarian, thyroid, endometrial, cervical and renal cancer (Ahn *et al.*, 2018; Hu *et al.*, 2017; Lee *et al.*, 2016; Lei *et al.*, 2017; Li *et al.*, 2017; Yeganeh & Hernandez, 2020). In agreement with the functional data found by Lei *et al.* (Lei *et al.*, 2017), we found an epigenetic de-repression of vtRNA2-1 promoter DNA methylation in renal carcinoma (KIRP). Likewise, a closer look at THCA samples, shows an average decrease in the DNA methylation of vtRNA2-1 promoter that supports the OG function described in this tissue (Lee *et al.*, 2016). Unfortunately, normal ovarian, cervical and endometrial tissues are not available at the TCGA. Finally, vtRNA1-2 has an oncogenic pattern of inferred expression between normal and tumor tissues for 13 of 16 tissues analyzed, raising a possible oncogenic function for this RNA.

Our study found also an association of DNA methylation at the promoter with a shorter overall survival for vtRNA1-1,

vtRNA1-2 and vtRNA2-1. This is surprising for vtRNA2-1, since it has both TSG and OG roles and TSG compatible expression profiles in three cancer types, inferred by DNA methylation of its promoter in tumor vs normal tissues and the average promoter DNA methylation is increased in tumor compared to normal tissue. A higher impact of vtRNA2-1 expression in patient survival in the oncogenic context may justify this discrepancy. On the contrary, the association between vtRNA1-2 low promoter DNA methylation and low patient survival is in agreement with its OG expression in cancer. The lack of survival association with the ATAC-seq values may be due to the small number of patients studied by this approach.

VtRNA expression in individual cancer types, inferred from the epigenetic status of their promoters, has been associated with patient survival in several studies of vtRNA2-1, one of vtRNA1-3 and none for vtRNA1-1 or vtRNA1-2. Low methylation or high expression of vtRNA2-1 promoter were associated with good prognosis or overall survival in lung (Cao *et al.*, 2013), esophageal (Lee *et al.*, 2014a), prostate (Fort *et al.*, 2018), AML (Treppendahl *et al.*, 2012), gastric (Lee *et al.*, 2014b) and liver (Yu *et al.*, 2020). Conversely, a worse prognosis or overall survival association of vtRNA2-1 was reported in thyroid (Lee *et al.*, 2016) and ovarian (Ahn *et al.*, 2018). From these reports, only prostate (Fort *et al.*, 2018) and gastric (Lee *et al.*, 2014b) studies used the TCGA data. Likewise, the methylation status of the vtRNA1-3 promoter associates with overall survival in the lower risk Myelodysplastic Syndrome patients (Helbo *et al.*, 2015). Additionally, vtRNA1-1 and vtRNA1-2 correlate to chemotherapeutic resistance by direct interaction with drugs (doxorubicin, etoposide and mitoxantrone) and the modulation of vtRNA1-1 confirmed this finding in osteosarcoma cell lines (Gopinath *et al.*, 2005; Gopinath *et al.*, 2010; Mashima *et al.*, 2008; van Zon *et al.*, 2001). Furthermore, increased levels of vtRNA1-1 were associated with increased proliferation and chemoresistance due to GAGE6 induction in MCF-7 cells (Chen *et al.*, 2018). Besides, Norbert Polacek group showed that vtRNA1-1 expression confers apoptosis resistance in several human cell lines (BL2, BL41, HS578T, HEK293, A549 and HeLa) and revealed its capacity to repressed intrinsic and extrinsic apoptosis pathway (Amort *et al.*, 2015; Bracher *et al.*, 2020). The later findings are in agreement with the lower patient survival associated with high levels of vtRNA1-1 expression cancer-wide observed in our analysis.

Seeking to get insights into the cancer related function of the vtRNAs we performed a genome wide search for genes co-regulated at the level of promoter chromatin accessibility. Remarkably, we identified immune and cytokine related terms for the genes co-regulated with vtRNA2-1. This association agrees with its proposed roles in innate immune modulation via PKR repression or OAS1 regulation and in cytokine production (Calderon & Conn, 2018; Golec *et al.*, 2019; Lee *et al.*, 2020; Li *et al.*, 2015). Furthermore, vtRNA2-1 has been associated with autoimmune disorders (Renauer *et al.*, 2015; Weeding & Sawalha, 2018) and tumor engraftment in prostate cancer (Ma *et al.*, 2020). This enrichment in viral infection pathways, together with the viral infection involvement of the core TFs common to all vtRNAs, reinforces the hypothesis of the re-utilization of a regulatory RNA induced upon viral response

in favor of cancer development at upstream and downstream regulatory steps. Additionally, we found a non-random clustering localization at chromosome arm 5q for genes co-regulated with vtRNA1-1. The chromosome arm 5q and regions chr5q31 and chr5q32-q33 were found frequently deleted in myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML) (Fuchs, 2012; Treppendahl *et al.*, 2012). Furthermore, the modulation of vtRNA2-1 and vtRNA1-3 was previously associated with human bone marrow CD34+ cells, AML and MDS (Helbo *et al.*, 2015; Treppendahl *et al.*, 2012). Yet, our analysis points to vtRNA2-1 and vtRNA1-2, and to a lesser extent to vtRNA1-1 and vtRNA1-3, as the most relevant drivers across cancer types.

Support for the involvement of the vtRNAs in the immune responses in the context of cancer came from the association of their chromatin status with the immune subtype categories defined by Thorsson *et al.* (Thorsson *et al.*, 2018). Our findings suggest a possible role of the vtRNAs in lymphocyte response in tumor samples, since the immunologically quiet subtype shows the lowest leukocyte fraction and vtRNAs promoter chromatin accessibility (Thorsson *et al.*, 2018). Meanwhile, the vtRNA2-1 and vtRNA1-2 expression distinguishes two groups among the remaining four subtypes. The participation of vtRNA2-1 in the induction of the IFN- γ and IL-2 expression in activated T cells through PKR modulation (Golec *et al.*, 2019) goes in agreement with this hypothesis. Although the same study demonstrated that vtRNA1-1 expression was unchanged during T cell activation, it is remarkable that the 5q31-q33 region, comprising the vtRNA1 locus and chromatin co-regulated genes, encodes several cytokines that regulate the differentiation of Th1 and Th2 lymphocytes and has been linked with susceptibility to infections (Jeronimo *et al.*, 2007; Lacy *et al.*, 2000; Naka *et al.*, 2009; Rodrigues *et al.*, 1999). Furthermore, vtRNA2-1 levels were recently associated with macrophages M1/M2 fates in prostate PC3 cell line mice xenografts via TGF- β (Ma *et al.*, 2020), two pathways that define the six immune subtypes (Thorsson *et al.*, 2018). Finally, we found that the proliferation rate and wound healing of the tumors are associated with the levels of vtRNA1-2 chromatin accessibility, which reinforces an OG role of vtRNA1-2 in cancer. Nonetheless, the participation of the vtRNAs in immune cells response inside the neoplastic niche needs to be further investigated.

Conclusion

Taken together our analysis reveals the pattern of chromatin accessibility and DNA methylation at the four vtRNA promoters, analyzed by tissue of origin in the TCGA cohort. The agreement between both dataset and previous related literature endorses the use of these variables as surrogates of the vtRNA transcripts expression. VtRNA genes are enriched in viral related TFs. The comparison of TFs and chromatin status of the vtRNAs suggest that vtRNA2-1 is likely the most independently regulated vtRNA locus. The pattern of chromatin accessibility and DNA promoter methylation in normal vs tumor tissue and the association with cancer cell pathways and patient survival suggest that vtRNA2-1 and vtRNA1-2 are possibly the

more relevant contributors to cancer. The results favor tissue specific TSG and OG roles for vtRNA2-1, a still not investigated oncogenic role of vtRNA1-2 and a limited cancer driver effect of vtRNA1-1/3 in specific tissue types. Lastly, we uncovered new evidence linking the vtRNAs with the immune response, cell proliferation and overall survival in cancer, which guarantees further investigation.

Data availability

Underlying data

All raw data used in this study can be downloaded from Xena Browser, a tool to explore functional genomic data sets (Goldman *et al.*, 2020) (<https://xenabrowser.net/>), and UCSC genome browser (Kent *et al.*, 2002) (<https://genome.ucsc.edu/>).

Extended data

Zenodo: Pan-Cancer chromatin analysis of the human vtRNA genes - Supplementary Figures, <http://doi.org/10.5281/zenodo.4784353> (Fort & Duhagon, 2021a)

This project contains the following extended data:

- Figure S1. Data type and tissue sample distribution.** Sample distribution per tissue type in the ATAC-seq, DNA methylation, ATAC-seq & DNA methylation and Normal adjacent tissue sample groups. **A.** Number of samples. **B.** Percentage of samples. The color reference, the number of samples of this tissue and the accounted percentage are presented to each tissue. Names references: ACC (Adrenocortical carcinoma), BLCA (Bladder Urothelial Carcinoma), BRCA (Breast invasive carcinoma), CESC (Cervical squamous cell carcinoma and endocervical adenocarcinoma), CHOL (Cholangiocarcinoma), COAD (Colon adenocarcinoma), DLBC (Lymphoid Neoplasm Diffuse Large B-cell Lymphoma), ESCA (Esophageal carcinoma), GBM (Glioblastoma multiforme), HNSC (Head and Neck squamous cell carcinoma), KICH (Kidney Chromophobe), KIRC (Kidney renal clear cell carcinoma), KIRP (Kidney renal papillary cell carcinoma), LGG (Brain Lower Grade Glioma), LIHC (Liver hepatocellular carcinoma), LUAD (Lung adenocarcinoma), LUSC (Lung squamous cell carcinoma), MESO (Mesothelioma), OV (Ovarian serous cystadenocarcinoma), PAAD (Pancreatic adenocarcinoma), PCPG (Pheochromocytoma and Paraganglioma), PRAD (Prostate adenocarcinoma), READ (Rectum adenocarcinoma), SARC (Sarcoma), SKCM (Skin Cutaneous Melanoma), STAD (Stomach adenocarcinoma), TGCT (Testicular Germ Cell Tumors), THCA (Thyroid carcinoma), THYM (Thymoma), UCEC (Uterine Corpus Endometrial Carcinoma), UCS (Uterine Carcinosarcoma) and UVM (Uveal Melanoma).
- Figure S2. Correlation between ATAC-seq values and promoter methylation values of vtRNAs in Pan-Cancer TCGA dataset.** A–D. The correlation between ATAC-seq and DNA methylation average normalized beta values was calculated for vtRNAs

(vtRNAs1-1 (A), vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D)), in 329 primary tumors samples. The box plots show the median line and lower and upper quartile and the whiskers the 2.5 and 97.5 percentile. Horizontal grey striped and red dotted lines are shown to denote unmethylated (average beta-value ≤ 0.2), 50% methylated (average beta-value = 0.5) and highly methylated (average beta-value ≥ 0.8) levels. Spearman correlation value and the best-fit line (red line) with 95% confidence bands (black dot lines) are shown.

- Figure S3. Genomic context and chromatin accessibility of human vtRNA3-1P, vtRNA2-2P and RNYs (RNY1, RNY3, RNY4 and RNY5).** A. Genomic view of the 1.5 kb region of human vtRNA3-1P, vtRNA2-2P and RNYs (RNY1, RNY3, RNY4 and RNY5) genes in UCSC Genome browser (GRCh37/hg19) centered at the 500bp of the ATAC-seq. Highlighted in yellow is the 500bp ATAC-seq region used in the posterior analyses. The following Gene annotation and ENCODE Project tracks for seven cell lines (GM12878, H1-hESC, HSMM, HUVEC, K562, NHEK, NHL) are displayed: DNA accessibility (DNase-seq hypersensitivity clusters (color intensity is proportional to the maximum signal strength)), DNA methylation (CpG islands length greater than 200 bp), histone modification (H3K27Ac, H3K4me1, H3K4me3 marks), conservation of the region in 100 Vertebrates (log-odds PhyloP scores). The vertical viewing range of the tracks displays the default settings of the browser for each variable. B. The distribution of ATAC-seq values for of human vtRNA3-1P, vtRNA2-2P and RNYs (RNY1, RNY3, RNY4 and RNY5) in Pan-Cancer TCGA dataset (385 tumors samples across 23 cancer types) expressed as log2 normalized values. The box plots show the median and the lower and upper quartile, and the whiskers the 2.5 and 97.5 percentile of the distribution. C. ATAC-seq values for of human vtRNA3-1P, vtRNA2-2P and RNYs (RNY1, RNY3, RNY4 and RNY5) in 21 different tumors with at least 5 tumor samples available in Pan-Cancer TCGA dataset (385 tumors samples) expressed as log2 normalized values. The chart shows the average and standard error for each vtRNAs in each tumor type.
- Figure S4. VtRNA promoter's chromatin accessibility (ATAC-seq and DNA methylation) average values for tumor tissues from Pan-Cancer TCGA dataset.** The average vtRNA promoters ATAC-seq and DNA methylation values for each primary tumor tissue type (21 tissues). VtRNA1-1 (A), vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D). The Spearman correlation between averages ATAC-seq and DNA methylation values was calculated for each vtRNA. The chart shows the average and standard deviation for each tissue with at least five samples available at Pan-Cancer TCGA.

- Figure S5. VtRNA promoter's DNA methylation for normal and tumor tissues from Pan-Cancer TCGA dataset.** The average promoter DNA methylation beta-values of normal adjacent (Normal) and primary tumor (Tumor) tissues (16 tissue types) for vtRNA1-1 (A), vtRNA1-2 (B), vtRNA1-3 (C) and vtRNA2-1 (D) is shown. The Spearman (r) correlation between normal and tumor values was calculated. The chart shows the average and standard deviation for each tissue with at least five samples available at Pan-Cancer TCGA.

Zenodo: Pan-Cancer chromatin analysis of the human vtRNA genes - Supplementary Tables, <http://doi.org/10.5281/zenodo.4784473> (Fort & Duhagon, 2021b)

This project contains the following extended data:

- Tables S1–S5. ATAC-seq and DNA methylation data for vtRNA promoters in primary tumors and normal adjacent tissues.** CSV spreadsheets: Table_S1_ATAC-seq_data_500bp: All ATAC-seq data of vtRNAs promoter (500bp) data for primary tumor samples; Table_S2_DNA_methylation_500bp: All DNA methylation data and ATAC-seq data of vtRNAs promoter (500bp) data for total primary tumor and normal adjacent samples; Table_S3_DNA_methylation_NORMAL: All DNA methylation data of vtRNAs promoter (500bp) data for normal adjacent samples; Table_S4_DNA_methylation_TUMOR: All DNA methylation data of vtRNAs promoter (500bp) data for primary tumor samples; Table_S5_Normal_&_Tumor_matched: All DNA methylation data of vtRNAs promoter (500bp) data for primary tumor and normal adjacent samples.
- Table S6. VtRNAs Transcription Factor Binding and KEGG enriched terms.** CSV spreadsheets: Table_S6_Binding_Factors: Transcription factors identified in the cell line K562 as ChIP-seq Peaks by ENCODE 3 project and KEGG_terms: enriched KEGG pathway terms (FDR < 0.05).
- Tables S7–S8. DNA methylation, ATAC-seq data and associated survival data for primary tumors.** CSV spreadsheets: Table_S7_DNA-methylation_Survival_data: All DNA methylation data of vtRNAs promoter (500bp) and survival data for primary tumor samples; Table_S8_ATAC-seq_Survival_data: ATAC-seq data of vtRNAs promoter (500bp) and survival data for primary tumor samples.
- Tables S9–S10. Correlation of ATAC-seq values between vtRNA and all genome promoters in primary tumor samples.** CSV spreadsheets: Table_S9_ATAC-seq_gene_promoter_spearman_correlation: Spearman correlation values of all promoter genes and vtRNAs in primary tumors samples; Table_S10_vtRNAs_pathway_enrichment_and_cluster_chromosome_localization_analysis: vtRNAs vtRNA1-1, vtRNA1-2,

vtRNA1-3 and vtRNA2-1 pathway enrichment and cluster chromosome localization data.

- **Tables S11–S14. ATAC-seq and DNA methylation data for vtRNA promoters in primary tumors and the associated Immune Subtypes data.** *CSV spreadsheets:* Table_S11_Immune_Subtypes_DNA_methylation_data: All DNA methylation data of vtRNAs promoter (500bp) and Immune Subtypes data for primary tumor samples; Table_S12_Spearman_corr_vtRNAs_Immune_Subtypes_DNA_methylation_data: Spearman correlation values of DNA methylation data of vtRNAs promoter (500bp) and Immune Subtypes data for primary tumor samples; Table_S13_Immune_Subtypes_ATAC-seq_data: All ATAC-seq data of vtRNAs promoter (500bp)

and Immune Subtypes data for primary tumor samples; Table_S14_Spearman_corr_vtRNAs_Immune_Subtypes_ATAC-seq_data: Spearman correlation values of ATAC-seq data of vtRNAs promoter (500bp) and Immune Subtypes data for primary tumor samples.

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgments

A previous version of this article is available on BioRxiv: <https://doi.org/10.1101/2020.10.05.324723>

References

- Aakula A, Kohonen P, Leivonen SK, *et al.*: **Systematic Identification of MicroRNAs That Impact on Proliferation of Prostate Cancer Cells and Display Changed Expression in Tumor Tissue.** *Eur Urol.* 2016; **69**(6): 1120–1128.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ahn JH, Lee HS, Lee JS, *et al.*: **Nc886 is induced by TGF- β and suppresses the microRNA pathway in ovarian cancer.** *Nat Commun.* 2018; **9**(1): 1166.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Amort M, Nachbauer B, Tuzlak S, *et al.*: **Expression of the vault RNA protects cells from undergoing apoptosis.** *Nat Commun.* 2015; **6**: 7030.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Attar N, Kurdistani SK: **Exploitation of EP300 and CREBBP lysine acetyltransferases by cancer.** *Cold Spring Harb Perspect Med.* 2017; **7**(3): a026534.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bi N, Cao J, Song Y, *et al.*: **A MicroRNA signature predicts survival in early stage small-cell lung cancer treated with surgery and adjuvant chemotherapy.** *PLoS One.* 2014; **9**(3): e91388.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bland JM, Altman DG: **Survival probabilities (the Kaplan-Meier method).** *BMJ.* 1998; **317**(7172): 1572.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Boivin V, Deschamps-Francoeur G, Couture S, *et al.*: **Simultaneous sequencing of coding and noncoding RNA reveals a human transcriptome dominated by a small number of highly expressed noncoding genes.** *RNA.* 2018; **24**(7): 950–965.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Boivin V, Faucher-Giguère L, Scott M, *et al.*: **The cellular landscape of mid-size noncoding RNA.** *Wiley Interdiscip Rev RNA.* 2019; **10**(4): e1530.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bracher L, Ferro I, Pulido-Quetglas C, *et al.*: **Human vtRNA1-1 levels modulate signaling pathways and regulate apoptosis in human cancer cells.** *Biomolecules.* 2020; **10**(4): 614.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buenrostro JD, Giresi PG, Zaba LC, *et al.*: **Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position.** *Nat Methods.* 2013; **10**(12): 1213–1218.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buenrostro JD, Wu B, Chang HY, *et al.*: **ATAC-seq: A method for assaying chromatin accessibility genome-wide.** *Curr Protoc Mol Biol.* 2015; **109**: 21.29.1–21.29.9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Burke JM, Kincaid RP, Nottingham RM, *et al.*: **DUSP11 activity on triphosphorylated transcripts promotes Argonaute association with noncanonical viral microRNAs and regulates steady-state levels of cellular noncoding RNAs.** *Genes Dev.* 2016; **30**(18): 2076–2092.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Büscher M, Horos R, Hentze MW: **‘High vault-age’: non-coding RNA control of autophagy.** *Open Biol.* 2020; **10**(2): 190307.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Calderon BM, Conn GL: **A human cellular noncoding RNA activates the antiviral protein 2’-5’-oligoadenylate synthetase 1.** *J Biol Chem.* 2018; **293**(41): 16115–16124.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Campbell KJ, White RJ: **MYC regulation of cell growth through control of transcription by RNA polymerases I and III.** *Cold Spring Harb Perspect Med.* 2014; **4**(5): a018408.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Canella D, Praz V, Reina JH, *et al.*: **Defining the RNA polymerase III transcriptome: Genome-wide localization of the RNA polymerase III transcription machinery in human cells.** *Genome Res.* 2010; **20**(6): 710–721.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cao J, Song Y, Bi N, *et al.*: **DNA methylation-mediated repression of miR-886-3p predicts poor outcome of human small cell lung cancer.** *Cancer Res.* 2013; **73**(11): 3326–3335.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Carpenter BL, Zhou W, Madaj Z, *et al.*: **Mother-child transmission of epigenetic information by tunable polymorphic imprinting.** *Proc Natl Acad Sci U S A.* 2018; **115**(51): E11970–E11977.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chen J, OuYang H, An X, *et al.*: **Vault RNAs partially induces drug resistance of human tumor cells MCF-7 by binding to the RNA/DNA-binding protein PSF and inducing oncogene GAGE6.** *PLoS One.* 2018; **13**(1): e0191325.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chen X, Shen Y, Draper W, *et al.*: **ATAC-seq reveals the accessible genome by transposase-mediated imaging and sequencing.** *Nat Methods.* 2016; **13**(12): 1013–1020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Corces MR, Granja JM, Shams S, *et al.*: **The chromatin accessibility landscape of primary human cancers.** *Science.* 2018; **362**(6413): eaav1898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dettmer MS, Perren A, Moch H, *et al.*: **MicroRNA profile of poorly differentiated thyroid carcinomas: New diagnostic and prognostic insights.** *J Mol Endocrinol.* 2014; **52**(2): 181–189.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Duren Z, Chen X, Jiang R, *et al.*: **Modeling gene regulation from paired expression and chromatin accessibility data.** *Proc Natl Acad Sci U S A.* 2017; **114**(25): E4914–E4923.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ertosun MG, Hapil FZ, Nidai OO: **E2F1 transcription factor and its impact on growth factor and cytokine signaling.** *Cytokine Growth Factor Rev.* 2016; **31**: 17–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Euskirchen GM, Rozowsky JS, Wei CL, *et al.*: **Mapping of transcription factor binding regions in mammalian cells by ChIP: Comparison of array- and sequencing-based technologies.** *Genome Res.* 2007; **17**(6): 898–909.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fendler A, Jung M, Stephan C, *et al.*: **miRNAs can predict prostate cancer biochemical relapse and are involved in tumor progression.** *Int J Oncol.* 2011; **39**(5): 1183–1192.
[PubMed Abstract](#) | [Publisher Full Text](#)

- Fort RS, Duhagon MA: **Pan-Cancer chromatin analysis of the human vtRNA genes - Supplementary Figures.** 2021a.
- Fort RS, Duhagon MA: **Pan-Cancer chromatin analysis of the human vtRNA genes - Supplementary Tables.** 2021b.
- Fort RS, Mathó C, Geraldo MV, *et al.*: **Nc886 is epigenetically repressed in prostate cancer and acts as a tumor suppressor through the inhibition of cell growth.** *BMC Cancer.* 2018; **18**(1): 127.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fort RS, Garat B, Sotelo-Silveira JR, *et al.*: **vtRNA2-1/nc886 Produces a Small RNA That Contributes to Its Tumor Suppression Action through the microRNA Pathway in Prostate Cancer.** *Noncoding RNA.* 2020; **6**(1): 7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fuchs O: **Important genes in the pathogenesis of 5q- syndrome and their connection with ribosomal stress and the innate immune system pathway.** *Leuk Res Treatment.* 2012; **2012**: 179402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gao W, Shen H, Liu L, *et al.*: **Mir-21 overexpression in human primary squamous cell lung carcinoma is associated with poor patient prognosis.** *J Cancer Res Clin Oncol.* 2011; **137**(4): 557–566.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gerber A, Ito K, Chu CS, *et al.*: **Gene-Specific Control of tRNA Expression by RNA Polymerase II.** *Mol Cell.* 2020; **78**(4): 765–778.e7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Goldman MJ, Craft B, Hastie M, *et al.*: **Visualizing and interpreting cancer genomics data via the Xena platform.** *Nat Biotechnol.* 2020; **38**(6): 675–678.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Goldman M, Craft B, Swatloski T, *et al.*: **The UCSC Cancer Genomics Browser: update 2013.** *Nucleic Acids Res.* 2013; **41**(Database issue): D949–D954.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Golec E, Lind L, Qayyum M, *et al.*: **The Noncoding RNA nc886 Regulates PKR Signaling and Cytokine Production in Human Cells.** *J Immunol.* 2019; **202**(1): 131–141.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gopinath SCB, Matsugami A, Katahira M, *et al.*: **Human vault-associated non-coding RNAs bind to mitoxantrone, a chemotherapeutic compound.** *Nucleic Acids Res.* 2005; **33**(15): 4874–4881.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gopinath SCB, Wadhwa R, Kumar PKR: **Expression of noncoding vault RNA in human malignant cells and its importance in mitoxantrone resistance.** *Mol Cancer Res.* 2010; **8**(11): 1536–1546.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Graczyk D, White RJ, Ryan KM: **Involvement of RNA Polymerase III in Immune Responses.** *Mol Cell Biol.* 2015; **35**(10): 1848–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Helbo AS, Lay FD, Jones PA, *et al.*: **Nucleosome Positioning and NDR Structure at RNA Polymerase III Promoters.** *Sci Rep.* 2017; **7**: 41947.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Helbo AS, Treppendahl M, Aslan D, *et al.*: **Hypermethylation of the VTRNA1-3 promoter is associated with poor outcome in lower risk myelodysplastic syndrome patients.** *Genes (Basel).* 2015; **6**(4): 977–990.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Horos R, Büscher M, Kleinendorst R, *et al.*: **The Small Non-coding Vault RNA1-1 Acts as a Riboregulator of Autophagy.** *Cell.* 2019; **176**(5): 1054–1067.e12.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Horsley V, Pavlath GK: **NFAT: Ubiquitous regulator of cell differentiation and adaptation.** *J Cell Biol.* 2002; **156**(5): 771–774.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hu Z, Zhang H, Tang L, *et al.*: **Silencing nc886, a Non-Coding RNA, Induces Apoptosis of Human Endometrial Cancer Cells-1A In Vitro.** *Med Sci Monit.* 2017; **23**: 1317–1324.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hussain S, Sajini AA, Blanco S, *et al.*: **NSun2-mediated cytosine-5 methylation of vault noncoding RNA determines its processing into regulatory small RNAs.** *Cell Rep.* 2013; **4**(2): 255–261.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Im WR, Lee HS, Lee YS, *et al.*: **A Regulatory Noncoding RNA, nc886, Suppresses Esophageal Cancer by Inhibiting the AKT Pathway and Cell Cycle Progression.** *Cells.* 2020; **9**(4): 801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jean D, Tellez C, Huang S, *et al.*: **Inhibition of tumor growth and metastasis of human melanoma by intracellular anti-ATF-1 single chain Fv fragment.** *Oncogene.* 2000; **19**(22): 2721–2730.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeon SH, Johnson BH, Lee YS: **A tumor surveillance model: A non-coding RNA senses neoplastic cells and its protein partner signals cell death.** *Int J Mol Sci.* 2012a; **13**(10): 13134–13139.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jeon SH, Lee K, Lee KS, *et al.*: **Characterization of the direct physical interaction of nc886, a cellular non-coding RNA, and PKR.** *FEBS Lett.* 2012b; **586**(19): 3477–3484.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeronimo SMB, Holst AKB, Jamieson SE, *et al.*: **Genes at human chromosome 5q31.1 regulate delayed-type hypersensitivity responses associated with *Leishmania chagasi* infection.** *Genes Immun.* 2007; **8**(7): 539–551.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Joo JE, Dowty JG, Milne RL, *et al.*: **Heritable DNA methylation marks associated with susceptibility to breast cancer.** *Nat Commun.* 2018; **9**(1): 867.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kedersha NL, Rome LH: **Isolation and characterization of a novel ribonucleoprotein particle: Large structures contain a single species of small RNA.** *J Cell Biol.* 1986; **103**(3): 699–709.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kedersha NL, Heuser JE, Chugani DC, *et al.*: **Vaults. III. Vault ribonucleoprotein particles open into flower-like structures with octagonal symmetry.** *J Cell Biol.* 1991; **112**(2): 225–235.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kent WJ, Sugnet CW, Furey TS, *et al.*: **The Human Genome Browser at UCSC.** *Genome Res.* 2002; **12**(6): 996–1006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kickhoefer VA, Rajavel KS, Scheffer GL, *et al.*: **Vaults are up-regulated in multidrug-resistant cancer cell lines.** *J Biol Chem.* 1998; **273**(15): 8971–8974.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kickhoefer VA, Poderycki MJ, Chan EKL, *et al.*: **The La RNA-binding protein interacts with the vault RNA and is a vault-associated protein.** *J Biol Chem.* 2002; **277**(43): 41282–41286.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Klemm SL, Shipony Z, Greenleaf WJ: **Chromatin accessibility and the regulatory epigenome.** *Nat Rev Genet.* 2019; **20**(4): 207–220.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kuleshov MV, Jones MR, Rouillard AD, *et al.*: **Enrichr: a comprehensive gene set enrichment analysis web server 2016 update.** *Nucleic Acids Res.* 2016; **44**(W1): W90–W97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kunkeaw N, Jeon SH, Lee K, *et al.*: **Cell death/proliferation roles for nc886, a non-coding RNA, in the protein kinase R pathway in cholangiocarcinoma.** *Oncogene.* 2012; **32**(32): 3722–3731.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kunkeaw N, Lee YS, Im WR, *et al.*: **Mechanism mediated by a noncoding RNA, nc886, in the cytotoxicity of a DNA-reactive compound.** *Proc Natl Acad Sci U S A.* 2019; **116**(17): 8289–8294.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Łabno A, Warkocki Z, Kulinski T, *et al.*: **Perlman syndrome nuclease DIS3L2 controls cytoplasmic non-coding RNAs and provides surveillance pathway for maturing snRNAs.** *Nucleic Acids Res.* 2016; **44**(21): 10437–10453.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lacy DA, Wang ZE, Symula DJ, *et al.*: **Faithful expression of the human 5q31 cytokine cluster in transgenic mice.** *J Immunol.* 2000; **164**(9): 4569–4574.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Landgraf P, Rusu M, Sheridan R, *et al.*: **A mammalian microRNA expression atlas based on small RNA library sequencing.** *Cell.* 2007; **129**(7): 1401–1414.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee YS: **A Novel Type of Non-coding RNA, nc886, Implicated in Tumor Sensing and Suppression.** *Genomics Inform.* 2015; **13**(2): 26–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee EK, Hong SH, Shin S, *et al.*: **Nc886, a non-coding RNA and suppressor of PKR, exerts an oncogenic function in thyroid cancer.** *Oncotarget.* 2016; **7**(46): 75000–75012.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee HS, Lee K, Jang HJ, *et al.*: **Epigenetic silencing of the non-coding RNA nc886 provokes oncogenes during human esophageal tumorigenesis.** *Oncotarget.* 2014a; **5**(11): 3472–3481.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee KS, Park JL, Lee K, *et al.*: **nc886, a non-coding RNA of anti-proliferative role, is suppressed by CpG DNA methylation in human gastric cancer.** *Oncotarget.* 2014b; **5**(11): 3944–3955.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee K, Kunkeaw N, Jeon SH, *et al.*: **Precursor miR-886, a novel noncoding RNA repressed in cancer, associates with PKR and modulates its activity.** *RNA.* 2011; **17**(6): 1076–1089.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee KS, Shin S, Cho E, *et al.*: **nc886, a non-coding RNA, inhibits UVB-induced MMP-9 and COX-2 expression via the PKR pathway in human keratinocytes.** *Biochem Biophys Res Commun.* 2019; **512**(4): 647–652.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee YS, Kunkeaw N, Lee YS: **Protein kinase R and its cellular regulators in cancer: An active player or a surveillant?** *Wiley Interdiscip Rev RNA.* 2020; **11**(2): e1558.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Legendre CR, Demeure MJ, Whitsett TG, *et al.*: **Pathway Implications of Aberrant Global Methylation in Adrenocortical Cancer.** *PLoS One.* 2016; **11**(3): e0150629.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lei J, Xiao JH, Zhang SH, *et al.*: **Non-coding RNA 886 promotes renal cell carcinoma growth and metastasis through the Janus kinase 2/signal**

- transducer and activator of transcription 3 signaling pathway.** *Mol Med Rep.* 2017; **16**(4): 4273–4278.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Li B, Carey M, Workman JL: **The Role of Chromatin during Transcription.** *Cell.* 2007; **128**(4): 707–719.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Li F, Chen Y, Zhang Z, *et al.*: **Robust expression of vault RNAs induced by influenza A virus plays a critical role in suppression of PKR-mediated innate immunity.** *Nucleic Acids Res.* 2015; **43**(21): 10321–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li JH, Wang M, Zhang R, *et al.*: **E2F1-directed activation of nc886 mediates drug resistance in cervical cancer cells via regulation of major vault protein.** *Int J Clin Exp Pathol.* 2017; **10**(9): 9233–9242.
[PubMed Abstract](#) | [Free Full Text](#)
- Liu G, Wang W, Hu S, *et al.*: **Inherited DNA methylation primes the establishment of accessible chromatin during genome activation.** *Genome Res.* 2018; **28**(7): 998–1007.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ma H, Wang M, Zhou Y, *et al.*: **Noncoding RNA 886 alleviates tumor cellular immunological rejection in host C57BL/C mice.** *Cancer Med.* 2020; **9**(14): 5258–5271.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mashima T, Kudo M, Takada Y, *et al.*: **Interactions between antitumor drugs and vault RNA.** *Nucleic Acids Symp Ser (Oxf).* 2008; (52): 217–218.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Meng C, Wei Z, Zhang y, *et al.*: **Regulation of cytochrome P450 3A4 by small vault RNA derived from the non-coding vault RNA1 of multidrug resistance-linked vault particle.** (D.A. Spandidos). *Mol Med Rep.* 2016; **14**(1): 387–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Miñones-Moyano E, Friedländer MR, Pallares J, *et al.*: **Upregulation of a small vault RNA (svtRNA2-1a) is an early event in Parkinson disease and induces neuronal dysfunction.** *RNA Biol.* 2013; **10**(7): 1093–1106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moran S, Arribas C, Esteller M: **Validation of a DNA methylation microarray for 850,000 CpG sites of the human genome enriched in enhancer sequences.** *Epigenomics.* 2016; **8**(3): 389–399.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Naka I, Nishida N, Patarapotikul J, *et al.*: **Identification of a haplotype block in the 5q31 cytokine gene cluster associated with the susceptibility to severe malaria.** *Malar J.* 2009; **8**: 232.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nandy C, Mrázek J, Stoiber H, *et al.*: **Epstein-Barr Virus-Induced Expression of a Novel Human Vault RNA.** *J Mol Biol.* 2009; **388**(4): 776–784.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nikitina TV, Tischenko LI, Schulz WA: **Recent insights into regulation of transcription by RNA polymerase III and the cellular functions of its transcripts.** *Biol Chem.* 2011; **392**(5): 395–404.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nomura M, Saito K, Aihara K, *et al.*: **DNA demethylation is associated with malignant progression of lower-grade gliomas.** *Sci Rep.* 2019; **9**(1): 1903.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nordentoft I, Birkenkamp-Demtroder K, Agerbæk M, *et al.*: **miRNAs associated with chemo-sensitivity in cell lines and in advanced bladder cancer.** *BMC Med Genomics.* 2012; **5**: 40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Okumura T, Kojima H, Miwa T, *et al.*: **The expression of microRNA 574-3p as a predictor of postoperative outcome in patients with esophageal squamous cell carcinoma.** *World J Surg Oncol.* 2016; **14**(1): 228.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Paliwal A, Temkin AM, Kerkel K, *et al.*: **Comparative Anatomy of Chromosomal Domains with Imprinted and Non-Imprinted Allele-Specific DNA Methylation.** *PLoS Genet.* 2013; **9**(8): e1003622.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Park JL, Lee YS, Kunkeaw N, *et al.*: **Epigenetic regulation of noncoding RNA transcription by mammalian RNA polymerase III.** *Epigenomics.* 2017a; **9**(2): 171–187.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Park JL, Lee YS, Song MJ, *et al.*: **Epigenetic regulation of RNA polymerase III transcription in early breast tumorigenesis.** *Oncogene.* 2017b; **36**(49): 6793–6804.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pazos Obregón F, Soto P, Lavín JL, *et al.*: **Cluster Locator, online analysis and visualization of gene clustering.** *Bioinformatics.* 2018; **34**(19): 3377–3379.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Persengiev SP, Green MR: **The role of ATF/CREB family members in cell growth, survival and apoptosis.** *Apoptosis.* 2003; **8**(3): 225–228.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Persson H, Kvist A, Vallon-Christersson J, *et al.*: **The non-coding RNA of the multidrug resistance-linked vault particle encodes multiple regulatory small RNAs.** *Nat Cell Biol.* 2009; **11**(10): 1268–1271.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Raha D, Wang Z, Moqtaderi Z, *et al.*: **Close association of RNA polymerase II and many transcription factors with Pol III genes.** *Proc Natl Acad Sci U S A.* 2010; **107**(8): 3639–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rechache NS, Wang Y, Stevenson HS, *et al.*: **DNA methylation profiling identifies global methylation differences and markers of adrenocortical tumors.** *J Clin Endocrinol Metab.* 2012; **97**(6): E1004–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Renauer P, Coit P, Jeffries MA, *et al.*: **DNA methylation patterns in naïve CD₄⁺ T cells identify epigenetic susceptibility loci for malar rash and discoid rash in systemic lupus erythematosus.** *Lupus Sci Med.* 2015; **2**(1): e000101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Richmond RC, Sharp GC, Herbert G, *et al.*: **The long-term impact of folic acid in pregnancy on offspring DNA methylation: follow-up of the Aberdeen Folic Acid Supplementation Trial (AFAST).** *Int J Epidemiol.* 2018; **47**(3): 928–937.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rodrigues V Jr, Piper K, Couissinier-Paris P, *et al.*: **Genetic control of schistosome infections by the SM1 locus of the 5q31-q33 region is linked to differentiation of type 2 helper T lymphocytes.** *Infect Immun.* 1999; **67**(9): 4689–4692.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rohini M, Haritha Menon A, Selvamurugan N: **Role of activating transcription factor 3 and its interacting proteins under physiological and pathological conditions.** *Int J Biol Macromol.* 2018; **120**(Pt A): 310–317.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Romanelli V, Nakabayashi K, Vizoso M, *et al.*: **Variable maternal methylation overlapping the nc886/vtRNA2-1 locus is locked between hypermethylated repeats and is frequently altered in cancer.** *Epigenetics.* 2014; **9**(5): 783–790.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sajini AA, Choudhury NR, Wagner RE, *et al.*: **Loss of 5-methylcytosine alters the biogenesis of vault-derived small RNAs to coordinate epidermal differentiation.** *Nat Commun.* 2019; **10**(1): 2550.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sallustio F, Serino G, Cox SN, *et al.*: **Aberrantly methylated DNA regions lead to low activation of CD₄⁺ T-cells in IgA nephropathy.** *Clin Sci (Lond).* 2016; **130**(9): 733–746.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schou JV, Rossi S, Jensen BV, *et al.*: **miR-345 in Metastatic Colorectal Cancer: A Non-Invasive Biomarker for Clinical Outcome in Non-KRAS Mutant Patients Treated with 3rd Line Cetuximab and Irinotecan.** *PLoS One.* 2014; **9**(6): e99886.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sender E, Johnson GD, Krawetz SA: **Local and global factors affecting RNA sequencing analysis.** *Anal Biochem.* 2011; **419**(2): 317–322.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shen J, Zhou W, Bi N, *et al.*: **MicroRNA-886-3P functions as a tumor suppressor in small cell lung cancer.** *Cancer Biol Ther.* 2018; **19**(12): 1185–1192.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Silver MJ, Kessler NJ, Hennig BJ, *et al.*: **Independent genomewide screens identify the tumor suppressor VTRNA2-1 as a human epiallele responsive to periconceptional environment.** *Genome Biol.* 2015; **16**(1): 118.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stadler PF, Chen JLL, Hackermüller J, *et al.*: **Evolution of vault RNAs.** *Mol Biol Evol.* 2009; **26**(9): 1975–1991.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stark R, Grzelak M, Hadfield J: **RNA sequencing: the teenage years.** *Nat Rev Genet.* 2019; **20**(11): 631–656.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Steitz J: **RNA-RNA base-pairing: Theme and variations.** *RNA.* 2015; **21**(4): 476–477.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sun Y, Miao N, Sun T: **Detect accessible chromatin using ATAC-seq, from principle to applications.** *Hereditas.* 2019; **156**: 29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Suojalehto H, Lindström I, Majuri ML, *et al.*: **Altered microRNA expression of nasal mucosa in long-term asthma and allergic rhinitis.** *Int Arch Allergy Immunol.* 2014; **163**(3): 168–178.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Szklarczyk D, Morris JH, Cook H, *et al.*: **The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible.** *Nucleic Acids Res.* 2017; **45**(D1): D362–D368.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tahiri A, Leivonen SK, Lüders T, *et al.*: **Deregulation of cancer-related miRNAs is a common event in both benign and malignant human breast tumors.** *Carcinogenesis.* 2014; **35**(1): 76–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Thorsson V, Gibbs DL, Brown SD, *et al.*: **The Immune Landscape of Cancer.** *Immunity.* 2018; **48**(4): 812–830.e14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thurman RE, Rynes E, Humbert R, *et al.*: **The accessible chromatin landscape of the human genome.** *Nature.* 2012; **489**(7414): 75–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Treppendahl MB, Qiu X, Sogaard A, *et al.*: **Allelic methylation levels of the noncoding *VTRNA2-1* located on chromosome 5q31.1 predict outcome in AML.** *Blood*. 2012; **119**(1): 206–216.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tsompana M, Buck MJ: **Chromatin accessibility: A window into the genome.** *Epigenetics Chromatin*. 2014; **7**(1): 33.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

van Dijk SJ, Peters TJ, Buckley M, *et al.*: **DNA methylation in blood from neonatal screening cards and the association with BMI and insulin sensitivity in early childhood.** *Int J Obes (Lond)*. 2018; **42**(1): 28–35.

[PubMed Abstract](#) | [Publisher Full Text](#)

van Zon A, Mossink MH, Schoester M, *et al.*: **Multiple human vault RNAs. Expression and association with the vault complex.** *J Biol Chem*. 2001; **276**(40): 37715–37721.

[PubMed Abstract](#) | [Publisher Full Text](#)

Vilalta A, Kickhoefer VA, Rome LH, *et al.*: **The rat vault RNA gene contains a unique RNA polymerase III promoter composed of both external and internal elements that function synergistically.** *J Biol Chem*. 1994; **269**(47): 29752–29759.

[PubMed Abstract](#)

Weeding E, Sawalha AH: **Deoxyribonucleic acid methylation in systemic lupus erythematosus: Implications for future clinical practice.** *Front Immunol*. 2018; **9**: 875.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Weinstein JN, Collisson EA, Mills GB, *et al.*: **The cancer genome atlas pan-cancer analysis project.** *Nat Genet*. 2013; **45**(10): 1113–1120.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Xiong Y, Zhang L, Holloway AK, *et al.*: **MiR-886-3p regulates cell proliferation and migration, and is dysregulated in familial non-medullary thyroid cancer.** *PLoS One*. 2011; **6**(10): e24717.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yeganeh M, Hernandez N: **RNA polymerase III transcription as a disease factor.** *Genes Dev*. 2020; **34**(13–14): 865–882.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yu Z, Chen D, Su Z, *et al.*: **miR-886-3p upregulation in clear cell renal cell carcinoma regulates cell migration, proliferation and apoptosis by targeting PITX1.** *Int J Mol Med*. 2014; **34**(5): 1409–1416.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yu MC, Lee CW, Lin CH, *et al.*: **Differential hypermethylation of the *VTRNA2-1* promoter in hepatocellular carcinoma as a prognostic factor: Tumor marker prognostic study.** *Int J Surg*. 2020; **79**: 282–289.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yu XF, Zou J, Bao ZJ, *et al.*: **miR-93 suppresses proliferation and colony formation of human colon cancer stem cells.** *World J Gastroenterol*. 2011; **17**(42): 4711–4717.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhang XO, Gingeras TR, Weng Z: **Genome-wide analysis of polymerase III-transcribed *Alu* elements suggests cell-type-specific enhancer function.** *Genome Res*. 2019; **29**(9): 1402–1414.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zheng G, Qin Y, Clark WC, *et al.*: **Efficient and quantitative high-throughput tRNA sequencing.** *Nat Methods*. 2015; **12**(9): 835–837.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 20 August 2021

<https://doi.org/10.5256/f1000research.57200.r87202>

© 2021 Hasler D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Daniele Hasler

Biochemistry Center Regensburg (BZR), Laboratory for RNA Biology, University of Regensburg, Regensburg, Germany

The authors addressed all the points meticulously or provided convincing arguments in their point-by-point response in case requested analyses were not performed. The only, very very little, point is that, despite claiming to have modified the legend to Figure 8, the abbreviation "PANCAN" is still used in the current version of the manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: RNA biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 24 Aug 2021

Maria Ana Duhagon

Dear Reviewer,

As we expressed before, your revision has been of great value for our manuscript, so we again thank you. We will amend the uncorrected error in the legend of Fig 8 in the final version of the manuscript, that will also incorporate the recommendations of the other two Reviewers.

Sincerely,

Maria Ana Duhagon

Competing Interests: No competing interests were disclosed.

Reviewer Report 16 July 2021

<https://doi.org/10.5256/f1000research.57200.r88586>

© 2021 Lee Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yong Sun Lee

Department of Cancer Biomedical Science, Graduate School of Cancer Science and Policy, National Cancer Center, Goyang, South Korea

In this manuscript, Fort and Duhagon examined the expression of vault RNAs (vtRNAs). vtRNAs were initially known to be a component of the vault complex, whose role is still elusive. However, recently, a growingly number of studies has begun to report interesting roles of vtRNAs, most of which are independent of the vault complex. These days, next generation sequencing techniques have provided a huge amount of information on gene expression profiles in diverse biological contexts. However, as the authors indicated, vtRNAs have been out of this benefit, because of their sizes and, allegedly by the authors, their secondary structures. So, the authors turned into ATAC-seq data and CpG methylation data to infer their expression levels and analyzed them in several different aspects.

The best of all, the value of this manuscript lies in the fact that it has provided comprehensive information about vtRNAs. This manuscript is useful, albeit not a breakthrough paper, for vtRNA researchers. Overall, this manuscript is well qualified for indexing. However, incorporation of my points below would improve this manuscript, I hope.

Major points

1. The manuscript is too lengthy and I recommend succinct description. Especially, the Discussion part is mostly repetition of what has been already stated in the Results section.

Minor points

1. Page 3, "SMAD5 and TGFB1 genes": not TGFB1, but TGFBI.
2. Page 4, "VtRNA2-1 has also been shown to act as a microRNA precursor in different tissues, serving both as TSG...": this description is followed by citation of many references. However, most of them simply deemed vtRNA2-1 to be miR-886-5p or -3p to go for subsequent experiments; for example, miRNA mimic transfection, target prediction, etc. All those experiments and analyses were nonsense, without prudent validation whether vtRNA2-1 generates miRNAs. The authors should mention this point and eliminate all references, except those that have provided experimental evidence for the presence of miRNA miR-886-5p or -3p.
3. Page 15, "there is evidence of vtRNA transcript regulation via RNA methylation by NSUN2, or via its association to other proteins like DUSP11, DIS3L2, SSB, SRSF2 and DICER": regarding vtRNA2-1 regulation by Dicer, this paper "nc886 is induced by TGF- β and suppresses the microRNA pathway in ovarian cancer"¹ should be included.

4. Page 16, "...Alternatively, the methods used to quantify the vtRNAs are not comparable among the studies": I believe that the levels of the four vtRNAs could not be compared among one another even within one study, unless they included a standard to normalize variation in efficiency between Northern probes or between qRT-PCR primers. This caveat must be mentioned. In addition, the authors should check each of the references (whether they did measurement with a standard) and cite selectively.

References

1. Ahn J, Lee H, Lee J, Lee Y, et al.: nc886 is induced by TGF- β and suppresses the microRNA pathway in ovarian cancer. *Nature Communications*. 2018; **9** (1). [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 06 July 2021

<https://doi.org/10.5256/f1000research.57200.r88408>

© 2021 Sengupta D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Deepanwita Sengupta

Department of Biology, Stanford University, Stanford, CA, USA

In this article, Fort *et al.* speculate the comprehensive function of vtRNAs in human health and disease. They analyze the vtRNA expression in TCGA Pan-Cancer cohort using dual approach: ATAC-seq to determine chromosome accessibility and genome-wide DNA methylation. Their analyses revealed the landscape of vtRNA chromatin status, and also identified pathways that the different vtRNA genes are part of, thereby shedding light into their diverse roles in cancer.

The paper is very well written, is easy to follow and has proper background and references. Methods have been described in sufficient details. The questions postulated by the authors are sound and the analyses performed by them directly test their hypothesis/question. Their conclusions are in accordance with their findings in most parts. I only have a few minor comments:

1. The authors used the ChIP-seq data of only K562 cell line to analyze TFs occupancy at a +/- 3000 bp region. It will be nice if they could validate their finding from this analysis with the ChIP-seq data of at least another cell line.
2. Its interesting that when the authors tested the clinical significance of the chromatin accessibility of the vtRNA promoters, they could only find a significant difference in OS when patients were stratified by promoter DNA methylation and not when stratification was done on basis of ATAC-seq. Can the authors comment on why this is happening? Also it will be really interesting to see if the results are different when instead of doing the analysis on all patients together, the analysis is performed on patient cohorts stratified on the basis of age, sex, disease type, etc.
3. In the discussion section, can the authors address the discrepancy in correlation between chromatin accessibility and promoter methylation between vtRNA1-2 and 2-1 with respect to the others?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epigenetics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 14 April 2021

<https://doi.org/10.5256/f1000research.31555.r82633>

© 2021 Hasler D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Daniele Hasler

¹ Biochemistry Center Regensburg (BZR), Laboratory for RNA Biology, University of Regensburg, Regensburg, Germany

² Biochemistry Center Regensburg (BZR), Laboratory for RNA Biology, University of Regensburg, Regensburg, Germany

The study by Fort and Duhagon estimates the expression of a class of non-coding (nc)RNAs termed 'vault RNAs' (vtRNAs) across different cancer types. Since vtRNAs might be underrepresented applying conventional RNA sequencing techniques, the authors infer vtRNAs expression by analyzing chromatin characteristics at promoter regions of vtRNA genes. In more detail, Fort and Duhagon use genome-wide chromatin accessibility data (ATAC-seq) as well as DNA methylation data sets from The Cancer Genome Atlas (TCGA) consortium to provide a comprehensive comparison between the four annotated vtRNA genes. By this approach, the authors determine the correlation between chromatin accessibility and DNA methylation of vtRNA promoters and compare their chromatin characteristics across different cancer types. Furthermore, Fort and Duhagon use publically available CHIP-seq datasets originating from a leukemia cell line to determine transcription factors (TFs) binding to vtRNA genes and identify a set of 23 core TFs common to all vtRNAs which might govern chromatin accessibility at these genomic sites.

Further analyses of the TCGA DNA methylation data sets provide evidence for vtRNA-specific deregulation upon malignant transformation in specific cancer types, as well as lower patient survival probability associated with reduced vtRNA promoter methylation. Finally, the authors determine genes that are co-regulated with vtRNAs in terms of chromatin accessibility and the also profile chromatin accessibility and DNA methylation of vtRNA promoters in six immune subtypes of the pan-cancer TCGA datasets.

VtRNAs are a class of ncRNAs with rather enigmatic functions, albeit they have been recently

implicated in important cellular processes^{1,2} and in cancer biology[ref-]. The work by Fort and Duhagon is the first pan-cancer study focusing on vtRNAs and it provides a nice and comprehensive overview for the potential function of the single vtRNA transcripts as tumor suppressors or oncogenes in specific cancer types.

The manuscript is well written, although some aspects could have been presented a bit more concisely. In particular, the authors dedicate special emphasis to the relative comparison between vtRNA genes (e.g. Figures 3A and 3B and result section related to Figures 4 and 8). The question arises whether a key message of the study is that specific vtRNA genes are up- or down-regulated in certain cancer conditions or whether a key point is that the relative expression levels of the four vtRNA genes are altered. If so, the authors should propose a model in the discussion section.

Additional major points:

The major limitation of this study is that vtRNA expression was not analyzed directly but was inferred from chromatin accessibility data (ATAC-seq) as well as DNA methylation data. Nevertheless, the authors honestly point out this limitation and provide convincing evidences from literature that support the validity of their study. Taken together, this approach sounds reasonable.

Nevertheless, the findings could be further strengthened by analyzing TCGA miRNA-seq datasets for the expression of vtRNA-derived small RNA fragments. This strategy has been applied in previous studies to infer the expression of snoRNAs⁴ and tRNAs⁵ across different cancer types, assuming that small processing products correlate with expression levels of the full-length transcripts. Indeed snoRNAs and in particular tRNAs are ncRNA with similar sequencing biases as vtRNAs.

Furthermore, since a small RNA derived from vtRNA-2-1 has been linked to different cancers (see references within the manuscript), this additional pan-cancer analysis would be informative not only concerning this specific fragment but might also uncover the relevance of additional vtRNA-derived small RNAs.

The analysis of TF occupancy at vtRNA loci is, in my opinion, rather speculative and vague. Since vtRNAs are transcribed by RNA polymerase III (pol III), is there any clear and direct evidence from literature that the identified TFs regulate Pol III activity (except for the TFIIIA, TFIIIB and TFIIIC complexes)? Could the binding of these TFs rather regulate transcription of adjacent RNA polymerase II genes and not of vtRNAs? The authors might at least briefly touch these aspects in the discussion.

It would be also worth to determine whether deregulation (e.g. of any of the 9 vtRNA1-2-specific TFs or any of the 4 vtRNA2-1-specific TFs) correlates with the changes of vtRNA promoter DNA methylation in normal vs tumor samples (Figure 6). This would be more informative than the current conclusions concerning shared TFs between vtRNA loci and their co-regulated expression. Indeed, this analysis might help to understand, e.g., how the expression of a single vtRNA locus is regulated within the vtRNA1 cluster.

Interestingly, the chromatin profile of vtRNAs in brain lower grade glioma (LGG) indicates that vtRNA expression is strongly reduced in this condition. Does this effect hold true also for other classes of pol III transcripts (e.g. Y RNAs)? If so, this finding might point towards a general

reduction of pol III activity in LGG.

Minor points:

The authors should reformulate their statement in the introduction “VtRNA2-1 has also been shown to act as a microRNA precursor”, as the vtRNA2-1-derived fragment was not convincingly proven to be a *bona fide* microRNA and the hsa-miR-886 entry has been retracted in the current miRBase release 22.1⁶.

Please correct the typo on page 3 replacing “TGFB1” by “TGFB1”; i.e. Transforming Growth Factor Beta Induced (TGFB1) and not Transforming Growth Factor Beta 1 (TGF1).

Since lower patient survival of cancer patients was associated with lower vtRNA promoter methylation and not with higher promoter accessibility, the paragraph “High chromatin accessibility at the promoter of vtRNA1-1, vtRNA1-2 and vtRNA2-1 is associated with low patient overall survival” should be renamed. In light of the statement “DNA methylation is a less direct and less accurate measurement of chromatin accessibility compared to ATAC-seq” (page 16), it is questionable whether the relationship between DNA methylation and patient survival is due to increased vtRNA expression, as a similar result would be expected also from the analysis of the ATAC-seq data. Is this simply due to a larger samples size of the DNA methylation dataset?

The authors do not mention that a pseudogene of vtRNA2-1 (vtRNA2-2P) has been annotated at chr2:65,555,432-65,555,534. Is there any reason for that? VtRNA2-2P could be included as proof of concept similarly to vtRNA3-1P on page 8.

Page 3: “...the presence of silencing mutations at the B box element of its promoter B-box”. Since the organization of RNA polymerase III promoters has not been explained before, it would be better to keep this statement more general without mentioning the B-box.

The abbreviation “PANCAN” in the title of the legend to Figure 8 was not used before in the text.

I am not sure whether the term “chromatin polymorphism” on page 10 is correct. The referenced publication refers to epigenetic variations.

References

1. Amort M, Nachbauer B, Tuzlak S, Kieser A, et al.: Expression of the vault RNA protects cells from undergoing apoptosis. *Nat Commun.* 2015; **6**: 7030 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Horos R, Büscher M, Kleinendorst R, Alleaume A, et al.: The Small Non-coding Vault RNA1-1 Acts as a Riboregulator of Autophagy. *Cell.* 2019; **176** (5): 1054-1067.e12 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Yeganeh M, Hernandez N: RNA polymerase III transcription as a disease factor. *Genes & Development.* 2020; **34** (13-14): 865-882 [PubMed Abstract](#) | [Publisher Full Text](#)
4. Gong J, Li Y, Liu CJ, Xiang Y, et al.: A Pan-cancer Analysis of the Expression and Clinical Relevance of Small Nucleolar RNAs in Human Cancer. *Cell Rep.* 2017; **21** (7): 1968-1981 [PubMed Abstract](#) | [Publisher Full Text](#)
5. Sun X, Yang J, Yu M, Yao D, et al.: Global identification and characterization of tRNA-derived RNA fragment landscapes across human cancers. *NAR Cancer.* 2020; **2** (4). [PubMed Abstract](#) | [Publisher Full Text](#)
6. Kozomara A, Birgaoanu M, Griffiths-Jones S: miRBase: from microRNA sequences to function. *Nucleic Acids Research.* 2019; **47** (D1): D155-D162 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: RNA biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 16 Jun 2021

Maria Ana Duhagon

Reviewer: Daniele Hasler, Biochemistry Center Regensburg (BZR), Laboratory for RNA Biology, University of Regensburg, Regensburg, Germany.

Reviewers' comment: The study by Fort and Duhagon estimates the expression of a class of non-coding (nc)RNAs termed 'vault RNAs' (vtRNAs) across different cancer types. Since vtRNAs might be underrepresented applying conventional RNA sequencing techniques, the authors infer vtRNAs expression by analyzing chromatin characteristics at promoter regions of vtRNA genes. In more detail, Fort and Duhagon use genome-wide chromatin accessibility data (ATAC-seq) as well as DNA methylation data sets from The Cancer Genome Atlas (TCGA) consortium to provide a comprehensive comparison between the four annotated vtRNA genes. By this approach, the authors determine the correlation between chromatin accessibility and DNA methylation of vtRNA promoters and compare their chromatin characteristics across different cancer types. Furthermore, Fort and Duhagon use publically available CHIP-seq datasets originating from a leukemia cell line to determine transcription factors (TFs) binding to vtRNA genes and identify a set of 23 core TFs common to all vtRNAs which might govern chromatin accessibility at these genomic sites.

Further analyses of the TCGA DNA methylation data sets provide evidence for vtRNA-specific deregulation upon malignant transformation in specific cancer types, as well as lower patient survival probability associated with reduced vtRNA promoter methylation. Finally, the authors determine genes that are co-regulated with vtRNAs in terms of chromatin accessibility and the also profile chromatin accessibility and DNA methylation of vtRNA promoters in six immune subtypes of the pan-cancer TCGA datasets.

VtRNAs are a class of ncRNAs with rather enigmatic functions, albeit they have been recently implicated in important cellular processes^{1,2} and in cancer biology[ref-]. The work by Fort and Duhagon is the first pan-cancer study focusing on vtRNAs and it provides a nice and comprehensive overview for the potential function of the single vtRNA transcripts as tumor suppressors or oncogenes in specific cancer types.

The manuscript is well written, although some aspects could have been presented a bit more concisely. In particular, the authors dedicate special emphasis to the relative comparison between vtRNA genes (e.g. Figures 3A and 3B and result section related to Figures 4 and 8). The question arises whether a key message of the study is that specific vtRNA genes are up- or down-regulated in certain cancer conditions or whether a key point is that the relative expression levels of the four vtRNA genes are altered. If so, the authors should propose a model in the discussion section.

Authors response: We are grateful for the evaluation of our manuscript entitled “Pan-Cancer chromatin analysis of the human vtRNA genes uncovers their association with cancer biology”, since we find these constructive comments rather pertinent and valuable. We tried to answer all the points raised by the Reviewer and hope it led us to the improved of the manuscript.

It is true that the article is long. We believe the length is justified by the type of information it provides. In particular, the description of the patterns of vtRNA inferred expression involves four genes in 20 cancer types assessed by two methods, plus normal tissues data for DNA methylation.

We think that the message of the study is fairly written in the abstract and conclusion, neither of which prioritize the inferred relative expression of the vtRNAs per se, but what in the inferred expression pattern is suggestive of the involvement of the four genes in cancer. The article aimed to use the chromatin status of the genes to withdraw their candidate roles in cancer tissue. In our view the figures mentioned by the Reviewer are about their pattern of expression in cancer and immune subtypes, regulation, and the derived functional implications of those findings. Nevertheless, the narrative requires a description of their relative abundance and change, while their comparison helps to visualize how different they are expressed and the implication for their function in normal and tumor tissue.

Additional major points:

Reviewers' comment: *The major limitation of this study is that vtRNA expression was not analyzed directly but was inferred from chromatin accessibility data (ATAC-seq) as well as DNA*

methylation data. Nevertheless, the authors honestly point out this limitation and provide convincing evidences from literature that support the validity of their study. Taken together, this approach sounds reasonable.

Nevertheless, the findings could be further strengthened by analyzing TCGA miRNA-seq datasets for the expression of vtRNA-derived small RNA fragments. This strategy has been applied in previous studies to infer the expression of snoRNAs⁴ and tRNAs⁵ across different cancer types, assuming that small processing products correlate with expression levels of the full-length transcripts. Indeed snoRNAs and in particular tRNAs are ncRNA with similar sequencing biases as vtRNAs.

Authors response: We agree the small RNA sequencing analysis proposed by the Reviewer would be valuable to provide further support for the expression of the vtRNAs inferred from the chromatin status. Indeed, we have done it previously while studying snc886-3p, a microRNA-like RNA derived from vtRNA2-1 (published in Fort *et al.*, 2020). There we showed that nc886 and snc886-3p expression (determined by qRT-PCR) negatively correlate with DNA methylation in six PrCa cell lines analyzed (r Spearman -0.71).

Nevertheless, since vtRNAs derived fragments are not quantified with the TCGA small RNA workflow, **we need to request access to protected controlled-access data** - primary sequencing data (BAM or FASTQ files) to identify them. We understand that this approach and the subsequent data analyses exceed the extent of the current manuscript.

Attempting to follow the Reviewer's recommendation using readily available data, we analyzed the NCI-60 cell lines panel, whose genome DNA methylation and small RNA-seq data is published. Indeed, the 4 vtRNA are known to be subject to posttranscriptional processing at some extent (Hussain *et al.*, 2013; Sajini *et al.*, 2019). We found that vtRNA2-1 produces more specific small RNAs than the other three vtRNAs, thus arguing in favor of a specialization of vtRNA2-1 as a Dicer substrate, which may explain the better correlation between the precursor and the small RNAs derived from it (see Panel A of the figure attached to this response; *Figure_reviewer_response-I*. Zenodo.

<http://doi.org/10.5281/zenodo.4784690>). In addition, vtRNA2-1 generates more than 30 times small RNAs than the other vtRNAs (see Panel B of the figure attached to this response). As anticipated, the expression of the small RNA fragments derived from vtRNA2-1 shows a better correlation with promoter methylation in the NCI-60 cell lines ($r = -0.77$). Meanwhile, the small RNAs derived from vtRNAs of cluster 1 are less abundant and less associated to promoter methylation ($r = -0.19$ - -0.24) (see Panel C of the figure attached to this response).

In addition, the average abundance of the small RNAs derived from the vtRNAs of cluster 1 correlates with the relative methylation of their promoters (small RNAs derived from vtRNA1-1 > vtRNA1-2 > vtRNA1-3 and promoter methylation vtRNA1-1 < vtRNA1-2 < vtRNA1-3) (see Panel B of the figure attached to this response), in agreement with the TCGA data presented in our manuscript. Since small RNAs derived from vtRNA2-1 are more abundant than those derived from the other three vtRNAs, the relative expression of its fragments does not correlate with the relative DNA methylation of the others.

In summary, the analysis of NCI-60 cell lines, a panel comprising all tissue types, confirms that the abundance of the small RNAs derived from vtRNAs negatively correlates with their promoter methylation but indicated it may be useful just for vtRNA2-1. The latter, together

with the low abundance of the small RNAs derived from the vtRNAs and their processing (Hussain *et al.*, 2013; Sajini *et al.*, 2019), suggest that the abundance of small RNAs in the TCGA small RNA-seq data may not be a good proxy for vtRNA expression.

As discussed above, the post-transcriptional regulation of vtRNA seems to represent an additional layer of complexity of vtRNA biology that includes the processing into small RNAs. We speculate that only some of the fragments derived from specific vtRNAs may have functional significance, but it is likely reached by mechanisms different from those in which their precursors vtRNAs participate. We believe small RNAs derived from vtRNA are an interesting subject of investigation, requiring an independent study, which incorporates the biological mechanism in which they play a part.

Reviewers' comment: *Finally, in order to be more accurate about the limitation of the study mentioned by the Reviewer, we substituted the term "expression" for "chromatin status" or related terms when was appropriate.*

The analysis of TF occupancy at vtRNA loci is, in my opinion, rather speculative and vague. Since vtRNAs are transcribed by RNA polymerase III (pol III), is there any clear and direct evidence from literature that the identified TFs regulate Pol III activity (except for the TFIID, TFIIB and TFIIC complexes)? Could the binding of these TFs rather regulate transcription of adjacent RNA polymerase II genes and not of vtRNAs? The authors might at least briefly touch these aspects in the discussion.

Authors response: We welcome the Reviewers comment since we had a similar concern about the significance of these observations. Yet, we reasoned that the identification of viral related transcription factors (TFs) near vtRNA genes was a provocative finding in the context of their involvement in viral infection and innate immune response. In addition, RNA Pol III (Pol III) has been linked to the innate immune response at different levels, thus it is tempting to speculate that the Pol III regulated genes are coordinately regulated by viral related TFs dependently or independently of RNA Pol II (Pol II) action.

The identified TFs that are shared by the vtRNA (23 TFs) comprise TFs known to be part of the Pol III basal machinery (3/23), or to be direct (6/23) or indirect interactors (3/23) of Pol III machinery. This observation endorses their participation in the vtRNA genes transcription.

There are no Pol II genes near or overlapping the vtRNA genes, thus this is an unlikely explanation for TF occupancy in the vicinity of the genes. Indeed, several lines of evidence have shown that Pol II and some of its specific TFs affect Pol III activity (White, 2011). The localization of the Pol II and its specific TFs (FOS, JUN, ETS1) near Pol III promoters (200bp upstream of TSS) in human cell lines, the global regulation by Pol II specific TFs (p53, MAF1, MYC, RB), the regulation of Pol III by Pol II activity, the sharing chromatin marks and the basal TFIIB component TBP. The direct interaction between specific TFs and TFIIB is also known to modulate Pol III transcription.

It has been proposed that Pol II may assist RNA pol III activity by either binding, transcription, or stalling, contributing to make the chromatin more permissive for transcription by Pol III. Curiously, Raha et al demonstrated a reduction of the three vtRNAs

of locus 1 after α -amanitin treatment and the co-localization of Pol III and Pol II near these genes (Raha *et al.*, 2010). Additional evidence of the Pol II regulation of specific Pol III transcribed genes has been published, including tRNAs (mediated by MAF1 (Gerber *et al.*, 2020), GAF1 (Rodríguez-López *et al.*, 2020), U6 RNA (OCT1, SP1, STAF, reviewed in (Hirsch *et al.*, 2000)) Alu elements (mediated by YY1, SP1, MEG2, FOS, EP300, AP-1, (Zhang *et al.*, 2019 and references therein). Finally, the crosstalk between the two polymerases seems to be tissue and gene specific, suggesting that Pol II-Pol III crosstalk may be finer than previously envisioned.

To amend the lack of discussion of this aspect we included the following sentence in the Discussion of the manuscript: *"It is worth to mention that RNA polymerase II TFs clustered near TSS of RNA polymerase III transcripts could modulate occupancy and transcription rate of RNA polymerase III through RNA pol II activity (Campbell and White, 2014; Gerber *et al.*, 2020; Raha *et al.*, 2010; Zhang *et al.*, 2019). Specifically, RNA polymerase II could assist chromatin opening, thus allowing accessibility of RNA polymerase III or its associated factors in the human genome (Raha *et al.*, 2010)."*

Reviewers' comment: *It would be also worth to determine whether deregulation (e.g. of any of the 9 vtRNA1-2-specific TFs or any of the 4 vtRNA2-1-specific TFs) correlates with the changes of vtRNA promoter DNA methylation in normal vs tumor samples (Figure 6). This would be more informative than the current conclusions concerning shared TFs between vtRNA loci and their co-regulated expression. Indeed, this analysis might help to understand, e.g., how the expression of a single vtRNA locus is regulated within the vtRNA1 cluster.*

Authors response: Following the Reviewer's suggestion, we determined the correlation between vtRNAs chromatin accessibility and the specific TFs associated with vtRNAs (Table attached to this revision, *Table_TFs_Binding_Factors.csv* Zenodo.

<http://doi.org/10.5281/zenodo.4784690>), but we did not find significant associations. It is worth to recall that TF activity is strongly regulated at the post-transcription level, thus TF abundance may not be a good indicator of TF activity. It might be also reasoned that TFs may be poised in an inactive state, to turn active upon the proper signalling, which might well be a viral invasion or an immune system response mediator. Whether this TF DNA occupancy is relevant in cancer remains to be elucidated.

While reviewing the literature to respond to this point it came to our attention that there is a link between RNAPol III and innate immune response (Graczyk *et al.*, 2015), that could be an alternative explanation of the viral TF enrichment at the vicinity of vtRNAs. Therefore, we decided to introduce this phrase at the end of the discussion of this aspect: *"Alternatively, since RNA polymerase III transcripts are involved in virus replication and immune response (Graczyk *et al.*, 2015), it is possible that the TF bound to vtRNA vicinity are part of a global and not a vtRNA specific viral response."*

In view of the above, in an attempt to reduce the conjectures derived from the TFs analysis we modified the abstract and the conclusion of the paper as explained below.

Abstract: We substituted the sentence "Although the vtRNAs are co-regulated by transcription factors related to viral infection, vtRNA2-1 is the most independently regulated homologue." for the following: *"VtRNAs promoters are enriched in transcription factors related*

to viral infection. VtRNA2-1 is likely the most independently regulated homologue."

Conclusion: We substituted the sentence "The four vtRNAs seem to be co-regulated at the transcriptional level, and TFs involved in viral infection are likely to take part in their coordinated transcription." for the following: *"VtRNA genes are enriched in viral related TFs. The comparison of TFs and chromatin status of the vtRNAs suggest that vtRNA2-1 is likely the most independently regulated vtRNA locus."*

Reviewers' comment: *Interestingly, the chromatin profile of vtRNAs in brain lower grade glioma (LGG) indicates that vtRNA expression is strongly reduced in this condition. Does this effect hold true also for other classes of pol III transcripts (e.g. Y RNAs)? If so, this finding might point towards a general reduction of pol III activity in LGG.*

Authors response: In order to answer the Reviewer, we determined the ATAC-seq chromatin accessibility of RNYs (RNY1, RNY3, RNY4 and RNY5). As seen in the new Supplementary Figure 3 and modified Supplementary Table 1, the RNYs showed a different chromatin accessibility profile within different tissues, among them and compared to vtRNAs. Interestingly, RNY3 and RNY4 show a conspicuously higher chromatin accessibility in LGG compared with other tissues, which is opposite to the lower accessibility observed for the vtRNAs. These observations raise a possible RNA pol III misregulation in LGG tumors, which must be gene specific to justify opposite outcomes for RNY and vtRNAs. The modified Supplementary Figure 3 and modified Supplementary Table 1 are now included in the manuscript.

To incorporate these findings in the text, the following sentence was included in the results section: *"The same approach was performed for the Y-RNAs genes RNY1 (Ave. 3.7, 0.4 SD), RNY3 (Ave. 2.1, 0.6 SD), RNY4 (Ave. 1.9, 0.6 SD) and RNY5 (Ave. 3.1, 0.4 SD) (Canella et al., 2010), which code for produce ubiquitously expressed RNA polymerase III transcripts (Extended Data: Figure S3 and Tables S1)."*

Minor points:

Reviewers' comment: *The authors should reformulate their statement in the introduction "VtRNA2-1 has also been shown to act as a microRNA precursor", as the vtRNA2-1-derived fragment was not convincingly proven to be a bona fide microRNA and the hsa-miR-886 entry has been retracted in the current miRBase release 22.1⁶.*

Authors response: Following the advice, we modified the sentences related to small RNAs derived from vtRNA2-1. *"VtRNA2-1 has also been shown to generate small RNAs that could act like microRNAs in different tissues,..."*

Reviewers' comment: *Please correct the typo on page 3 replacing "TGFB1" by "TGFB1"; i.e. Transforming Growth Factor Beta Induced (TGFB1) and not Transforming Growth Factor Beta 1 (TGF1).*

Authors response: We are sorry for the mistake; we have corrected the name of the gene.

Reviewers' comment: *Since lower patient survival of cancer patients was associated with lower vtRNA promoter methylation and not with higher promoter accessibility, the paragraph "High chromatin accessibility at the promoter of vtRNA1-1, vtRNA1-2 and vtRNA2-1 is associated with low patient overall survival" should be renamed. In light of the statement "DNA methylation is a less direct and less accurate measurement of chromatin accessibility compared to ATAC-seq" (page 16), it is questionable whether the relationship between DNA methylation and patient survival is due to increased vtRNA expression, as a similar result would be expected also from the analysis of the ATAC-seq data. Is this simply due to a larger samples size of the DNA methylation dataset?*

Authors response: We amended the title of the paragraph to "Low DNA methylation at the promoter of vtRNA1-1, vtRNA1-2 and vtRNA2-1 is associated with low patient overall survival."

Despite no statistically significant differences in overall survival were observed using ATAC-seq patient data the curves were like those observed using the promoter DNA methylation. The lack of significance is likely due to the smaller ATAC-seq sample size as anticipated by the Reviewer. In addition, the association between OS and vtRNA promoter DNA methylation is weak per se, so sample size is likely critical to sense significant differences.

Reviewers' comment: *The authors do not mention that a pseudogene of vtRNA2-1 (vtRNA2-2P) has been annotated at chr2:65,555,432-65,555,534. Is there any reason for that? VtRNA2-2P could be included as proof of concept similarly to vtRNA3-1P on page 8.*

Authors response: We have previously looked for vtRNA2-2P in ATAC-seq data but there is no ATAC-seq region determined for this gene in the TSS, so we decided to do not include it. However, we now extended the analysis to the ATAC-seq region nearest to the TSS of vtRNA2-2P. As can be seen in modified Supplementary Figure 3 and modified Supplementary Table 1, the vtRNA2-2P has a very low and stable pattern of ATAC-seq chromatin accessibility across the different tissues that agrees with its pseudogene annotation. These results were now included in the manuscript and the text was modified accordingly. In the introduction section we added: "The vtRNA3-1P and vtRNA2-2P are two additional vault RNA genes situated in chromosome X and 2 respectively, but were classified as a pseudogenes due to the absence of expression in cell lines and the presence of silencing mutations at the promoter (Büscher et al., 2020; van Zon et al., 2001).". In the results section we added: "The low ATAC-seq value of the pseudogenes vtRNA3-1P (Ave. -1.2, 0.6 SD) and vtRNA2-2P (Ave. 0.1, 0.6 SD), represents a proof of concept of the analyses (Extended Data: Figure S3 and Tables S1-S5). The same approach was performed for the Y-RNAs genes RNY1 (Ave. 3.7, 0.4 SD), RNY3 (Ave. 2.1, 0.6 SD), RNY4 (Ave. 1.9, 0.6 SD) and RNY5 (Ave. 3.1, 0.4 SD) (Canella et al., 2010), which code for produce ubiquitously expressed RNA polymerase III transcripts (Extended Data: Figure S3 and Tables S1)." and "VtRNA2-2P reaches a maximum and minimum promoter chromatin accessibility in Lung Adenocarcinoma (LUAD) (1.0) and ACC (-0.34) respectively (Extended Data: Figure S3C)."

Reviewers' comment: *Page 3: "...the presence of silencing mutations at the B box element of its promoter B-box". Since the organization of RNA polymerase III promoters has not been explained before, it would be better to keep this statement more general without mentioning the B-box.*

Authors response: To attend the Reviewer's suggestion, we modified the sentence to *"the presence of silencing mutations at the promoter"*.

Reviewers' comment: *The abbreviation "PANCAN" in the title of the legend to Figure 8 was not used before in the text.*

Authors response: Due the Reviewer's comment, we modified "PANCAN" for *"Pan-Cancer"*.

Reviewers' comment: *I am not sure whether the term "chromatin polymorphism" on page 10 is correct. The referenced publication refers to epigenetic variations.*

Authors response: Due the Reviewer's comment, we modified the term "chromatin polymorphism" for *"epigenetic variation"*.

References:

- Fort, R.S., Garat, B., Sotelo-Silveira, J.R., & Duhagon, M.A. (2020). *vtRNA2-1/nc886 Produces a Small RNA That Contributes to Its Tumor Suppression Action through the microRNA Pathway in Prostate Cancer*. Non-Coding RNA 6, 7.
- Gerber, A., Ito, K., Chu, C.S., & Roeder, R.G. (2020). *Gene-Specific Control of tRNA Expression by RNA Polymerase II*. Mol. Cell 78, 765–778.
- Graczyk, D., White, R.J., & Ryan, K.M. (2015). *Involvement of RNA Polymerase III in Immune Responses*. Mol. Cell. Biol. 35, 1848–1859.
- Hirsch, H.A., Gu, L., & Henry, R.W. (2000). *The Retinoblastoma Tumor Suppressor Protein Targets Distinct General Transcription Factors To Regulate RNA Polymerase III Gene Expression*. Mol. Cell. Biol. 20, 9182–9191.
- Hussain, S., Sajini, A.A., Blanco, S., Dietmann, S., Lombard, P., Sugimoto, Y., Paramor, M., Gleeson, J.G., Odom, D.T., Ule, J., *et al.* (2013). *NSun2-mediated cytosine-5 methylation of vault noncoding RNA determines its processing into regulatory small RNAs*. Cell Rep. 4, 255–261.
- Raha, D., Wang, Z., Moqtaderi, Z., Wu, L., Zhong, G., Gerstein, M., Struhl, K., & Snyder, M. (2010). *Close association of RNA polymerase II and many transcription factors with Pol III genes*. Proc. Natl. Acad. Sci. U. S. A. 107, 3639–3644.
- Rodríguez-López, M., Gonzalez, S., Hillson, O., Tunnacliffe, E., Codlin, S., Tallada, V.A., Bähler, J., & Rallis, C. (2020). *The GATA Transcription Factor Gaf1 Represses tRNAs, Inhibits Growth, and Extends Chronological Lifespan Downstream of Fission Yeast TORC1*. Cell Rep. 30, 3240–3249.e4.
- Sajini, A.A., Choudhury, N.R., Wagner, R.E., Bornelöv, S., Selmi, T., Spanos, C., Dietmann, S., Rappsilber, J., Michlewski, G., & Frye, M. (2019). *Loss of 5-methylcytosine alters the biogenesis of vault-derived small RNAs to coordinate epidermal differentiation*. Nat. Commun. 10.
- White, R.J. (2011). *Transcription by RNA polymerase III: More complex than we thought*. Nat. Rev. Genet. 12, 459–463.
- Zhang, X.O., Gingeras, T.R., & Weng, Z. (2019). *Genome-wide analysis of polymerase III-transcribed Alu elements suggests cell-type-specific enhancer function*. Genome Res. 29, 1402–1414.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research