SYSTEMATIC REVIEW

REVISED The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis [version 2; peer review: 1 approved, 1 approved with reservations]

Osfar Sjöfjan¹, Danung Nur Adli ^{1,2}, Rakhmad Perkasa Harahap^{2,3}, Anuraga Jayanegara^{2,4}, Dicky Tri Utama⁵, Ainun Pizar Seruni⁶

¹Department of Nutrition and Feed Technology, University of Brawijaya, Malang, East Java, 65145, Indonesia

²Animal Feed and Nutrition Modelling (AFENUE) Research Group, IPB university, Bogor, West Java, 16680, Indonesia

³Study Program of Animal Science, University of TanjungPura, Pontianak, Borneo, 78124, Indonesia

⁴Department of Nutrition and Feed Technology, IPB University, Bogor, West Java, 16680, Indonesia

⁵Department of Animal Product Technology, Faculty of Animal Science, Malang, East Java, 65145, Indonesia

⁶Graduate School, University of Missouri, Missouri, Columbia, 65211, USA

v2 First published: 05 Mar 2021, 10:183
<https://doi.org/10.12688/f1000research.51219.1>
 Second version: 19 Aug 2021, 10:183
<https://doi.org/10.12688/f1000research.51219.2>
 Latest published: 13 Oct 2021, 10:183
<https://doi.org/10.12688/f1000research.51219.3>

Abstract

Introduction: The number of publications in Scopus on this topic increased from less than 50 in 1995 to more than 250 in 2015. In other hand, inconsistency in results about the correlation between yeast and lactic acid bacteria as probiotics has been evident since the early publications on use in broilers.

Methods: A meta-analysis was conducted to determine relationship between lactic acid bacteria and yeast as probiotics to broiler diets on the growth performance, relative organ weight, blood parameters, and immune response of the broiler. A database was designed based on published data that reported the use of probiotics on the broiler. The method used for selecting articles was based on the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) method. Articles selected were taken from PubMed, Web of science, Scopus, Google Scholar, and Science direct databases as well as individual.

Results: The final database consists of 49 *in vivo* articles, 93 studies, and 225 treatments. The analysis statement in the system was a PROC MIXED procedure of SAS software. The level of probiotic increased ($p < 0.001$) body weight, body weight gain, and feed intake of broiler.

Open Peer Review

Approval Status

	1	2
version 3		
(revision)
13 Oct 2021		view
version 2
(revision)
19 Aug 2021	view	view
version 1
05 Mar 2021	view	

- Hazem Mohammed Ebraheem Shaheen**
 , Damanhour University, Damanhour, Egypt
- Youssef A. Attia**, King Abdulaziz University,
 Jeddah, Saudi Arabia

Any reports and responses or comments on the

There was a reduction ($p < 0.01$) on feed conversion ratio and mortality on the level probiotic given to broiler. Supplementation of probiotics in broiler diet increased ($p < 0.001$) the weight of liver, spleen, gizzard, bursa of fabricius and carcass yield, while reduced ($p < 0.001$) abdominal fat weight. The probiotic given increased the total of red and white blood cells (both at $p < 0.001$) but did not affect lymphocyte.

Discussion: It can be concluded the yeast act as supporting agent that serves lactic acid bacteria as probiotic increases the growth performance, relative organ weight, blood parameters, and immune response of the broiler.

Keywords

broiler, lactic acid bacteria, meta-analysis, probiotic, yeast.

article can be found at the end of the article.

Corresponding author: Osfar Sjöfjan (osfar@ub.ac.id)

Author roles: Sjöfjan O: Conceptualization, Funding Acquisition, Supervision, Writing – Review & Editing; Adli DN: Data Curation, Project Administration, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; Harahap RP: Formal Analysis, Software, Validation, Visualization; Jayanegara A: Conceptualization, Data Curation, Investigation, Methodology, Software, Supervision, Validation, Writing – Review & Editing; Utama DT: Resources, Writing – Review & Editing; Seruni AP: Investigation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Sjöfjan O *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sjöfjan O, Adli DN, Harahap RP *et al.* **The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis [version 2; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:183 <https://doi.org/10.12688/f1000research.51219.2>

First published: 05 Mar 2021, 10:183 <https://doi.org/10.12688/f1000research.51219.1>

REVISED Amendments from Version 1

We followed the suggestions of the reviewer. Some of the improvements are detailed as follows:

On database development, correction was:

- 'After careful evaluation, the parameters included were growth performance, relative organ weight, carcass quality, blood parameters, and immune responses. While the economical parameters are not included due to insufficient data in the paper. The strains recorded on the raw database were Ross308, which dominated at 63.26%; Arbor Acres at 32.65%; and others at 4.09%.'

On the growth performance section, we added:

- 'Thus, utilization of feed is also reflected on the nutrient digestibility.'

We also edited the references.

Any further responses from the reviewers can be found at the end of the article

Introduction

In 1997, the use of antibiotics in livestock was first addressed in Denmark with *avoparcin* as an antibiotic growth promoter (AGP). The trend continued and a European Union (EU)-wide ban on AGPs in animal feed (poultry) took effect in 2006 (EC Regulation No 1831/2003)¹. Since then, this type of regulation has spread to developing countries, including Indonesia, which has been banning antibiotics and imported poultry feed products since the most recent regulation, PERMENTAN/14/16/2017, was put in place². The EU introduced probiotics as an alternative to antibiotics and this has subsequently become an area of great interest for researchers worldwide³. Probiotics are living microorganisms that when ingested in sufficient amounts, may positively improve growth, intestinal health and animal productivity. Probiotics are commonly sourced from lactic acid bacteria, namely, *Lactobacillus* and *Bifidobacterium*, which are usually found in the intestine⁴.

Earlier studies have reported an active role for probiotics in reducing or eliminating the pathogen bacteria in the intestine. In recent research^{5,6} probiotic mixtures have also been found to have beneficial effects against a wide range of disorders, although evidence that mixtures are more effective than their component strains is more limited. Nevertheless, in the future, a further potential advantage of multi-strain probiotics, in addition to exerting additive or synergistic effects, is that the strain-specific effects of individual probiotic components could together exert a broader spectrum of activity^{5,6}. Probiotics can be given in both powder and liquid form and positively modulate the composition of broiler intestinal microflora via the stimulation of potentially beneficial bacterial populations and the reduction of pathogenic bacteria¹⁻⁴. The interaction between probiotics and micro biota added to diet influences the microbial population's stability and the health of the host. The gut micro biota plays a crucial role in host metabolism and fundamentally influences physiology, health and well-being, functionality and performance⁵.

Yeasts have been reported to act as supporting agents for lactic acid bacteria but also as having the potential to reduce avian bacterial in the gut micro biota of poultry^{1,4}. Inconsistency in results about the correlation between yeast and lactic acid bacteria as probiotics has been evident since the early publications on use in broilers⁴. Accordingly, the current study aims to determine the relationship between lactic acid bacteria and yeasts as probiotics in broiler diets on growth performance, meat quality, blood parameters, and immune responses, through a meta-analysis using data from published articles.

Methods

Database development

A database was constructed based on peer-reviewed and published research articles which reported the use of probiotics in the broiler diet. Articles were selected based on the Systematic Review Center for Laboratory Animal Experimentation (SYRCLE') method⁷ and Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA)⁸. Articles selected were taken from PubMed, Web of science, Scopus, Google Scholar, and Science direct databases as well as individual journals such as World Poultry Journal Science, British Poultry Science Journal, and International Journal of Poultry Science using the keywords 'probiotic', 'broiler', 'performance', 'organ weight', 'carcass', and 'blood serum'. In each article evaluated, the reference list was also searched for relevant articles. The raw database information from articles, authors, year of study, broiler (strain and sex), diet used in trial, length of trial, level of treatment, form and dosage of probiotic contained in the study was recorded in a spreadsheet following the referenced method⁷. After careful evaluation, the parameters included were growth performance, relative organ weight, carcass quality, blood parameters, and immune responses. While the economical parameters are not included due to insufficient data in the paper. The strains recorded on the raw database were Ross308, which dominated at 63.26%; Arbor Acres at 32.65%; and others at 4.09%.

Criteria for an article to be included in database were as follows: (a) article was published in a peer-reviewed with range 2008–2020, this paper length was chosen as related to journals aged last 10–12 years are often good⁹ (b) the broiler were modern-controlled-trial environment and management, (c) performed directly on broiler *in vivo* as the experimental animals, (d) The log concentration of lactic acid bacteria and yeast both powder and liquid form on the trial was transformation into 10¹⁰ in the database development, (e) non-probiotic treatment excluded from the database, (e) the articles written consistently in English were considered in studies, (f) the average duration of the study was minimum 0–21 days and the maximum at 0–53 days, (g) dosages given at the range 0–10 g/kg from total feed formulation. Moreover, the dependent and independent variables were selected with the aim of lactic acid bacteria and yeast related as probiotic on the broiler. Likewise, data extraction was completed in accordance with the task analysis to obtain the exact values from graphical data, the relevant figure from the papers were subjected to an online tool, WebplotDigitizer 4.4 (<https://automeris.io/WebPlotDigitizer/>), following the method¹⁰.

The final database consisted of 49 *in vivo* articles, 93 studies, and 225 n-total (3,375 n-total of total in this experiment). The details for the study selection included in this meta-analysis are provided in [Figure 1](#). The search strategy is presented in [Table 1](#). The summary of the final database is presented in [Table 2](#), and PICOS criteria presented in [Table 6](#).

Data analysis

Statistical dataset analysis using a mixed-model approach was applied^{11–14} with statement analysis in the system using the MIXED procedure of SAS (version 9.1, SAS Institute Inc., 2008), the following model was applied: The findings of a study was then taken as a random effect, while the supplementation concentration was taken as the fixed effect as follows^{15–17}:

$$Y_{ij} = B_0 + B_1 X_{ij} + B_2 X_{ij}^2 + s_i + b_i X_{ij} + e_{ij}$$

where Y_{ij} = the expected output for dependent variable Y at level j from the variable X as a continuous variable in the study i , B_0 = overall intercept across all studies (fixed effect), B_1 = linear regression coefficient of Y on X (fixed effect), B_2 = quadratic regression coefficient of Y on X (fixed effect), X_{ij} = value of the continuous predictor variable (probiotic supplementation level), s_i = random effect of study i , b_i = random effect of study i on the regression coefficient of Y on X in study i and e_{ij} = unexplained residual error. In the statement CLASS, the “study” variable was declared. Data were weighted by the number of replicates in each study. Additionally, an unstructured variance – covariance matrix (type = un) was performed at the random effect part of the model to avoid a positive correlation between intercepts and slopes. Significance of an effect was stated at the probability level of $p < .05$, and $p < .1$ was considered as a tendency of significance. In case that the quadratic model

Figure 1. Diagram flow of article selection included in meta-analysis adapted from PRISMA Method.

Table 1. Search strategy.

Databases	Search strategy
PubMed	((("probiotics" [MeSH Terms] OR "probiotics" [All Fields] OR "probiotic" [All Fields]) OR (serum [All Fields] AND blood [All Fields]) AND blood [All Fields])) OR
Scopus	("broiler" [MeSH Terms] OR "broiler" [All Fields]) OR (organ [All Fields] AND ("weight" [Journal] OR "weight" [All Fields])) AND ("chickens" [MeSH Terms] OR "carcass" [All Fields] OR "carcass" [All Fields]).
Google Scholar	<p>'Probiotic'</p> <p>'Growth Performances'</p> <p>'Relative organ weight'</p> <p>'Blood performances'</p> <p>'Carcasses'</p> <p>'broiler'</p> <p>'Cited by' 'related articles'</p> <p>'Since year'</p> <p>'2008–2020'</p> <p>Search in full text Search in title</p> <p>Result 1 Result 1</p> <p>Result 2 Result 2</p> <p>Result 3 Result 3</p> <p>Result 4</p> <p>Result 5 Result 5</p> <p>Result 6 Result 6</p> <p>Result 7 Result 7</p> <p>Result 8</p>

Table 2. Studies included in the meta-analyses of the relationship between lactic acid bacteria and yeast as probiotics on the growth performance, relative organ weight, blood parameters, and immune response of broiler.

No	References	Kind of Probiotic	Form	Dosage (g/kg)	Periods (d)
1	Chen <i>et al.</i> ¹⁸	Lactic acid bacteria	Powder	0-2	0-28
2	Park and Kim ¹⁹	Lactic acid bacteria	Powder	0-1	0-28
3	Zhang <i>et al.</i> ²⁰	Lactic acid bacteria	Powder	0-1	0-35
4	Jamshidparvar <i>et al.</i> ²¹	Lactic acid bacteria	Liquid	0-2	0-42
5	Khan <i>et al.</i> ²²	Lactic acid bacteria	Liquid	0-1	0-39
6	Gheisar <i>et al.</i> ²³	Lactic acid bacteria	Powder	0-0.50	0-35
7	Hussein <i>et al.</i> ²⁴	Yeast	Powder	0-0.50	0-35
8	Nosrati <i>et al.</i> ²⁵	Lactic acid bacteria	Powder	0-0.18	0-42
9	Javandel <i>et al.</i> ²⁶	Lactic acid bacteria	Powder	0-0.9	0-42
10	Sun and Kim ²⁷	Yeast	Powder	0-0.02	0-35
11	Sugiharto <i>et al.</i> ²⁸	Lactic acid bacteria	Powder	0-0.05	0-42
12	Ghasemi <i>et al.</i> ²⁹	Lactic acid bacteria	Powder	0-0.04	0-42
13	Abdel-Hafeez <i>et al.</i> ³⁰	Lactic acid bacteria	Powder	0-1.5	0-42
14	Toghyani <i>et al.</i> ³¹	Lactic acid bacteria	Powder	0-0.15	0-42
15	Salah <i>et al.</i> ³²	Lactic acid bacteria	Powder	0-2	0-42

No	References	Kind of Probiotic	Form	Dosage (g/kg)	Periods (d)
16	Paryad and Mahmoud ³³	Yeast	Powder	0-0.02	0-42
17	Koc <i>et al.</i> ³⁴	Yeast	Powder	0-2	0-21
18	Zhou <i>et al.</i> ³⁵	Lactic acid bacteria	Powder	0-0.4	0-35
19	Rezaei-pour ³⁶	Yeast	Powder	0-7.5	0-42
20	Cho <i>et al.</i> ³⁷	Lactic acid bacteria	Powder	0-0.2	0-35
21	Priya and Babu ³⁸	Yeast	Powder	0-1.5	0-36
22	Lan <i>et al.</i> ³⁹	Lactic acid bacteria	Powder	0-0.05	0-35
23	Sun and Kim ⁴⁰	Yeast	Powder	0-0.2	0-35
24	Elnagar ⁴¹	Yeast	Powder	0-0.2	0-53
25	Mashayekhi <i>et al.</i> ⁴²	Lactic acid bacteria	Powder	0-0.5	0-42
26	Attia <i>et al.</i> ⁴³	Yeast	Powder	0-1	0-35
27	Pournazari <i>et al.</i> ⁴⁴	Lactic acid bacteria	Powder	0-2	0-42
28	Riyazi <i>et al.</i> ⁴⁵	Lactic acid bacteria	Powder	0-0.15	0-42
29	Sugiharto <i>et al.</i> ⁴⁶	Yeast	Powder	0-0.4	0-35
30	Manafi <i>et al.</i> ⁴⁷	Yeast	Powder	0-0.01	0-42
31	Ashayerizadeh <i>et al.</i> ⁴⁸	Lactic acid bacteria	Powder	0-0.9	0-42
32	Zahirian <i>et al.</i> ⁴⁹	Yeast	Powder	0-4	0-42
33	Sugiharto <i>et al.</i> ⁵⁰	Lactic acid bacteria	Powder	0-0.2	0-38
34	Sugiharto <i>et al.</i> ⁵¹	Lactic acid bacteria	Powder	0-0.15	0-35
35	Sugiharto <i>et al.</i> ⁵²	Lactic acid bacteria	Powder	0-0.01	0-35
36	Miah <i>et al.</i> ⁵³	Lactic acid bacteria	Liquid	0-0.5	0-21
37	Sherief <i>et al.</i> ⁵⁴	Yeast	Powder	0-0.5	0-42
38	Reisinger <i>et al.</i> ⁵⁵	Yeast	Powder	0-0.2	0-35
39	Attia <i>et al.</i> ⁵⁶	Yeast	Powder	0-0.05	0-35
40	Sjofjan and Adli ⁵⁷	Lactic acid bacteria	Combination	0-0.8	0-35
41	Makled <i>et al.</i> ⁵⁸	Lactic acid bacteria	Powder	0-5	0-42
42	Vase <i>et al.</i> ⁵⁹	Lactic acid bacteria	Liquid	0-0.03	0-42
43	Waqas <i>et al.</i> ⁶⁰	Yeast	Powder	0-0.06	0-35
44	Caruk <i>et al.</i> ⁶¹	Lactic acid bacteria	Powder	0-0.1	0-42
45	Yalçin <i>et al.</i> ⁶²	Yeast	Powder	0-3	0-42
46	Yalçinkaya <i>et al.</i> ⁶³	Yeast	Powder	0-1	0-42
47	Shokaiyan <i>et al.</i> ⁶⁴	Lactic acid bacteria	Powder	0-0.5	0-42
48	Salehizadeh <i>et al.</i> ⁶⁵	Lactic acid bacteria	Powder	0-0.1	0-42
49	Khajeh <i>et al.</i> ⁶⁶	Lactic acid bacteria	Powder	0-0.5	0-42

above was not significant, the model was changed into its corresponding linear model. The variable of the study was declared in the class statement as it did not contain any

quantitative information. The regression equations are also presented with *p*-value, and root mean square error (RMSE).

Furthermore, to determine (1) interaction between lactic acid bacteria and yeast; (2) interaction of type probiotic (powder and liquid) according to the following model¹⁶:

$$Y_{ij} = \mu + S_i + \tau_j + S\tau_{ij} + e_{ij}$$

Where Y_{ij} = the expected output for dependent variable Y , μ = overall mean, S_i = random effect of i study, τ_j = fixed effect of the j level, $S\tau_{ij}$ = random interaction between i study and the j level, and e_{ij} = residual error. A significant effect was declared at $p < 0.05$ or there is a tendency when the p -value was between 0.05 and 0.10.

Results

Table 3 presents the effects of probiotics on broiler performance. The meta-analysis results show the level of probiotic ($p < 0.001$) body weight, body weight gain, and feed intake of broilers. In contrast, there was a reduction ($p < 0.01$) on feed conversion ratio (FCR) and mortality on the level probiotic given to broiler. Furthermore, the analysis also shows that the

form of probiotic in the feed does not create any significant difference in broiler performance.

The weight of abdominal organs and carcass yield of broilers were affected by the supplementation of probiotics in the diet (Table 4). Supplementation of probiotics in broiler diet increased ($p < 0.001$) the weight of liver, spleen, gizzard, bursa of Fabricius and carcass yield, while reduced ($p < 0.001$) abdominal fat weight. Different types of probiotic, i.e., powder or liquid, influenced the weight of liver ($p = 0.001$), spleen ($p < 0.005$), gizzard ($p = 0.045$) and bursa of Fabricius, ($p < 0.001$). In contrast, abdominal fat and carcass yield were not affected by the type of probiotics supplemented in the diet. Further, different culture type, i.e., lactic acid bacteria or yeast, had no significant effect on the abdominal organs weight and carcass yield of broilers.

The effects on blood parameters of lactic acid bacteria and yeasts as probiotics are presented in Table 5. The probiotic given increased the total of red and white blood cells (both at

Table 3. Effect of probiotics on performance of broiler.

Response parameter	Unit	n	Parameter estimates				Model statistics			
			Intercept	SE intercept	Slope	SE slope	p-Value	RMSE	P x L	B x Y
BW	gram	225	2071	62.1	13.1	2.22	<.001	314	0.523	0.166
BWG	gram	225	1567	61.7	13.8	1.76	<.001	248	0.506	0.847
FCR	-	225	1.85	0.031	-0.010	0.00	<.001	0.193	0.630	0.310
FI	gram	225	3163	120	4.74	2.7	<.001	380	0.609	0.361
Mortality	%	225	0.665	0.179	-0.002	0.01	<.001	1.80	0.83	0.474

Note: P = powder; L=liquid; B=LAB; Y=yeast; Slope: The respond when the probiotic at the zero level, SE intercept: standard error intercept; BW: body weight; BWG: body weight gain; FCR: feed conversion ratio; FI: feed intake; root mean square error (RMSE)

Table 4. Effect of probiotics on carcass and organ weight of broiler.

Response parameter	Unit	n	Parameter estimates				Model statistics			
			Intercept	SE intercept	Slope	SE slope	p-Value	RMSE	P x L	B x Y
Liver	g/kg	225	2.44	0.06	0.003	0.004	<.001	0.60	0.0001	0.27
Spleen	g/kg	225	0.34	0.07	0.004	0.002	<.001	0.30	0.0005	0.11
Gizzard	g/kg	225	1.50	0.06	0.003	0.002	<.001	0.26	0.004	0.52
Bursa of fabricius	g/kg	225	0.34	0.055	0.0005	0.002	<.001	0.34	<.001	0.51
Abdominal fat	g/kg	225	1.58	0.07	-0.015	0.004	<.001	0.62	0.37	0.50
Carcass yield	%	225	67.6	0.73	0.067	0.02	<.001	3.36	0.38	0.23

Note: P= powder; L=liquid; B=LAB; Y=yeast; Slope: The respond when the probiotic at the zero level, SE intercept: standard error intercept; root mean square error (RMSE)

Table 5. Effect of probiotics on blood and immune responses of broiler.

Response parameter	Unit	n	Parameter estimates				Model statistics			
			Intercept	SE intercept	Slope	SE slope	p-Value	RMSE	P x L	B x Y
RBC	/ μ L	225	2.25	0.05	0.003	0.002	<.001	0.30	0.38	0.50
WBC	/ μ L	225	289	9.40	0.42	0.41	<.001	58	0.21	0.03
Lymphocyte	%	225	53	1.55	-0.02	0.06	<.001	8.73	0.47	0.004
Hemoglobin	mg/dL	225	8.24	0.38	0.003	0.01	<.001	1.63	0.70	0.97

Note: RBC: red blood cell; WBC: white blood cell; P= powder; L=liquid; B=LAB; Y=yeast; Slope: The respond when the probiotic at the zero level, SE intercept: standard error intercept; root mean square error (RMSE)

Table 6. PICOS Criteria.

	Search strategy	Exclusion criteria
Participant	Broiler	Non-broiler
Interventions	Probiotic (Lactic acid bacteria and yeast)	Irrelevant treatment
Comparison	Control group (Maize – Soya bean diet [basal])	
Outcomes	LAB, BW, BWG, FCR, FI, RBC, WBC, RMSE, SE, PXL	
Study design	Modern controlled environment during <i>in-vivo</i>	

P = powder; L=liquid; B=LAB; Y=yeast; Slope: The respond when the probiotic at the zero level, SE intercept: standard error intercept; BW: body weight; BWG: body weight gain; FCR: feed conversion ratio; FI: feed intake; RBC: red blood cell; WBC: white blood cell root mean square error (RMSE); PICOS: population, intervention, comparison, outcomes and study

$p < 0.001$) but did not affect lymphocyte. Furthermore, the immune response hemoglobin results were not significantly influenced by the delivery of different lactic acid bacteria and yeast forms such as powder or liquid.

Discussion

The effect of probiotics on growth performance

Our meta-analysis shows that probiotics positively affect growth performance. In terms of growth performance, we suggest that this finding is related to the ability of probiotics to induce intestinal mechanisms, resulting in a reduction in pathogenic bacteria. In the digestive system, intestinal pH, intestinal bacteria composition, and digestive activity are improved when probiotics are present in diets¹⁸. Some probiotics are known to produce enzymes, amylase, protease, and lipase to optimize nutrients' breakdown^{19,20}. They can also increase specific enzymes in the host digestive tract to enhance nutrient absorption in the diet. In the poultry industry, probiotics are supplemented into the diet to maintain health by enhancing gut health, modulating the immune system, lowering glycaemic response, and improving various performances parameters^{21–23}. Moreover, the administering of probiotics has several ways in practice. The administering of probiotics can be included in the basal diet or combined with raw materials that contain prebiotics to enhance its effect^{20,30,50–54,67}. The probiotic can be given alone or with

another additive without any negative effect such as acidifiers and phytogenics^{25,26,32,42,44}. Furthermore, probiotics can contain one or multiple microorganism strains that can be added to animals' diets^{24,28,37}.

There are previous studies that report that cell-wall components of yeast in dietary supplementation to lactic acid bacteria improve the growth rate, feed consumption, and feed efficiency in broilers^{51,55}. These positive and consistent results were due to yeast activating spores to reduce and remove potential pathogens in the gut which possibly increases body weight⁴⁴. Linearly, the factors related to synergism between yeast and lactic acid bacteria reported in the research could be related to environmental conditions in various experiments^{45,46}.

One study from 46 showed that rearing the broiler with a heat-stress environment at 35°C was more low weight than heat stress exposed. Thus, yeast failed to alleviate heat-stress on the performance of broiler⁴⁶. The lower temperature in the chicken house may help increase feed intake to eat more of the experimental diets⁵⁷. In addition, other factors related due to being reared under a stocking density stress of 43 kg live weight per m² floor space⁵⁷. Another result⁴⁵ explains in more detail that yeasts in powder form significantly increase the body weight of broilers starting at 21 days old, with an increase

in line with increasing feed intake and reducing FCR. These consistent results^{57–59} explain that the use of lactic acid bacteria as probiotics in powder form help to increase the body weight of broilers, as a result of digestibility and metabolic process improvement caused by the bacteria, affecting energy partition and putting more energy into growth than maintenance⁴⁶.

Moreover, the positive use of probiotics both as powders and liquids was in line with the increasing level of treatment in broiler⁵⁷. The probiotics enhance liquid lactic acid bacteria synergism with yeast in the feed but suggested at an optimized level⁵⁷ of 0.8%. However, the dose-response relationship of probiotics in animal trials is rarely studied¹. At low doses a probiotic may be specific, for example *bifidobacteria*, due to the high specificity of *bifidobacteria* for that particular probiotic¹. In other hand, if the dose increased, this would leave some substrate for other probiotic strains able to ferment it. The outcome of high dose would show less specificity than that of the low dose¹. Treatment with both powder and liquid forms increases body weight and feed intake and reduces feed conversion^{31,57,61}. Moreover, probiotics for farm animals have positive effects on growth, efficiency of feed utilization⁶⁸. Thus, utilization of feed is also reflected on the nutrient digestibility⁶⁹. In addition, the consistent result in studies⁴⁵ vs 50–52 show the relationship between both yeast and lactic acid bacteria working together to reduce potential pathogens in the gut of broilers but dose is dependent.

The effect of probiotic on the relative organ weight and carcass quality

The meta-analysis results show limited effects on the carcass and organ weight of broilers. In agreement from 52,62 the carcass quality shows no significant difference after administering probiotics of both lactic acid bacteria and yeast. The one factor can be caused reduced of percentage carcass are heat-stress environmental. The carcass heat-stress was associated with the reduced of carcass quality⁵². Apart from physiological adjustment derived from depressed feed intake, the increased *Corticosterone* level may be responsible lower percentage of carcass⁵². Carcass percentage was reported to increase by one study⁴⁵, with the saleable product in terms of edible portions. Reported⁴⁵ the carcass quality can be affected from physiological and genetic potential, feed formulated, strain of the broiler rearing⁶⁹. The excess fat deposition in carcass of broiler is undesirable to producers because of reduced carcass yield and to consumers that prefer a leaner product¹.

Likewise, one study⁶² the use of yeast as probiotic reduced abdominal fat of broiler. Reported⁶² yeast help to reduce fat deposition because, modern-broiler-farming were intensive feeding (*ad-libitum*), fatter caused limb defects, and sudden death syndrome. The probiotics reported reduced fat of broiler compared without probiotic⁶⁵. The mode of action was that probiotics decreased the activity of acetyl-CoA carboxylase⁶⁶. Acetyl-CoA carboxylase has been widely suggested as the rate-limiting enzyme in fatty acid synthesis⁶⁶. The decline in the synthesis of fatty acids, in turn, would decrease their availability for esterification to triglycerides for deposited in the adipose tissue⁶⁶. Furthermore, the minimum dose of probiotic to

stimulate fatty acids are currently unknown^{63,64,66}. Differences in the broiler line/breed and conditions, as well as microorganism strains (highly species-and strain specific), origin species, concentrations, and methods of administration of the probiotic bacteria, may explain these results⁶⁵. However, our study can't exactly suggest the dose optimum for using this probiotic.

In one study⁴², the effects of probiotics on relative weights of liver and spleen were not significant ($P > 0.05$), while bursa of Fabricius relative weight increased ($p < 0.05$). Supplementing diets with probiotics could help to prevent necrotic enteritis which associated with degeneration of hepatocytes and immune system of the broiler⁷⁰. The smaller liver in broiler may indicate a higher resistance to pathogen microorganism such as *Clostridium perfringens*⁷¹. The IGF-1 can produce short-fatty acids (SCFAs), which act either directly or indirectly on the liver and adipose tissue to promote growth of organ and skeletal development⁶⁵. The report from 72 at the end of the feeding trial showed that the development of gizzard weight was decreased, dateable irregularities in the gizzard are a sensitive index to reduce anti-nutritional factors in the basal diet after exposure to toxic substance not to amount of lactic bacteria⁷². Moreover, increased weight of this lymphoid organ may indicate a higher immunity achieved in treated broiler, which could be explained by probiotic anti-microbial activity^{18,42}. The factor affected by significant differences in the relative organ weight is the ability to absorb substances from probiotics^{18,43,44}. 18 stated that variances between the broilers result from impacts on absorption and other capacities of the relative organ weight. The growth factors correlate with age, while the broiler's uses in the relative age cause the same internal organ's growth. In instances, an increased relative organ weight may be in line with an increase in lymphocyte concentration¹⁸.

The effect of probiotic on the blood parameters and immune responses

The meta-analysis of different probiotic levels on some blood parameters showed red and white blood cell concentration increased ($p < 0.05$) with increasing probiotic supplementation levels in the feed. The increased of the white blood cells had correlated with yeast reduce the uric acid (UA) content in the blood⁷³. The uric acid is a metabolite of protein that has an antioxidant function, but is converted to a pro-oxidant in the cell or cytoplasm⁷³. In contrast, lymphocyte, and hemoglobin were not significantly different ($p > 0.05$)¹⁵. Linearly, blood serum rose in line with probiotic increase. Additionally, one¹⁸ shows that, the lactic acid bacteria that help reduce avian-pathogenic bacteria were *Escherichia coli* and *Clostridium perfringens*. The beneficial action indicates that lactic acid bacteria produced extracellular enzymes to enhance the nutrient digestibility of feed and synthesize immune function using endogenous anti-microbial²⁰. In terms of negative linear response in studies²¹ there was no positive result on red and white blood cells.

Moreover, probiotics could be related to a lowered recycling of bile salts in the gut or inhibited hepatic 3-hydroxy-3-methylglutaryl coenzyme A reductase activity²¹. The mechanism operating in lactic acid bacteria, as probiotics to elicit

their hypocholesterolemic effect is interference with intestinal bile acid transport and absorption, leading to an increase in bile acid excretion²¹. The potential pathogens reduce but are not eliminated, thus, probiotics balance the intestinal environment to enhance the broiler's immune systems^{45,50–52}. Although, the lactic acid bacteria do not produce butyric acid themselves, they stimulate the proliferation of butyric acid and cell-wall of yeast in the blood circulation by the mechanism that is called cross-feeding⁶⁵. Continued research⁴² shows probiotics help to increase the white blood cell count as level of probiotic is increased. One study⁴² stated that the increase of white blood cells and immune response was due to the level increase of B and T lymphocyte production. In line with the 41 studies the amount of red blood cells, hemoglobin, and white blood cells consistently tends to increase compared to controls. The positive effect from yeast as a probiotic could derive from its outer cell wall components namely: chitin, mannan, and glucan which have an immunostimulant effect. Moreover, these outer wall components promote lactic acid bacteria activity, which is activated by producing enzymes that cause disintegration of bile salts, making them unconjugated⁶⁸. The yeast can enhance the immune response by promoting growth of lactic acid bacteria and thus simultaneously producing antibacterial substances and stimulating the production of immunoglobulin³³. Thus, yeast acts as a supporting agent of lactic acid bacteria, which adhere to the endogenous epithelial cells to initiate colonization³³.

Conclusions

The results provided by this meta-analysis demonstrates the enhancement of overall performance of broilers supplemented with lactic acid bacteria and yeast as probiotics. Effects of the probiotics on blood parameters are dose dependent, where areas, the additives have limited effects on organ weight and carcass percentage. Both powder and liquid forms of probiotics do not affect the results differently. The future research

trends are to determine the dose optimum of probiotic for broiler.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Extended data

Figshare: Extended data for ‘The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis’. <https://doi.org/10.6084/m9.figshare.14060414>⁷⁴.

This project contains extracted data of outcome measures (BW: body weight; BWG: body weight gain; FCR: feed conversion ratio; FI: feed intake; RBC: red blood cell; WBC: white blood cell).

Reporting guidelines

Figshare: PRISMA checklist for ‘The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis’. <https://doi.org/10.6084/m9.figshare.14060501>⁷⁵.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC BY 4.0)

Acknowledgements

Thanks to Animal Feed and Nutrition Modelling (AFENUE) Research Group, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia

References

1. Venema K, do Carmo AP: **Probiotics and prebiotics: current research and future trends**. Academic Press, 1st ed, 2015.
[Publisher Full Text](#)
2. Maron DF, Smith TJS, Nachman KE: **Restrictions on antimicrobial use in food animal production: an international regulatory and economic survey**. *Global Health*. 2013; 9(1): 48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Mehdi Y, Létourneau-Montminy MP, Gaucher ML, et al.: **Use of Antibiotics in Broiler Production: Global Impacts and Alternatives**. *Anim Nutr*. 2018; 4(2): 170–178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Park YH, Hamidon F, Rajangan C, et al.: **Application of probiotics for the production of safe and high-quality poultry meat**. *Korean J Food Sci Anim Resour*. 2016; 36(5): 567–576.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Śliżewska K, Chlebicz-Wójcik A, Nowak A: **Probiotic Properties of New *Lactobacillus* Strains Intended to Be Used as Feed Additives for Monogastric Animals**. *Probiotics Antimicrob Proteins*. 2021; 13(1): 146–162.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Żbikowski A, Pawłowski K, Śliżewska K, et al.: **Comparative Effects of Using New Multi-Strain Synbiotics on Chicken Growth Performance, Hematology, Serum Biochemistry and Immunity**. *Animals (Basel)*. 2020; 10(9): 1555.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. de Vries RBM, Hooijmans CR, Langendam MW, et al.: **A protocol format for the preparation, registration and publication of systematic reviews of animal intervention studies**. *Evi Pre Med*. 2015; 2(1): 1–9.
[Publisher Full Text](#)
8. Liberati A, Altman DG, Tetzlaff J, et al.: **The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration**. *BMJ*. 2009; 339: b2700.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Pautasso M: **Ten simple rules for writing a literature review**. *PLoS Comput Biol*. 2013; 9(7): e1003149.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Drevon D, Fursa SR, Malcolm AL: **Intercoder reliability and validity of WebPlotDigitizer in extracting graphed data**. *Behav Modif*. 2017; 41(2): 323–339.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Hidayat C, Sumiati, Jayanegara A, et al.: **Effect of zinc addition on the immune response and production performance of broilers: a meta-analysis**. *Asian-Australas J Anim Sci*. 2019; 33(3): 465–479.
[PubMed Abstract](#) | [Free Full Text](#)
12. Patra AK: **The effect of dietary fats on methane emissions, and its other effects on digestibility, rumen fermentation and lactation performance in cattle: A meta-analysis**. *Livest Sci*. 2013; 155(2–3): 244–254.
[Publisher Full Text](#)

13. St-Pierre NR: **Invited Review: Integrating quantitative findings from multiple studies using mixed model methodology.** *J Dairy Sci.* 2001; **84**(4): 741–755.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Sauviant D, Schmidely P, Daudin JJ, *et al.*: **Meta-analyses of experimental data in animal nutrition.** *Animal.* 2008; **2**(8): 1203–1214.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Prihambodo TR, Sholikin MM, Qomariyah N, *et al.*: **Effects of dietary flavonoids on performance, blood constituents, carcass composition and small intestinal morphology of broilers: A meta-analysis.** *Anim Biosci.* 2021; **34**(3): 434–442.
[PubMed Abstract](#) | [Free Full Text](#)
16. Sjöfjan O, Adli DN, Sholikin MM, *et al.*: **The effects of probiotics on the performance, egg quality and blood parameters of laying hens: A meta-analysis.** *J Anim Feed Sci.* 2021; **30**(1): 11–18.
[Publisher Full Text](#)
17. Jayanegara A, Wina E, Takahashi J: **Meta-analysis on methane mitigating properties of saponin-rich sources in the rumen: influence of addition levels and plant sources.** *Asian-Australas J Anim Sci.* 2014; **27**(10): 1426–35.
[PubMed Abstract](#) | [Free Full Text](#)
18. Chen W, Wang JP, Yan L, *et al.*: **Evaluation of probiotics in diets with different nutrient densities on growth performance, blood characteristics, relative organ weight and breast meat characteristics in broilers.** *Br Poult Sci.* 2013; **54**(5): 635–641.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Park JH, Kim IH: **Supplemental effect of probiotic *Bacillus subtilis* B2A on productivity, organ weight, intestinal *Salmonella* microflora, and breast meat quality of growing broiler chicks.** *Poult Sci.* 2014; **93**(8): 2054–2059.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Zhang ZF, Zhou TX, Ao X, *et al.*: **Effects of β -glucan and *Bacillus subtilis* on growth performance, blood profiles, relative organ weight and meat quality in broilers fed maize-soybean meal based diets.** *Livest Sci.* 2012; **150**(1–3): 419–424.
[Publisher Full Text](#)
21. Jamshidparvar A, Javandel F, Seidavi A, *et al.*: **Effects of golpar (*Heracleum persicum* Desf.) and probiotics in drinking water on performance, carcass characteristics, organ weights, blood plasma constituents, and immunity of broilers.** *Environ Sci Pollut Res Int.* 2017; **24**(30): 23571–23577.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Khan SH, Yousaf B, Mian AA, *et al.*: **Assessing the effect of administering different probiotics in drinking water supplement on broiler performance, blood biochemistry and immune response.** *J Appl Anim Res.* 2011; **39**(4): 418–428.
[Publisher Full Text](#)
23. Gheisar MM, Hosseindoust A, Kim IH: **Effects of dietary *Enterococcus faecium* on growth performance, carcass characteristics, faecal microbiota, and blood profile in broilers.** *Vet Med.* 2016; **61**(1): 28–34.
[Publisher Full Text](#)
24. Hussein E, Selim S: **Efficacy of yeast and multi-strain probiotic alone or in combination on growth performance, carcass traits, blood biochemical constituents, and meat quality of broiler chickens.** *Livest Sci.* 2018; **216**: 153–159.
[Publisher Full Text](#)
25. Nosrati M, Javandel F, Camacho LM, *et al.*: **The effects of antibiotic, probiotic, organic acid, vitamin C, and *Echinacea purpurea* extract on performance, carcass characteristics, blood chemistry, microbiota, and immunity of broiler chickens.** *J Appl Poult Res.* 2017; **26**(2): 295–306.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Javandel F, Nosrati M, van den Hoven R, *et al.*: **Effects of Hogweed (*Heracleum persicum*) Powder, Flavophospholipol, and Probiotics as Feed Supplements on the Performance, Carcass and Blood Characteristics, Intestinal Microflora, and Immune Response in Broilers.** *J Poult Sci.* 2019; **56**(4): 262–269.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Sun HY, Kim IH: **Dietary Supplementation of Mixed Yeast Culture Derived from *Saccharomyces cerevisiae* and *Kluyveromyces fragilis*: Effects on Growth Performance, Nutrient Digestibility, Meat Quality, Blood Parameters, and Gut Health in Broilers.** *J Poult Sci.* 2019; **56**(2): 140–147.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Sugiharto S, Isroli I, Yudiarti T, *et al.*: **The effect of supplementation of multistrain probiotic preparation in combination with vitamins and minerals to the basal diet on the growth performance, carcass traits, and physiological response of broilers.** *Vet World.* 2018; **11**(2): 240–247.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Ghasemi-Sadabadi M, Ebrahimnezhad Y, Shaddel-Tili A, *et al.*: **The effects of fermented milk products (kefir and yogurt) and probiotic on performance, carcass characteristics, blood parameters, and gut microbial population in broiler chickens.** *Arch Anim Breed.* 2019; **62**(1): 361–374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Abdel-Hafeez HM, Saleh ESE, Tawfeek SS, *et al.*: **Effects of probiotic, prebiotic, and synbiotic with and without feed restriction on performance, hematological indices and carcass characteristics of broiler chickens.** *Asian-Australas J Anim Sci.* 2017; **30**(5): 672–682.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Toghyani M, Mosavi SK, Modaresi M, *et al.*: **Evaluation of kefir as a potential probiotic on growth performance, serum biochemistry and immune responses in broiler chicks.** *Anim Nutr.* 2015; **1**(4): 305–309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Salah AS, El-Tarabany MS, Ali MA: **Impact of dietary supplementation with a synbiotic, organic acids or their combination on growth performance, carcass traits, economic efficiency, jejunum histomorphometry and some blood indices of broiler chickens.** *Anim Prod Sci.* 2018; **59**(7): 1318–1326.
[Publisher Full Text](#)
33. Paryad A, Mahmoudi M: **Effect of different levels of supplemental yeast (*Saccharomyces cerevisiae*) on performance, blood constituents and carcass characteristics of broiler chicks.** *Afr J Agric Res.* 2008; **3**(12): 835–842.
[Reference Source](#)
34. Koc F, Samli H, Okur A, *et al.*: **Effects of *Saccharomyces cerevisiae* and/or mannanoligosaccharide on performance, blood parameters and intestinal microbiota of broiler chicks.** *Bulgarian J Agr Sci.* 2010; **16**(5): 643–650.
[Reference Source](#)
35. Zhou TX, Chen YJ, Yoo JS, *et al.*: **Effects of chitoooligosaccharide supplementation on performance, blood characteristics, relative organ weight, and meat quality in broiler chickens.** *Poult Sci.* 2009; **88**(3): 593–600.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Rezaeiipour V, Fononi H, Irani M: **Effects of dietary L-threonine and *Saccharomyces cerevisiae* on performance, intestinal morphology and immune response of broiler chickens.** *S Afr J Anim Sci.* 2012; **42**(3).
[Publisher Full Text](#)
37. Cho JH, Zhang ZF, Kim IH: **Effects of single or combined dietary supplementation of beta-glucan and kefir on growth performance, blood characteristics and meat quality in broilers.** *Br Poult Sci.* 2013; **54**(2): 216–221.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Priya BS, Babu SS: **Effect of different levels of supplemental probiotics (*Saccharomyces cerevisiae*) on performance, haematology, biochemistry, microbiology, histopathology, storage stability and carcass yield of broiler chicken.** *Int J Pharm Biol Arch.* 2013; **4**(1): 201–207.
[Reference Source](#)
39. Lan RX, Lee SI, Kim IH: **Effects of *Enterococcus faecium* SLB 120 on growth performance, blood parameters, relative organ weight, breast muscle meat quality, excreta microbiota shedding, and noxious gas emission in broilers.** *Poult Sci.* 2017; **96**(9): 3246–3253.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Sun HY, Kim IH: **Dietary supplementation of mixed yeast culture derived from *saccharomyces cerevisiae* and *Kluyveromyces fragilis*: effects on growth performance, meat quality, blood parameters, and gut health in broilers.** *J Poult Sci.* 2019; **56**(2): 140–147.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. ElNaggar AS, Abdelkhalek E: **Productive Performance, Blood Biochemical Traits And Immune Response of Sasso Chickens Supplemented With *Saccharomyces Cerevisiae* or Mannan Oligosaccharide As Natural Growth Additives.** *Egypt Poult Sci J.* 2017; **37**(2): 505–522.
42. Mashayekhi H, Mazhari M, Esmaeilipour O: **Eucalyptus leaves powder, antibiotic and probiotic addition to broiler diets: effect on growth performance, immune response, blood components and carcass traits.** *Anim.* 2018; **12**(10): 2049–2055.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Attia YA, Zewell HS, Alsaffar AA, *et al.*: **Effect of non-antibiotic feed additives as an alternative to flavomycin on productivity, meat quality and blood parameters in broilers.** *Arch Geflügelk.* 2011; **75**(1): 40–48.
[Reference Source](#)
44. Pournazari M, AA-Qotbi A, Seidavi A, *et al.*: **Prebiotics, probiotics and thyme (*Thymus vulgaris*) for broilers: performance, carcass traits and blood variables.** *Revista Colombiana de Ciencias Pecuarias.* 2017; **30**(1): 3–10.
[Publisher Full Text](#)
45. Riyazi SR, Ebrahimnezhad Y, Hosseini SA, *et al.*: **Comparison of the effects of basil (*Ocimum basilicum*) essential oil avilamycin and protexin on broiler performance, blood biochemistry and carcass characteristics.** *Arch Anim Breed.* 2015; **58**(2): 425–432.
[Publisher Full Text](#)
46. Sugiharto S, Yudiarti T, Isroli I, *et al.*: **Effect of dietary supplementation with *Rhizopus oryzae* or *Chrysonilia crassa* on growth performance, blood profile, intestinal microbial population, and carcass traits in broilers exposed to heat stress.** *Arch Anim Breed.* 2017; **60**(3): 347–356.
[Publisher Full Text](#)
47. Manafi M: **Comparison study of a natural non-antibiotic growth promoter and a commercial probiotic on growth performance, immune response and biochemical parameters of broiler chicks.** *J of Poult Sci.* 2015; **52**(4): 274–281.
[Publisher Full Text](#)
48. Ashayerizadeh O, Dastar B, Shargh MS, *et al.*: **Influence of antibiotic, prebiotic and probiotic supplementation to diets on carcass characteristics, hematological indices and internal organ size of young broiler chickens.** *J Anim Vet Adv.* 2009; **8**(9): 1772–1776.
[Reference Source](#)
49. Zahirian M, Seidavi A, Solka M, *et al.*: **Dietary supplementation of *Aspergillus oryzae* meal and its effect on performance, carcass characteristics, blood**

- variables, and immunity of broiler chickens. *Trop Anim Health Prod.* 2019; **51**(8): 2263–2268.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Sugiharto S, Yudiarti T, Isroli I: **Growth performance, haematological parameters, intestinal microbiology, and carcass characteristics of broiler chickens fed two-stage fermented cassava pulp during finishing phase.** *Trop Anim Sci J.* 2019; **42**(2): 113–120.
[Publisher Full Text](#)
 51. Sugiharto S, Yudiarti T, Isroli I, *et al.*: **Growth performance, haematological responses, intestinal microbiology and carcass traits of broiler chickens fed finisher diets containing two-stage fermented banana peel meal.** *Trop Anim Health Prod.* 2020; **52**(3): 1425–1433.
[PubMed Abstract](#) | [Publisher Full Text](#)
 52. Sugiharto S, Yudiarti T, Isroli I, *et al.*: **The effect of fungi-origin probiotic *Chrysonilia crassa* in comparison to selected commercially used feed additives on broiler chicken performance, intestinal microbiology, and blood indices.** *J Adv Vet Anim Res.* 2018; **5**(3): 332–342.
[Publisher Full Text](#)
 53. Miah MA, Sarker MMH, Das MR, *et al.*: **The influence of probiotics and antibiotic growth promoter on growth performance and hemato-biochemical parameters in Broilers.** *Curr Res in Agr Sci.* 2014; **1**(4): 103–109.
[Reference Source](#)
 54. Sherief MAR, Sherief MSA: **The Effect of single or combined dietary supplementation of mannan oligosaccharide and probiotics on performance and slaughter characteristics of broilers.** *Int J of Poult Sci.* 2011; **10**(11): 854–862.
[Reference Source](#)
 55. Reisinger N, Ganner A, Masching S, *et al.*: **Efficacy of a yeast derivative on broiler performance, intestinal morphology and blood profile.** *Livest Sci.* 2012; **143**(2–3): 195–200.
[Publisher Full Text](#)
 56. Attia YA, Al-Hamid AEA, Ibrahim MS, *et al.*: **Productive performance, biochemical and hematological traits of broiler chickens supplemented with propolis, bee pollen, and mannan oligosaccharides continuously or intermittently.** *Livest Sci.* 2014; **164**: 87–95.
[Publisher Full Text](#)
 57. Sjoftan O, Adli DN: **Effect of dietary of supplementation mannan-rich fraction (mrf) and probiotic-enhanced liquid acidifier on the growth performance, serum blood biochemistry, and intestinal properties of broilers.** *IOP Conf Ser Earth and Environ Sci.* 2020; **478**(1): 12–66.
[Reference Source](#)
 58. Makled MN, Abouelezz KFM, Gad-Elkareem AEG, *et al.*: **Comparative influence of dietary probiotic, yoghurt, and sodium butyrate on growth performance, intestinal microbiota, blood hematology, and immune response of meat-type chickens.** *Trop Anim Health Prod.* 2019; **51**(8): 2333–2342.
[PubMed Abstract](#) | [Publisher Full Text](#)
 59. Vase KK, Mortezaei SH, Rasouli B, *et al.*: **The effect of three tropical medicinal plants and superzist probiotic on growth performance, carcass characteristics, blood constituents, immune response, and gut microflora of broiler.** *Trop Anim Health Prod.* 2019; **51**(1): 33–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Waqas M, Mehmood S, Mahmud A, *et al.*: **Effect of yeast based mannan oligosaccharide (Actigen™) supplementation on growth, carcass characteristics and physiological response in broiler chickens.** *Indian J of Anim Res.* 2018; **53**(11): 1476–1479.
[Publisher Full Text](#)
 61. Kuprys-Caruk M, Michalczyk M, Chabłowska B, *et al.*: **Comparison of the effect of lactic acid bacteria added to feed or water on growth performance, health status and gut microbiota of chickens broilers.** *Anim Sci.* 2019; **58**(1): 55–67.
[Publisher Full Text](#)
 62. Yalçın S, Yalçın S, Eser H, *et al.*: **Effects of dietary yeast cell wall supplementation on performance, carcass characteristics, antibody production and histopathological changes in broilers.** *Kafkas Universitesi Veteriner Fakültesi Dergisi.* 2014; **20**(5): 757–764.
[Reference Source](#)
 63. Yalçinkaya I, Çınar M, Yildirim E, *et al.*: **The effect of prebiotic and organic zinc alone and in combination in broiler diets on the performance and some blood parameters.** *Ital J of Anim Sci.* 2016; **11**(3): e55.
[Publisher Full Text](#)
 64. Shokaiyan M, Ashayerizadeh O, Shargh MS, *et al.*: **Algal crude fucoidan alone or with *Bacillus subtilis* DSM 17299 in broiler chickens diet: growth performance, carcass characteristics, blood metabolites, and morphology of intestine.** *Poult Sci J.* 2019; **7**(1): 87–94.
[Publisher Full Text](#)
 65. Salehizadeh M, Modarressi MH, Mousavi SN, *et al.*: **Effects of probiotic lactic acid bacteria on growth performance, carcass characteristics, hematological indices, humoral immunity, and IGF-I gene expression in broiler chicken.** *Trop Anim Health Prod.* 2019; **51**(8): 2279–2286.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Khajeh BM, Afsharmanesh M, Ebrahimnejad H: **Effect of dietary *bacillus coagulans* and different forms of zinc on performance, intestinal microbiota, Carcass and Meat Quality of Broiler Chickens.** *Probiotics Antimicrob Proteins.* 2020; **12**(2): 461–472.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Adli DN, Sjoftan O, Natsir MH, *et al.*: **The effect of replacing maize with fermented palm kernel meal (FPKM) on broiler performance.** *Livest Res Rural Dev.* 2020; **32**(7): 120.
[Reference Source](#)
 68. Min Y, Choi Y, Choe J, *et al.*: **Effects of dietary mixture of protease and probiotics on growth performance, blood constituents, and carcass characteristics of growing-finishing pigs.** *J Anim Sci Technol.* 2019; **61**(5): 272–277.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. Sjoftan O, Adli DN, Natsir MH, *et al.*: **Effect of dietary modified-banana-tuber meal substituting dietary corn on growth performance, carcass trait and dietary-nutrients digestibility of coloured-feather hybrid duck.** *JITV.* 2021; **26**(1): 39–48.
[Reference Source](#)
 70. Hussein EO, Ahmed SH, Abudabos AM, *et al.*: **Effect of antibiotic, phytobiotic and probiotic supplementation on growth, blood indices and intestine health in broiler chicks challenged with *clostridium perfringens*.** *Animals (Basel).* 2020; **10**(3): 507.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Ramlucken U, Ramchuran SO, Moonsamy G, *et al.*: **A novel *Bacillus* based multi-strain probiotic improves growth performance and intestinal properties of *Clostridium perfringens* challenged broilers.** *Poult sci.* 2020; **99**(1): 331–341.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Qorbanpour M, Fahim T, Javandel F, *et al.*: **Effect of dietary ginger (*Zingiber officinale* Roscoe) and multi-strain probiotic on growth and carcass traits, blood biochemistry, immune responses and intestinal microflora in broiler chickens.** *Animals (Basel).* 2018; **8**(7): 117.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Chuang WY, Lin WC, Hsieh YC, *et al.*: **Evaluation of the combined use of *saccharomyces cerevisiae* and *aspergillus oryzae* with phytase fermentation products on growth, inflammatory, and intestinal morphology in broilers.** *Animals (Basel).* 2019; **9**(12): 1051.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Sjoftan O, Adli DN, Harahap RP, *et al.*: **Extended data for “The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis”.** 2021.
<http://www.doi.org/10.6084/m9.figshare.14060414>
 75. Sjoftan O, Adli DN, Harahap RP, *et al.*: **PRISMA checklist for “The effects of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis”.** 2021.
<http://www.doi.org/10.6084/m9.figshare.14060501>

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 23 September 2021

<https://doi.org/10.5256/f1000research.76610.r93793>

© 2021 Attia Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Youssef A. Attia

Department of Agriculture, Faculty of Environmental Sciences, King Abdulaziz University, Jeddah, Saudi Arabia

This MS has an updated review on the use of lactic acid bacteria and yeasts as probiotics on the growth performance, relative organ weight, blood parameters, and immune responses of broiler: A meta-analysis of value to broilers industry. The MS has provided helpful information about the concept of probiotics and their role in enhancing performance, health and product quality.

Major concern

- To go one step further, the dose of lactic acid bacteria and yeasts should be estimated based on this work.
- The update and renewable application of lactic acid bacteria and yeasts in broilers should given considered attention such as: compact of heat stress; control of infectious diseases; and detoxification of nitrate and aflatoxin. I have included some of my works which are good examples:
 - Control of *Salmonella enteritidis* infection in male layer chickens by acetic acid and/or prebiotics, probiotics, and antibiotics¹.
 - The detoxication of nitrate by two antioxidants or a probiotic, and the effects on blood and seminal plasma profiles and reproductive function of New Zealand White rabbit bucks².
 - Enhancing Tolerance of Broiler Chickens to Heat Stress by Supplementation with Vitamin E, Vitamin C and/or Probiotics³.
 - Nitrate detoxification using antioxidants and probiotics in the water for rabbits⁴.
 - [Health Benefits of Probiotics in Human and Animal Nutrition and Prospective Research](#).
 - Effects of different dietary levels of blue lupine (*Lupinus angustifolius*) seed meal with or

without probiotics on the performance, carcass criteria, immune organs, and gut morphology of broiler chickens⁵.

Minor concern

- Improve the language as the MS needs an English language and grammar edit.
- Enhancing the MS with figures and infographics can improve readability of the MS.

References

1. Attia YA, Ellakany HF, Abd El-Hamid AE, Bovera F, et al.: Control of Salmonella enteritidis infection in male layer chickens by acetic acid and/or prebiotics, probiotics and antibiotics. *Arch.Geflügelk.* 2012; **76** (4).
2. Attia YA, Abd El Hamid EA, Ismaiel AM, El-Nagar A: The detoxication of nitrate by two antioxidants or a probiotic, and the effects on blood and seminal plasma profiles and reproductive function of New Zealand White rabbit bucks. *Animal.* 2013; **7** (4): 591-601 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Attia Y, Al-Harathi M, El-Shafey A, Rehab Y, et al.: Enhancing Tolerance of Broiler Chickens to Heat Stress by Supplementation with Vitamin E, Vitamin C and/or Probiotics. *Annals of Animal Science.* 2017; **17** (4): 1155-1169 [Publisher Full Text](#)
4. King Abdulaziz University, Attia Y, Hamid E A, King Abdulaziz University, et al.: Nitrate detoxification using antioxidants and probiotics in the water for rabbits. *Revista Colombiana de Ciencias Pecuarias.* 2018; **31** (2): 130-138 [Publisher Full Text](#)
5. Al-Sagan AA, Al-Yemni AH, Al-Abdullatif AA, Attia YA, et al.: Effects of Different Dietary Levels of Blue Lupine (*Lupinus angustifolius*) Seed Meal With or Without Probiotics on the Performance, Carcass Criteria, Immune Organs, and Gut Morphology of Broiler Chickens. *Front Vet Sci.* 2020; **7**: 124 [PubMed Abstract](#) | [Publisher Full Text](#)

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Poultry production and nutrition

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 26 August 2021

<https://doi.org/10.5256/f1000research.76610.r92326>

© 2021 Shaheen H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hazem Mohammed Ebraheem Shaheen

Department of Pharmacology, Faculty of Veterinary Medicine, Damanhour University, Damanhour, 22511, Egypt

No additional comments.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Partly

Are sufficient details of the methods and analysis provided to allow replication by others?

Partly

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Veterinary Pharmacology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 26 July 2021

<https://doi.org/10.5256/f1000research.54362.r90146>

© 2021 Shaheen H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hazem Mohammed Ebraheem Shaheen

Department of Pharmacology, Faculty of Veterinary Medicine, Damanhour University, Damanhour, 22511, Egypt

Please specify what is the object of this study.

You could add an Economic interpretation of this study.

Data in the Results section were insufficient and need more clear for discussion.

This paragraph is not understandable:

"Criteria for an article to be included in database were as follows: (a) article was published in a peer-reviewed with range 2008–2020, this paper length was chosen as related to journals aged last 10–12 years are often good9 (b) the broiler were modern-controlled-trial environment and management, (c) performed directly on broiler in vivo as the experimental animals, (d) The log concentration of lactic acid bacteria and yeast both powder and liquid form on the trial was transformation into 1010 in the database development, (e) non-probiotic treatment excluded from the database, (e) the articles written consistently in English were considered in studies, (f) the average duration of the study was minimum 0–21 days and the maximum rom total feed formulation. Moreover, the dependent and independent variables were selected with the aim of lactic acid bacteria and yeast related as probiotic on the broiler. Likewise, data extraction was completed in accordance with the task analysis to obtain the exact values from graphical data, the relevant figure from the papers were subjected to an online tool, WebplotDigitizer 4.4 (<https://automeris.io/WebPlotDigitizer/>), following the method10".

I suggest an English language edit for the revision of the manuscript.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Partly

Are sufficient details of the methods and analysis provided to allow replication by others?

Partly

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Veterinary Pharmacology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 26 Jul 2021

Danung Nur Adli, University of Brawijaya, Malang, Indonesia

We would like to express our sincere thanks and appreciation for comprehensively reviewed our manuscript that helped us to improve our manuscript.

Regarding the reviewer suggestion and question

1. Please specify what is the object of this study.

Answer: We already mentioned in the introduction that "Accordingly, the current study aims to determine the relationship between lactic acid bacteria and yeasts as probiotics in broiler diets on growth performance, meat quality, blood parameters, and immune responses, through a meta-analysis using data from published articles".

2. You could add an Economic interpretation of this study.

Answer: Thank you for the suggestion. Regarding this suggestion the economic parameters, initially we input this parameter but in final paper we don't put down this parameter due to lack of data provided in the paper.

3. Data in the Results section were insufficient and need more clear for discussion.

Answer: We mentioned clearly in the result section.

4. **Answer:** The statement that you are asking for are the requirements when we select the paper and criteria when we put down the information, to develop the data.

5. I suggest an English language edit for the revision of the manuscript.

Answer: Authors would like say thank to second reviewer. The manuscript has been edited for English language usage by a native British proof-reader of Britannia Proofreading Service.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research