Check for updates

RESEARCH ARTICLE

Genome-wide regulation of CpG methylation by ecCEBP α in acute myeloid leukemia [version 1; peer review: 1 approved, 1 approved with reservations]

Adewale J. Ogunleye¹ , Ekaterina Romanova², Yulia A. Medvedeva^{1,2}

¹Department of Biological and Medical Physics, Moscow Institute of Physics and Technology, Moscow, Russian Federation

²Research Center of Biotechnology, Institute of Bioengineering, Institute of bioengineering, Russian Academy of Sciences, Moscow, Russian Federation

V1 First published: 11 Mar 2021, 10:204
<https://doi.org/10.12688/f1000research.28146.1>

Latest published: 31 Aug 2021, 10:204
<https://doi.org/10.12688/f1000research.28146.2>

Abstract

Background: Acute myeloid leukemia (AML) is a hematopoietic malignancy characterized by genetic and epigenetic aberrations that alter the differentiation capacity of myeloid progenitor cells. The transcription factor *CEBP α* is frequently mutated in AML patients leading to an increase in DNA methylation in many genomic locations. Previously, it has been shown that *ecCEBP α* (extra coding *CEBP α*) - a lncRNA transcribed in the same direction as *CEBP α* gene - regulates DNA methylation of *CEBP α* promoter in *cis*. Here, we hypothesize that *ecCEBP α* could participate in the regulation of DNA methylation in *trans*.

Method: First, we retrieved the methylation profile of AML patients with mutated *CEBP α* locus from The Cancer Genome Atlas (TCGA). We then predicted the *ecCEBP α* secondary structure in order to check the potential of *ecCEBP α* to form triplexes around CpG loci and checked if triplex formation influenced CpG methylation, genome-wide.

Results: Using DNA methylation profiles of AML patients with a mutated *CEBP α* locus, we show that *ecCEBP α* could interact with DNA by forming DNA:RNA triple helices and protect regions near its binding sites from global DNA methylation. Further analysis revealed that triplex-forming oligonucleotides in *ecCEBP α* are structurally unpaired supporting the DNA-binding potential of these regions. *ecCEBP α* triplexes supported with the RNA-chromatin co-localization data are located in the promoters of leukemia-linked transcriptional factors such as MLF2.

Discussion: Overall, these results suggest a novel regulatory mechanism for *ecCEBP α* as a genome-wide epigenetic modulator through triple-helix formation which may provide a foundation for sequence-specific engineering of RNA for regulating methylation of specific genes.

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
31 Aug 2021	view

version 1
11 Mar 2021	view	view

1. **Amrita Singh** , CSIR Institute of Genomics and Integrative Biology (CSIR-IGIB), New Delhi, India
2. **Oleg N. Demidov** , University of Bourgogne-Franche Comté, Dijon, France

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

Acute myeloid leukemia, Triplex, DNA methylation, long non coding RNA, extra-coding CEBPα

This article is included in the **Genomics and Genetics** gateway.

This article is included in the **Cell & Molecular Biology** gateway.

Corresponding author: Adewale J. Ogunleye (ogunleie.ad@phystech.edu)

Author roles: **Ogunleye AJ:** Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Romanova E:** Conceptualization, Data Curation, Supervision; **Medvedeva YA:** Conceptualization, Funding Acquisition, Methodology, Resources, Software, Supervision, Validation, Visualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The work has been partially supported by Russian Science Foundation (grant 18-14-00240) to YAM and the Ministry of the Sciences and Higher Education of the Russian Federation.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Ogunleye AJ *et al.* This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ogunleye AJ, Romanova E and Medvedeva YA. **Genome-wide regulation of CpG methylation by ecCEBPα in acute myeloid leukemia [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:204 <https://doi.org/10.12688/f1000research.28146.1>

First published: 11 Mar 2021, 10:204 <https://doi.org/10.12688/f1000research.28146.1>

Introduction

Acute myeloid leukemia (AML) is a malignant tumor characterized by the proliferation of undifferentiated myeloblasts^{1,2}. It is the most prevalent form of leukemia in older adults (>60 years) with an annual mortality rate of 50% and a 5-year survival rate of 24%^{2,3}. With the combined effects of the global increase in average life expectancy and AML drug inefficiency, the number of patients is expected to significantly increase in the coming years^{4,5}.

The current understanding of the molecular interplay in AML has been defined under two distinct categories; (i) genetic abnormalities and (ii) non-random chromosomal rearrangements. Cases of AML with chromosomal rearrangements as t(15;17) [*PML-RARA*], t(9;22) [*BCR-ABL*], inv(16) [*CBFB-MYH11*], t(8;21) [*RUNX1-ETO*] are called cytogenetically abnormal (CA-AML), while cases with genetic abnormalities (including frequent mutations in *DNMT3A*, *NPM1*, *CEBPα*, *IDH1/2*, *TET2*, *FLT3-ITD*) are called cytogenetically normal (CN-AML)^{1,4}. The former accounts for 50–55%, while the latter accounts for 45–50% of diagnosed AML cases^{6,7}. Even though these mutations and chromosomal alterations are crucial for initiating AML, they are not sufficient to explain AML progression, heterogeneity, and relapse⁸.

Recently, studies have identified the role of non-coding RNAs, especially long non-coding RNAs (lncRNA) in the initiation and progression of cancers^{9–11}. lncRNAs are emerging functional transcriptional products of at least 200 nucleotides lacking an open reading frame. Although they account for a large proportion of transcriptional products in mammals (about 58,000 loci)¹², only a small number of lncRNAs have been well characterized. Although lncRNAs are mostly not conserved evolutionarily, they are heavily regulated suggesting their functional role^{12,13}. They may function either as signal transducers, protein guides, or molecular scaffolds to regulate transcriptional and epigenetic events^{14–16}. Some lncRNAs perform these functions in *cis*- by modulating transcription of nearby genes (*Dum*)¹⁷ or act in *trans*-, by modulating genes at multiple distant loci (*MALAT1*)¹⁸, while some can do both (*HOTAIR*)^{9,19}.

Recent studies have identified lncRNAs for their remarkable role in regulating major epigenetic processes such as DNA methylation and chromatin remodeling. DNA methylation in mammals is coordinated by one of the three DNA methyltransferases (DNMT); *DNMT1*, *DNMT3A*, and *DNMT3B*^{7,17,20}. lncRNAs have been identified in recent studies as important agents that can modulate DNA methylation, either by activating or repressing DNMTs. For example, the lncRNA *Dum* was discovered to repress a nearby gene *Dppa2* by recruiting multiple DNMTs leading to methylation of a promoter region, thus promoting myoblast differentiation¹⁷. Conversely, the lncRNA *H19* represses the activity of *DNMT3B* by interacting with *SAHH* which hydrolyzes *SAH*, a step required for *DNMT3B* activation²¹. *ecCEBPα*, which is transcribed from the *CEBPα* locus, directly blocks *DNMT1* to prevent methylation of proximal and distal located promoters, thus promoting *CEBPα*-mediated

granulocyte differentiation²⁰. Mechanistic studies via reduced representation bisulfite sequencing and RNA immunoprecipitation sequencing shows that *ecCEBPα* suppresses DNA methylation in *cis*- by acting as a shield that sequesters *DNMT1* from the *CEBPα* promoter. We speculate that *ecCEBPα* could regulate *DNMT1* activities in distant DNA regions (*in trans*-) as well. The mechanism of this potential interaction is to be determined.

Methods

ecCEBPα sequence

In the recent version of GENCODE, *ecCEBPα* is not annotated most likely due to an overlap with a protein-coding gene *CEBPα* on the same strand. We retrieved the complete *ecCEBPα* sequence from the human genome (hg19, chr19: 33298573–33303358) based on information reported in the work of Di Ruscio *et al.*²⁰. *ecCEBPα* is approximately 4.8kb and it overlaps with the intronless *CEBPα* gene (~2.6kb) on the same strand. *ecCEBPα* does not share either the same transcription start site (TSS) or transcription end site (TES) with *CEBPα*, starting ~0.89kb upstream and ending ~1.46kb downstream of the *CEBPα* gene.

DNA methylation data processing

CpG methylation (Illumina 450K array) and *CEBPα* mutation data for 186 AML patients were retrieved from the Cancer Genome Atlas (TCGA: <http://firebrowse.org>). CpG methylation levels were measured in 307796 unique loci. We split all the AML patients into two groups based on *CEBPα* mutation status (13 patients with a *CEBPα* mutation and 173 patients without a mutation). We classified a CpG position as hypermethylated (HM) in patients with a *CEBPα* locus mutation if DNA methylation level was significantly increased in the case of a *CEBPα* mutation (t-test, FDR ≤ 0.05 and Δ-value ≥ 0.1) and all non-hypermethylated (NHM) CpG in the case of a *CEBPα* mutation were classified as non-hypermethylated CpGs (t-test, FDR > 0.05 and |Δ-value| < 0.1). As a result, we obtained 11955 HM and 261433 NHM CpGs.

Secondary structure and triplex prediction

As suggested in a previous study²², unpaired RNA nucleotides are more likely to form triplexes with DNA. We predicted RNA secondary structure using RNAplfold (V 2.4.14), from the Vienna suite using a cut-off for pairing probability (-c) of 0.1^{23,24}. To search for potential interactions between *ecCEBPα* and DNA target regions we used only unpaired nucleotides, while the nucleotides predicted to pair were replaced with 'N'.

DNA target regions were defined as 100 nucleotides centered at each CpG. To predict *ecCEBPα* triplex formation with DNA target regions, we used Triplexator (V 1.3.2)²⁵, since it has higher accuracy of prediction¹⁴, with the following optimization parameters suggested in 22: minimum length = 10 nucleotides, error rate = 20%, G-C content = 70%, and filter-repeats = off. Using these parameters out of 307796 unique CpG loci, we predicted 272131 loci with at least one triplex and 35715 without any. Among them, 10351 and 222105 potential triplex targets were predicted in the HM and NHM regions respectively.

To estimate the statistical significance of predicted triplexes we used Triplex domain finder (TDF v 0.12.3), which clusters RNA triplex-forming oligonucleotide (TFO) into DNA binding domains (DBD)²⁶. Briefly, all 272131 CpG loci with at least one predicted triplex were taken as input target regions. By predicting triplexes in the background regions, TDF is capable of estimating the statistical significance of *ecCEBPa* binding between target regions and other non-target CpGs regions. Since TDF allows only to mask regions in the genomic background rather than to select the background explicitly we had to prepare a special mask for the non-target regions. To do so we removed 35715 CpG loci with zero triplex predictions from the human genome using BEDtools subtract (BEDTools v2.29.2). TDF was implemented with a minimum triplex length (-l) of 10 nucleotides, an error rate (-e) of 20%, and (-f) to mask background loci in 100 random samplings (-n).

RNA:chromatin colocalization analysis

To validate the predicted interactions we used RNA:chromatin interactome obtained with iMARGI method capturing chromatin-associated RNA (caRNA) and their genomic interaction loci²⁷. The data was downloaded from GEO (GSM3478205). The iMARGI dataset was mapped to the hg38 genome assembly. We used UCSC Liftover to convert *ecCEBPa* sequence coordinates from hg19 to hg38 sequences²⁸. We expanded the DNA coordinates of CpGs by 3.0kb nucleotides upstream and downstream. IntersectBed from BEDTools was used to check the co-location of predicted triplexes and experimentally validated interactions of *ecCEBPa*²⁹. Fisher's exact test was calculated for the number of confirmed *ecCEBPa* interactions between TDF and iMARGI data.

GO enrichment analysis

Finally, since Illumina 450K array probes are located close to genes, we performed functional enrichment using BiNGO (v 3.0.3) (binomial test)³⁰ to infer the biological significance of the genes potentially affected by *ecCEBPa* binding.

All statistical analyses were performed using R 4.0 or SciPy v1.5.1 library. Visualization was done in Cytoscape 3.2.0³¹ and Python 3.7. Code is available at <https://zenodo.org/record/4385259>³².

Results

ecCEBPa forms triplexes with promoter regions and affects promoter methylation

The current study explores the potential of the lncRNA *ecCEBPa* in the modulation of global CpG methylation status in *trans* via direct interaction with DNA regions. *ecCEBPa* (extra coding *CEBPa*), reported in work by Di Russo *et al.*²⁰, is located on chromosome 19 and transcribed from the *CEBPa* locus (Figure 1a).

Mutations in *CEBPa* locus are a common feature of AML leading to whole genome hypermethylation (Figure 1b). Since TCGA is focused on protein-coding genes, all reported mutations are located within the *CEBPa* gene and could affect both *CEBPa* and *ecCEBPa*. To investigate if *ecCEBPa* could affect

DNA methylation in *trans*, first we checked if it is capable of interaction via forming triple helices (triplexes) with its binding sites and if such interactions affect DNA methylation. We observed that regions capable of forming triplexes with *ecCEBPa* remain protected from global DNA hypermethylation observed in case of a mutation in a *CEBPa* locus (Figure 1c, Fisher's exact test, p-value <0.001). This result suggests a negative relationship between DNMT access to promoter sites and *ecCEBPa* binding.

ecCEBPa binding is not affected by the mutation in the *CEBPa* locus

To investigate deeper the potential of *ecCEBPa* to form triplexes we used Triplex Domain Finder (TDF) - a triplex prediction tool that refines the resolution of predicted TFOs in RNA into DNA binding domains (DBD) and calculates the significance of the number of predicted triplexes for each DBD. Overall, 17 significant DBDs were identified within *ecCEBPa*, interspaced between sequences 4086 and 4968 towards the 3' end of the lncRNA (Figure 1d). These DBDs form triplexes with the majority of regions protected from hypermethylation in patients with a *CEBPa* mutation (Figure 1e, Extended data: Supplementary Table 1). The DBD region is located downstream from the *CEBPa* gene suggesting that *ecCEBPa* binding region is not affected by the mutation in the *CEBPa* locus. The predicted secondary structure of the *ecCEBPa* sequence showed that more than 95% of sequence positions from 4087-4987 (~0.5kb from *CEBPa* TES) (Figure 1d) were unpaired and potentially capable of forming triple helices with the target DNA region (Figure 2a).

ecCEBPa interacts with predicted binding sites

Since we use relatively relaxed thresholds for triplex prediction, we decided to validate the predicted RNA:DNA triplexes using experimental data obtained with the iMARGI method, allowing detection of RNA-chromatin interactions. We identified 157 *ecCEBPa* contacts within the iMARGI dataset and 29 of them contained predicted triplexes. Altogether, these 29 iMARGI interactions were made up of 182 predicted TTS (Fisher's exact test, p-value < 2.2E-16) located *in cis* and *in trans* on 14 chromosomes (Figure 2b). Chromosomes 19 (the native chromosome for *ecCEBPa*) are accounted for by all predictions.

Genes protected from methylation by *ecCEBPa* are involved in transcription factor activities

We performed gene ontology analysis on the genes located nearby 182 *ecCEBPa* triplexes supported by iMARGI. Key gene ontology categories such as nucleic acid binding and transcription factor activities were enriched among putative *ecCEBPa* targets (Figure 2c). Representative genes among transcription factors include *MLF2*, *SUV39H2*, *RBM5*, *UBTF*, and among sequence-specific DNA binding proteins include *POU2F2*, *MED12L*, and *DNASE1L1*. The enrichment in transcription factors (TF) suggests that triplex formation may represent a possible mechanism employed by *ecCEBPa* to regulate TF methylation and as a consequence, their expression.

Figure 1. (a) Schematic diagram for transcriptional products in the *CEBPA* locus; *CEBPA* is represented by an orange arrow and *ecCEBPA* is represented by the blue line. (b) Global change in DNA methylation (c) Difference in DNA methylation levels between patients with and without *CEBPA* mutation in the regions of *ecCEBPA* predicted binding (non parametric t-test, p-value >1E-10) (d) Number of DNA Triplex Target Site (y-axis) and a location of the corresponding TFO on the *ecCEBPA* (TFO: RNA; x-axis). (e) Number of Triplex Target sites per TFO predicted for NHM and HM CpG regions.

Discussion

Unlike other forms of cancers, AML progression is often mutation-independent but may be explained by altered epigenetic regulation, DNA methylation specifically^{1,8}. In this study, we elucidate a putative mechanism for the regulation of DNA methylation by *ecCEBPA*. In a previous study, *ecCEBPA*, which accompanies the transcription of *CEBPA* on the same locus, was shown to protect the promoter of *CEBPA* from DNA methylation leading to active expression²⁰. We speculated that *ecCEBPA* might perform a similar function in trans. We demonstrated that *ecCEBPA*-DNA triplex formation might provide the molecular basis of this interaction. *ecCEBPA* binding

presumably protects the region from genome-wide hypermethylation induced by *CEBPA* mutation in AML patients. *ecCEBPA* contains a TFO/DBD-rich region at its 3' end, with low pairing probability, suggesting that it is capable of triplex formation. Several of the predicted *ecCEBPA* binding sites, which include transcription factors such as MLF2, SUV39H2, RBM5, UBTF, and sequence-specific DNA binding proteins include *POU2F2*, *MED12L*, and *DNASE1L1*, were supported by experimental iMARGI RNA:chromatin interactions data.

Currently, the understanding of lncRNA function and mechanisms of action is limited to a few dozen well-annotated lncRNA

Figure 2. (a) Predicted secondary structure of *ecCEBPα*. Nucleotide color represents base pairing probabilities as predicted by RNAfold. (b) Experimentally validated *ecCEBPα* triplexes per chromosome. The x-axis represents the chromosome and y-axis represents the triplex potential relative to TTS. Blue points represent all TDF predictions that are present in the iMARGI dataset (c) Functional enrichment of genes located nearby CpG with predicted triplexes. (d) Schematic representation of *ecCEBPα*:DNA interactions in *trans* and its implication on DNA methylation. The presence of *ecCEBPα* inhibits DNA methylation process.

transcripts. A few functional characterization attempts are based on the ‘guilt by association’ hypothesis, which may not resonate well with the ability of lncRNA to interact in *trans*³³. As thoroughly reviewed previously, lncRNAs such as *ecCEBPα*, *Dum*, *Dali*, *Dacor1*, and *LINCRA-P21* interact with DNA in *trans* to regulate DNA methylation³⁴. The results presented herein further demonstrate that triplex formation between *ecCEBPα* and CpG containing DNA regions could indeed be regulatory and protect CpG sites from DNMT activity.

Unfortunately, RNA:chromatin interaction protocols are relatively new and the data is available only for a few cell types. Since RNA:chromatin interactions are highly cell-type specific³⁵ and lowly expressed, it is not surprising that we could validate only a few of the predicted interactions. Nevertheless, based on our results, we suggest a model of potential *ecCEBPα* chromatin interaction in *trans* (Figure 2d). In this model, *ecCEBPα* uses its unpaired regions to directly bind to specific DNA sequences by forming triplexes and in this way prevents

DNA methylation in the region of binding. *ecCEBPα* binding to distant regions could be mediated either by 3-dimensional chromatin organization¹⁷ which brings them close to *ecCEBPα*.

Recent studies have observed that promoter or transcription start sites (TSS) regions, which tend to be rich in CpG dinucleotides, are TTS-rich and potential triplex-forming hotspots^{36,37}. Through functional enrichment analysis, we observed that transcription factors might be preferential targets of *ecCEBPα*. Interestingly, previous studies have shown that the suppression of a myeloid leukemia factor (*MLF2*), an oncogene in breast cancer and myeloid leukemia^{38,39} as well as UBTF which controls rDNA expression^{40,41} contributes significantly to cancers upon promoter hypermethylation^{40,42}. The suppressor of variegation 3-9 homolog 2 (*SUV39H2*), a histone-lysine-N-methyltransferase which regulates the hypermethylation H3K9 has also been reported to indirectly influence over 450 promoters in AML⁴³. Having in mind that *ecCEBPα* is transcribed from *CEBPα* locus - a key transcription factor of hematopoiesis - this lncRNA

could participate in the formation of a hub in the hematopoiesis regulatory network.

Conclusion

In conclusion, we have shown that ecCEBP α could serve as a trans-acting regulatory agent protecting its binding sites from genome-wide CpG methylation, and its dysregulation could contribute to aberrant methylation profile in AML patients. These results suggest a novel regulatory mechanism for ecCEBP α as a modulator of DNA methylation through triplex formation providing a foundation for sequence-specific engineering of RNA for regulating methylation of specific genes.

Data availability

Underlying data

Complete ecCEBP α sequence retrieved from the human genome (hg19, chr19: 33298573-33303358): https://www.ncbi.nlm.nih.gov/assembly/GCF_000001405.13/

CEBP α mutation data for 186 AML patients retrieved from the Cancer Genome Atlas (TCGA): <http://firebrowse.org>.

GEO: Embryonic kidney that expresses SV40 large T antigen, Accession number GSM3478205: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM3478205>

Zenodo: josoga2/eccebp-alpha-project: F1000 Code Release, <https://doi.org/10.5281/zenodo.4385259>³²

This project contains the following underlying data:

- DNA_BINDING_DOMAINS_ID.tsv
- predicted_secondary_structure_of_ecCEBP α .fa
- probes.csv (main data)

Code used for analysis available from: <https://github.com/josoga2/eccebp-alpha-project/tree/f1000>

Archived code as at time of publication: <https://doi.org/10.5281/zenodo.4385259>³²

Extended data

Zenodo: Supplementary Data for Secondary structure and DNA binding domain prediction, <https://doi.org/10.5281/zenodo.4433222>⁴⁴.

This project contains the following extended data:

- Supplementary Table 1: Summary table of DNA binding domains (DBD), the counts of target regions within the genome and statistical analysis.
- ecCEBP α secondary structure prediction with RNAplfold

Data and code are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

1. Goldman SL, Hassan C, Khunte M, et al.: **Epigenetic Modifications in Acute Myeloid Leukemia: Prognosis, Treatment, and Heterogeneity.** *Front Genet.* 2019; **10**: 133. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Gambacorta V, Gnani D, Vago L, et al.: **Epigenetic Therapies for Acute Myeloid Leukemia and Their Immune-Related Effects.** *Front Cell Dev Biol.* 2019; **7**: 207. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. van Oosterwijk JG, Buelow DR, Drenberg CD, et al.: **Hypoxia-induced upregulation of BMX kinase mediates therapeutic resistance in acute myeloid leukemia.** *J Clin Invest.* 2018; **128**(1): 369–380. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Zhang J, Gu Y, Chen B: **Mechanisms of drug resistance in acute myeloid leukemia.** *Onco Targets Ther.* 2019; **12**: 1937–1945. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Zueva I, Dias J, Lushchekina S, et al.: **New evidence for dual binding site inhibitors of acetylcholinesterase as improved drugs for treatment of Alzheimer's disease.** *Neuropharmacology.* 2019; **155**: 131–141. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Hackl H, Astanina K, Wieser R: **Molecular and genetic alterations associated with therapy resistance and relapse of acute myeloid leukemia.** *J Hematol Oncol.* 2017; **10**(1): 51. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. O'Brien EC, Prideaux S, Chevassut T: **The epigenetic landscape of acute myeloid leukemia.** *Adv Hematol.* 2014; **2014**: 103175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Mehdi pour P, Santoro F, Minucci S: **Epigenetic alterations in acute myeloid leukemias.** *FEBS J.* 2015; **282**(9): 1786–1800. [PubMed Abstract](#) | [Publisher Full Text](#)
9. Hajjari M, Salavaty A: **HOTAIR: an oncogenic long non-coding RNA in different cancers.** *Cancer Biol Med.* 2015; **12**(1): 1–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Gao Y, Wang JW, Ren JY, et al.: **Long noncoding RNAs in gastric cancer: From molecular dissection to clinical application.** *World J Gastroenterol.* 2020; **26**(24): 3401–3412. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Zhang T, Hu H, Yan G, et al.: **Long Non-Coding RNA and Breast Cancer.** *Technol Cancer Res Treat.* 2019; **18**: 1533033819843889. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Hon CC, Ramilowski JA, Harshbarger J, et al.: **An atlas of human long non-coding RNAs with accurate 5' ends.** *Nature.* 2017; **543**(7644): 199–204. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Alam T, Medvedeva YA, Jia H, et al.: **Promoter Analysis Reveals Globally Differential Regulation of Human Long Non-Coding RNA and Protein-Coding Genes.** *PLoS One.* 2014; **9**(10): e109443. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Antonov IV, Mazurov E, Borodovsky M, et al.: **Prediction of lncRNAs and their interactions with nucleic acids: benchmarking bioinformatics tools.** *Brief Bioinform.* 2019; **20**(2): 551–564. [PubMed Abstract](#) | [Publisher Full Text](#)
15. Cruz-Miranda GM, Hidalgo-Miranda A, Bárcenas-López DA, et al.: **Long Non-Coding RNA and Acute Leukemia.** *Int J Mol Sci.* 2019; **20**(3): 735. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Marchese FP, Raimondi I, Huarte M: **The multidimensional mechanisms of long noncoding RNA function.** *Genome Biol.* 2017; **18**(1): 206. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Wang L, Zhao Y, Bao X, et al.: **lncRNA Dum interacts with Dnmts to regulate Dppa2 expression during myogenic differentiation and muscle regeneration.** *Cell Res.* 2015; **25**(3): 335–350. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. He B, Peng F, Li W, et al.: **Interaction of lncRNA-MALAT1 and miR-124 regulates HBx-induced cancer stem cell properties in HepG2 through PI3K/Akt signaling.** *J Cell Biochem.* 2019; **120**(3): 2908–2918. [PubMed Abstract](#) | [Publisher Full Text](#)
19. Botti G, Scognamiglio G, Aquino G, et al.: **lncRNA HOTAIR in Tumor**

- Microenvironment: What Role?** *Int J Mol Sci.* 2019; **20**(9): 2279.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Di Ruscio A, Ebralidze AK, Benoukrat T, et al.: **DNMT1-interacting RNAs block gene-specific DNA methylation.** *Nature.* 2013; **503**(7476): 371–376.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Zhou J, Yang L, Zhong T, et al.: **H19 lncRNA alters DNA methylation genome wide by regulating S-adenosylhomocysteine hydrolase.** *Nat Commun.* 2015; **6**: 10221.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Matveishina E, Antonov I, Medvedeva YA: **Practical Guidance in Genome-Wide RNA:DNA Triple Helix Prediction.** *Int J Mol Sci.* 2020; **21**(3): 830.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Lorenz R, Bernhart SH, Siederdisen CHZ, et al.: **ViennaRNA Package 2.0.** *Algorithms Mol Biol.* 2011; **6**: 26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Bernhart SH, Hofacker IL, Stadler PF: **Local RNA base pairing probabilities in large sequences.** *Bioinformatics.* 2006; **22**(5): 614–615.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. Buske FA, Bauer DC, Mattick JS, et al.: **Triplexator: detecting nucleic acid triple helices in genomic and transcriptomic data.** *Genome Res.* 2012; **22**(7): 1372–1381.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Kuo CC, Hänzelmann S, Cetin NS, et al.: **Detection of RNA-DNA binding sites in long noncoding RNAs.** *Nucleic Acids Res.* 2019; **47**(6): e32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Wu W, Yan Z, Nguyen TC, et al.: **Mapping RNA-chromatin interactions by sequencing with iMARGI.** *Nat Protoc.* 2019; **14**(11): 3243–3272.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Kuhn RM, Haussler D, Kent WJ: **The UCSC genome browser and associated tools.** *Brief Bioinform.* 2013; **14**(2): 144–161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.** *Bioinformatics.* 2010; **26**(6): 841–842.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Maere S, Heymans K, Kuiper M: **BINGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks.** *Bioinformatics.* 2005; **21**(16): 3448–3449.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Shannon P, Markiel A, Ozier O, et al.: **Cytoscape: a software environment for integrated models of biomolecular interaction networks.** *Genome Res.* 2003; **13**(11): 2498–2504.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Ogunleye AJ: **josoga2/cecbp-alpha-project: F1000 Code Release (Version f1000).** *Zenodo.* 2020.
<http://www.doi.org/10.5281/zenodo.4385259>
 33. Jalali S, Singh A, Maiti S, et al.: **Genome-wide computational analysis of potential long noncoding RNA mediated DNA:DNA:RNA triplexes in the human genome.** *J Transl Med.* 2017; **15**(1): 186.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Zhao Y, Sun H, Wang H: **Long noncoding RNAs in DNA methylation: new players stepping into the old game.** *Cell Biosci.* 2016; **6**: 45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Bonetti A, Agostini F, Suzuki AM, et al.: **RADICL-seq identifies general and cell type-specific principles of genome-wide RNA-chromatin interactions.** *Nat Commun.* 2020; **11**(1): 1018.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Sridhar B, Rivas-Astroza M, Nguyen TC, et al.: **Systematic Mapping of RNA-Chromatin Interactions In Vivo.** *Curr Biol.* 2017; **27**(4): 602–609.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Antonov I, Medvedeva YA: **Purine-rich low complexity regions are potential RNA binding hubs in the human genome [version 2; peer review: 3 approved].** *F1000Res.* 2018; **7**: 76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Dave B, Granados-Principal S, Zhu R, et al.: **Targeting RPL39 and MLF2 reduces tumor initiation and metastasis in breast cancer by inhibiting nitric oxide synthase signaling.** *Proc Natl Acad Sci U S A.* 2014; **111**(24): 8838–8843.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Gobert V, Haenlin M, Waltzer L: **Myeloid leukemia factor: a return ticket from human leukemia to fly hematopoiesis.** *Transcription.* 2012; **3**(5): 250–254.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Nguyen LXT, Raval A, Garcia JS, et al.: **Regulation of ribosomal gene expression in cancer.** *J Cell Physiol.* 2015; **230**(6): 1181–1188.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Santoro R, Grummt I: **Molecular mechanisms mediating methylation-dependent silencing of ribosomal gene transcription.** *Mol Cell.* 2001; **8**(3): 719–725.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Leshchenko VV, Kuo PY, Shakhovich R, et al.: **Genomewide DNA methylation analysis reveals novel targets for drug development in mantle cell lymphoma.** *Blood.* 2010; **116**(7): 1025–1034.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Monaghan L, Massett ME, Bunschoten RP, et al.: **The Emerging Role of H3K9me3 as a Potential Therapeutic Target in Acute Myeloid Leukemia.** *Front Oncol.* 2019; **9**: 705.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Ogunleye AJ: **Supplementary Data for Secondary structure and DNA binding domain prediction [Data set].** *Zenodo.* 2021.
<http://www.doi.org/10.5281/zenodo.4433222>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 10 June 2021

<https://doi.org/10.5256/f1000research.31133.r81320>

© 2021 Demidov O. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oleg N. Demidov

INSERM U1231, Faculty of Medicine and Pharmacy, University of Bourgogne-Franche Comté, Dijon, France

The manuscript by Ogunleye AF and co-authors describes a novel phenomenon of lncRNA ecCEBP α -dependent regulation of methylation in several genes. The authors analyzed the open datasets and showed that mutations in CEBPA gene (coding both, CEBPA transcriptional factor and lncRNA ecCEBP α) changed the global CpG methylation status in AML cells. The manuscript is nicely written, conclusions are logically justified. This work will be interesting to the readers in the epigenetic regulation and cancer research field.

Still, I have several concerns about the analysis. There are a lot of predicted triplex target regions for ecCEBP α genome-wide. Some of them could be false-positive predictions. Is there any way to confirm the observed effect only of experimentally validated contacts? Or in regions where predictions are most reliable?

Also, it is unclear other genes, not only transcription factors, predicted to be regulated by ecCEBP α are related to hematopoiesis. It would be helpful to have at least some hypotheses in the discussion. If there is some link found between ecCEBP α targets and regulation of hematopoiesis, it would really strengthen the conclusions of the work. Probably, KEGG pathway enrichment analysis may help with that.

Minor comments:

It is unclear on the figure with a secondary structure of ecCEBP α where the triplex-forming region is located.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Models of human diseases, cancer research, fibrosis, paging

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 05 Aug 2021

ADEWALE OGUNLEYE

Response to Reviewer Two

Comment one: There are a lot of predicted triplex target regions for ecCEBPA genome-wide. Some of them could be false-positive predictions. Is there any way to confirm the observed effect only of experimentally validated contacts? Or in regions where predictions are most reliable?

Response one: We are grateful to the reviewer for this suggestion. We performed a validation for the most reliable predicted triplexes using only experimentally validated ecCEBPA:DNA contacts. Overall, a mean methylation delta score of only 0.01 (p-value < 1E-09) was observed which further emphasizes the protective effect of ecCEBPA.

The following text and images were added to the main body of the paper:

"Since these ecCEBPA contacts have been experimentally validated, we suggested that they are the most reliable regions where ecCEBPA might have a protective effect from DNA methylation. Overall, a mean methylation delta score of experimentally validated binding sites is significantly lower than that of the non-binding sites (p-value < 1E-09) (Figure 2C)."

New figure panels (1&2) here: <https://drive.google.com/file/d/1iynm-t0DfGkxLF5vsumG4iCcfo2u2xCJ/view?usp=sharing>

Comment two: Also, it is unclear if other genes, not only transcription factors, predicted to be regulated by ecCEBPA are related to hematopoiesis. It would be helpful to have at least some hypotheses in the discussion. If there is some link found between ecCEBPA targets and regulation of hematopoiesis, it would really strengthen the conclusions of the work.

Probably, KEGG pathway enrichment analysis may help with that.

Response two: We thank the reviewer for this important comment. Indeed, we discovered that other non-transcription factor genes have links to acute myeloid leukemia via literature mining. Unfortunately, this could not be enriched through KEGG. We suggest including a supplementary table that details the different roles of the identified genes.

In the main text, we added:

"Out of the 33 genes we identified to be targets of ecCEBPA, 16 genes (ICAM1, PDXK, etc.) are clearly related to hematopoiesis and various leukemias. A summary of the genes and their function is provided in Table 1."

Table 1: AML specific function of nearby genes that are protected by ecCEBPA (added to the paper).

Competing Interests: No competing interests were disclosed.

Reviewer Report 15 April 2021

<https://doi.org/10.5256/f1000research.31133.r81314>

© 2021 Singh A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Amrita Singh

CSIR Institute of Genomics and Integrative Biology (CSIR-IGIB), New Delhi, Delhi, India

The article has addressed an important question of the ability of a lncRNA to act in trans and regulate a multitude of genes, which might be affecting the progression of acute myeloid leukemia. The authors have also highlighted the possibility of a triplex structure for defining the binding of lncRNA at its target gene CpG region. However, there are certain parts wherein the authors haven't been able to convey their points properly.

1. In Figure 1(b), a global DNA hypermethylation has been represented in AML patients with CEBPα mutation, but the numbers of the HM and NHM sites in the methods section do not reflect the same. It shows higher sites for NHM, than the HM sites, kindly check if some mislabeling is there at the authors part.
2. Also, the reviewer has failed to understand Figure 1(c). In both the x-axis and the side legend, binding and non-binding have been depicted, but which one is the AML patients with and without mutation isn't clear from the figure.
3. There are few other key points that were not clear;
(a) what is the level of ecCEBPα expression in AML patients with and without mutation.

(b) Additionally, if the expression of the lncRNA remains the same in both cases, how does one explain the increase in binding of the lncRNA and subsequent higher NHM sites, in the case of AML patients with CEBPα mutation?

(c) Moreover, a previous report (Di Ruscio *et al.*¹) on ecCEBPα had also analyzed genome-wide methylation, in which they report that methylation levels remain unchanged even when ecCEBPα was overexpressed. This is in contrast with the major theme of the paper i.e., methylation of genes in trans is affected by the ecCEBPα. The authors should comment on this in the discussion part.

References

1. Di Ruscio A, Ebralidze AK, Benoukraf T, Amabile G, et al.: DNMT1-interacting RNAs block gene-specific DNA methylation. *Nature*. 2013; **503** (7476): 371-6 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Structure function relationship of noncoding RNA, with major focus on lncRNA function via triple helical structures.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Aug 2021

ADEWALE OGUNLEYE

Response to Reviewer One

Comment One: In Figure 1(b), a global DNA hypermethylation has been represented in AML patients with CEBPα mutation, but the numbers of the HM and NHM sites in the methods section do not reflect the same. It shows higher sites for NHM, than the HM sites, kindly check if some mislabeling is there at the authors part.

Response Two: We believe that there is a confusion here. The plot was correct and reflected the change in DNA methylation in all CpG positions including those that were not statistically differentially methylated. We counted hypermethylated CpG (t-test, $FDR \leq 0.05$ and Δ -value ≥ 0.1) and those that were not hypermethylated (t-test, $FDR > 0.05$ and $|\Delta$ -value| < 0.1). In this way, we refer to non-hypermethylated (NHM) to any CpG with DNA methylation change less than 0.1, which includes either not changing or hypomethylated and not only hypomethylated. To avoid this confusion we suggest replacing the plot with the one below. We specified a 0.1 mid-point to aid visualization. (Figure 1b).

Comment Two: Also, the reviewer has failed to understand Figure 1(c). In both the x-axis and the side legend, binding and non-binding have been depicted, but which one is the AML patients with and without mutation isn't clear from the figure.

Response Two: Each violin in Figure 1(c) is a representation of the differential methylation score between CpGs of AML patients with and without CEBPα mutation. The blue violin represents all CpGs that are located in ecCEBPα binding sites, while the orange violin represents CpGs that do not bind with ecCEBPα. To provide clarity, we corrected the labels on the X and Y axes and added explicit explanation in the legend of the figure (Figure 1c). Difference in methylation = Mean methylation of CpGs in AML patients with CEBPα mutation - Mean methylation of CpGs in AML patients without CEBPα mutation.

Comment 3: There are few other key points that were not clear;

- (a) what is the level of ecCEBPα expression in AML patients with and without mutation.
- (b) Additionally, if the expression of the lncRNA remains the same in both cases, how does one explain the increase in binding of the lncRNA and subsequent higher NHM sites, in the case of AML patients with CEBPα mutation?
- (c) Moreover, a previous report (Di Ruscio et.al1) on ecCEBPα had also analyzed genome-wide methylation, in which they report that methylation levels remain unchanged even when ecCEBPα was overexpressed. This is in contrast with the major theme of the paper i.e., methylation of genes in trans is affected by the ecCEBPα. The authors should comment on this in the discussion part.

Response:

(a&b): It is relatively difficult to estimate the expression level of ecCEBPα since this gene is not in the annotation used by TCGA. The raw data is not freely available in TCGA. ecCEBPα gene also overlaps with the CEBPα gene making the estimation of expression of each of the genes even more complicated. Since the mutation in AML patients happens in the body of CEBPα gene, we do not expect a significant change in the expression of either CEBPα or ecCEBPα.

We explained the relationship between ecCEBPα binding and methylation in the first result

section (Figures 1c,d&e). We suggested that the mutation(s) in CEBPA does not affect the expression of the lncRNA, but rather the ability to bind its targets.

We further emphasize this point in the paper with this line (discussion):

"Di Ruscio et al. confirmed that most mutations in CEBPa do not influence the expression of ecCEBPa but rather, its ability of the RNA to fold properly. We suggest that the inability to fold properly even in overexpressed cases may affect its structural ability to bind its targets."

(c): We are very grateful to the reviewer for this valuable comment. In response to this, we retrieved the DNA methylation data (RRBS) for wild-type HL-60 cells ([ENCODE Dataset](#)) and overexpressed HL-60 cells ([Di Ruscio et al.](#)). We then calculated the methylation difference between the two cell states and compared it between the ecCEBPA binding sites versus the non-binding sites. Non ecCEBPA binding sites were more methylated in comparison to ecCEBPA binding sites (Fisher Test: $p\text{-val} = 1.24\text{E-}09$). Bearing in mind that ecCEBPA targets are located genome-wide, it is tempting to suggest that "enforced overexpression" (which was achieved with the R1 variant of ecCEBPA; comprise of downstream ecCEBPA sequences) of ecCEBPA strongly protects local CpGs from methylation while the rest of the genome gain some methylation.

We added the following statement to the results in the main text:

"Furthermore, the overexpression of ecCEBPA in HL-60 cells (Wang et al) lead to ecCEBPA binding sites stay unmethylated while non ecCEBPA binding sites gain methylation (Fisher Test: $p\text{-val} = 1.24\text{E-}09$) , suggesting a protective effect on ecCEBPA's binding to its target locations (Fig 1d)."

Competing Interests: No competing interests were disclosed.

Comments on this article

Version 1

Reader Comment 14 Apr 2021

Olumide Inyang, moscow institute of physics and technology (MIPT), Russian Federation

This report gave a clear understanding of ecCEBPa as a genome-wide epigenetic modulator via triple-helix stacks for RNA(engineered) in regulating methylation of specific genes...great work from the authors.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research