RESEARCH ARTICLE

REVISED Spatial distribution analysis of dentists, dental technicians, and dental therapists in Indonesia [version 2; peer review: 2 approved]

Nanda Rachmad Putra Gofur ¹, Achmad Zam Zam Aghasy²,
Aisyah Rachmadani Putri Gofur³

¹Department of Health, Faculty of Vocational Studies, Universitas Airlangga, Surabaya, Indonesia

²Indonesian Health Collaboration and Innovation Institute, Surabaya, Indonesia

³Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

v2 First published: 18 Mar 2021, 10:220
<https://doi.org/10.12688/f1000research.50869.1>
Latest published: 10 Jun 2021, 10:220
<https://doi.org/10.12688/f1000research.50869.2>

Abstract

Background: Access to health services is needed around the world, from healthcare providers to doctors. One of the needs in public health is a system that is accessible for everyone, but, unequal distribution of healthcare provider and health workers, especially in dentistry fields is still a main problem in several countries, including Indonesia. The aim of this study is to analyze the spatial distribution of dentists, dental technicians, and dental therapists.

Methods: This spatial analysis study was conducted after obtaining secondary data in Indonesia. All data were collected between September 1st, 2020 and October 1st, 2020 from open access sources of de-identified data. The data of dentists per area, dental technicians per area, and dental therapists per area were calculated for analysis. A spatial distribution map was prepared using the Quantum Geographic Information System (QGIS Desktop, version 3.10.6).

Results: The results of this study found a ratio of dentists to members of the population in Indonesia of 1:17,105. The average number of dental technicians that work in the public health centers in each province (dental technicians per area) in Indonesia was calculated to be 0.13. The average number of dental therapists that work in the public health centers in each province (dental therapists per area) in Indonesia was calculated to be 0.40. This spatial autocorrelation illustrates that there is a relationship between values of dentists per area and dental therapists per area between provinces in Indonesia, and shows geographic clustering relationships or patterns that are grouped and have similar characteristics in adjacent locations. This spatial autocorrelation did not occur in the value of dental technicians.

Conclusions: From this study we can conclude that there is an unequal distribution of dental personnel in Indonesia.

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
10 Jun 2021		view
version 1		
18 Mar 2021	view	view

1. **Reny Purnama Hadi** , Brawijaya
University, Malang, Indonesia

2. **Alessandro Leite Cavalcanti** , UEPB -
Universidade Estadual da Paraíba, Campina
Grande, Brazil

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

Healthcare, Public health, Health service, Health worker density

Corresponding author: Nanda Rachmad Putra Gofur (nanda.gofur@gmail.com)

Author roles: **Gofur NRP:** Conceptualization, Formal Analysis, Funding Acquisition, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Aghasy AZZ:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Resources, Software, Visualization; **Gofur ARP:** Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Gofur NRP *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Gofur NRP, Aghasy AZZ and Gofur ARP. **Spatial distribution analysis of dentists, dental technicians, and dental therapists in Indonesia [version 2; peer review: 2 approved]** F1000Research 2021, 10:220 <https://doi.org/10.12688/f1000research.50869.2>

First published: 18 Mar 2021, 10:220 <https://doi.org/10.12688/f1000research.50869.1>

REVISED Amendments from Version 1

There is no differences in the abstract of this new version article and previously published. We only change the keywords of the article.

Any further responses from the reviewers can be found at the end of the article

Introduction

Indonesia is a country in Southeast Asia located between Indian and Pacific oceans. It has more than 17,000 islands, but the biggest islands are Sumatra, Java, Borneo or Kalimantan, Sulawesi and Papua. Indonesia is the 14th largest country in the world with a land area of 1,904,569 square kilometers or 735,358 square miles and is the world's largest island country. Indonesia is bordered by several country such as Malaysia, East Timor, Papua New Guinea, Singapore, Vietnam, Philippines, Palau and Australia¹.

Based on the Global Burden of Disease (GBD) study in 2016², dental and oral health problems, especially dental caries, affects nearly half of the world's population (3.58 billion people). Gum disease (periodontal) is the 11th most common oral health disease in the world. Meanwhile, in Asia Pacific, oral cancer is the third most common type of cancer. The results of the Basic Health Research (Riskesdas) in 2018 stated that 57.6% of the Indonesian population had dental and oral problems during the last 12 months, but only 10.2% received treatment by dental medical personnel (dental nurses, dentists or specialist dentists), while the rest received no treatment^{2,3}. The largest proportion of dental problems in Indonesia are dental caries, necrosis tooth, and toothache complaints (45.3%). Meanwhile, the majority of oral health problems experienced by Indonesians are swollen gums (14%)^{3,4}.

Effective Medical Demand (EMD) is defined as the percentage of the population who had oral health problems in the last 12 months multiplied by the percentage of the population who received dental care or treatment from dental medical personnel (specialist, dentists, dental nurses/dental therapists) and dental technicians⁵. The EMD in Indonesia is only 8.1%, it means that only few of the population receive dental treatments when they have dental problems. The overall ability to get services from dental medical personnel (EMD) in Indonesia is only 8.1%^{5,6}. Three provinces of Indonesia in 2013, as South Sulawesi, South Kalimantan and Central Sulawesi had quite high levels of oral and dental problems (>35%), with EMD of 10.3%, 8.0%, and 6.4%, respectively⁷. Based on this finding, a clear picture of the gap between dental problems and the EMD that occur in society can be seen.

A public health system is needed, whereby everyone can access public health services, including dental services, at an affordable rate including affordability of health services. Private health-care providers might help affordability of health services beside government health services. The main focus which needs attention is on curative care in single-practitioner and

group practices in dentistry fields. But, unfortunately, these private health services are driven by market demands in society and are business-oriented services. The Geographic Information System (GIS) is a method commonly used in health research. This method could analyze various healthcare variables related to other fields such as physical, social and cultural environments^{8,9}. Access to health services is very important and needed around the world. The correlation between the service provider (the health practitioner) and the consumer (the patient) is still a main problem in many other countries including Indonesia^{10,11}.

As of 2012, from 8,975 public health centers, there are 5,439 public health centers with dentists and 3,536 public health center without dentists. Nationally, 47.4% of public health centers had one dentist and 13.2% of public health centers had more than two dentists^{12,13}. Public health centers that have the highest efforts in dental and oral health service is in the Bali province, which is 100% and the lowest efforts in dental and oral health service in Indonesia is in Papua Province, which is 24%; the national rate for this efforts of dental and oral health services is 84%. This means, Bali is the province that is best at providing dental services and appropriately considers dentistry fields, meanwhile Papua is the worst^{14,15}.

The residents ease to reach services and facilities based on the distance and travel time to a resource is called geographic accessibility. Optimal delivery of dental health services should take into consideration availability and accessibility, which together are referred to as spatial accessibility¹⁶⁻¹⁸. The aim of this study was to analyze the spatial distribution of dentists, dental technicians, and dental therapists in Indonesia.

Methods

This spatial analysis study was conducted after obtaining secondary data in Indonesia. All data were collected between 1st September 2020 and 1st October 2020 from open access sources of de-identified data; therefore, no ethical approval was required. We sought to identify the average number of dentists, dental technicians and dental therapists working in public sectors and the main building of public health centers in each province.

This study retrieved data of the number of dentists in each province in 2018 from the national Indonesian Health Facility Research report (Riset Fasilitas Kesehatan), accessed via the [Ministry of Health](#), Indonesia.

The geographic administrative area data of Indonesia were retrieved from the Indonesia [Geospatial Portal](#), which is publicly available. The provincial maps of the Indonesian database were processed with the Geospatial Portal for further geographic information system (GIS) based analysis to identify spatial distribution of dentists, dental technicians, and dental therapists per area by obtaining the province level polygon map that contains information regarding latitudes and longitudes of each province. We used the map of all 34 provinces of Indonesia for the analysis. Data on the distribution of dentists,

dental technicians, and dental therapists was collected from the Ministry of Health and Geospatial Portal.

We created a map using the **Quantum Geographic Information System** (QGIS Desktop, version 3.10.6). Global and local Moran's indices were calculated to determine the autocorrelation value and local indicator of spatial autocorrelation (LISA). LISA was analyzed using **GeoDa** software, version 1.10.0.8. The level of significance was set at p-value of <0.05, $Z_{w/2}$ at 1.96, and randomization run to 999 permutations. We used automatic Euclidean weight distance, which matched with the assumption that each province has at least four neighboring provinces. Interpretation of the LISA significance map includes the following categories:

- “high–high” indicates a clustering of high value rates (positive spatial autocorrelation)
- “low–high” indicates that the low value rates are adjacent to high value rates (negative spatial autocorrelation)
- “low–low” indicates clustering of low value rates (positive spatial autocorrelation)
- “high–low” indicates that high value rates are adjacent to low value rates (negative spatial autocorrelation)
- “not significant” indicates that there is no spatial autocorrelation.

The outcome of Moran's I identifies the intensity of spatial autocorrelation along with the result of statistically significant test, that is, the p value. The following mathematical representation exhibits the computation of Moran's I:

$$I = \frac{N}{S_0} \frac{\sum_{i=1}^N \sum_{j=1}^N W_{ij} (x_i - \bar{x})(x_j - \bar{x})}{\sum_{i=1}^N (x_i - \bar{x})(x_i - \bar{x})^2}$$

$$S_0 = \sum_{i=1}^N \sum_{j=1}^N W_{ij}$$

Where W_{ij} is the spatial weight between the parameters in provinces i and j . The parameters are the number of dentists per area, dental technicians per area, and dental therapists per area. N is the total number of spatial units; S_0 is the aggregate of all spatial weights; x_i and x_j are the amount of each dental health personnel in provinces i and j , respectively.

Results and discussion

The total number of public health centers in Indonesia were counted at 9,831. The total number of dentists, dental technicians, and dental therapists that work in public sectors of public health center main buildings in each province was 15,833, 1,214, and 3,834, respectively (Table 1).

Figure 1 shows the distribution of dental health personnel in several major islands in Indonesia. The distribution is still unequal due to the big difference in the amount of dental personnel in each island. The average number of dentists that

work in public sectors and public health centers in each province (dentist per area) in Indonesia was calculated at 1.61 (Figure 2). The ratio of dentist to members of the population in Indonesia was calculated at 1:17,105. The average number of dental therapists that work in main buildings of public health centers in each province (dental therapist per area) in Indonesia was calculated at 0.40 (Figure 3). The average number of dental technicians that work in main buildings of public health centers in each province, (dental technicians per area) in Indonesia were calculated at 0.13 (Figure 4).

There are 13 provinces that have a value of dentists per area below 1, which indicates that there are areas that do not have dentists at all in public health centers. All provinces in Indonesia have a value of dental technicians per area below 1, which shows that all regions of Indonesia do not have sufficient numbers of dental technicians for each public health center area. There are 31 provinces that have a value of dental therapists per area below 1, which indicates that there are areas that do not have dental therapists at all in a health center area (Figure 1–Figure 4).

The highest number of dentists per distribution area is in the province of Bali, Java-Bali, and the lowest is in the province of West Papua, Papua. The highest number of dental technicians per distribution area is in the province of Bangka Belitung, Sumatra, and the lowest is in the province of North Maluku, Maluku. The highest number of dental therapists per distribution area is in the province of Bali, Java-Bali, and the lowest is in the province of Papua. Visualization maps of dentists, dental technicians, and dental therapist per area were developed and are presented in Figure 5.

Indonesia had both a shortage of and an unequal distribution of dental health personnel in public health center areas. The distribution of dentists is similar to that in other studies, which have indicated maldistribution of both general practitioners (GPs) and specialists¹⁹. Maldistribution indicates inequality associated with area. Dentist and dental nurses are two health workers who play an important role in providing dental health services. The distribution of the two health workers was the highest in the Java-Bali region (60.1%), and the lowest in Papua region (10.3%). However, there are a high number of dentists and dental therapists in the Maluku Islands region (57.9%)^{11,12}. Based on this, it appears that there is a gap that occurs between regions related to the availability of dentists and dental therapists on duty at the health centers. These results are also supported by a previous study that found an unequal distribution^{19–21}.

The poor distribution of health workers is not only present in dental health service personnel, but in other types of health workers too. Although there is a health worker placement policy in Indonesia which uses a temporary employee system for medical personnel (doctors and dentists), the distribution is not equal especially in remote areas. From the results of the analysis conducted by Ihsan Husain, *et al.* (2006), on average there are more dentists in Java-Bali compared to the other regions^{10,15}.

Table 1. Distribution of dental personnel in each province in Indonesia.

Province	Total Public health center (n)	Dentist		Dental Technician		Dental Therapist	
		Total	Average of each public health center work area	Total	Average of each public health center work area	Total	Average of each public health center work area
Aceh	347	414	1,19	92	0,27	145	0,42
North Sumatera	571	900	1,58	64	0,11	60	0,11
West Sumatera	271	544	2,01	38	0,14	139	0,51
Riau	216	555	2,57	25	0,12	38	0,18
Jambi	193	287	1,49	29	0,15	100	0,52
South Sumatera	328	362	1,10	36	0,11	259	0,79
Bengkulu	179	129	0,72	15	0,08	7	0,04
Lampung	299	238	0,80	27	0,09	87	0,29
Bangka Belitung	63	113	1,79	24	0,38	50	0,79
Kepulauan Riau	80	138	1,73	4	0,05	31	0,39
DKI Jakarta	313	1416	4,52	15	0,05	115	0,37
West Java	1069	1915	1,79	168	0,16	451	0,42
Central Java	876	1517	1,73	89	0,10	428	0,49
DI Yogyakarta	121	390	3,22	28	0,23	132	1,09
East Java	964	2318	2,40	112	0,12	324	0,34
Banten	233	535	2,30	23	0,10	24	0,10
Bali	120	640	5,33	13	0,11	206	1,72
West Nusa Tenggara	161	198	1,23	49	0,30	103	0,64
East Nusa Tenggara	374	210	0,56	27	0,07	252	0,67
West Kalimantan	241	197	0,82	39	0,16	240	1,00
Central Kalimantan	197	147	0,75	38	0,19	81	0,41
South Kalimantan	232	264	1,14	49	0,21	118	0,51
East Kalimantan	178	341	1,92	8	0,04	51	0,29
North Kalimantan	55	85	1,55	9	0,16	11	0,20
North Sulawesi	193	128	0,66	20	0,10	82	0,42
Central Sulawesi	196	171	0,87	31	0,16	21	0,11
South Sulawesi	452	844	1,87	57	0,13	203	0,45
Southeast Sulawesi	281	219	0,78	41	0,15	31	0,11
Gorontalo	93	89	0,96	5	0,05	4	0,04
West Sulawesi	94	108	1,15	12	0,13	14	0,15
Maluku	199	90	0,45	9	0,05	10	0,05
North Maluku	129	102	0,79	2	0,02	3	0,02
West Papua	157	54	0,34	5	0,03	12	0,08
Papua	356	175	0,49	11	0,03	2	0,01
Indonesia	9831	15833	1,61	1214	0,13	3834	0,40

Figure 1. Distribution of dentists (red), dental therapists (orange) and dental technicians (grey) in Indonesia (x axis is island name in Indonesia; y axis is the average number of dental health personnel per public health center).

Figure 2. Average number of dentists at every Province in each public health center.

Unequal distribution of dentists can be seen from comparing the actual number versus ideal number of dentists with the total population. The ideal ratio is one dentist for 9,090 residents. In the year of 2010, based on dentist registration data from the Indonesian Medical Council (KKI), there were 22,237 registered dentists consisting of 20,665 general dentists and 1,582 specialist dentists⁷. However, not all dentists that were registered at the Indonesia database work in a public health center. That means only 60.6% of dentists work for an Indonesian public health center, and the rest works in private sector. The amount of Dentists who work in the private sector

is 49.4% and they might not be distributed equally around Indonesia⁹.

Many factors cause this inequality, including the placement policy of health workers in each region (province and district / city), the quality and number of health service facilities, shifting disease patterns, especially in urban regions, high disparity in health status community between regions, and characteristics of the geographic area. From the results of a study conducted by Bappenas, there is a gap between the number of workers and public health centers who need health workers, in that the

Figure 3. Average number of dental therapists at every Province in each public health center.

Figure 4. Average number of dental technicians at every Province in each public health center.

number of workers is higher than public health structure itself. Furthermore, to ensure health workers are evenly distributed, it is necessary to develop an even infrastructure¹⁰⁻¹².

Health worker placement policy is largely determined collectively by health offices and regional civil service agencies. According to a result from a study in two districts in the province of Gorontalo, the unequal distribution of healthcare workers occurs due to lack of healthcare facilities. Increasing the number of health workers every year was not followed by an equal distribution to health facilities. The study also found that there were three factors that affected the policies and development of health facilities, namely disparities of health

status, migration of population, and mutual geographic characteristics^{10,15}.

Based on the results of the spatial autocorrelation test with Moran's I, there was a significant spatial autocorrelation of dentists per area (I: 0.272, z-value: 3.20) and dental therapists per area (I: 0.238, z-value: 2.85) in Indonesia. Meanwhile, the value of dental technicians per area (I: 0.002, z-value: 0.47) did not show significant spatial autocorrelation (Figure 6).

According to the World Health Organization (WHO), the ideal ratio of dentists for an area is 1:8,000 people, while Indonesia currently stands at 1:17,105. Dentists per area in Indonesia

Figure 5. Visualization maps of dentists per area (a), dental therapists per area (b), and dental technicians per area (c). Orange = <1; red = ≥1.

have spatial autocorrelation at the provincial level and are concentrated only in the Java-Bali region, with very few in the Papua region. For example, Saudi Arabia is one of the countries that has improved the ratio of dentists to population. In 1987 the ratio of dentists to population was 1:8,906, but in 2016 this number increased to 1:1,880. The ratio of dentists to population in Saudi Arabia recently was 5.3 per 10,000 people. This number is higher than all the developing countries in Asia-Pacific region. Also, in this region, China reported the lowest ratio of dentists to population, with 0.12 per 10,000 people. And on the other side, Japan has the highest ratio of dentists to population ratio in Asia-Pacific region with 7.7 per 10,000 people^{18,22,23}.

Other studies reported that the OECD member countries, excluding Scandinavian countries and Greece, have varied dentists-to-population ratios ranging from 5 to 8 with the average of 6.1 per 10,000 people. In addition, most of the European countries have dentists-to-population ratios ranging from 5.07

to 7.3 per 10,000 people. Among Middle Eastern countries, Bahrain has the lowest dentists-to-population ratio of 1.5 per 10,000 people, and Qatar has the highest dentists-to-population ratio of 5.8 per 10,000 people^{19,22,24,25}.

Based on the distribution pattern that we found, we assumed that the distribution of dentists in Indonesia occurred due to the large population, the number of dental institutions, as well as the equitable distribution of development in each region. Java island is one of the areas with an equal distribution of dentists in Indonesia, though it is also the most populated island in Indonesia^{20,21}. In line with the large population growth, Java Island is an area that has 18 dental institutions out of a total of 31 dental institutions throughout Indonesia. Issues of development and availability of public facilities and infrastructure are the main reasons why dentists prefer to practice in the Java area. Dental therapists per area also have spatial autocorrelation at the provincial level, where the Nusa Tenggara region has the highest distribution and the Papua region has the least^{26,27}.

Figure 6. Spatial autocorrelation with Moran's index of dentists per area (a), dental therapists per area (b), and dental technicians per area was developed (c). The X axis is referring to the number of observations and is also known as the response axis. The Y axis is referring to the average or spatial lag of the corresponding observation of dental health personnel per area of the X axis.

This spatial autocorrelation illustrates that there is a correlation between the number of dentists per area and dental therapists per area between provinces in Indonesia and shows geographic clustering relationships or patterns that are grouped and have similar characteristics in adjacent locations. This spatial autocorrelation did not occur in the value of dental technicians. The results of Moran's I test show that there is an autocorrelation or spatial relationship of the number of dentists per area and dental therapists per area in Indonesia. Furthermore,

Moran's scatterplot also illustrates the pattern of relationships between the existing provinces (Figure 7).

Based on the distribution patterns, similar to what happened in the maldistribution of dentists, the maldistribution that occurs in dental therapists may also be due to the population number, the number of dental institutions, and the equal distribution of welfare in each region according to this result. In addition, we assumed that the geographical grouping of dental therapists

Figure 7. Spatial autocorrelation illustration of dentists per area (a), dental therapists per area (b), and dental technicians per area (c). The plots show the spatial autocorrelation between the number of dentists and dental therapists per area in every province in Indonesia. But the dental technician is not showing the spatial autocorrelation.

in the Nusa Tenggara region was due to the process of substituting the role of dentists by dental therapists where the distribution of dentists in the area was low. Meanwhile, a very minimal distribution was seen in the value of dental technicians per area. In this study, it was found that all regions of Indonesia did not have a sufficient number of dental technicians for each public health center, with a mean value of less than 1 per public health center. The factor that most influences this is the very limited number of schools for dental technicians; only 10 institutions throughout Indonesia. Several conditions above then describe the minimum distribution of dental personnel in Indonesia²⁸.

A sufficient number of dental health personnel, such as dentists, dental technicians, and dental therapists in a Province is very important in efforts to ensure health service. The Indonesian government has made regulations regarding the minimum number of dentists in public health centers (minimum 1), but there is not a distribution policy for dental technicians and dental therapists. Indonesia has also implemented various policies, such as increased numbers of dental students to be trained in faculties of dentistry nationwide and pre contract for dentists to work in rural areas after graduation through the Nusantara Sehat and PTT Daerah programs, to solve the shortage of dentists. However, the inequality of dental personnel distribution is still being found²⁹.

We determined the significance of local spatial autocorrelation through LISA. From this test, the significance of the relationship in each province was obtained. The pattern of Moran's local significance levels is presented in Figure 8. Interpretation of the LISA significance map includes the following categories:

“High-high” indicates a clustering of high-value rates (positive spatial autocorrelation)

“Low-high” indicates that the low-value rates are adjacent to high-value rates (negative spatial autocorrelation)

“Low – low” indicates clustering of low-value rates (positive spatial autocorrelation)

“High – low” indicates that high-value rates are adjacent to low-value rates (negative spatial autocorrelation)

“Not significant” indicates that there is no spatial autocorrelation.

Spatial analysis was used to identify dental health personnel shortages and geographical distribution. The traditional methods use administrative boundaries such as counties as the basic spatial units as well as dental health personnel to identify shortages based on those numbers^{30–32}. Such approaches have been criticized for their inability to account for either the spatial variations of population demand and dental health personnel supply within those boundaries or for population–dental health personnel interactions across them. Spatial analysis can assist current and future dental health personnel, dental school administrators, and policymakers in making informed decisions to determine suitable practice locations, dental school admissions criteria, and target areas for public health initiatives^{33,34}.

The problem of unequal distribution of dental health personnel should use spatial analysis to understand the issues. This study included number of dental health personnel in each public health center for the accuracy of this research. This

Figure 8. Autocorrelation local indicator of spatial autocorrelation (LISA) map of dentists per area (a), dental therapists per area (b), and dental technicians per area (c). Grey = not significant, red = high-high, blue = low-low, purple = low-high, peach = high-low.

research can be accurate because the characteristics of public health centers in each region are similar^{35–37}. Future studies are recommended to use data of dental health personnel both in the public sector and in the private sector.

Conclusion

The number of dentists in Indonesia has increased due to the increase of dental students who have been trained in faculties of dentistry which are also done the pre-contracts for dentists to work in rural areas after graduation through the Nusantara Sehat and PTT Daerah programs, to solve the shortage of dentists. However, the inequality of dental

personnel distribution is still found, there is no distribution policy for dental technicians and dental therapists, and the dentist-to-population ratios in Indonesia have not improved. Spatial analysis might help identify dental health personnel shortage and geographical distribution.

Data availability

Source data

The distribution data of the number of dentists in each province is available at <https://www.litbang.kemkes.go.id/> (Ministry of Health) and geographic area data is available from <https://tanahair.indonesia.go.id/portal-web> (Geospatial Portal).

References

1. Azwar A: **Pengantar Administrasi Kesehatan**. Edisi Ketiga. Jakarta: Binarupa Aksara; 1996.
2. Kurcz R, Kruger E, Tennant M: **Using GIS to analyse dental practice distribution in Indiana, USA**. *Community Dent Health*. 2013; **30**(3): 155–60. [PubMed Abstract](#)
3. Widström E, Tiira H, Tillberg A: **Public dental service personnel facing a major health care reform in Finland**. *BDJ Open*. 2019; **5**: 5. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Liu DL, Xie YF, Shu R: **Statistical analysis of current oral health care and dental education resources in China**. *Chin J Dent Res*. 2019; **22**(1): 37–43. [PubMed Abstract](#) | [Publisher Full Text](#)
5. Gabriel M, Cayetano MH, Galante ML, et al.: **A global overview of the geographical distribution of dentists: A scoping review**. *JDR Clin Trans Res*. 2018; **3**(3): 229–37. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Kruger E, Whyman R, Tennant M: **High acuity GIS mapping of private practice dental services in New Zealand: does service match need?** *Int Dent J*. 2012; **62**(2): 95–99. [PubMed Abstract](#) | [Publisher Full Text](#)
7. Almado H, Kruger E, Tennant M: **Application of spatial analysis technology to the planning of access to oral health care for at-risk populations in Australian capital cities**. *Aust J Prim Health*. 2015; **21**(2): 221–226. [PubMed Abstract](#) | [Publisher Full Text](#)
8. Tennant M, Kruger E, Shiyha J: **Dentist-to-population and practice-to-population ratios: in a shortage environment with gross mal-distribution what should rural and remote communities focus their attention on?** *Rural Remote Health*. 2013; **13**(4): 2518–2524. [PubMed Abstract](#) | [Publisher Full Text](#)
9. Evans B, Sabel CE: **Open-Source web-based geographical information system for health exposure assessment**. *Int J Health Geogr*. 2012; **11**: 2. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Kruger E, Whyman R, Tennant M: **High acuity GIS comparison of dentist and doctor surgery locations in Auckland, New Zealand**. *Community Dent Health*. 2013; **30**(2): 83–7. [PubMed Abstract](#)
11. Badan Litbangkes: **Laporan Riset Kesehatan Dasar 2013**. Jakarta: Lembaga Penerbit Badan Litbangkes; 2014.
12. Badan Litbangkes: **Laporan Riset Fasilitas Kesehatan 2011**. Jakarta : Lembaga Penerbit Badan Litbangkes; 2012.
13. **Kementerian Kesehatan Peraturan Menteri Kesehatan No. 75/2014 tentang Pusat Kesehatan Masyarakat**. Jakarta : Kementerian Kesehatan RI; 2014.
14. Pusat Perencanaan dan Pendayagunaan Sumber Daya Manusia Kesehatan: **Analisis Kebutuhan Tenaga Kesehatan Sebagai Dasar Pengembangan Prodi Pendidikan Tenaga Kesehatan**. Jakarta : Pusrengun Kemenkes; 2013. [Reference Source](#)
15. Blas E, Kurup AS: **Equity, Social Determinants and Public Health Programmes**. World Health Organization. 2010; 159–176. [Reference Source](#)
16. Husain I, Hasanbasri M, Soetjipto HP: **Kualitas dan Kuantitas Tenaga Kesehatan Public health center – Studi Distribusi Desa/Kota dan Regional, Analisis Data SAKERTI 2000**. Working Paper Series No. 18, Juli 2006; KMPK Universitas Gadjah Mada, Yogyakarta. 1–62. [Reference Source](#)
17. Roeger LS, Reed RL, Smith BP: **Equity of access in the spatial distribution of GPS within an Australian metropolitan city**. *Aust J Prim Health*. 2010; **16**(4): 284–90. [PubMed Abstract](#) | [Publisher Full Text](#)
18. Krishnan V: **A macro model of change in specialty and spatial distribution of physicians in Canada, 1971-1981**. *Socioecon Plann Sci*. 1992; **26**(2): 111–27. [PubMed Abstract](#) | [Publisher Full Text](#)
19. Gächter M, Schwazer P, Theurl E, et al.: **Regional density of private dentists: Empirical evidence from Austria**. *Community Dent Oral Epidemiol*. 2014; **42**(1): 20–9. [PubMed Abstract](#) | [Publisher Full Text](#)
20. Luo W: **Using a GIS-based floating catchment method to assess areas with shortage of physicians**. *Health Place*. 2004; **10**(1): 1–11. [PubMed Abstract](#) | [Publisher Full Text](#)
21. Lamster IB, Eaves K: **A model for dental practice in the 21st century**. *Am J Public Health*. 2011; **101**(10): 1825–1830. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Gallagher JE, Manickam S, Wilson NHF: **Sultanate of Oman: building a dental workforce**. *Hum Resour Health*. 2015; **13**: 50. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Huang CS, Cher TL, Lin CP, et al.: **Projection of the dental workforce from 2011 to 2020, based on the actual workload of 6762 dentists in 2010 in Taiwan**. *J Formos Med Assoc*. 2013; **112**(9): 527–536. [PubMed Abstract](#) | [Publisher Full Text](#)
24. Koletsi-Kounari H, Papaioannou W, Stefanitis T: **Greece's high dentist to population ratio: comparisons, causes, and effects**. *J Dent Educ*. 2011; **75**(11): 1507–1515. [PubMed Abstract](#) | [Publisher Full Text](#)
25. Krause D, Frate DA, May WL: **Demographics and distribution of dentists in Mississippi: A dental work force study**. *J Am Dent Assoc*. 2005; **136**(5): 668–77. [PubMed Abstract](#) | [Publisher Full Text](#)
26. Thanakanjanaphakdee W, Laohasirirong W, Puttanapong N: **Spatial distribution of dentists in Thailand**. *J Int Oral Health*. 2019; **11**: 340–6. [Publisher Full Text](#)
27. Susi L, Mascarenhas AK: **Using a geographical information system to map the distribution of dentists in Ohio**. *J Am Dent Assoc*. 2002; **133**(5): 636–42. [PubMed Abstract](#) | [Publisher Full Text](#)
28. **Konsil Kedokteran Indonesia (Indonesia Medical Council): Standar Kompetensi Dokter Gigi Jakarta: Konsil Kedokteran**. 2006. [Reference Source](#)
29. **Kementerian Kesehatan RI Keputusan Menteri Kesehatan No. 378/Menkes/SKIII/2007 tentang Standar Profesi Perawat Gigi**. Jakarta: Kemenkes RI; 2007. [Reference Source](#)
30. Wright SE: **The spatial distribution and geographic analysis of endodontic office locations at the national scale**. *J Endod*. 1994; **20**(10): 500–505. [PubMed Abstract](#) | [Publisher Full Text](#)
31. Perera I, Kruger E, Tennant M: **GIS as a decision support tool in health informatics: spatial analysis of public dental care services in Sri Lanka**. *J Health Inform Dev Ctries*. 2012; **6**: 422–434. [Reference Source](#)
32. Jaafar N, Hakim H, Mohd Nor NA, et al.: **Is the burden of oral diseases higher in urban disadvantaged community compared to the national prevalence?** *BMC Public Health*. 2014; **14**(Suppl 3): S2. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Esa R, Ong AL, Humphris G, et al.: **The relationship of dental caries and dental fear in Malaysian adolescents: a latent variable approach**. *BMC Oral Health*. 2014; **14**: 19. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Petersen PE, Kwan S: **Equity, social determinants and public health programmes—the case of oral health**. *Commun Dent Oral Epidemiol*. 2011; **39**(6): 481–487. [PubMed Abstract](#) | [Publisher Full Text](#)
35. Watt RG: **Social determinants of oral health inequalities: implications for action**. *Commun Dent Oral Epidemiol*. 2012; **40** Suppl 2: 44–48. [PubMed Abstract](#) | [Publisher Full Text](#)
36. Hosny G, Sayed S, Tantawi M, et al.: **Evaluation of accessibility of dental services using Geographic Information System**. *Sci Afric J Sci Issues*. 2015; **3**: 763–767.
37. Omogunloye OG, Tijani OA, Abiodun EO, et al.: **Geospatial distribution and utilization of dental facilities in Lagos State**. *J Biodivers Endanger Species*. 2016; **4**: 1–18. [Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 11 June 2021

<https://doi.org/10.5256/f1000research.57637.r87316>

© 2021 Leite Cavalcanti A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alessandro Leite Cavalcanti

Department of Dentistry, UEPB - Universidade Estadual da Paraíba, Campina Grande, Brazil

No further comments.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health and Epidemiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 25 May 2021

<https://doi.org/10.5256/f1000research.53958.r81866>

© 2021 Leite Cavalcanti A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alessandro Leite Cavalcanti

Department of Dentistry, UEPB - Universidade Estadual da Paraíba, Campina Grande, Brazil

I congratulate the authors for the excellent quality of the manuscript. The study is relevant and describes the profile of dental professionals in Indonesia. Some minor corrections are suggested:

1. Page 1: ABSTRACT: It is not necessary to repeat the name of the country in the results, since it is already in the objective.

2. Page 3, left column, 2nd paragraph, line 11: I suggest excluding "while the rest received no treatment".
3. Page 3, right column, 1st paragraph, sentence 1: New wording "As of 2012, from 8,975 public health centers, there are 5,439 public health centers with dentists (60.6%)." You must delete "and 3,536 public health center without dentists".
4. Page 6, left column, 2nd paragraph: I suggest that the authors inform how many dental schools there are in the country and how many dentists are trained each year. The same applies to dental therapists and technicians.
5. Page 10: I recommend that the authors describe the limitations of the study.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public Health and Epidemiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 17 May 2021

<https://doi.org/10.5256/f1000research.53958.r85400>

© 2021 Hadi R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Reny Purnama Hadi

Master Program in Biomedical Science, Faculty of Medicine, Brawijaya University, Malang, Indonesia

First of all the article is unique, never discussed before and adequate. Moreover, there are some additional suggestions:

1. The abstract background needs to focus on spatial distribution; 3 professions.
2. The method is adequate and clear.
3. The first sentence of the conclusion should be answering on the background.
4. The discussion is clear and adequate.

After all, the overall is very good.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biomedical, Public Health of Medicine

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research