RESEARCH ARTICLE

Cytotoxic T cells response with decreased CD4/CD8 ratio during mammary tumors inhibition in rats induced by non-contact electric fields [version 1; peer review: 1 approved, 1 approved with reservations]

Firman Alamsyah ¹, Rarastoeti Pratiwi ², Nisrina Firdausi ², Jessica Irene Mesak Pello², Subekti Evi Dwi Nugraheni ², Ahmad Ghitha Fadhlurrahman ², Luthfi Nurhidayat ², Warsito Purwo Taruno¹

¹Center for Medical Physics and Cancer Research, Ctech Labs Edwar Technology, Tangerang, Banten, 15143, Indonesia

²Faculty of Biology, Universitas Gadjah Mada, Sleman, DI Yogyakarta, 55281, Indonesia

V1 First published: 19 Jan 2021, 10:35
<https://doi.org/10.12688/f1000research.27952.1>
 Latest published: 04 May 2021, 10:35
<https://doi.org/10.12688/f1000research.27952.2>

Abstract

Background: Breast cancer is the most common cancer in women worldwide and is the leading cause of death in women with cancer. One novel therapy used for breast cancer treatment is non-contact electric fields called electro-capacitive cancer therapy (ECCT) with intermediate frequency (100 kHz) and low intensity (18 Vpp). The objective of this study was to examine the effect of ECCT on mammary tumors growth in rats and observing the immune responses that play a role in fighting the tumor.

Methods: Female SD rats were used and divided into four groups, namely control (NINT), placebo (NIT), non- therapy (INT), and therapy (IT) groups with 6 biological replicates in each group. Rats in INT and IT groups were treated with 7,12-dimethylbenz[a]anthracene for mammary tumor induction. Only rats in NIT and IT groups were exposed to ECCT individually for 10 hours per day for 21 days. The size of all tumors was measured with a digital caliper. The distributions of PCNA, ErbB2, caspase-3, CD68, CD4 and CD8-positive cells were observed with immunohistochemistry and scoring with ImageJ.

Results: The growth rate of mammary tumors in IT group was significantly lower ($p < 0.05$) than that in the INT group. The number of mitotic figures and the percentage of PCNA, caspase-3, and CD68-positive cells in IT group were significantly lower ($p < 0.05$) than those in INT group. Conversely, the percentage of CD8-positive T cells in IT group was significantly higher ($p < 0.05$) than that in INT group. Moreover, the CD4/CD8 ratio in IT group was decreased. Some tumor tissues were blackened and detached from the surrounding tissue,

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
04 May 2021	view	
version 1		
19 Jan 2021	view	view

1. **Bilgin Keserci**, Universiti Sains Malaysia, Kelantan, Malaysia

2. **Elfahmi Elfahmi**, Bandung Institute of Technology, Bandung, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

resulting in an open wound which then healed up upon exposure.

Conclusions: Non-contact electric fields exposure showed inhibition on mammary tumor growth in rats while inducing CD8+ T cells that lead to tumor cells death and potentially helps wound healing.

Keywords

non-contact electric fields, ECCT, mammary tumors, CD8+ T cells, CD4/CD8 ratio

This article is included in the **Cell & Molecular Biology** gateway.

Corresponding author: Firman Alamsyah (firman.alamsyah@c-techlabs.com)

Author roles: **Alamsyah F:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation; **Pratiwi R:** Conceptualization, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Original Draft Preparation; **Firdausi N:** Data Curation, Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation; **Irene Mesak Pello J:** Data Curation, Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation; **Evi Dwi Nugraheni S:** Data Curation, Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation; **Ghitha Fadhlurrahman A:** Data Curation, Investigation, Project Administration, Writing – Original Draft Preparation; **Nurhidayat L:** Methodology, Supervision, Validation, Writing – Original Draft Preparation; **Purwo Taruno W:** Funding Acquisition, Resources, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by Ctech Labs Edwar Technology for the in vivo experiment and a grant from the Indonesian Ministry of Research, Technology and Higher Education (Contract number: 11/PPK/E/E4/2018) to RP for the immunohistochemistry application.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Alamsyah F *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Alamsyah F, Pratiwi R, Firdausi N *et al.* **Cytotoxic T cells response with decreased CD4/CD8 ratio during mammary tumors inhibition in rats induced by non-contact electric fields [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:35 <https://doi.org/10.12688/f1000research.27952.1>

First published: 19 Jan 2021, 10:35 <https://doi.org/10.12688/f1000research.27952.1>

Introduction

Breast cancer is the most common cancer among women in the world, in which 1 in 8 women in the world is at risk of developing breast cancer¹. Furthermore, about 2.1 million new cancer cases in 2018 are breast cancer cases and this type of cancer is the leading cause of death in women with cancer². Commonly, the treatment of breast cancer is carried out in two ways. The first, termed local therapy, controls or removes the tumor in specific areas, such as through breast-conserving therapy and mastectomy that may be followed by radiotherapy. The second, called systemic therapy, uses the circulatory system in the body to inhibit cancer cells growth that spread in parts of the body. This type of treatment includes chemotherapy, targeted therapy, and endocrine therapy³. However, breast cancer therapy applied so far often causes negative effects, such as the presence of pathophysiological mechanisms that cause chronic pain⁴, or the mechanism of multidrug resistance (MDR) in breast cancer cells as a reaction to chemotherapy^{5,6}. Moreover, breast cancer therapies may attack certain normal cells or tissues that result in toxicity symptoms⁷. A novel cancer therapy based on electric field exposure with intermediate frequency and low intensity called Tumor Treating Fields (TTFields) has been developed to prevent such negative effects of cancer therapy and still inhibit cancer growth. No toxic side effects or adverse events were found on animal tumor models nor patients under the exposure to this electric field-based cancer therapy, except for skin toxicity or dermatitis of low severity^{8,9}.

Since skin toxicity or dermatitis arises in electric field therapy due to prolonged direct contact between the electrodes and the skin¹⁰, we have developed a type of non-contact electric field therapy using capacitive modality to avoid such injuries. In our previous study in preclinical setting, cancer therapy namely Electro-Capacitive Cancer Therapy (ECCT) using the non-contact electric fields with intermediate frequency (100 kHz) and low intensity (18 peak-to-peak Voltage/Vpp) could reduce the size of breast tumors in mice significantly without causing histological damage to mammary and skin tissues. In addition, lymphocyte and macrophage infiltration was found around the tumor area through the blood vessels¹¹. One of the mechanisms of tumor cells death can be carried out with the act of anti-tumor mechanism of various immune cells, including helper and cytotoxic T cells of lymphocytes, as well as macrophages¹². Cytotoxic CD8⁺ T cells which is activated by helper CD4⁺ T cells, may enter the tumor microenvironment and induce apoptosis, resulting in shrinkage of the tumor lesions^{12,13}. Macrophages with the M1 phenotype play an important role in the recognition and destruction of cancer cells and their presence usually indicates a favorable prognosis. Whereas macrophages with the M2 phenotype are involved in anti-inflammation, helping the angiogenesis process and the expression of scavenger receptors, and have a significant role in tumor development and metastasis^{12,13}, including breast tumor^{14,15}. One of the marker proteins to observe the infiltration of macrophages in a tissue, including mammary tumor tissue, is the CD68 protein^{14–16}.

Cancer therapy that can re-induce immune response to destroy and clean up tumor cells is a therapy that is being developed

because of its positive effects. By activating immune cells, the immune cells may reach the tumor-specific areas, including metastatic area, and specifically attacking only tumor cells, not the normal cells in the vicinity¹⁷. Petri *et al.*¹⁸ reported that static electric fields increase the immune response in mice. Moreover, electric fields modulate the activation and polarisation of some immune cells, including T cells¹⁹ and macrophages²⁰. Voloshin *et al.*²¹ reported that TTFields exposure combined with anti-PD-1 therapy induced immunogenic cell death in lung and colon tumors. In addition, Pratiwi *et al.*²² reported the increased expression of CD68 and caspase-3 in rat breast tumor tissues under the exposure to 150 kHz non-contact electric fields while reducing the expression of PCNA and ErbB2 proteins. However, the study by Pratiwi *et al.*²² showed an increase of the tumor size during therapy. Using different intermediate frequency (100 kHz), we examined the effect of non-contact electric fields exposure on tumor growth in mammary tumors-induced rats and observed the activation of immune cells in fighting the tumor cells. We hypothesized that the increase in the size of mammary tumors would not happen under exposure to this non-contact electric fields with the induction of immune cells, especially lymphocytes and macrophages.

Methods

Ethics

This study was conducted at the Integrated Research and Testing Laboratory (LPPT) of Universitas Gadjah Mada (UGM) and at the Animal Structure and Development Laboratory of Faculty of Biology, UGM. LPPT UGM has been awarded ISO/IEC 17025:2000 accreditation for the competence of testing and calibration²². Experimental protocol in this study was performed following approval by the Ethical Clearance Committee of LPPT UGM with ethical clearance number: 00015/4/LPPT/IV/2017. The Ethical Clearance Committee stated that this study has been declared to meet the ethical requirements for research on experimental animals and the Ethical Clearance Committee has the right to conduct monitoring during the research.

Experimental design

A total of 40 5-week-old healthy female Sprague Dawley (SD) rats (*Rattus norvegicus*, Berkenhout 1769) weighing 50–80 gram were used for this study. SD rats have been used as animal tumor model to study human breast cancer with various induction agents, as well as with DMBA administration^{23,24}, since rats have 98% genetic homology with humans²⁵. The rats were supplied by LPPT UGM laboratory, Yogyakarta, Indonesia, and never used for any other experiment. Rats showing any symptoms of illness or abnormalities were excluded from the experiment. The rats were acclimated for 1 week in polypropylene cages as communal home cages with each size: 50 × 40 cm² that covered with rice hulls bedding at the bottom. One communal cage consisted of 5 rats. The temperature of the animal room was maintained at between 23–26°C with 81.09% average relative humidity to avoid dehydration during electric fields exposure, lighting condition was light from lamps during the day and total darkness during the night (12L:12D photoperiod). The rats were fed with AD2 pellet animal diet and tap water *ad libitum*. However, during treatment, cucumber slices were given as

a substitute for water, to avoid the animals being shocked with electricity, since the tip of the drink bottle contains metal which can permit electric fields. Cage cleaning was carried out every day by cleaning the animal waste from the cage and changing food and water. Individual marking was conducted by giving picric acid to rat fur, while group marking was conducted by labelling the cage with a paint marker to avoid potential confounders.

The SD rats were divided into four groups, namely control (non-induction and non-therapy/NINT), placebo (non-induction and therapy/NIT), DMBA-induced mammary tumors without therapy (induction and non-therapy/INT), and DMBA-induced mammary tumors with therapy (induction and therapy/IT) groups. The sample size in each group was calculated according to the Federer formula, where 6 biological replicates were used for each group²². The animals were selected randomly and assigned to control and treatment groups. Rats were picked randomly for mammary tumors induction with a single dose of DMBA (20 mg/kg body weight) administration two times per week for 5 weeks. The first 6 rats to develop tumors were assigned to IT group, and then 6 more rats were assigned to INT group. Not all rats grew tumors after DMBA administration. All healthy, normal rats in the placebo (NIT) group were given corn oil as DMBA (7,12-dimethylbenz[a]anthracene) solvent. DMBA compound is a neuroendocrine disruptor that commonly used for carcinogenesis in specific organs, including breast tumors in rats^{26,27}. All administrations were performed by the same researcher (AGF). Each researcher in the team had different task. Besides the researcher who administered DMBA, other researchers measured the size of the nodules, recorded the morphology of mammary tumors, dissected the euthanized rats and analysed the data statistically. One researcher (FA) controlled and monitored all the different steps in the experiment.

Non-contact electro-capacitive cancer therapy for mammary tumors in rats

Following the development of mammary tumors up to 1.5 cm in length, all rats in the IT group were exposed to intermediate frequency (100 kHz) and low intensity (18 Vpp) non-contact electric fields generated between pairs of capacitive electrodes embedded in individual cages that had been modified to an ECCT device (Figure 1a). The device is called non-contact because its electrodes do not directly attach to the animal skin (Figure 1b). All individual cages were placed on the same shelf at the same height. Non-contact electric fields therapy was performed 10 hours per day for 21 days with 2 hours rest after first 5 hours therapy with the same starting time of therapy each day. The therapy was stopped when the mammary tumors has increased to 2.25 cm² in size or it has reached the last day of therapy. All rats in placebo (NIT) group were also exposed to non-contact electric fields in the same period, 3 weeks after the last corn oil administration. The duration of electric fields exposure for 10 hours per day was shorter than the previous exposure in our preliminary study¹¹ to minimize any possible side effects due to exposure to electric fields^{8,10}. After treatment, all animals were returned to their communal cages.

During therapy, all tumors were palpated once every 2 days and their size (cm²) was measured with a digital caliper and tabulated. All nodule measurements were conducted by the same researcher (NF). In this study, we did not measure nodule size in volume due to tool limitations. The tumor size data were used to calculate the tumor surface area (TSA) using the formula for circle and ellipse areas, and subsequently the TSA data were used to calculate the tumor growth rates (TGR) of mammary tumors. The TGR was calculated by using specific growth rates (SGR) formula²⁸ with modification as follows: $SGR (\%/day) = \ln (A2/A1)/(t2 - t1)$, where A1 and A2 are the tumor surface area before (t1) and after therapy (t2), respectively.

Figure 1. Electro-Capacitive Cancer Therapy (ECCT) device in individual cage. (a) ECCT with oscillators connected to electricity, (b) A rat inside an individual cage of ECCT. The dimension of the cage was 23 cm × 18 cm × 19 cm. The arrangement of the electrodes has been described in our previous study¹¹.

Besides tumor size, the morphology of mammary tumors was also recorded by the same researcher (JIMP).

Necropsy and tissue harvesting

After therapy was completed, all rats were euthanized by an overdose of ketamine (150 mg/kg of body weight) for deep anesthesia using intramuscular injection. The rats were pinned ventral side up on a dissected box and dissected with the same surgeon (AGF). Nodules and normal breast tissues were taken and washed using physiological saline. All tissues were fixed using neutral buffered formalin (NBF). Three nodules from three different rats were collected from each INT and IT groups, while one normal breast tissue was collected from each NINT and NIT groups. These six nodules were used for histological examination with immunohistochemical staining (IHC) of the target antigens. While the normal breast tissues were used for comparison of the overall histological structure without IHC.

Histological and immunohistochemical examination

Histological examination was performed by using paraffin-embedded Hematoxylin and Eosin (H&E)-stained histology slides followed by IHC²⁹ using Starr Trek Universal HRP Detection Kit Biocare. Observations and counting were performed on the mitotic figures on H&E preparations. Six antibodies purchased from Abcam Cambridge, United Kingdom, were used for IHC, namely anti-PCNA polyclonal antibody (Cat# ab18197, RRID:AB_444313), anti-ErbB2 monoclonal antibody (Cat# ab16901), anti-caspase-3 polyclonal antibody (Cat# ab13847), anti-CD68 monoclonal antibody (Cat# ab201340), anti-CD4 polyclonal antibody (Cat# ab203034), and anti-CD8 alpha monoclonal antibody (Cat# ab33786, RRID:AB_726709). The dilution for anti-CD4 and anti-CD8 antibodies was 1:750, where 1 µl of antibody was diluted with 750 µl of BSA-PBST solution (Bovine Serum Albumin 1% - Phosphate Buffered Saline with Tween® 20). While the dilution for anti-PCNA, anti-ErbB2, anti-caspase-3 and anti-CD68 antibodies was 1:500. As much as 100 µl diluted primary antibody was applied on the appropriate slide covered with a parafilm and placed in a humidity chamber. Then, the slide was removed to a refrigerator (4°C) overnight. One to two drops (50 to 100 µl) of biotinylated secondary antibody Trekkie Universal (Biocare Medical, STUHRP700 H L10) were applied on the slides covered with a parafilm and incubated for 30 minutes at room temperature. The concentration of antibodies given was followed the manufacturer's instructions.

Data analysis

All measured data were analysed with appropriate methods without any exclusion. Semi-quantitative IHC scoring was performed using the percentage of positive area of stained cells (CD4 and CD8-positive T cells; caspase-3, ErbB2, PCNA-positive tumor cells) and the number of stained cells (CD68-positive macrophages). IHC scoring was performed in ImageJ ver.1.51, a freely available software²². The comparison between positive area of stained cells and total area showed the distributions of targeted antigens and cells³⁰. Scoring was carried out in 50 fields per group (INT and IT groups), in which 17–18 random fields were examined in one slide. In addition, CD68+

macrophages counting was performed by using 60 fields per group with 20 random fields examined in a slide. In total, three slides from three different nodules were examined from each group.

All data were tested for normality (Shapiro-Wilk test, $\alpha = 0.05$). The data of mammary tumors growth rate from 12 animals (6 rats each from IT and INT groups) and the number of mitotic figures were analysed using T-test ($\alpha = 0.05$) to find statistical significance. The data of PCNA, ErbB2, caspase-3, CD68, CD4, and CD8-positive cells and CD4/CD8 ratio were analysed quantitatively using non-parametric Mann-Whitney test, since the data were not normally distributed. All data were analysed statistically using GraphPad Prism ver.8.4.3 software²² for Windows by the same researcher (NF). Qualitative data (morphology of mammary tumors) were analysed descriptively.

Results

The outcome of this study was the growth rate of mammary tumors under exposure to non-contact electric fields. Comparison of histological characteristics between breast tissue and breast cancer in the animal tumor model, the distribution of molecular markers that described the proliferation and the mechanism of cell death, and the response of immune cells under exposure, as will be explained coherently below.

The growth rate of rat mammary tumors under exposure to non-contact electric fields

The growth rate of rat mammary tumors in IT group (-0.043 ± 0.065) was significantly lower ($p < 0.05$) than that in INT group (0.108 ± 0.078) and showing negative tumor growth rate as shown in Figure 2a. There was no tumor growth in the placebo (NIT) group. Mammary tumors morphology in INT and IT groups showed differences, in which mammary tumors in the INT group were generally more compact as previously reported by Pratiwi *et al.*²² and appear bruised (Figure 2b) due to thrombocytopenia from the malignancy³¹, whereas mammary tumors in the IT group were generally softer because they contained fluid²² and some were blackened (Figure 2c). The blackened mammary tumors in the IT group were released from the surrounding tissue, as shown in Figure 3a. These tumors initially darkened, then hardened like a scab as occurs in a dry wound, and slowly sloughed off from the surrounding tissue, causing an open wound which then dried up (Figure 3b) and shrank upon exposure to non-contact electric field (Figure 3c).

Histological characteristics of breast tissue and breast cancer in rats

The characteristics of normal breast tissue and the pathogenesis of actively dividing cells in DMBA-induced breast cancer in rats can be seen through H&E staining (Figure 4). Figures 4a, b show microscopical sections of breast tissue from rats of NINT and NIT groups, respectively, having normal histological features with the presence of adipose tissue and other connective tissues, and in the absence of actively proliferating cells. In addition, a myoepithelial cell layer was clearly visible in the glandular epithelium³². Conversely, Figure 4c, d show the level of malignancy of breast cancer cells from rats of INT and IT groups, respectively. They show histological

Figure 2. Comparison of specific growth rates (SGR) and morphology of tumors in INT and IT groups. (a) SGR of mammary tumors in INT and IT groups, (b) morphology of a mammary tumor in INT group with the appearance of bruises, (c) morphology of a mammary tumor in IT group with a blackened condition. Different letters (a, b) on top of the charts indicate significant differences ($p < 0.05$).

Figure 3. An open wound after the release of darkened tumor and its size (cm²) upon exposure. (a) open wound on day 14 of therapy, (b) open wound on day 21 (final day) of therapy, (c) the reduction of wound size from day 14 to day 21 of therapy.

feature of papillary carcinoma with the characteristics of malignant epithelial proliferation protruding into the lumen of the glandular duct that made it constricted. However, breast cancer in rats of the IT group tend to be less malignant than that in the INT group, indicated by the clear glandular duct and

glandular lobule lumens, and the prominent epithelial structure as well (Figure 4d). In addition, the lumens in IT group still maintained an intact layer of cells (Figure 4d), while most of the lumens in INT group did not (Figure 4c), due to the pressure exerted by the breast cancer cells. Moreover, the

region of adipose tissue and other connective tissues were also pressed by the mammary tumor cells (Figure 4d). On the other hand, the tumor tissue itself undergoing necrosis (Figure 4c, d) which was characterized by inflammation^{33,34} and the necrotic areas with inflammatory environment in the INT group was larger than that in the IT group. Figure 4e and Figure 4f show mitotic figures and apoptotic bodies in INT and IT groups, respectively.

Distribution of PCNA, ErbB2 and caspase-3 in breast cancer with and without non-contact electric field exposure

Immunohistochemical staining with anti-PCNA, anti-ErbB2 and anti-caspase-3 detected PCNA, ErbB2, and caspase-3-positive cells, respectively, in rat mammary tumors. The expression of PCNA in INT and IT groups were detected in the nucleus of tumor cells (Figure 5a and Figure 5b, respectively), while the expression of ErbB2 in both groups was found on the cell membranes (Figure 5c and Figure 5d, respectively). Caspase-3 was expressed in different locations in mammary tumors of INT and IT groups (Figure 5e and Figure 5f, respectively). The expression of caspase-3 in INT group was found in the cell cytoplasm (Figure 5e), while in IT group, caspase-3

expression was found in the nucleus (Figure 5f) and resulted in chromatin condensation and DNA fragmentation, indicating that apoptosis has reached a late stage^{35,36}. In addition, the expression of caspase-3 in IT group was also found at hollow regions in mammary tumors (Figure 5f), whereas caspase-3 expression in INT group was rarely found in the same region (Figure 5e).

Breast cancer cells that actively proliferated in rats of the INT group were seen to be relatively greater than those in the IT group, indicated by the higher number of mitotic figures per field in the INT group (10.14 ± 3.80) compared to that in the IT group (5.43 ± 1.81) as shown in Figure 6a. In addition, the percentage of positively stained areas of PCNA in INT group ($36.16 \pm 5.16\%$) was significantly higher than that in INT group ($23.65 \pm 3.84\%$, Figure 6b). Figure 6c and Figure 6d show that INT group had a greater percentage of positively stained areas of ErbB2 ($0.045 \pm 0.029\%$) and caspase-3 ($8.90 \pm 5.48\%$) than those in IT group (0.036 ± 0.029 for

Figure 4. Histological features of mammary glands, mammary tumors and mitotic figures in rats with H&E staining. (a) Control (NINT) group, (b) Placebo (NIT) group, (c) Non-therapy (INT) group, (d) Therapy (IT) group, (e) Mitotic figures (indicated by arrows) and apoptotic bodies (in circles) in INT group, and (f) Mitotic figures and apoptotic bodies in IT group. CT= Connective tissue, NA= Necrotic area, DEC= Ductal epithelial cells, MEC= Myoepithelial cells, LU=Lumen, AC= Adipose Cells, BV = Blood Vessel.

Figure 5. Immunohistochemical staining (3,3'-diaminobenzidine) of rat mammary tumors with PCNA, ErbB2, and caspase-3-positive cells. (a) Expression of PCNA in the nucleus of tumor cells in non-therapy (INT) group, (b) Expression of PCNA in the nucleus of tumor cells in therapy (IT) group, (c) Expression of ErbB2 on the cell membranes of tumor cells in INT group, (d) Expression of ErbB2 on the cell membranes of tumor cells in IT group, (e) Expression of caspase-3 in the cytoplasm of tumor cells in INT group, (f) Expression of caspase-3 in the nucleus and Hollow regions (HR) of tumor cells in IT group. Brown color chromogen and arrows indicates positively stained cells.

Figure 6. Number of mitotic figures and percentage of PCNA-, ErbB2-, caspase-3-positive cells in mammary tumors. (a) Number of mitotic figures, (b) percentage of PCNA-positive cells, (c) percentage of ErbB2-positive cells, (d) percentage of caspase-3-positive cells. Error bars represent the standard error of the mean. Different letters (a, b) on top of the charts indicate significant differences ($p < 0.05$).

ErbB2 and 3.01 ± 1.52% for caspase-3). However, the difference of percentage of positively stained areas of caspase-3 between the two groups was considered statistically significant ($p < 0.05$), while the difference in ErbB2 expression was not significant ($p > 0.05$).

Distribution of CD68⁺ macrophages, CD4⁺ and CD8⁺ T cells in breast cancer with and without non-contact electric field exposure

Immunohistochemical staining with anti-CD68, anti-CD4 and anti-CD8 detected CD68⁺ macrophages, CD4⁺ and CD8⁺ T cells, respectively, in rat mammary tumors. The expression of CD68, CD4 and CD8 were detected on the cell membrane and in cytoplasm (Figure 7). In addition, CD68⁺ macrophage infiltration was found in mammary tumors of INT and IT groups, particularly in necrotic areas and hollow regions (Figure 7a and Figure 7b, respectively). The hollow regions in IT group were seen to be more numerous and larger in size than those found in INT group. In addition, CD68⁺ macrophage infiltration into the hollow regions of mammary tumors in IT group were also seen to be greater than that in INT group. Conversely, CD68⁺ macrophages infiltration in necrotic areas was seen to be higher in INT group (Figure 7a) than that in IT group (Figure 7b), since the distribution of necrotic areas in INT group was also wider than that in IT group. Similarly, CD4⁺ and CD8⁺ T cells were also

found around the proliferating tumor cells that may respond to certain antigens in the mammary tumor cells, within blood vessels, and in the necrotic areas^{12,13} which were characterized by inflammation^{33,34} (Figure 7c and Figure 7d for CD8, Figure 7e and Figure 7f for CD4).

Significantly higher number ($p < 0.05$) of CD68⁺ macrophages was found in the INT group (8.20 ± 0.52) than that in IT group (4.87 ± 0.30 , Figure 8a) with lower growth rate (Figure 2a). The percentage of positively stained areas of CD4 in INT group ($0.42 \pm 0.08\%$) was higher than that in IT group ($0.32 \pm 0.08\%$), but the difference was not statistically significant ($p > 0.05$, Figure 8b). In contrast, the percentage of positively stained areas of CD8 in INT group ($0.21 \pm 0.04\%$) was significantly lower ($p < 0.05$) than those in IT group ($0.64 \pm 0.11\%$, Figure 8c). Consequently, the ratio of CD4/CD8 in INT group (7.16 ± 2.60) was higher than that in IT group (2.44 ± 1.08 , Figure 8d).

Discussion

In the present study, the significantly lower growth rate of rat mammary tumors in the IT group with a negative mean value of SGR compared to INT group with a positive mean value of SGR (Figure 2a) showed the inhibitory effect of non-contact electric fields of ECCT against mammary tumor cells that are actively dividing in the animals, as we have previously reported

Figure 7. Immunohistochemical staining (DAB) of mammary tumors with CD68-positive macrophages, CD4 and CD8-positive T cells. (a) CD68⁺ macrophages in non-therapy (INT) group, (b) CD68⁺ macrophages in therapy (IT) group, (c) CD8⁺ T cells in INT group, (d) CD8⁺ T cells in IT group, (e) CD4⁺ T cells in INT group, (f) CD4⁺ T cells in IT group. Brown color chromogen indicates positively stained cells. MC: Macrophage cells, CT: Connective Tissue, BV: Blood Vessel, DEC: ductal epithelial cells, NA: Necrotic area, CTL: Cytotoxic T Lymphocyte, and HTL: Helper T Lymphocyte.

in our preliminary study in mice¹¹. In addition, the development of mammary tumors exposed to 100 kHz non-contact electric fields of ECCT was decreased, whereas in the study of Pratiwi *et al.*²² which used a frequency of 150 kHz, it was increased. This may suggest that differences in the frequency of the electric fields can give different therapeutic results in cancer^{37,38} and the 100 kHz was the appropriate frequency of non-contact electric fields in inhibiting the proliferation of mammary tumor cells.

The absence of tumors in the placebo (NIT) group (Figure 4b) demonstrated the safety of non-contact electric fields of ECCT to healthy or normal animal models, as we have previously reported¹¹. In addition to its safety, this electric fields exposure did not produce acute damage to rat skin tissue or chronic pain which usually occurs after radiation therapy, including breast cancer radiation therapy^{4,39,40}. Instead, it helped wound healing after cancer treatment in rats of IT group, indicated by the drying wound and the reduction of wound size (Figure 3b and Figure 3c). Endogenous electric fields are known to play as a signal that direct cell migration in epithelial

wound healing and external electric fields exposure may accelerate wound healing by directing the migration of T cells¹⁹ and fibroblast, as well as their proliferation and transdifferentiation^{41,42}. In addition, Hoare *et al.*²⁰ reported that external electric fields may contribute to the coordination and regulation of macrophage functions, including wound healing. The effect of non-contact electric fields of ECCT in directing immune cells was also seen in the fluid contained in the mammary tumors of the IT group as indicated by the presence of macrophages and lymphocytes infiltration into the tumor area (Figure 7) as we have previously reported^{11,22}. This immune cell infiltration can promote the destruction of cancer cells^{12,21,43} as demonstrated in the IT group (Figure 2a and Figure 2c).

The cellular effects of electric fields exposure on PCNA expression during S-phase may indicate the incidence and the growth rate of mammary tumors⁴⁴. Since PCNA is a S-phase marker in interphase that has a central role in DNA replication and repair⁴⁵, the decrease of PCNA expression in mammary tumor cells in IT group (Figure 6b) may indicate a decrease in the incidence and the growth rate of mammary tumors⁴⁴. In addition, alternating electric fields exposure with intermediate frequency and low intensity used in this study can cause tumor cells that are dividing to experience uneven segregation of chromosomes and mitotic catastrophe which followed by programmed cell death⁴⁶. This is supported by the study of Mujib *et al.*⁴⁷ which reported that p53 protein can serve as a marker for cancer cell death via apoptosis due to non-contact electric fields exposure. Moreover, the hollow cavities, also called dead cell islands, with cells debris, may be the spaces that used to be the place for mammary tumor cells that then died through apoptosis^{48,49}, since caspase-3 was found in this place in the IT group (Figure 5f). In addition, within these dead cell islands, the expression level of caspase-3 was very low⁴⁹, as we found in the IT group (Figure 6d).

The lower expression of caspase-3 in the IT group (Figure 6d) may suggest the mechanism of mammary tumor cell death under exposure to non-contact electric fields of ECCT through caspase-independent cell death. Meggyeshaz *et al.*⁵⁰ reported tumor destruction by electric fields exposure and concomitant heat with elevated DNA fragmentation and low level of caspase-3 expression localized to inflammatory cells which dominantly follows a caspase-independent pathway. Meanwhile, higher caspase-3 expression in INT group (Figure 6d) suggested that higher apoptosis events may have occurred in this group, resulted from oxidative stress due to elevated reactive oxygen species in mammary tumors^{51,52}. Although apoptosis occurred in the mammary tumors, the tumor cell death rate was relatively much lower than the proliferation rate, so that mammary tumor size increased⁵³, as seen in the INT group (Figure 2a and Figure 2b).

The type of receptor possessed by breast cancer cells in this study was the ErbB2 receptor (Figure 5c and Figure 5d). The generation of mammary tumors with ErbB2 receptors were the results of carcinogenesis due to DMBA administration into the animal tumor model⁵⁴. Despite such insignificant results

Figure 8. Number of CD68-, percentage of CD4- and CD8-positive cells, and CD4/CD8 ratio in mammary tumors. (a) Number of CD68-positive macrophages, (b) percentage of CD4-positive T cells, (c) percentage of CD8-positive T cells, (d) ratio of CD4/CD8. Error bars represent the standard error of the mean. Different letters (a, b) on top of the charts indicate significant differences ($p < 0.05$).

in the expression level of ErbB2 between INT and IT groups (Figure 6c), tumor size in the INT group continued to increase from baseline, and its tumor growth rate was higher than that in the IT group (Figure 2a). In addition, the number of mitotic figures and the percentage of PCNA-positive tumor cells in the INT group were also higher than those in the IT group (Figure 6a and Figure 6b, respectively). This suggests that INT group experienced higher tumor cell proliferation than IT group. Moreover, the higher number of CD68⁺ macrophages as one of inflammatory cells seen in the tumor microenvironment of INT group (Figure 7a and Figure 8a) was capable to induce tumor growth by contributing to cellular migration of tumor cells and metastasis^{12,14,55}. Ong *et al.*⁵⁶ also reported that rat mammary carcinoma had significantly higher percentage of CD68⁺ macrophages compared to benign mammary tumors with a lower growth rate. CD68⁺ macrophages that infiltrate to the necrotic areas of INT group (Figure 7a) may be considered as M2 macrophages which play a role in tumor growth and provide a poor prognosis^{12,56,57}. In addition, increased M2 macrophages or Tumor-associated macrophages (TAMs) infiltration in mammary tumors would be accompanied by an increase in tumor size^{14,15,58}, as shown in the INT group (Figure 2a and Figure 8a). Conversely, the lower the number of infiltrating TAMs in rat mammary tumors, the smaller the size of the tumors^{12,15}, as shown in the IT group (Figure 2a and Figure 8a). Moreover, the expressions of chemokine (C-C motif) ligand 2 (CCL2) and interleukin 18

(IL18) that have main roles in the development of metastatic breast cancer and excessive angiogenesis by amplifying M2 macrophage polarization^{59,60} were down-regulated under the exposure to non-contact electric fields of ECCT at 150 kHz frequency²². Similarly, non-contact electric field (100 kHz) exposure in our study may also inhibit the amplification of M2 macrophage polarization resulted in the negative mammary tumor growth rate in rats of IT group (Figure 2a). The infiltration of CD68⁺ macrophages in mammary tumors of IT group suggested that exposure to non-contact electric fields of ECCT may direct macrophages, likely of M1 phenotype, to the tumor areas^{11,21} and then engulf and digest dead cell, debris and tumor cells^{12,61} that eventually produced hollow cavities^{48,49} as seen in Figure 7b. This macrophage-mediated clearance of dead cells was occurred in caspase-3-independent death⁶², since the IT group had lower expression of caspase-3 than the INT group (Figure 6d).

Immunogenic tumor cell death induced by the electric fields exposure in the IT group can also be caused by the interaction of lymphocyte with receptors on membrane of mammary tumor cells with the production of cytokines^{21,46}. The distribution of CD4⁺ and CD8⁺ T cells within blood vessels and mammary tumors of IT group revealed that non-contact electric fields of ECCT may direct the spreading of these lymphocytes from blood vessels to tumor areas^{11,19,21}, including

necrotic areas in it (Figure 7d and Figure 7f). The presence of CD4⁺ T cells in mammary tumors would induce the activation of CD8⁺ T cells to kill the tumors^{12,63,64} as an adaptive immune response^{12,13,65}. However, since the percentage of CD4⁺ T cells in INT and IT groups was not significantly different (Figure 8b), the presence of CD4⁺ T cells in tumor areas may not always indicate tumor elimination, instead they may indicate tumor progression^{12,30,66}. Moreover, electric field exposure can decrease CD4⁺ T cells polarisation¹⁹, resulted in the lower percentage of CD4⁺ T cells in IT group compared to INT group (Figure 8b). Conversely, the presence of CD8⁺ T cells in the necrotic areas which characterized by inflammation^{33,34}, indicated their role in the inflammatory response to clear dead cells^{12,67} recognized as foreign agents¹³. In addition, a higher percentage of CD8⁺ T cells in IT group (Figure 8c) indicated increased anti-tumor immunity and immunological ability against tumor progression^{66,68,69}. CD8⁺ T cells may release granzymes during granule exocytosis when tumor cells are marked for elimination^{12,13,70}, and the granzymes may induce caspase-independent apoptotic pathways^{71,72} which support the presence of hollow cavities in the tumor areas with low caspase-3 expression⁴⁹ as seen in the IT group (Figure 5f and Figure 6d).

The ratio of CD4/CD8 is correlated to clinical aspects and could be used as prognostic factor for breast cancer, where higher CD4/CD8 ratio in INT group (Figure 8d) was correlated to mammary tumor progression⁷³. Conversely, the higher positive area of CD8⁺ T cells and the lower ratio of CD4/CD8 in IT group suggested a good prognosis in rat breast cancer^{12,66,73} with non-contact electric field exposure. Further study will be conducted to investigate the potency of non-contact electric fields exposure in wound healing upon breast cancer treatment by observing the presence of cytokines, as well as the migration of immune cells and fibroblast that regulate wound healing^{20,41,42} by using tissue samples from the IT group. Based on the evidence of its efficacy and safety on preclinical studies, the 100 kHz non-contact electric field has been used in phase I clinical trials of ECCT for healthy volunteers.

Conclusions

In summary, low intensity (18 Vpp) and intermediate frequency (100 kHz) non-contact electric fields exposure showed inhibition of mammary tumor growth in rats by inducing CD8⁺ T cells that lead to tumor cells death supported by decreased CD4/CD8 ratio. It also showed inhibition of M2 macrophage polarization resulted in the negative tumor growth rate. The proposed therapy gives good prognosis in rats with mammary tumors and potentially helps in wound healing of the tumor aftermath during the treatment.

Data availability

Underlying data

Open Science Framework: Cytotoxic T cells response with decreased CD4/CD8 ratio during mammary tumors inhibition in rats induced by non-contact electric fields, <https://doi.org/10.17605/OSF.IO/3P794>⁷⁴ (registered on 11th December 2020: <https://osf.io/b6d4m>).

This project contains the following underlying data:

- Ethical Clearance document
- Growth rate of mammary tumors and wound healing data
- IHC scoring data
- Raw images for IHC and hematoxylin figures
- Statistical analysis with GraphPad Prism ver.8.4.3

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

We would like to thank Mrs. Ardaning Nurliani from Faculty of Biology of Universitas Gadjah Mada for helping the antibodies procurement.

References

1. Siegel R, Naishadham D, Jemal A: **Cancer Statistics, 2012**. *CA Cancer J Clin*. 2012; **62**(1): 10–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Bray F, Ferlay J, Soerjomataram I, et al.: **Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries**. *CA Cancer J Clin*. 2018; **68**(6): 394–424.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Moo TA, Sanford R, Dang C, et al.: **Overview of Breast Cancer Therapy**. *PET Clin*. 2018; **13**(3): 339–354.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Andersen K, Kehlet H: **Persistent pain after breast cancer treatment: a critical review of risk factor and strategies for prevention**. *J Pain*. 2011; **12**(7): 725–746.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Bukowski K, Kciuk M, Kontek R: **Mechanisms of Multidrug Resistance in Cancer Chemotherapy**. *Int J Mol Sci*. 2020; **21**(9): 3233.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Conze D, Weiss L, Regen P, et al.: **Autocrine production of interleukin 6 causes multidrug resistance in breast cancer cells**. *Cancer Res*. 2001; **61**(24): 8851–8858.
[PubMed Abstract](#)
7. Kumar M, Nagpal R, Hemalatha R, et al.: **Targeted cancer therapies: the future of cancer treatment**. *Acta Biomed*. 2012; **83**(3): 220–233.
[PubMed Abstract](#)
8. Bokstein F, Blumenthal D, Limon D, et al.: **Concurrent Tumor Treating Fields (TTFields) and Radiation Therapy for Newly Diagnosed Glioblastoma: A Prospective Safety and Feasibility Study**. *Front Oncol*. 2020; **10**: 411.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Kirson ED, Schneiderman RS, Dbalý V, et al.: **Chemotherapeutic treatment efficacy and sensitivity are increased by adjuvant alternating electric fields (TTFields)**. *BMC Med Phys*. 2009; **9**: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Jo Y, Hwang SG, Jin YB, et al.: **Selective toxicity of tumor treating fields to melanoma: an in vitro and in vivo study**. *Cell Death Discov*. 2018; **4**: 46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Alamsyah F, Ajrina IN, Dewi FN, et al.: **Antiproliferative effect of electric fields**

- on breast tumor cells *in vitro* and *in vivo*. *Indones J Cancer Chemoprev*. 2015; 6(3): 71–77.
[Publisher Full Text](#)
12. Gonzalez H, Hagerling C, Werb Z: **Roles of the immune system in cancer: from tumor initiation to metastatic progression.** *Genes Dev*. 2018; 32(19–20): 1267–1284.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Maimela NR, Liu S, Zhang Y: **Fates of CD8+ T cells in Tumor Microenvironment.** *Comput Struct Biotechnol J*. 2019; 17: 1–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Yang M, Li Z, Ren M, *et al.*: **Stromal Infiltration of Tumor-Associated Macrophages Conferring Poor Prognosis of Patients with Basal-Like Breast Carcinoma.** *J Cancer*. 2018; 9(13): 2308–2316.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. Medrek C, Ponten F, Jirstrom K, *et al.*: **The Presence of Tumor Associated Macrophages in Tumor Stroma as A Prognostic Marker for Breast Cancer Patients.** *BMC Cancer*. 2012; 12: 306.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Donzelli S, Milano E, Pruszkowski M, *et al.*: **Expression of ID4 protein in breast cancer cells induces reprogramming of tumour associated macrophages.** *Breast Cancer Res*. 2018; 20(1): 59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Dimberu PM, Leonhardt RM: **Cancer immunotherapy takes a Multi-Faceted Approach to Kick the immune System into Gear.** *Yale J Biol Med*. 2011; 84(4): 371–380.
[PubMed Abstract](#) | [Free Full Text](#)
 18. Petri AK, Schmiedchen K, Stunder D, *et al.*: **Biological Effects of Exposure to Static Electric Fields in Humans and Vertebrates: A Systematic Review.** *Environ Health*. 2017; 16(1): 41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Arnold CE, Rajnicek AM, Hoare JJ, *et al.*: **Physiological strength electric fields modulate human T cell activation and polarisation.** *Sci Rep*. 2019; 9(1): 17604.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Hoare JJ, Rajnicek AM, McCaig CD, *et al.*: **Electric fields are novel determinants of human macrophage functions.** *J Leukoc Biol*. 2016; 99(6): 1141–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Voloshin T, Kaynan N, Davidi S, *et al.*: **Tumor treating fields (TTFields) induce immunogenic cell death resulting in enhanced antitumor efficacy when combined with anti PD 1 therapy.** *Cancer Immunol Immunother*. 2020; 69(7): 1191–1204.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Pratiwi R, Antara NY, Fadliansyah LG, *et al.*: **CCL2 and IL18 expressions may associate with the anti-proliferative effect of noncontact electro capacitive cancer therapy *in vivo*.** *F1000 Res*. 2020; 8: 1770.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Abbasalipourkabir R, Dehghan A, Salehzadeh A, *et al.*: **Induction of mammary gland tumor in female Sprague Dawley rats with LA7 cells.** *Afr J Biotechnol*. 2010; 9(28): 4491–4498.
[Reference Source](#)
 24. Akla B, Monteil J, Paraf F, *et al.*: **A New Orthotopic Model of Human Breast Cancer in Immunocompetent Rats.** *Anticancer Res*. 2003; 23(5A): 3761–6.
[PubMed Abstract](#)
 25. Costa E, Ferreira-Gonçalves T, Chasqueira G, *et al.*: **Experimental Models as Refined Translational Tools for Breast Cancer Research.** *Sci Pharm*. 2020; 88(32): 1–29.
[Publisher Full Text](#)
 26. Kerdelhue B, Forest C, Coumoul X: **Dimethyl-Benz(a)anthracene: A mammary carcinogen and a neuroendocrine disruptor.** *Biochimie Open*. 2016; 3: 49–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Liu Y, Yin T, Feng Y, *et al.*: **Mammalian models of chemically induced primary malignancies exploitable for imaging-based preclinical theragnostic research.** *Quant Imaging Med Surg*. 2015; 5(5): 708–729.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Lee SH, Kim YS, Han W, *et al.*: **Tumor growth rate of invasive breast cancers during wait times for surgery assessed by ultrasonography.** *Medicine (Baltimore)*. 2016; 95(37): e4874.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Bancroft JD, Cook HC: **Manual of Histology Techniques and Their Diagnostic Application.** Churchill Livingstone, London, 1994.
[Reference Source](#)
 30. Sawe RT, Kerper M, Badve S, *et al.*: **Aggressive breast cancer in western Kenya has early onset, high proliferation, and immune cell infiltration.** *BMC Cancer*. 2016; 16: 204.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Ghanavat M, Ebrahimi M, Rafieemehr H, *et al.*: **Thrombocytopenia in solid tumors: Prognostic significance.** *Oncol Rep*. 2019; 13(1): 413.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Feng M, Feng C, Yu Z, *et al.*: **Histopathological Alterations During Breast Carcinogenesis in A Rat Model Induced by 7,12-Dimethylbenz(a)anthracene and Estrogen-Progestogen Combinations.** *Int J Clin Exp Med*. 2015; 8(1): 346–357.
[PubMed Abstract](#) | [Free Full Text](#)
 33. Koerner F: **Diagnostic Problems in Breast Pathology: 1st Edition.** Boston: Saunders, 2009.
[Reference Source](#)
 34. Davidovich P, Kearney CJ, Martin SJ: **Inflammatory outcomes of apoptosis, necrosis and necroptosis.** *Biol Chem*. 2014; 395(10): 1163–1171.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Elmore S: **Apoptosis: A Review of Programmed Cell Death.** *Toxicol Pathol*. 2007; 35(4): 495–516.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Trisciuglio L, Bianchi ME: **Several Nuclear Events during Apoptosis Depend on Caspase-3 Activation but Do Not Constitute a Common Pathway.** *PLoS One*. 2009; 4(7): e6234.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Kirson ED, Gurvey Z, Schneiderman R: **Disruption of cancer cell replication by alternating electric fields.** *Cancer Res*. 2004; 64(9): 3288–3295.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Swanson KD, Lok E, Wong ET: **An Overview of Alternating Electric Fields Therapy (NovoTTF Therapy) for the Treatment of Malignant Glioma.** *Curr Neurol Neurosci Rep*. 2016; 16(1): 8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Kole AJ, Kole L, Moran MS: **Acute radiation dermatitis in breast cancer patients: challenges and solutions.** *Breast Cancer (Dove Med Press)*. 2017; 9: 313–323.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Payne WG, Naidu DK, Wheeler CK: **Wound Healing in Patients with Cancer.** *J Plastic Surg*. 2008; 68–90.
 41. Funk RHW: **Endogenous electric fields as guiding cue for cell migration.** *Front Physiol*. 2015; 6: 143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Long Y, Wei H, Li J: **Effective Wound Healing Enabled by Discrete Alternative Electric Fields from Wearable Nanogenerators.** *ACS Nano*. 2018; 12(12): 12533–12540.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Gun SY, Lee SWL, Sieow JL, *et al.*: **Targeting immune cells for cancer therapy.** *Redox Biol*. 2019; 25: 101174.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Zuccari DAP, Pavam MV, Terzian ACB, *et al.*: **Immunohistochemical evaluation of e-cadherin, Ki-67 and PCNA in canine mammary neoplasias: Correlation of prognostic factors and clinical outcome.** *Pesq Vet Bras*. 2008; 28(4): 207–215.
[Publisher Full Text](#)
 45. Essers J, Theil AF, Baldeyron C, *et al.*: **Nuclear Dynamics of PCNA in DNA Replication and Repair.** *Mol Cell Biol*. 2005; 25(21): 9350–9359.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Giladi M, Schneiderman RS, Voloshin T, *et al.*: **Mitotic Spindle Disruption by Alternating Electric Fields Leads to Improper Chromosome Segregation and Mitotic Catastrophe in Cancer Cells.** *Sci Rep*. 2015; 5: 18046.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Mujib SA, Alamsyah FA, Taruno WP: **Cell Death and Induced p53 Expression in Oral Cancer, HeLa, and Bone Marrow Mesenchyme Cells under the Exposure to Noncontact Electric Fields.** *Integr Med Int*. 2017; 4: 161–170.
[Publisher Full Text](#)
 48. Debnath J, Mills KR, Collins NL, *et al.*: **The Role of Apoptosis in Creating and Maintaining Luminal Space within Normal and Oncogene-Expressing Mammary Acini.** *Cell*. 2002; 111(1): 29–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Liao DJ, Dickson RB: **Cell Death in MMTV-c-myc Transgenic Mouse Mammary Tumors May Not Be Typical Apoptosis.** *Lab Invest*. 2003; 83(10): 1437–1449.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Meggyeshazi N, Andocs G, Balogh L, *et al.*: **DNA fragmentation and caspase-independent programmed cell death by modulated electrohyperthermia.** *Strahlenther Onkol*. 2014; 190(9): 815–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Liou GY, Storz P: **Reactive oxygen species in cancer.** *Free Radic Res*. 2010; 44(5): 479–496.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Olsson M, Zhivotovsky B: **Caspases and cancer.** *Cell Death Differ*. 2011; 18(9): 1441–1449.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Parton M, Dowsett M, Smith I: **Studies of Apoptosis in Breast Cancer.** *BMJ*. 2001; 322(7301): 1528–1532.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. Ma Z, Kim YM, Howard EW, *et al.*: **DMBA promotes ErbB2 mediated carcinogenesis via ErbB2 and estrogen receptor pathway activation and genomic instability.** *Oncol Rep*. 2018; 40(3): 1632–1640.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. Vano YA, Oudard S, By MA, *et al.*: **Optimal cut-off for neutrophil-to-lymphocyte ratio: Fact or Fantasy? A prospective cohort study in metastatic cancer patients.** *PLoS One*. 2018; 13(4): e0195042.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. Ong CB, Brandenberger C, Kiupel M, *et al.*: **Immunohistochemical Characterization and Morphometric Analysis of Macrophages in Rat**

- Mammary Tumors.** *Vet Pathol.* 2015; **52**(2): 414–418.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Martinez FO, Gordon S: **The M1 and M2 Paradigm of Macrophage Activation: Time for Reassessment.** *F1000Prime Rep.* 2014; **6**(13): 1–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Yang J, Li X, Liu X, *et al.*: **The Role of Tumor-Associated Macrophages in Breast Carcinoma Invasion and Metastasis.** *Int J Clin Exp Pathol.* 2015; **8**(6): 6656–6664.
[PubMed Abstract](#) | [Free Full Text](#)
 59. Ruytinx P, Proost P, Van Damme J, *et al.*: **Chemokine-induced macrophage polarization in inflammatory conditions.** *Front Immunol.* 2018; **9**: 1930.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Kobori T, Hamasaki S, Kitaura A, *et al.*: **Interleukin-18 Amplifies Macrophage Polarization and Morphological Alteration, Leading to Excessive Angiogenesis.** *Front Immunol.* 2018; **9**: 334.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Hirayama D, Iida T, Nakase H: **The Phagocytic Function of Macrophage-Enforcing Innate Immunity and Tissue Homeostasis.** *Int J Mol Sci.* 2018; **19**(1): 92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Turner C, Devitt A, Parker K, *et al.*: **Macrophage-mediated clearance of cells undergoing caspase-3-independent death.** *Cell Death Differ.* 2003; **10**(3): 302–312.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Ahrends T, Borst J: **The opposing roles of CD4⁺ T cells in anti-tumour immunity.** *Immunology.* 2018; **154**(4): 582–592.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Tay RE, Richardson EK, Toh HC: **Revisiting the role of CD4⁺ T cells in cancer immunotherapy-new insights into old paradigms.** *Cancer Gene Ther.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. Vestweber D: **Lymphocyte trafficking through blood and lymphatic vessels: more than just selectins, chemokines and integrins.** *Eur J Immunol.* 2003; **33**(5): 1361–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Huang Y, Ma C, Zhang Q, *et al.*: **CD4⁺ and CD8⁺ T cells have opposing roles in breast cancer progression and outcome.** *Oncotarget.* 2015; **6**(19): 17462–17478.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 67. Rock KL, Lai JJ, Kono H: **Innate and adaptive immune responses to cell death.** *Immunol Rev.* 2011; **243**(1): 191–205.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 68. Shindo G, Endo T, Onda M, *et al.*: **Is the CD4/CD8 ratio an effective indicator for clinical estimation of adoptive immunotherapy for cancer treatment?** *J Cancer Therapy.* 2013; **4**(8): 1382–1390.
[Publisher Full Text](#)
 69. Tsukumo SI, Yasutomo K: **Regulation of CD8⁺ T cells and antitumor immunity by Notch signaling.** *Front Immunol.* 2018; **9**: 101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 70. Chowdhury D, Lieberman J: **Death by a Thousand Cuts: Granzyme Pathways of Programmed Cell Death.** *Annu Rev Immunol.* 2008; **26**: 389–420.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Martinvalet D, Zhu P, Lieberman J: **Granzyme A Induces Caspase-Independent Mitochondrial Damage, a Required First Step for Apoptosis.** *Immunity.* 2005; **22**(3): 355–370.
[PubMed Abstract](#) | [Publisher Full Text](#)
 72. Martinvalet D: **Mitochondrial Entry of Cytotoxic Proteases: A New Insight into the Granzyme B Cell Death Pathway.** *Oxid Med Cell Longev.* 2019; **2019**: 9165214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Yang X, Ren H, Sun Y, *et al.*: **Prognostic significance of CD4/CD8 ratio in patients with breast cancer.** *Int J Clin Exp Pathol.* 2017; **10**(4): 4787–4793.
[Reference Source](#)
 74. Alamsyah F: **Cytotoxic T cells response with decreased CD4/CD8 ratio during mammary tumors inhibition in rats induced by non-contact electric fields.** 2020.
<http://www.doi.org/10.17605/OSF.IO/3P794>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 03 March 2021

<https://doi.org/10.5256/f1000research.30917.r79324>

© 2021 Elfahmi E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Elfahmi Elfahmi

School of Pharmacy, Bandung Institute of Technology, Bandung, Indonesia

The article consists a quite good scientific background and methodology and has an innovation to be contributed in the treatment of cancer, however other clinical studies should be conducted. The quality of works and article is suitable for indexing, however, some comments should be addressed by author for the statement or question as follow:

1. Some data presented in the results showed a quite big deviation, are there any outliers in the 6 replicates data which contributed for this deviation? If it's possible that the authors could explain this clearly, it would then strengthen the paper.
2. Although the electrical field is not contacted with the animal models, it is placed in the closed chamber. Could the authors explain in scientific view whether there are no side effects of the electric field?
3. The condition parameter of electric field used in this study is only point. Is it an optimum condition? If yes, the proven study for the optimalization should be included.
4. If there is toxicity data of the instrument published, it would be better if included.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 30 Mar 2021

Firman Alamsyah

1. The deviation value that is quite large in some data shows a quite different response from each individual rat to the electric field exposure. For example, in the SGR data (Figure 2), there is indeed 1 outlier data in each of the IN and INT groups. This may occur because the estrous cycle that stimulates prolactin can be different in each individual mouse which can affect differences in the growth rate of mammary tumors in rat (Nandi et al., 1995; Schedin et al., 2000). In addition, different levels of stress due to therapy in individual cages could affect differences in the development of mammary tumors in individual rat (Hermes et al., 2009; De la Roca-Chiapas, et al., 2016). We have added this argument in the revision of our article.

2. We have explained the safety of electric field of ECCT to mammary and skin tissues of animal models in the second paragraph of the Introduction as reported in our previous publication. In addition, ECCT has passed the phase I clinical trials for healthy volunteers as stated in the last sentence of the Discussion. The safety of ECCT in normal tissue is determined by the membrane potential of the tissue which is higher than that in cancer cells (Yang and Brackenbury, 2013; Kadir et al., 2018). External electric field exposure cannot penetrate normal cell membranes due to the higher membrane potential (Kirson et al., 2004; Yang and Brackenbury, 2013; Kadir et al., 2018) and only causes rotational motion of the cell organelles (Kirson et al., 2004). On the other hand, external electric field exposure can penetrate cancer cell membranes with lower cell membrane potential, thereby interfering with mitosis (Kirson et al., 2004; Yang and Brackenbury, 2013; Kadir et al., 2018). However, not all normal cells have a higher membrane potential than cancer cells. For example, liver cell has the same membrane potential as MCF-7 cancer cell (Yang and Brackenbury, 2013; Kadir et al., 2018). Even so, we found that liver function was not impaired based on alanine transferase activity and bilirubin levels in blood plasma. The data of the safety of ECCT exposure in rat and healthy volunteers will be published separately this year.

3. We have tried various frequencies (100, 150, and 200 kHz) and intensities (10 Vpp, 18 Vpp and 30 Vpp) of the ECCT electric field in preclinical studies and found that a frequency of 100 kHz and an intensity of 18 Vpp are the optimum electric field exposure for breast cancer

therapy.

4. ECCT toxicity data on the liver, kidney and hematological profiles will be published separately this year.

Competing Interests: No competing interests were disclosed.

Reviewer Report 23 February 2021

<https://doi.org/10.5256/f1000research.30917.r79321>

© 2021 Keserci B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bilgin Keserci

School of Medical Sciences, Universiti Sains Malaysia, Kelantan, Malaysia

Even though the study results are quite impressive, I have some additional issues to be considered/corrected before this paper is indexed.

First, the manuscript needs serious language editing. I strongly advise them to get a professional editing service when revising.

Second, the below concerns must be answered and explained well in the manuscript before being indexed:

On page 4 it says, *"The first 6 rats to develop tumors were assigned to IT group, and then 6 more rats were assigned to INT group. Not all rats grew tumors after DMBA administration. All healthy, normal rats in the placebo (NIT) group were given corn oil as DMBA (7,12-dimethylbenz[a]anthracene) solvent."*

Questions:

- What does this mean? How many rats developed tumors in the IT and INT groups? Why were all the rats that developed the tumor first included in the IT group rather than equally distributed to INT and IT?
- Were the rats in the NIT group taken from the non tumor growing induced group after being given corn oil?

On page 5 it says, *"Three nodules from three different rats were collected from each INT and IT groups, while one normal breast tissue was collected from each NINT and NIT groups."*

Question:

- Why not all? How are the rats chosen?

On page 7 it says, *"In addition, the percentage of positively stained areas of PCNA in INT group ($36.16 \pm 5.16\%$) was significantly higher than that in INT group ($23.65 \pm 3.84\%$, Figure 6b)."*

Comment:

- 'INT' in bold above is supposed to be 'IT' group. Please correct it.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical imaging, cancer therapy, MRI, image guided therapies

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 27 Feb 2021

Firman Alamsyah

1. We will consider getting a professional language editing service to revise our manuscript.
2. The rats were randomly selected for mammary tumor induction with DMBA administration. One month after DMBA administration, the tumors began to grow with a different time of appearance which was between 38-63 days. Since we used 6 biological replicates for each group based on Federer formula, 12 rats were used for IT and INT groups. Then, we selected the first 6 rats that developed tumors to be assigned to the IT group, so that it was technically easier to carry out the research and avoided mistakes in operating the ECCT device. We did not equally distribute the induced rats to both groups to avoid human errors in operating the ECCT, since we had to turn on and off the device twice every day for the IT group, while not for the INT group.
3. The rats in the NIT group were normal rats with corn oil administration. They did not

experience any DMBA administration.

4. We used 3 nodules from 3 rats for each IT and INT group, since this sample size was sufficiently representative for the IHC method. We randomly selected the rats with nodules for histopathological analysis.

5. Yes, we made a mistake for that sentence. We will correct it.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research