STUDY PROTOCOL

Stage 1 Registered Report: Testing mediumship accuracy with a triple-blind protocol [version 1; peer review: awaiting peer review]

Fernando Sinesio¹, Laura Liberale¹, Ines Testoni², Patrizio E. Tressoldi³

¹Gruppo di Ricerca Italiano sulla Medianità, Padova, Italy

²FISPPA, Padova University, Padova, Italy

³Science of Consciousness Research Group - Studium Patavinum, Padova University, Padova, Italy

V1 First published: 06 May 2021, 10:351
<https://doi.org/10.12688/f1000research.52208.1>

Latest published: 21 May 2021, 10:351
<https://doi.org/10.12688/f1000research.52208.2>

Abstract

Even if mediumship is a practice that probably originated at the outset of human history and is embedded in many cultural and religious traditions, scientific investigations into mediumship are quite recent. In this study, we aim to investigate if self-defined mediums can retrieve information about specific deceased persons in an unconventional way, that is, without retrieving the information from the parents or friends of the deceased person or by other conventional means. To our knowledge, this is the first registered report related to this phenomenon.

Our experimental design will be triple-blinded: the mediums and their interviewer will only know the first name of the deceased person and of the requesting person. The parents or the close friends of the deceased person will not interact in any way with the medium. Furthermore, they will be blind about which of two lists of information (readings) paired with deceased persons of the same gender is related to their parents or friend. Accuracy of information will be analysed by three criteria: (1) percentage of correct identification of the readings related to the requested person; (2) comparison of the global scores (on a scale 1 to 6) assigned to the intended readings with that of control readings; and (3) comparison of the difference between the percentage of correct minus incorrect information identified in the intended readings with that of the control readings.

Keywords

registered report, mediumship, triple-blind protocol, anomalous cognition

Open Peer Review

Approval Status ?

1

version 2

(revision)

21 May 2021

version 1

06 May 2021

1. **Alexander Moreira-Almeida**, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora, Brazil

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Patrizio E. Tressoldi (patrizio.tressoldi@unipd.it)

Author roles: **Sinesio F:** Data Curation, Investigation, Resources, Validation; **Liberale L:** Conceptualization, Data Curation, Funding Acquisition, Project Administration, Validation, Writing – Review & Editing; **Testoni I:** Funding Acquisition, Writing – Review & Editing; **Tressoldi PE:** Conceptualization, Data Curation, Formal Analysis, Methodology, Project Administration, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Fundação Bial.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Sinesio F *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sinesio F, Liberale L, Testoni I and Tressoldi PE. **Stage 1 Registered Report: Testing mediumship accuracy with a triple-blind protocol [version 1; peer review: awaiting peer review]** F1000Research 2021, 10:351 <https://doi.org/10.12688/f1000research.52208.1>

First published: 06 May 2021, 10:351 <https://doi.org/10.12688/f1000research.52208.1>

Introduction

The scientific study of mediumship, notwithstanding its very old origin and its widespread diffusion in the Eastern and Western society, is relatively recent (Beischel, 2018). From a scientific point of view, the main questions are related to the level of accuracy of information given by mediums and its origin.

Consultation with a medium typically involves the presence of a person, defined as a 'sitter', usually a close relative or friend of the deceased person, who wants to know if the deceased person is still living after his/her death. In this setting, it is quite likely that the source of the information is the sitter him/herself, who could consciously or unconsciously suggest it to the medium through what is called 'cold reading' (Roe & Roxburgh, 2013), that is, the transmission of verbal and non-verbal information to the medium.

In order to eliminate this source of information, research protocols must ensure a complete separation and visually and auditory isolation between the sitter and the medium.

Given that the only person who can judge the accuracy of information is the sitter, in order to prevent confirmatory bias, i.e. judging if particular information is correct even if it is wrong because of their strong need of proof that their loved one is still alive (Snyder & Swann, 1978; Rabin & Schrag, 1999), research protocols must require the sitter to be given at least two anonymous lists of information (readings) related to two deceased persons for evaluation: one related to their requested parent or friend and the second one related to another person paired with the same gender.

A meta-analysis of all available studies completed between 2001 and 2019 that used research protocols with different levels of blindness, revealed an accuracy of 6-14% more than expected by chance with better performance of mediums who were preliminary screened about their abilities (Sarraf, Woodley of Menie and Tressoldi, 2020).

In this study, we aim to add a further contribution to this line of investigation by adopting the best research practices in order to prevent contamination of the results by so-called questionable research practices (John, Loewenstein & Prelec, 2012; Banks et al., 2016).

Hypotheses

The main hypotheses are:

- The percentage of correctly identified intended readings, that is those related to the requested person, will exceed the control ones at a statistical level (see section *Statistical analyses*);
- The global scores assigned to the intended readings will be higher with respect to the control ones at a statistical level;
- The difference between the percentage of correct information minus the percentage of wrong information of the intended readings will be higher with respect to the control ones at a statistical level.

Methods

Ethical approval

This study has been approved by the Comitato etico della ricerca psicologica (Ethical Committee of Psychological Research) of Padova University (Protocol no. 3670). Both mediums and sitters will be requested to read and confirm orally the protocol as informed consent.

Participants

Eight professional self-defined mediums will be included in the study. All have been purposively selected by the authors after testing their accuracy skills in a previous study (Tressoldi et al. 2020).

Eight sitters will be contacted by email mailing lists by the authors, from people with a serious interest in having a reading with respect to a deceased relative or friend, regardless of cause of death or length of time since the death. The only exclusion criteria will be a not well-defined relationship with the requested deceased person that could preclude a complete knowledge of him/her and consequently a difficulty in the evaluation of information given by the mediums.

Each medium will be asked to contribute a reading for each of the eight deceased persons for a total of 64 readings.

Procedure

Each trial, that is each consultation with the medium and each task assigned to the sitters, will comprise two readings related to deceased individuals of the same gender, male or female, as the only common characteristic.

The experimental procedure will be similar to that adopted by Tressoldi, Liberale, Sinesio et al. (2020) which was a reduced version of that presented by Beischel (2007), with some variants aimed at further excluding a mental connection with the sitter.

Furthermore, in order to keep the task difficulty as similar as possible for all mediums, they were requested to obtain response to a set of ten identical questions (see *Extended data*) plus two or three further questions requested by the sitters to increase their confidence of the authenticity of the contact with the requested deceased.

The steps of the procedure of each session are the follow:

1. To a pair of sitters interested in having a free consultation for a deceased loved parent or friend of the same gender, they will be asked to provide the deceased first name, and the time since the deceased passed.
2. This information will be kept by one of the co-authors, who will be called 'research assistant B' (raB).
3. The medium will be contacted for the consultation by another co-author, who will be called 'research assistant A' (raA).
4. On the day of consultation, raA will contact the medium either via Skype or WhatsApp and for each of the two paired deceased persons will give the medium only the deceased and the sitter's first name (without the family name) sent by raB.
5. For each of the two paired deceased persons, the medium will be required to respond either in oral or written form to the ten fixed questions (*Extended data*) and to provide all information pertinent to the deceased's identification as asked by the sitter, as well as anything the purported deceased wishes to communicate to the sitter. For oral form, the information will be recorded by raA using an audio recorder.
6. At the conclusion of the reading, raA will transfer all information into a vertical list, excluding generic statements, e.g. "I love you" or "Don't worry about me, I'm well", and will send them to raB.
7. Once raB receive the two readings from raA, the information will be listed vertically for evaluation of each piece of information and for global evaluation (see the example in *Extended data*).
8. Afterwards, both reading A and B will be anonymously sent to the requesting sitter, providing assistance, if needed, for the evaluation procedure, i.e. sitters will be asked to go through the two lists of information and mark as 'perfectly correct', 'clearly wrong', 'somewhat correct' or 'I do not have information for evaluation'.

In summary, the research protocol will comprise three levels of blindness: for the deceased's identity: the medium, raA; and for the sitters which of the two readings is that of their deceased.

Evaluation of information from the two readings

All information listed will be classified in two categories: those comprising very detailed information, e.g. "he (the deceased person) got a car accident when very young"; "she said to you (to the sitter) do not sell her gold ring left in her bedroom", and those comprising less detailed information e.g. "he was a very nice person"; "she was very in love with her husband".

- Each item of information from the reading list related to the "detailed" category evaluated by the sitters as "Perfectly correct" or "Clearly wrong" will be given a value of 1 or -1, respectively.
- Each item of information from the reading list related to the "less detailed" category evaluated by the sitters as "Perfectly correct" or "Clearly wrong" will be given a value of .5 or -.5, respectively.
- Finally, each item evaluated by the sitters as "somewhat correct" will be given a value of 0.5 regardless of being in the detailed or less detailed category.

Subsequently, excluding information marked as “I don’t have information for evaluation”, the percentages of correct and incorrect information (ratio of this information with their total number) and their difference will be calculated. These differences could range from –100% if all wrong, to 100% if all correct.

The reliability of this scoring will be tested by an independent judge who will analyze all the readings following the same criteria.

As well as evaluating each item of information from the two readings, the sitter will also be asked to give a global evaluation of each reading using a scale from 0 (totally wrong) to 6 (excellent: effectively free of errors and containing compelling evidence of authentic communication), identical to the scale used by [Beischel \(2007\)](#). If both readings are given identical scores, the sitter will be asked to choose the one considered to be closest to their own deceased person.

Statistical analyses

Statistical Power: from the results observed by [Tressoldi, Liberale, Sinesio et al. \(2020\)](#), 64 readings are sufficient to detect a global score difference of 1.5 on a scale of 0-6, a 15% above chance of readings identification and 25% of correct information difference with respect to the incorrect one, that we consider to be the smallest effect size of interest, setting the statistical power to .80 and a directional hypothesis with $\alpha = .05$.

Data collection will end when 64 readings are evaluated by the sitters without any missing values.

In order to test the robustness of results, we will adopt both a frequentist and a Bayesian statistical approach with the statistical package JASP v.0.14.1 ([JASP Team, 2020](#)).

The percentage of correct readings identification will be analyzed using a frequentist binomial test with chance probability equal to $\frac{1}{2}$ and a Bayesian binomial test with beta prior parameter $a = 1$ and parameter $b = 1$.

The mean of the global score assigned to correctly identified readings will be compared with that of the incorrectly identified readings by using a one-sided permutation t -test estimated with 5000 bootstrap samples and the confidence intervals will be bias-corrected following the procedure of [Ho et al. \(2019\)](#) and with a Bayesian independent sample t -test with a Cauchy prior of .707.

The same statistics will be used for:

- the comparison of the global scores assigned to the correctly identified readings with that of the control readings;
- the comparison of the difference between the percentage of correct minus incorrect information identified in the correctly identified readings with that of the control readings.

The syntax of the statistical analyses will be available at Figshare.

Dissemination of information

Raw data and statistical analyses scripts will be made available open access in the extended data repository

Study status

At present we have completed the recruiting of the mediums and sitters.

Conclusion

With this study, we aim to contribute to the investigation of mediumship accuracy and the source of their information with a triple-blind protocol free of questionable research practice.

Even if our results cannot be generalized beyond the participants (mediums and sitters/deceased) enrolled in the study given their large individual differences, they can be an example of how it is possible to study a controversial phenomenon, such as mediumship, with the best methodological scientific tools.

Data availability

Underlying data

No data is associated with this article.

Extended data

Figshare: Mediumship, <https://doi.org/10.6084/m9.figshare.13311710.v4> (Tressoldi et al, 2020).

This project contains the following extended data:

- List of common questions
- Excerpt of the reading format

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgments

We thank the Proof Readings Service for English revisions.

References

- Banks GC, Rogelberg SG, Woznyj HM, et al.: **Evidence on questionable research practices: The good, the bad, and the ugly.** *Journal of Business and Psychology*. 2016; **31**(3): 323–338.
- Beischel J: **Contemporary Methods used in laboratory-based mediumship research.** *J Parapsychol*. 2007; **71**: 37–68.
- Beischel J: **Mental Mediumship Research** | *Psi Encyclopedia*. 2018. Retrieved November 19, 2020.
[Reference Source](#)
- Ho J, Tumkaya T, Aryal S, et al.: **Moving beyond P values: Everyday data analysis with estimation plots.** *Nat Methods*. 2019; 1548–7105.
[PubMed Abstract](#) | [Publisher Full Text](#)
- JASP Team: **JASP (Version 0.14.1).** [Computer software]. 2020.
[Reference Source](#)
- John LK, Loewenstein G, Prelec D: **Measuring the prevalence of questionable research practices with incentives for truth telling.** *Psychol Sci*. 2012; **23**(5): 524–532.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rabin M, Schrag J: **First impressions matter: a model of confirmatory bias.** *Quarterly J Econ*. 1999; **114**(1): 37–82.
- Roe CA, Roxburgh E: **An overview of cold reading strategies.** In: Moreman C (Ed.) *The Spiritualist Movement: Speaking with the Dead in America and around the World: Volume 2, Belief, Practice, and Evidence for Life after Death*. 2013. (pp. 177–203). Santa Barbara.
- Sarraf M, Woodley of Menie MA, Tressoldi P: **Anomalous information reception by mediums: A meta-analysis of the scientific evidence.** *Explore*. 2020.
[Publisher Full Text](#)
- Snyder M, Swann WB: **Behavioral confirmation in social interaction: from social perception to social reality.** *J Experi. Soc. Psychol*. 2020; **14**: 148–162.
- Tressoldi P, Liberale L, Sinesio F, et al.: **Mediumship accuracy: a quantitative and qualitative study with a triple-blind protocol.** *PsyArxiv*. 2020.
[Publisher Full Text](#)
- Tressoldi P, Pederzoli L, Testoni I: **Mediumship.** *figshare. Dataset*. 2020.
[Publisher Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research