RESEARCH ARTICLE

REVISED Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology [version 2; peer review: 1 approved, 1 approved with reservations]

Mauro Petrillo ¹, Carlo Brogna ², Simone Cristoni³, Maddalena Querci ¹, Ornella Piazza ⁴, Guy Van den Eede^{1,5}

¹European Commission, Joint Research Centre (JRC), Ispra, Italy

²CranioMed group srl, Montemiletto, Italy

³ISB Ion Source & Biotechnologies Srl, Bresso, Italy

⁴Department of Medicine and Surgery, University of Salerno, Baronissi, Italy

⁵European Commission, Joint Research Centre (JRC), Geel, Belgium

v2 First published: 11 May 2021, 10:370
<https://doi.org/10.12688/f1000research.52540.1>

Second version: 23 Jun 2021, 10:370
<https://doi.org/10.12688/f1000research.52540.2>

Latest published: 01 Jul 2021, 10:370
<https://doi.org/10.12688/f1000research.52540.3>

Abstract

Background

Scientific evidence for the involvement of human microbiota in the development of COVID-19 disease has been reported recently. SARS-CoV-2 RNA presence in human faecal samples and SARS-CoV-2 activity in faeces from COVID-19 patients have been observed.

Methods

Starting from these observations, an experimental design was developed to cultivate *in vitro* faecal microbiota from infected individuals, to monitor the presence of SARS-CoV-2, and to collect data on the relationship between faecal bacteria and the virus.

Results

Our results indicate that SARS-CoV-2 replicates *in vitro* in bacterial growth medium, that the viral replication follows bacterial growth and it is influenced by the administration of specific antibiotics. SARS-CoV-2-related peptides have been detected in 30-day bacterial cultures and characterised.

Discussion

Our observations are compatible with a 'bacteriophage-like' behaviour of SARS-CoV-2, which, to our knowledge has not been observed or described before. These results are unexpected and hint towards a novel hypothesis on the biology of SARS-CoV-2 and on the COVID-19 epidemiology. The discovery of possible new modes of action of SARS-

Open Peer Review

Approval Status

	1	2
version 3		
(revision)	view	
01 Jul 2021		
version 2		
(revision)	view	view
23 Jun 2021		
version 1		
11 May 2021	view	view

1. Kourosh Honarmand Ebrahimi ,
University of Oxford, Oxford, UK

2. Margarita Aguilera , University of
Granada, Granada, Spain

Any reports and responses or comments on the article can be found at the end of the article.

CoV-2 has far-reaching implications for the prevention and the treatment of the disease.

Keywords

SARS-CoV-2, COVID-19, gut microbiota

This article is included in the **Cell & Molecular Biology** gateway.

This article is included in the **Emerging Diseases and Outbreaks** gateway.

This article is included in the **Pathogens** gateway.

This article is included in the **Coronavirus** collection.

Corresponding author: Mauro Petrillo (mauro.petrillo@ext.ec.europa.eu)

Author roles: **Petrillo M:** Conceptualization, Data Curation, Formal Analysis, Methodology, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Brogna C:** Conceptualization, Formal Analysis, Methodology, Resources, Supervision, Writing – Review & Editing; **Cristoni S:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Resources, Validation, Writing – Review & Editing; **Querci M:** Conceptualization, Data Curation, Methodology, Project Administration, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Piazza O:** Conceptualization, Writing – Review & Editing; **Van den Eede G:** Conceptualization, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Funding was provided by the European Commission Joint Research Centre.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Petrillo M *et al.* This is an open access article distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Petrillo M, Brogna C, Cristoni S *et al.* **Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology [version 2; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:370 <https://doi.org/10.12688/f1000research.52540.2>

First published: 11 May 2021, 10:370 <https://doi.org/10.12688/f1000research.52540.1>

REVISED Amendments from Version 1

This version contains text additions following the suggestions and comments from Dr. Ebrahimi and Dr. Margarita Aguilera in their referee reports. These additions include:

- A new version of Figure 3.
- A paragraph in the conclusions discussing the relevance of identifying taxa, species or consortia that can be prone to act as interactor/receptor of virus, including those present in the upper respiratory tract.

Any further responses from the reviewers can be found at the end of the article

Introduction

Recent scientific articles and reviews^{1–3} discuss the relationship between gastrointestinal microbiota and COVID-19 disease. In particular, the prolonged presence of SARS-CoV-2 RNA in human faecal samples from COVID-19 patients has been reported⁴ and the potential role of orofecal transmission of SARS-CoV-2 has been examined in systematic reviews^{5,6} and open evidence briefs.^{7,8} Cases of SARS-CoV-2 detection in faecal samples from patient with typical symptoms of COVID-19 but negative to multiple SARS-CoV-2 real-time reverse transcription polymerase chain reaction (rRT-PCR) tests on oropharyngeal and nasopharyngeal swabs have been reported.⁹ SARS-CoV-2 faecal viral activity was depicted in association with gut microbiota composition in patients with COVID-19,¹⁰ and the replicating virus was detected in faeces.¹¹ At the same time, Wölfel *et al.*¹² observed high viral RNA concentration in stool samples, but reported isolation of infectious virus only from throat- and lung-derived samples, while Yao *et al.*¹³ had indication of viable SARS-CoV-2 particles in stool samples, denoting that the detailed biology of SARS-CoV-2 is not yet fully elucidated. Moreover, the interaction between SARS-CoV-2 and individual variable microbiota composition could drive the differential pathophysiological effects and severity of symptoms (Yeoh YH *et al.* 2021, Zuo T *et al.* 2021, Zuo T *et al.* 2020).

Our experiments further explored the relationship between COVID-19 disease and SARS-CoV-2 infected faeces to provide data relevant for pandemic understanding and disease management. The results however did not correspond with current thinking of the epidemiology of SARS-CoV-2 and, therefore, we believe a quick sharing with the scientific community of our findings is imperative.

Methods

The experimental design included:

- 1) the inoculation of NutriSelect™ Plus nutrient broth at 37°C, fit for the growth of more fastidious bacteria, with a faecal sample (stool) from one subject positive to SARS-CoV-2 and from a healthy individual (here called sample A and sample B, respectively) following written informed consent.
- 2) The assessment of the presence of SARS-CoV-2 RNA in both samples, after seven days of culture, using the Luminex technology; with confirmation of the presence of SARS-CoV-2 RNA in sample A, and of its absence in sample B.
- 3) Inoculation of sample B with the supernatant of sample A, obtained after centrifugation (hereafter called sample B_(A+)) and resuspension of the formed pellet (sample C).
- 4) Incubation of all samples (A, B, B_(A+) and C) for 30 days under the same conditions in NutriSelect™ Plus nutrient broth at 37°C with measurement of the viral RNA load in each sample at days 1, 2, 3, 7, 14, 21, and 30, following the date of inoculation (day 0).
- 5) Antibiotic treatment on 18 aliquots derived from sample B_(A+) at day 21, consisting in the addition of a specific antibiotic (each of the following: metronidazole, clindamycin, lincomycin, piperacillin+tazobactam, vancomycin, amoxicillin, ampicillin, cefixime, ceftriaxone, meropenem, rifaximin, azithromycin, erythromycin, gentamicin, ciprofloxacin, colistin, levofloxacin, and teicoplanin) to each aliquot. SARS-CoV-2 RNA load was measured by Luminex technology in each aliquot before and 3 days after antibiotic administration.
- 6) In parallel, additional analyses were performed to evaluate and monitor over time the bacterial growth and metabolic activity of all samples and all aliquots of sample B_(A+), using SANIST Biotyper according to the method described by Cristoni *et al.*¹⁴

7) Purification and analysis of the peptides present in sample B_(A+) at day 30.

The details of the procedures and protocols used are presented in the extended data, together with a schematic representation of the experimental design (Graphical abstract).³⁵

Results

The experimental design included a series of analyses (performed on all samples A, B, B_(A+) and C) aimed at verifying: 1) the permanence/survival over time and the eventual multiplication of SARS-CoV-2 RNA *in vitro*; 2) the presence/synthesis of SARS-CoV-2 peptides in the cultures having confirmed SARS-CoV-2 RNA presence; 3) the effect of antibiotics administration in sample B_(A+); 4) the concomitant presence of other metabolites; and 5) the characterisation of the bacterial samples, including the verification of the presence of eukaryotic cells.

Presence of SARS-CoV-2 RNA

The results presented and discussed here, carried out over a period of 30 days, confirmed the extra-corporal multiplication of SARS-CoV-2 RNA: viral load highly increased over time in sample B_(A+), slightly increased in sample A, decreased in sample C while, as expected, sample B was found constantly negative (Figure 1).

In order to verify the reproducibility of our results, the whole experiment was repeated independently three times using the same infected and healthy samples (with the exception that the repetition experiments were stopped at day 14 instead of day 30). The results of the SARS-CoV-2 RNA load measurements in the repetitions are reported in Figure 2, where results are depicted as the average of the measurements of the three repetitions, together with the calculated standard deviations. The trend was confirmed, with the increase over time of measured viral RNA load in sample A and sample B_(A+). Decrease in sample C and no detection in sample B were also confirmed, but they are not reported in Figure 2.

In addition, three new couples of faecal samples from different “infected donors” (*i.e.* sources of A) and “healthy recipients” (*i.e.* sources of B) have been recruited, and subject to the same experimental procedure. Samples were collected from anonymous donors, and no information (*i.e.* age, sex, blood serotype, severity of the disease, time of the collection,

Figure 1. SARS-CoV-2 RNA load variation over time. SARS-CoV-2 RNA load measurements (reported as AU, see extended data) of samples A (blue bars), B (orange bars), B_(A+) (red bars), and C (azure bars) grown, all under the same conditions for thirty days from inoculation (day 0). SARS-CoV-2 RNA load in sample B_(A+) had a power increase trend over time (as shown in the small frame on top-left), slightly increased in sample A, and decreased in sample C. As expected, sample B was found constantly negative.

Figure 2. Average SARS-CoV-2 RNA load variation in repetitions. The graph reports average results of three repetitions of the experiment conducted using the same starting material as described in Figure 1 (with the exception that the repetitions were stopped at day 14 instead of day 30). To normalise the measurements, all values at day 0 were used as denominator (at day 0 all values = 1), i.e. for each sample, at day X, the ratio between LuminexCountAtDayX/LuminexCountAtDay0 was calculated. Each bar represents the average of the SARS-CoV-2 RNA load ratio of samples A (blue bars) and B_(A+) (red bars), together with the calculated standard deviations.

Figure 3. SARS-CoV-2 RNA load increase in different donor and recipient samples. Results of experiments combining samples from three “infected donor” sources (A1, A2 and A3) and from three “healthy recipient” sources (B1, B2 and B3). A) The graphs report results of nine combinations. To normalise the measurements, all values at day 0 were used as denominator (at day 0 all values = 1), i.e. for each sample, at day X, the ratio between LuminexCountAtDayX/LuminexCountAtDay0 was calculated. Each bar represents the SARS-CoV-2 RNA load ratio. Although with certain differences, the observed trends are similar, confirming the increase over time of SARS-CoV-2 RNA load in samples of type A and samples of type B_(A+), independently from the sources of A and B. B) Each line represents the average of the SARS-CoV-2 RNA load ratio of samples B1 (green line), B2 (yellow line), and B3 (azure line) infected each one with three different A donor sources. To normalise the measurements, all values at day 0 were normalised as described in panel A).

Figure 3. (continued)

Figure 4. Effect of antibiotics on viral load. SARS-CoV-2 RNA load measurements (reported as AU, see Supplementary Material) of eighteen aliquots pre- (red) and post- (three days, dashed) treatment with the following selection of antibiotics (ABX): Metronidazole (class: Azoles); Clindamycin, Lincomycin, Piperacillin+Tazobactam, Vancomycin (class: Carboxylic acids and derivatives); Amoxicillin, Ampicillin, Cefixime, Ceftriaxone, Meropenem (class: Lactams); Rifaximin (class: Macrolactams); Azithromycin, Erythromycin, Gentamicin (class: Organooxygen compounds); Ciprofloxacin, Colistin, Levofloxacin (class: Quinolones and derivatives); Teicoplanin (semisynthetic glycopeptide antibiotic). SARS-CoV-2 RNA load is reported as preABX-postABX variation in percentage.

Table 1. Examples of 12 peptides (named as in the first column) mapping on different SARS-CoV-2 proteins (column "Match") are here reported. Amino acids highlighted in red represent changes with respect to the translations of CDS regions reported in the reference severe acute respiratory syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome sequence (GenBank LOCUS: NC_045512.2). Pep51, pep121, and pep230 were found with a different mass spectrometry approach using the Q Exactive HF Hybrid Quadrupole-Orbitrap with an ultra-high-field analyser (Broga, personal communication).

Peptide ID	Length (AA)	Fragment	Match	From	To
pep51	27	ESDDYIKNGPLTVGGSCLLSGHNIAK		268	294
pep121	82	LILSVCLGSLIYSTAALGVLMNSNLGMPSTGYREGYLNSTNVTIATYCTGSIPCSVCLSLDSDTYPSTLETIQTISFFK	NSP3	1416	1497
pep20	77	CLGSLYSTAALGVLMNSNLGMPSTGYREGYLNSTNVTIATYCTGSIPCSVCLSLDSDTYPSTLETIQTISFFK		1421	1497
pep230	62	WVLNNDYYRSLPGVFCGVDVAVNLLTNMFTPLIQPIGALDISASIVAGGINAVITCLAYFIM	NSP4	241	302
pep199	62	TDAVRDPQTLEILDITPCSFSGGVSVITPGTNTSNQVAVLYQGVNCTEVPVAIHADQLTPTWR		573	634
pep33	69	DPQTLEILDITPCSFSGGVSVITPGTNTSNQVAVLYQDVNCTEVPVAIHADQLTPTWVSTGNSNVFQTR		578	646
pep190	74	SVASQSIAYTMLLGAENSVAYSNNSIAIPTNFTISVTTEILPVSMTKTSVDCTMYICGDSSTECNLLLLQYGSF	S	686	759
pep181	75	SIAYTMSLGAENSVAYSNNSIAIPTNFTISVTTEILPVSMTKTSVDCTMYICGDSSTECNLLLLQYGSFCTQLNR		691	765
pep103	31	IQDLSSTASALGKLQDVVNQNAQALNTLVK		934	964
pep25	77	DCWLHSYFTSDYYQLYSTQSLTDTGVEHVTFFIYNKIVDEPEEHVQIHTIDGSSGVNPNVMEPICDDEPTTTTSVPL	NS3	199	275
pep68	13	GISPGRMAGNGGDAALALLLDR	N	204	216
pep38	31	EAVGTNLPLQLGFSTGVNLVAVPTGVVDTPN	NSP14	99	129

Table 2. Amino acid changes reported in Table 1 have been checked for their existence among the observed variations in SARS-CoV-2 sequenced isolates available GISAID at time of writing. All of them, except one, have been already reported in humans, and only two in Italy. For each amino acid change, the number of occurrences in GISAID isolates is reported, together with details of the first human isolate recorded in GISAID with reported collection date. AA change NSP3:A274K of pep51 has never been reported in human SARS-CoV-2 sequences, but it has been found in beta-CoV genome sequences from bats (isolate hCoV-19/bat/Yunnan/RmYN01/2019, collection date 25-06-2019).

Peptide ID	AA change	Observed in human?	#Occurrence in human	Observed in Italy?	Observed in other than human?	First human isolate recorded in GISAID with reported collection date	Collection date
pep51	NSP3:A274K	No	0	No	Yes	-	-
pep51	NSP3:K280T	Yes	2	No	No	hCoV-19/Finland/HEL-18-471/2021	23/01/2021
pep51	NSP3:V286L	Yes	1	No	Yes	hCoV-19/USA/TX-HMH-MCoV-25096/2021	20/01/2021
pep121	NSP3:L141I	Yes	4	No	No	hCoV-19/USA/WA-UW163/2020	13/03/2020
pep199	S:D614G	Yes	728,982	Yes	Yes	hCoV-19/Australia/NSW2153/2020	25/01/2020
pep33	S:R634S	Yes	2	No	Yes	hCoV-19/India/MH-NIV-815-3/2020	07/04/2020
pep190	S:S698L	Yes	398	Yes	No	hCoV-19/USA/AZ-TG666166/2020	25/03/2020
pep25	NS3:Y264C	Yes	106	No	No	hCoV-19/Canada/ON-5738/2020	09/04/2020
pep68	N:T205I	Yes	25,665	Yes	No	hCoV-19/Beijing/Wuhan_IME-BJ07/2020	29/01/2020
pep68	N:A208G	Yes	514	No	No	hCoV-19/USA/MD-HP00076/2020	11/03/2020

fatality, *etc.*) is available. All combinations of “infected donors” sources (A1, A2 and A3) and “healthy recipients” donors (B1, B2 and B3) were subject to the same experimental procedure. Although with certain differences, the observed trends are similar (Figure 3A), confirming the increase over time of SARS-CoV-2 RNA load in samples of type A and in samples of type B_(A+). Lastly, for each “recipient”, we measured SARS-CoV-2 RNA load (Figure 3B).

Effect of antibiotics administration

Aliquots of sample B_(A+) tested after three days of culture in the presence of the single different antibiotics belonging to different classes were analysed and the SARS-CoV-2 RNA load measured in each of them. SARS-CoV-2 RNA load was found to be influenced by the presence of antibiotics in different ways (Figure 4):

- SARS-CoV-2 RNA load was reduced to undetectable levels in the four aliquots treated with metronidazole, vancomycin, amoxicillin and azithromycin, respectively.
- SARS-CoV-2 RNA load decreased by 20% to 85% in the aliquots treated with piperacillin+tazobactam, ampicillin, cefixime, ceftriaxone, meropenem, gentamicin, ciprofloxacin and teicoplanin. For example, cefixime induced a decrease of viral RNA load of 85%, ciprofloxacin of 61% and teicoplanin of 56%.
- SARS-CoV-2 RNA load was not substantially affected by the presence of clindamycin, lincomycin, rifaximin, erythromycin, colistin and levofloxacin.

Presence of SARS-CoV-2 peptides

After 30 days of bacterial growth in culture, aliquots of samples B_(A+) (from couple 0) were collected and tested for the presence of SARS-CoV-2-related peptides using mass spectrometry (details are described in extended data³⁵). Several peptides found in the aliquot from sample B_(A+) were assigned to SARS-CoV-2 proteins. As shown in Table 1, the sequence of some of the peptides (pep51 and pep121, matching on NSP3; pep199, matching on the spike protein; pep25 and pep68, matching on NS3 and N, respectively) have one or more amino acid (AA) changes (highlighted in red) with respect to the translations of CDS regions reported in the reference ‘Severe acute respiratory syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome sequence’ (GenBank LOCUS: NC_045512.2). No SARS-CoV-2-related peptide was identified in the aliquot of sample B.

The identified AA changes have been checked for their existence among the observed variations in SARS-CoV-2 sequenced isolates available in GISAID¹⁵ at time of writing. As shown in Table 2, all of them except NSP3:A274K in pep51 have been already reported in humans; the majority of them have been never reported in the country of origin of the samples (Italy), the remaining ones have been observed in samples sequenced in Italy, but after the time of collection of the infected sample A (February 2020). Some of the found peptides mapping on the SARS-CoV-2 spike protein are shown in Figure 5.

Presence of other metabolites

We have already described the detection of toxin-like peptides in plasma, urine and faecal samples from COVID-19 affected individuals (Cristoni *et al.*,¹⁶ *under review*). The evaluation on the potential release of toxic-like peptides in aliquots from sample B_(A+) has been assessed by performing the same analyses. Toxin-like peptides have been observed, but their presence

Figure 5. Peptides identified in sample B_(A+) mapped on SARS-CoV-2 spike protein. Local alignments of peptides identified in sample B_(A+) mapping on three different regions of SARS-CoV-2 reference spike protein (NCBI protein LOCUS YP_009724390.1). Amino acids highlighted in red correspond to changes described in Table 1 and Table 2.

Figure 6. Bacteria genera. The presence of bacteria genera was monitored over time by looking at their metabolic activity as described by Cristoni et al.¹⁴ Measures on Y-axis are reported as “detection frequency” (range 0-10). The two charts report the most metabolically active genera identified together with the “generic bacterial gut microbiota” (representing other bacterial genera not classified by the instrument) at day 0, 1, 7, 14, 21, and 30 for samples A, and B_(A+). Other microbial organisms were observed at low levels (2 or less, at day 7) and not reported in the figure: *Mycobacterium*, *Actinobacteria*, *Bacteroidetes*, *Blautia*, *Brevibacterium*, *Brevundimonas*, *Candida* (*C. albicans*), *Collinsella*, *Enterococcus*, *Eubacterium*, *Klebsiella*, *Lactonifactor*, *Microbacterium*, *Porphyromonas*, *Propionibacterium*, *Sphingomonas*, *Stenotrophomonas*, *Streptococcus gordonii*, *Xanthomonas*.

was completely reduced to negligible levels in the aliquots treated with metronidazole and vancomycin administration (data not shown). These results need to be carefully interpreted, taking into account the different antimicrobials kinetics.

Presence of eukaryotic cells and virus-like particles

Samples A and B_(A+) were found to contain some bacterial genera particularly abundant and metabolically active during the whole experiment, as shown in Figure 6.

Aliquots of samples A, B, and B_(A+), collected at different times, were analysed with both transmission electron microscope (TEM) and scanning electron microscope (SEM) to verify the presence of eukaryotic cells. More than 30 different preparations (including at 30 days of culture) have been observed: none was found to contain any structure resembling cells with nuclei, but only bacterial cells. Analyses of images on samples A and B_(A+) revealed the presence of virus-like particles interacting with bacterial cells. Immune electron microscopy is ongoing for confirming that these particles are of SARS-CoV-2 origin (*in preparation*).

Discussion

Our results indicate that the SARS-CoV-2 genome, in addition to its known interactions with eukaryotic cells, is additionally capable of replicating outside the human body, suggesting a possible ‘bacteriophage-like’ mode of action. It is not clear yet whether the SARS-CoV-2 genome could just be replicated by its RNA polymerase (which would correspond to a bacteriophage pseudo-lysogenic mechanism), or if the production of full-blown SARS-CoV-2 replicating particles within the bacteria occur (which would correspond to the typical lytic cycle of bacteriophages). In either case, according to our knowledge, this is a novel observation and has never described before for SARS-CoV-2.

The experiment here described (Figure 1) was repeated three additional times using the same samples (Figure 2). In addition, independent replications were performed in a 3×3 design using different starting material (Figure 3). In all of them, very similar trends were observed and the increase of SARS-CoV-2 RNA load in sample A and sample B_(A+) was confirmed in all experiments.

In the cases of replications with different starting material (*i.e.* faecal samples from different “infected donors” as sources of A, and “healthy recipients” as sources of B), the trends are similar and confirm that the experiment is reproducible. In terms of relevance, we noticed that SARS-CoV-2 RNA load was particularly high in one combination (A2×B2). It is thus plausible that viral RNA load depends on the gut eubiotic/dysbiotic condition met by the virus. This is also a plausible explanation of why in the initial couple of A-B sources (Figure 1) the difference between viral RNA load measurements in A and B_(A+) is notable. SARS-CoV-2 is considered as a respiratory virus, and many bacteria reside in the upper respiratory tract (URT) interacting with different viruses like influenza (see Schenk *et al.*, 2016 for an overview). With this respect, our observations are in line with the hypothesis formulated by Shah, who has recently proposed the existence of a gut-lung equilibrium mediated by multiple mechanisms of action, including the abundance of certain microorganisms in the gut microbiota as responsible for determining the sensitivity and severity of SARS-CoV-2 infections.

Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (Ebrahimi, 2020, Budding *et al.*, 2020). In particular, Ebrahimi identified *in silico* a series of serine protease TMPRSS2 and peptidyl peptidases with high similarity to the ACE2 peptidase domain (ACE2-PD) in members of *Proteobacteria* phylum. It can't be excluded that these or other similar proteins act as the cellular receptors for SARS-CoV-2 in bacteria. Looking for taxa, species or consortia that can be prone to act as receptor of virus is imperative, and, with this respect, a whole-genome metagenomic sequencing on the samples is ongoing, aimed at characterising further which bacterial species are candidate target(s) of the observed behaviour of SARS-CoV-2.

Whereas our experimental design was intended to grow bacterial cells, the possibility that SARS-CoV-2 RNA increase could be due to replication in human cells present in the original faecal samples, was considered. The human cells most abundantly present in faecal samples are colonic epithelial cells (colonocytes). Loktionov,¹⁶ reported that cell exfoliation events from colonic epithelium are rare under normal conditions, while they dramatically increase in cases of uncontrolled growth of cells not under physiologic control (like in neoplasia), when cell removal by apoptosis does not function properly. In addition, Iyengar *et al.*¹⁷ reported that colonic epithelial cells terminally differentiated are devoid of proliferative activity. More recently, Nair *et al.*¹⁸ and Chandel *et al.*¹⁹ developed specific methodologies to recover viable colonocytes from stool. In our case, both sample A and B originated from adult individuals with no diagnosis of cancer. In addition, it is unlikely that human cells potentially present in samples A and B are able to:

- grow in a culture medium typically formulated for bacteria and not containing growth factors, serum, nor other important components for eukaryotic cell sustainment;

- survive in such a medium for 30 days, and in co-occurrence with an event of SARS-CoV-2 infection;
- multiply in the absence of specific CO₂ concentration conditions.

Also, the possibility of interaction between SARS-CoV-2 and other eukaryotic organisms potentially present in the cultures, as *e.g.* parasitic nematodes and fungal cells, has been considered.

During the whole experiment, parasitic nematodes were not noted at visual inspections by eye. In addition, stool of sample B was independently analysed and certified to be free of known parasites and microbial pathogens (certification provided by the Italian diagnostic laboratory [Biomolecular Diagnostic Srl](#)). Parasitic nematodes are usually not able to survive outside the host and many intestinal roundworms (like those of genus *Ascaris*) release antimicrobial factors that interfere with bacterial growth,²⁰ in contrast with the found high increase of metabolic activities of some bacterial genera. In the used medium, chemical elements relevant for (parasitic and not) nematodes (*e.g.* cholesterol and traces of metals) are missing.

The possibility of involvement of the mycobiome fraction present in the stool was considered. As highlighted by Chin *et al.*,²¹ multifaceted and multidisciplinary approaches are necessary to identify uncultivable, low-abundance, permanent and transient fungal species residing in the gut, confirming that the human mycobiome is not yet fully characterised. Accordingly, while the ability of unknown fungi to grow in the used culture medium cannot be excluded, no significant metabolic activity of *Candida albicans*, most commonly found in the microbiota, was observed.

Finally, inspections of images from TEM and SEM on more than 30 different preparations did not reveal presence of eukaryotic cells (*in preparation*). If on the one hand, the possibility that a nematode or another unknown eukaryotic cell is able grow in the medium cannot be excluded, the used conditions make this possibility very unlikely. Anyhow, the ability of SARS-CoV-2 to interact either with nematodes or with fungal cells has never been observed before and would be a novel and surprising observation as well.

As indicated above, several peptides matching to SARS-CoV-2 proteins were found in the aliquot from sample B_(A+). The identification of peptides with amino acid changes, compared to the translations of CDS regions of the reference SARS-CoV-2 genome, is intriguing but is compatible with the mechanism of viral replication in bacteria. RNA viruses such as SARS-CoV-2 inhabit the host as a population of variants called *quasispecies*, *i.e.* a group made of different variants that are genetically linked through mutation events, and contribute collectively to the characteristics of the whole (viral) population in the host.²² Recent studies highlighted the significant amount of intra-host genomic diversity in SARS-CoV-2 samples.^{23,24} In a 'bacteriophage-like' mode of action, as bacteria were grown for 30 days, it can't be excluded that the observed amino acid changes represent viral *quasispecies* emerged through replication events in bacterial hosts. In relation to this, recent studies²⁵⁻²⁷ evidenced hypermutations occurrences in SARS-CoV-2 genomes and suggested APOBEC and ADAR deaminases as the possible responsible of these phenomena. The APOBEC family is related to bacterial, yeast, and plant deaminases all possessing highly conserved amino acid motifs responsible for coordination of zinc in the active site.²⁸ As no sequencing was performed on the original infected stool sample, the presence of SARS-CoV-2 haplotypes in the initial SARS-CoV-2 population used to perform the experiments, therefore justifying the amino acid changes observed, cannot be excluded. However, all the amino acid changes found have been reported in sequences of SARS-CoV-2 found for the first time after the date of collection of the infected sample A (February 2020), and some of them have never been reported in Italy, the country of origin of the samples.

On the other hand, other mechanisms like those at the basis of diversity-generating retroelements (DGR) systems²⁹ have that could contribute to SARS-CoV-2 hypermutation phenomena have recently been described in bacteria, and could therefore be responsible of the AA changes found. Additional experiments aimed to verify increase of viral peptides similar to viral RNA over time are planned.

These results can potentially provide new insights in the epidemiology of SARS-CoV-2. Considering the possible impact and implications that such relationship has on the manifestation, therapy and control of COVID-19 disease, some questions immediately arise like *e.g.*:

- Can this 'bacteriophage-like' behaviour of SARS-CoV-2 explain the long-term presence of SARS-CoV-2 observed in some recovered patients?³⁰
- Can antibiotics and/or bacteriophage-based therapies play a role in the treatment of COVID-19 affected patients?³¹

- How would the (antecedent) administration of antibiotics to patients, influencing the microbiota population, impact the clinical course of the disease?³²
- Can the involvement of bacteria in COVID-19 epidemiology help to explain clinical observations, like the elevated serum C-reactive protein, procalcitonin, D-dimer, and ferritin associated with poor outcomes in COVID-19?³³

These questions are only examples of the plethora of questions to be addressed. Our results support the way to tackle COVID-19 pandemic proposed by Mushi,³⁴ *i.e.* by using the One Health holistic approach. If individuals are considered not only human bodies, but as 'holobionts', *i.e.* discrete ecological units that need to be studied and treated as such, a deeper understanding of the role of the microbial community living in the human body is fundamental to tackle COVID-19 disease.

Consent

Faecal samples were collected and handled by CranioMed S.R.L. from anonymous donors who agreed to participate in this study by signing informed consent, as foreseen by Italian legislation. No personal information (*i.e.* age, sex, blood serotype, severity of the disease, time of the collection, fatality, *etc.*) were collected.

The study is compliant with the JRC Scientific Integrity and Research Ethics guidance.

Declarations

The scientific output expressed does not imply a policy position of the European Commission. Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use that might be made of this publication.

Data availability

Underlying data

Zenodo: Underlying data for 'Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology'. <https://doi.org/10.5281/zenodo.4723549>.³⁵

The project contains the following underlying data:

Mass spectrometry raw data of the peptides.

NCBI Genome: Severe acute respiratory syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome. Accession number: NC_045512.2; https://www.ncbi.nlm.nih.gov/nuccore/NC_045512.2.

NCBI Protein: Surface glycoprotein [Severe acute respiratory syndrome coronavirus 2]. Accession number: YP_009724390.1; <https://www.ncbi.nlm.nih.gov/protein/1796318598>.

Extended data

Zenodo: Extended data for 'Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology'. <https://doi.org/10.5281/zenodo.4723549>.³⁵

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

We gratefully acknowledge the Authors and the Originating laboratories where the clinical specimen or virus isolate was first obtained and the Submitting laboratories, where sequence data have been generated and submitted to GISAID.

Authors' notes

The efforts and commitment put into this research work are dedicated to all EU citizens who suddenly left us because of COVID-19, and to their families.

References

1. Trottein F, Sokol H: **Potential Causes and Consequences of Gastrointestinal Disorders during a SARS-CoV-2 Infection.** *Cell Rep.* 2020; **32**: 107915.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Zuo T, et al.: **Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization.** *Gastroenterology.* 2020; **159**: 944–955. e8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. He L-H, et al.: **Intestinal Flora as a Potential Strategy to Fight SARS-CoV-2 Infection.** *Front. Microbiol.* 2020; **11**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Wu Y, et al.: **Prolonged presence of SARS-CoV-2 viral RNA in faecal samples.** *Lancet Gastroenterol. Hepatol.* 2020; **5**: 434–435.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Heneghan C, Spencer E, Brassey J, et al.: **SARS-CoV-2 and the Role of Orofecal Transmission: Systematic Review.** *medRxiv* 2020.08.04.20168054. 2020.
[Publisher Full Text](#)
6. Jefferson T, Spencer EA, Brassey J, et al.: **Viral cultures for COVID-19 infectious potential assessment – a systematic review.** *Clin. Infect. Dis.* 2020;
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Jefferson T, Spencer E, Brassey J, et al.: **SARS-CoV-2 and the Role of Orofecal Transmission: Evidence Brief.**
8. O'Keefe J, Freeman S, Nicol A: **The Basics of SARS-CoV-2 Transmission.** *Vancouver.* 2021.
9. Brogna B, et al.: **SARS-CoV-2 Detection in Faecal Sample from a Patient with Typical Findings of COVID-19 Pneumonia on CT but Negative to Multiple SARS-CoV-2 RT-PCR Tests on Oropharyngeal and Nasopharyngeal Swab Samples.** *Medicina (B. Aires).* 2021; **57**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Zuo T, et al.: **Depicting SARS-CoV-2 faecal viral activity in association with gut microbiota composition in patients with COVID-19.** *Gut.* 2020; gutjnl-2020-322294.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Wang W, et al.: **Detection of SARS-CoV-2 in Different Types of Clinical Specimens.** *JAMA.* 2020;
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Wölfel R, et al.: **Virological assessment of hospitalized patients with COVID-2019.** *Nature.* 2020; **581**: 465–469.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Yao H, et al.: **Patient-derived SARS-CoV-2 mutations impact viral replication dynamics and infectivity in vitro and with clinical implications in vivo.** *Cell Discov.* 2020; **6**: 76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Cristoni S, et al.: **Predicting and preventing intestinal dysbiosis on the basis of pharmacological gut microbiota metabolism.** *Rapid Commun. Mass Spectrom.* 2019; **33**: 1221–1225.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Elbe S, Buckland-Merrett G: **Data, disease and diplomacy: GISAID's innovative contribution to global health.** *Glob. Challenges.* 2017; **1**: 33–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Loktionov A: **Cell exfoliation in the human colon: Myth, reality and implications for colorectal cancer screening.** *Int. J. Cancer.* 2007; **120**: 2281–2289.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Iyengar V, Albaugh GP, Lohani A, et al.: **Human stools as a source of viable colonic epithelial cells.** *FASEB J.* 1991; **5**: 2856–2859.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Nair P, et al.: **Coprocytobiology.** *J. Clin. Gastroenterol.* 2003; **36**: S84–S93.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Chandel DS, Braileanu GT, Chen J-HJ, et al.: **Live Colonocytes in Newborn Stool: Surrogates for Evaluation of Gut Physiology and Disease Pathogenesis.** *Pediatr. Res.* 2011; **70**: 153–158.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Midha A, et al.: **The Intestinal Roundworm *Ascaris suum* Releases Antimicrobial Factors Which Interfere With Bacterial Growth and Biofilm Formation.** *Front. Cell. Infect. Microbiol.* 2018; **8**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Chin VK, et al.: **Mycobiome in the Gut: A Multiperspective Review.** *Mediators Inflamm.* 2020; **2020**: 1–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Luring AS, Andino R: **Quasispecies Theory and the Behavior of RNA Viruses.** *PLoS Pathog.* 2010; **6**: e1001005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Jary A, et al.: **Evolution of viral quasispecies during SARS-CoV-2 infection.** *Clin. Microbiol. Infect.* 2020; **26**: 1560.e1–1560.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Wang Y, et al.: **Intra-host variation and evolutionary dynamics of SARS-CoV-2 populations in COVID-19 patients.** *Genome Med.* 2021; **13**: 30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Di Giorgio S, Martignano F, Torcia MG, et al.: **Evidence for host-dependent RNA editing in the transcriptome of SARS-CoV-2.** *Sci. Adv.* 2020; **6**: eabb5813.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Klimczak LJ, Randall TA, Saini N, et al.: **Similarity between mutation spectra in hypermutated genomes of rubella virus and in SARS-CoV-2 genomes accumulated during the COVID-19 pandemic.** *PLoS One.* 2020; **15**: e0237689.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Simmonds P: **Rampant C→U Hypermutation in the Genomes of SARS-CoV-2 and Other Coronaviruses: Causes and Consequences for Their Short- and Long-Term Evolutionary Trajectories.** *mSphere.* 2020; **5**: e00408–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Ratcliff J, Simmonds P: **Potential APOBEC-mediated RNA editing of the genomes of SARS-CoV-2 and other coronaviruses and its impact on their longer term evolution.** *Virology.* 2021; **556**: 62–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Wu L, et al.: **Diversity-generating retroelements: natural variation, classification and evolution inferred from a large-scale genomic survey.** *Nucleic Acids Res.* 2018; **46**: 11–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. D'Ardes D, et al.: **Long-term Positivity to SARS-CoV-2: A Clinical Case of COVID-19 with Persistent Evidence of Infection.** *Eur. J. Case Reports Intern. Med.* 2020; **7**: 1707.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Goncalves Mendes Neto A, et al.: **Bacterial infections and patterns of antibiotic use in patients with COVID-19.** *J. Med. Virol.* 2020; jmv.26441.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Zhang L, et al.: **Retrospective analysis of clinical features in 134 coronavirus disease 2019 cases.** *Epidemiol. Infect.* 2020; **148**: e199.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Huang I, Pranata R, Lim MA, et al.: **C-reactive protein, procalcitonin, D-dimer, and ferritin in severe coronavirus disease-2019: a meta-analysis.** *Ther. Adv. Respir. Dis.* 2020; **14**: 175346662093717.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Mushi V: **The holistic way of tackling the COVID-19 pandemic: the one health approach.** *Trop. Med. Health.* 2020; **48**: 69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Petrillo M, Brogna C, Cristoni S, et al.: **Underlying and extended data for 'Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology'.**
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 25 June 2021

<https://doi.org/10.5256/f1000research.57680.r88114>

© 2021 Honarmand Ebrahimi K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Kourosh Honarmand Ebrahimi

Department of Chemistry, University of Oxford, Oxford, UK

The authors have fully addressed my original concerns. I found their hypothesis very interesting, however, they should note and somewhere in the paper they mention that ACE2 receptor is expressed in other tissues like the small intestine, and what they have observed could be due to the replication of the virus in the gastrointestinal tract tissue (Han *et al.* (2020¹)).

References

1. Han T, Kang J, Li G, Ge J, et al.: Analysis of 2019-nCoV receptor ACE2 expression in different tissues and its significance study. *Ann Transl Med.* 2020; **8** (17): 1077 [PubMed Abstract](#) | [Publisher Full Text](#)

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Virology and immunology and bioinorganic chemistry

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 25 Jun 2021

Mauro Petrillo

Dear Dr Kourosh Honarmand Ebrahimi,

Thanks a lot for your comment.

Your point is relevant and we modified the Discussion section accordingly:

Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (Ebrahimi, 2020, Budding et al., 2020). In particular, Ebrahimi identified *in silico* a series of serine protease TMPRSS2 and peptidyl peptidases with high similarity to the angiotensin-converting enzyme 2 (ACE2) receptor peptidase domain (ACE2-PD) in members of Proteobacteria phylum. It can't be excluded that these or other similar proteins act as the cellular receptors for SARS-CoV-2 in bacteria. **At the same time, ACE2 receptor gene, whose protein is known to be critical in SARS-CoV-2 transmission [Yan 2020], is expressed in various tissues and organs of the human body, including the small intestine [Han 2020]. It is thus very likely that SARS-CoV-2 found in faecal samples of infected individuals is from infections occurring in human body cells. Our observations are not in contrast, and they suggest that, in the gastrointestinal tract, human cells, like those of small intestine, are not the sole SARS-CoV-2 target.**

We hope that the quality of the manuscript, thanks to your comment, has been improved and you consider it suitable for publication.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

Reviewer Report 25 June 2021

<https://doi.org/10.5256/f1000research.57680.r88113>

© 2021 Aguilera M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Margarita Aguilera

Department of Microbiology, Faculty of Pharmacy, Campus of Cartuja, University of Granada, Granada, Spain

Authors have fully improved the manuscript according to all the suggestions proposed. No further actions are needed.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbiota, Probiotics, taxonomy, Microbiology, Molecular biology, Biotechnology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 25 Jun 2021

Mauro Petrillo

Dear Prof. Aguilera,

Thanks a lot for your valuable comments and suggestions that you provided. They surely improved the quality of the manuscript.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

Version 1

Reviewer Report 07 June 2021

<https://doi.org/10.5256/f1000research.55836.r85802>

© 2021 Aguilera M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Margarita Aguilera

¹ Department of Microbiology, Faculty of Pharmacy, Campus of Cartuja, University of Granada, Granada, Spain

² Department of Microbiology, Faculty of Pharmacy, Campus of Cartuja, University of Granada, Granada, Spain

The work titled "*Increase of SARS-CoV-2 RNA load in faecal samples prompts for rethinking of SARS-CoV-2 biology and COVID-19 epidemiology*" has been well designed and soundly performed.

After the urgent needs to invest efforts on clinical research for handling the COVID-19, new questions should be posed and the present work has done innovatively. The hypothesis is relevant due to the high differential effects within SARS-CoV-2 patients, based on the individual health statuses that determine variable host responses. The Authors have shown experimental data to demonstrate that SARS-CoV-2 could have a bacteriophage activity through evaluating SARS-CoV-2 long-term survival, proliferation and the potential interaction between the virus and gut microbiota taxa. Multiple methodologies and experimental data support appropriately the results and conclusions: viral RNA by Luminex technology; impact of antibiotic treatments; Biotyper for cultured microbiota predominant taxa over the time, eukaryotic cells, peptides, other metabolites molecular alignments, etc.

Therefore, I consider that Petrillo *et al.* have carried out an excellent and holistic approach; **I would suggest its publication after minor corrections/explanations:**

- Please use the term gut microbiota instead gut flora along the document.
- Please add the underlined phrase at the end of this paragraph: *At the same time, Wölfel et al. 12 observed high viral RNA concentration in stool samples, but reported isolation of infectious virus only from throat- and lung-derived samples, while Yao et al. had indication of viable SARS-CoV-2 particles in stool samples, denoting that the detailed biology of SARS-CoV-2 is not yet fully elucidated. Moreover, the interaction between SARS-CoV-2 and individual variable microbiota composition could drive the differential pathophysiological effects and severity of symptoms.*
- Please explain if the medium NutriSelect™Plus nutrient used for the inoculation and cultivation of microorganisms could have an impact on the specific taxa cultured as well as metabolites measured. Have you tried other media?
- Please explain if you have been used only aerobic conditions for all experimental culturing data.
- Please specify the concentration of antibiotics added to the experiment.
- Figure 1: Please give a plausible explanation for the data shown in this figure - Could it be that microbiota of healthy individuals contain eubiotic taxa that allow a better proliferation of SARS-CoV-2?
- Figure 3: Please explain the differences observed between the three graphics. The microbiota composition of the sample receptors seems to have an impact. Please give a plausible explanation.
- Future research is directly derived from these results: to look for the taxa, species or consortia that can be prone to act as receptor of virus. Please comment on that.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbiota, Probiotics, taxonomy, Microbiology, Molecular biology, Biotechnology.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 07 Jun 2021

Mauro Petrillo

Dear Prof. Aguilera,

Thanks a lot for your valuable comments and suggestions that you have provided in the report.

We will address all of them, and will provide a revised version of the manuscript.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

Author Response 09 Jun 2021

Mauro Petrillo

Dear Dr Aguilera,

Thanks a lot for your valuable comments and suggestions that you have provided in the report.

As anticipated, we have addressed all your points, and provided a new version of the manuscript:

- *Please use the term gut microbiota instead gut flora along the document.*
- **Response:** Done, thanks.
- *Please add the underlined phrase at the end of this paragraph: 'At the same time, Wölfel et al.12 observed high viral RNA concentration in stool samples, but reported isolation of infectious virus only from throat- and lung-derived samples, while Yao et al. had indication of viable SARS-CoV-2 particles in stool samples, denoting that the detailed biology of SARS-CoV-2 is not yet fully elucidated. Moreover, the interaction between SARS-CoV-2 and individual variable microbiota composition could drive the differential pathophysiological effects and severity of symptoms.'*
- **Response:** Very relevant point. We added it, together with references.

- *Please explain if the medium NutriSelect™ Plus nutrient used for the inoculation and cultivation of microorganisms could have an impact on the specific taxa cultured as well as metabolites measured. Have you tried other media?*
- *Please explain if you have been used only aerobic conditions for all experimental culturing data.*
- *Please specify the concentration of antibiotics added to the experiment.*
- **Response:** With respect to these points, the information has been added in the supplementary material, points 1, 3, 8). The new link for get the supplementary material is <https://doi.org/10.5281/zenodo.4723549>
- *Figure 1: Please give a plausible explanation for the data shown in this figure - Could it be that microbiota of healthy individuals contain eubiotic taxa that allow a better proliferation of SARS-CoV-2?*
- *Figure 3: Please explain the differences observed between the three graphics. The microbiota composition of the sample receptors seems to have an impact. Please give a plausible explanation.*
- *Future research is directly derived from these results: to look for the taxa, species or consortia that can be prone to act as receptor of virus. Please comment on that.*
- **Response:** Your points are relevant and we modified the Discussion section accordingly: 'In terms of relevance, we noticed that SARS-CoV-2 RNA load was particularly high in one combination (A2×B2). It is thus plausible that viral RNA load depends on the gut eubiotic/dysbiotic condition met by the virus. This is also a plausible explanation of why in the initial couple of A-B sources (Figure 1) the difference between viral RNA load measurements in A and B(A+) is notable. SARS-CoV-2 is considered as a respiratory virus, and many bacteria reside in the upper respiratory tract (URT) interacting with different viruses like influenza (see Schenk et al., 2016 for an overview). With this respect, our observations are in line with the hypothesis formulated by Shah, who has recently proposed the existence of a gut-lung equilibrium mediated by multiple mechanisms of action, including the abundance of certain microorganisms in the gut microbiota as responsible for determining the sensitivity and severity of SARS-CoV-2 infections. Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (Ebrahimi, 2020, Budding et al., 2020). In particular, Ebrahimi identified in silico a series of serine protease TMPRSS2 and peptidyl peptidases with high similarity to the ACE2 peptidase domain (ACE2-PD) in members of Proteobacteria phylum. It can't be excluded that these or other similar proteins act as the cellular receptors for SARS-CoV-2 in bacteria. Looking for taxa, species or consortia that can be prone to act as receptor of virus is imperative, and, with this respect, a whole-genome metagenomic sequencing on the samples is ongoing, aimed at characterising further which bacterial species are candidate target(s) of the observed behaviour of SARS-CoV-2.'

We believe that, thanks to your comments, the quality of the manuscript has improved a lot.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

Reviewer Report 17 May 2021

<https://doi.org/10.5256/f1000research.55836.r85393>

© 2021 Honarmand Ebrahimi K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Kourosh Honarmand Ebrahimi ¹ Department of Chemistry, University of Oxford, Oxford, UK² Department of Chemistry, University of Oxford, Oxford, UK

Petrillo and colleagues report that SARS-CoV-2 could have a bacteriophage activity in faecal samples. They measure the presence of viral RNA using Luminex technology and viral proteins using mass spectrometry. The experimental design is sound and their findings are exciting, which support the publication of this work. However, I have few concerns that must be addressed before publication. Therefore, I recommend publication after minor revisions.

- SARS-CoV-2 is a respiratory virus. Many bacteria reside in the upper respiratory tract (URT) and interact with different viruses like influenza (Schenk *et al.*, 2016¹). Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (e.g. (Ebrahimi, 2020²); (Budding *et al.*, 2020³)). The authors should cite these literatures and discuss their findings with respect to a similar bacteriophage behaviour of SARS-CoV-2 in URT.
- In Figure 3, the graph for donor A2: Why in B2(A2+) sample the amount of RNA is hugely different than the other samples? The authors need to explain.
- The authors suggest that the presence of SARS-CoV-2 peptides is compatible with the mechanism of viral replication in bacteria. If this is true, shouldn't the authors observe an increase in viral peptides similar to viral RNA?

References

1. Schenck LP, Surette MG, Bowdish DM: Composition and immunological significance of the upper respiratory tract microbiota. *FEBS Lett.* 2016; **590** (21): 3705-3720 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Honarmand Ebrahimi K: SARS-CoV-2 spike glycoprotein-binding proteins expressed by upper respiratory tract bacteria may prevent severe viral infection. *FEBS Lett.* **594** (11): 1651-1660 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Budding A, Sieswerda E, Wintermans B, Bos M: An Age Dependent Pharyngeal Microbiota Signature Associated with SARS-CoV-2 Infection. *SSRN Electronic Journal.* 2020. [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Virology and immunology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 May 2021

Mauro Petrillo

Dear Dr Kourosh Honarmand Ebrahimi,

Thanks a lot for your valuable comments and suggestions that you have provided in the report.

We will address all of them, and wait for those of other reviewers, in order to provide a fully revised version of the manuscript.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

Author Response 09 Jun 2021

Mauro Petrillo

Dear Dr Kourosh Honarmand Ebrahimi,

Thanks a lot for your valuable comments and suggestions that you have provided in the report.

As anticipated, we have addressed all your points, and provided a new version of the

manuscript:

- *SARS-CoV-2 is a respiratory virus. Many bacteria reside in the upper respiratory tract (URT) and interact with different viruses like influenza (Schenk et al., 2016). Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (e.g. (Ebrahimi, 2020); (Budding et al., 2020). The authors should cite these literatures and discuss their findings with respect to a similar bacteriophage behaviour of SARS-CoV-2 in URT.*
- **Response:** Your point is relevant and we modified the Discussion section accordingly: 'SARS-CoV-2 is considered as a respiratory virus, and many bacteria reside in the upper respiratory tract (URT) interacting with different viruses like influenza (see Schenk et al., 2016 for an overview). With this respect, our observations are in line with the hypothesis formulated by Shah, who has recently proposed the existence of a gut-lung equilibrium mediated by multiple mechanisms of action, including the abundance of certain microorganisms in the gut microbiota as responsible for determining the sensitivity and severity of SARS-CoV-2 infections. Moreover, recent reports suggest an interaction of URT microbiota with SARS-CoV-2 (Ebrahimi, 2020, Budding et al., 2020). In particular, Ebrahimi identified in silico a series of serine protease TMPRSS2 and peptidyl peptidases with high similarity to the ACE2 peptidase domain (ACE2-PD) in members of Proteobacteria phylum. It can't be excluded that these or other similar proteins act as the cellular receptors for SARS-CoV-2 in bacteria. Looking for taxa, species or consortia that can be prone to act as receptor of virus is imperative, and, with this respect, a whole-genome metagenomic sequencing on the samples is ongoing, aimed at characterising further which bacterial species are candidate target(s) of the observed behaviour of SARS-CoV-2.'
- *In Figure 3, the graph for donor A2: Why in B2(A2+) sample the amount of RNA is hugely different than the other samples? The authors need to explain.*
- **Response:** We have modified Figure 3, by adding an additional panel. In addition, we modified the Discussion section accordingly: 'In terms of relevance, we noticed that SARS-CoV-2 RNA load was particularly high in one combination (A2×B2). It is thus plausible that viral RNA load depends on the gut eubiotic/dysbiotic condition met by the virus. This is also a plausible explanation of why in the initial couple of A-B sources (Figure 1) the difference between viral RNA load measurements in A and B(A+) is notable.'
- *The authors suggest that the presence of SARS-CoV-2 peptides is compatible with the mechanism of viral replication in bacteria. If this is true, shouldn't the authors observe an increase in viral peptides similar to viral RNA?*
- **Response:** Your point is relevant and we modified the Discussion section accordingly: 'Additional experiments aimed to verify increase of viral peptides similar to viral RNA over time are planned.'

We hope that the quality of the manuscript, thanks to your comments, has been improved and you consider it suitable for publication.

Best regards,

Mauro Petrillo, on behalf of the authors.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research