Check for updates

CASE REPORT

Case Report: unilateral condylar hyperplasia [version 1; peer review: 2 approved]

Shishir Shetty¹, Shrihari Guddadarangiah²

¹Oral and Craniofacial Health sciences, University of Sharjah, Sharjah, 27272, United Arab Emirates

²Oral Medicine and Radiology, Krishnadevaraya Dental College and Hospital, Bangalore, Karnataka, 562157, India

V1 First published: 25 Jan 2021, 10:46
<https://doi.org/10.12688/f1000research.48499.1>
 Latest published: 25 Jan 2021, 10:46
<https://doi.org/10.12688/f1000research.48499.1>

Abstract

Case: This report describes a clinical case of unilateral condylar hyperplasia (CH) with unique, atypical morphology. An important feature of this report is the documentation of a series of clinical photographs of the patient, showing a gradual increase in facial asymmetry associated with the CH. The main symptom reported in this case was facial asymmetry. The main intraoral clinical features observed in the patient were contralateral crossbite and ipsilateral open bite associated with CH. Surgical reshaping of the condyle was the treatment plan for this case.

Conclusions: The main take away point from this case is the importance of obtaining previous photographs of the patient at different ages during case diagnosis, which helps the clinician to determine the approximate time of commencement of CH. This case also highlights the imaging features of rarely observed atypical shape of the hyperplastic condyle.

Keywords

Mandibular condyle, hyperplasia, panoramic radiography, computed tomography

Open Peer Review

Approval Status

	1	2
version 1
25 Jan 2021	view	view
1. Prashant M. Battepati , University of Taif, Taif, Saudi Arabia		
2. Renita Lorina Castelino , AB Shetty Memorial Institute of Dental Sciences, Mangalore, India		

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Shishir Shetty (shishirshettyomr@gmail.com)

Author roles: Shetty S: Conceptualization, Investigation, Writing – Original Draft Preparation, Writing – Review & Editing; Guddadarangiah S: Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Shetty S and Guddadarangiah S. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Shetty S and Guddadarangiah S. **Case Report: unilateral condylar hyperplasia [version 1; peer review: 2 approved]** F1000Research 2021, 10:46 <https://doi.org/10.12688/f1000research.48499.1>

First published: 25 Jan 2021, 10:46 <https://doi.org/10.12688/f1000research.48499.1>

Introduction

Condylar hyperplasia (CH) is a rare condition associated with excessive condylar bone growth¹. Adams first described CH in the year 1836². CH often occurs unilaterally and manifests clinically as facial asymmetry³. Apart from facial asymmetry, occlusal discrepancies, chin deviation and temporomandibular joint discomfort are commonly associated with CH⁴. CH is known to be self-limiting in nature, normally commences during puberty, progresses gradually and sometimes may be only recognized at the age of 25–30 years⁵. Diagnosis of CH is usually made using a combination of clinical findings and imaging features⁶. The aim of this case report is to present clinical and imaging features of unilateral CH. One of the significant points of the present report is that the progression of CH-associated facial asymmetry has been described using a series of photographs of the patient. The unique finding is the atypical shape of the hyperplastic condyle observed.

Case report

A 31-year-old Asian male mechanic reported to the dental clinic in January 2013 with complaint of asymmetry of the face over the past five years. The patient felt that the asymmetry had increased during the first three years (of the five-year duration) but had remained constant over the next two years. There was no history of pain, discomfort or clicking sounds in the temporomandibular joints (TMJ). However, the patient had a history of lower jaw trauma during a sports event at the age of 15 years. No other family member had a similar condition.

Clinical examination of the patient revealed facial asymmetry due to the deviation of the chin to the left side of the face (Figure 1). Examination of the right TMJ revealed a bony swelling in the right preauricular area. Intraoral examination revealed posterior open bite on the right side (Figure 2) and posterior crossbite on the left side (Figure 3). A series of photographs of the patient at the age of 18 years (Figure 4a), 24 years (Figure 4b), and 27 years (Figure 4c) was evaluated. No evidence of facial asymmetry was noticed at 18 years. Mild features

Figure 1. Extraoral photograph of the patient showing facial asymmetry.

Figure 2. Intraoral photograph showing posterior open bite on the right side.

Figure 3. Intraoral photograph showing posterior cross bite on the left side.

Figure 4a. Extraoral photograph of the patient at the age of 18 years showing no evidence of facial asymmetry.

of asymmetry were noticed at 24 years and obvious features of asymmetry were noticed at 27 years. A panoramic radiograph revealed the presence of a beak shaped hyperplastic right condyle (Figure 5a). The posteroanterior skull view revealed increased length of the condylar neck on the right side (Figure 5b). A coronal computed tomography (CT) scan showed enlargement of the right condyle with beak like projection on the medial aspect (Figure 6a). An axial CT scan revealed the antero-medial projection of the beak like enlargement (Figure 6b).

Figure 4b. Extraoral photograph of the patient at the age of 24 years showing initial evidence of facial asymmetry.

Figure 5b. Posteroanterior view radiograph showing increase in length of condylar neck on the right side.

Figure 4c. Extraoral photograph of the patient at the age of 27 years showing obvious evidence of facial asymmetry.

Figure 6a. Coronal computed tomography (CT) scan showing enlargement of right condyle with beak like projection on the medial aspect.

Figure 5a. Panoramic radiograph showing enlarged right condyle with beak like projection.

Figure 6b. Axial computed tomography (CT) scan showing the antero-medial projection of the beak like enlargement.

Based on the patient's history, clinical features and imaging findings a diagnosis of hyperplasia of the right condyle was made. The patient was advised surgical treatment of the CH. Unfortunately, the patient was not willing to undergo surgical correction and long term follow up was not possible.

Discussion

CH is characterized by unilateral or bilateral increase in the volume of the mandibular condyle, often leading to facial asymmetry, jaw deviation and malocclusion⁷. The exact etiological factor of CH is still unclear, although endocrine alterations,

metabolic hyperactivity, trauma, and genetic factors⁸ have been implicated. In our patient there was history of trauma at the age of 15 years which could be the possible etiological factor. CH occurs predominantly in women⁹ with a recently published meta-analysis revealing that 64% of cases occurred in females¹⁰. In our report the patient was a 31-year-old male. The female predominance has been attributed to hormonal factors, particularly estrogen^{10,11}. Estrogen regulates bone growth and is found in the articular cartilage and growth plates^{11,12}.

CH usually occurs between the ages of 10 and 30 years and most cases occur between adolescence until the end of pubertal growth⁹. However, some cases of CH also occur after puberty. In our patient the CH seems to have occurred after puberty, as evident in the photographs taken at 24 years. The use of serial photographs of patients with CH at different ages helps physicians to estimate the approximate time of occurrence of the condition. This method was used to estimate the time of occurrence of CH in our patient.

Another important finding that depends on the time of occurrence of CH is posterior open bite. It has been observed that if CH occurs during puberty the occlusal plane usually inclines as a result of dental compensation, but if CH occurs after puberty posterior open bite may be evident^{13,14}. In our case posterior open bite was observed, suggesting that the CH must have occurred after the growth phase ended.

The main clinical feature of unilateral CH is enlargement of the same side of the face and flattened appearance of the contralateral side¹⁵. These clinical features were observed in our case.

Although the combination of clinical findings and imaging features is required for the diagnosis of CH, a radiological examination showing elongation of the neck and head of the condyle is necessary for a definitive diagnosis¹⁶. Osteoma, osteochondroma and resorption of the contralateral condyle are the important differential diagnoses for unilateral CH¹⁷. Condylar osteomas are extremely rare in occurrence. Condylar osteomas can be differentiated from CH radiographically, since osteomas tend to exhibit a mixed radiolucent-radiopaque appearance, unlike CH which are radiopaque¹⁸.

Condylar osteochondromas can be differentiated using CT imaging. In the case of condylar osteochondroma the coronal and sagittal CT sections tend to reveal a growth arising from the morphologically normal condyle in contrast to the uniform enlargement of condylar head which is characteristic of CH¹⁹.

Panoramic radiographs are useful for comparing both the condyles in a single image although the view is two dimensional²⁰. Panoramic radiographs are good for screening condyles but not considered suitable for the quantitative analysis of condyles and follow up of patients with unilateral CH²¹. We used a panoramic radiograph to screen our patient and evaluation of the panoramic view revealed hyperplasia of the right condyle. CT imaging aids in multiplanar imaging of the condyles²². A recently conducted retrospective CT based study revealed a significant increase in condylar length and other dimensions on the hyperplastic side when compared to the normal side²³. CH characteristically appears as a uniform enlargement of the condylar head^{19,24}. In our patient a beak like projection in the anteromedial direction was observed in the axial CT section which was atypical of CH.

Growth activity of the CH can be assessed using single-photon emission computed tomography (SPECT)²⁵. In SPECT the unilateral hyperplastic condyle is quantitatively compared to the normal contralateral side²⁶. A 0–5% difference in activity is usually observed between normal condyles. If the difference in activity is greater than 10% between two condyles, CH is suspected in the condyle with increased activity^{27,28}. SPECT could not be performed in our patient because of financial constraints. Prior to initiating treatment for patients with CH several factors such as the level of facial asymmetry, psycho-social consequences of the facial change, functional changes and malocclusion have to be considered⁴. Treatment options for CH include high condylectomy with or without orthognathic surgery and orthodontic treatment²⁹. Unfortunately, our patient was not willing to undergo surgical treatment of CH.

Conclusion

Prompt diagnosis is very important for successful management of CH. Apart from a patient's history and clinical findings, serial photographs of the patient from the past 10 to 15 years also provides vital information about the approximate time of occurrence and progression of the CH. Hence it is advisable to study serial photographs of patients with CH during the diagnostic stage.

Informed consent

Written informed consent for publication of their clinical details and clinical images was obtained from the patient.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

References

- Almeida LE, Zacharias J, Pierce S: **Condylar hyperplasia: An updated review of the literature.** *Korean J Orthod.* 2015; **45**(6): 333–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Iannetti G, Cascone P, Belli E, et al.: **Condylar hyperplasia: Cephalometric study, treatment planning, and surgical correction (our experience).** *Oral Surg Oral Med Oral Pathol.* 1989; **68**(6): 673–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hovell JH: **Condylar hyperplasia.** *Br J Oral Surg.* 1963; **1**: 105–111.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Olate S, Netto HD, Rodriguez-Chessa J, et al.: **Mandible condylar hyperplasia:**

- a review of diagnosis and treatment protocol.** *Int J Clin Exp Med.* 2013; **6**(9): 727–37.
[PubMed Abstract](#) | [Free Full Text](#)
5. Bharathi SC, Senthilnathan S, Kumar LD, *et al.*: **Unilateral condylar hyperplasia: A case report and review of literature.** *J Int Soc Prev Community Dent.* 2014; **4**(1): 67–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 6. Shankar U, Chandra S, Raju BH, *et al.*: **Condylar hyperplasia.** *J Contemp Dent Pract.* 2012; **13**(6): 914–7.
 7. Gelada K, Halli R, Hebbale M, *et al.*: **Unilateral Condylar Hyperplasia: A Case Report.** *Ann Clin Case Rep.* 2018; **3**: 1524.
[Reference Source](#)
 8. Hansson T, Oberg T, Carlsson GE, *et al.*: **Thickness of the soft tissue layers and the articular disk in the temporomandibular joint.** *Acta Odontol Scand.* 1977; **35**(2): 77–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
 9. Nitzan DW, Katsnelson A, Bermanis I, *et al.*: **The clinical characteristics of condylar hyperplasia: experience with 61 patients.** *J Oral Maxillofac Surg.* 2008; **66**(2): 312–318.
[PubMed Abstract](#) | [Publisher Full Text](#)
 10. Rajmakers PG, Karssemakers LH, Tuinzing DB: **Female predominance and effect of gender on unilateral condylar hyperplasia: a review and meta-analysis.** *J Oral Maxillofac Surg.* 2012; **70**(1): e72–e76.
[PubMed Abstract](#) | [Publisher Full Text](#)
 11. Talwar RM, Wong BS, Svoboda K, *et al.*: **Effects of estrogen on chondrocyte proliferation and collagen synthesis in skeletally mature articular cartilage.** *J Oral Maxillofac Surg.* 2006; **64**(4): 600–609.
[PubMed Abstract](#) | [Publisher Full Text](#)
 12. Yu S, Xing X, Liang S, *et al.*: **Locally synthesized estrogen plays an important role in the development of TMD.** *Med Hypotheses.* 2009; **72**(6): 720–722.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Mehrotra D, Dhasmana S, Kamboj M, *et al.*: **Condylar hyperplasia and facial asymmetry: report of five cases.** *J Maxillofac Oral Surg.* 2011; **10**(1): 50–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Norman JE, Painter DM: **Hyperplasia of the mandibular condyle. A historical review of important early cases with a presentation and analysis of twelve patients.** *J Maxillofac Surg.* 1980; **8**(3): 161–75.
[PubMed Abstract](#)
 15. López DF, Corral CM: **Condylar hyperplasia: characteristics, manifestations, diagnosis and treatment. A topic review.** *Rev Fac Odontol Univ Antioq.* 2015; **26**(21): 425–446.
[Reference Source](#)
 16. Wen B, Shen Y, Wang CY: **Clinical Value of 99Tcm-MDP SPECT Bone Scintigraphy in the Diagnosis of Unilateral Condylar Hyperplasia.** *Sci World J.* 2014; **2014**: 256256.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Tripathi T, Srivastava D, Neha, *et al.*: **Differential diagnosis and treatment of condylar hyperplasia.** *J Clin Orthod.* 2019; **53**(1): 29–38.
[PubMed Abstract](#)
 18. Valente L, Tieghi R, Mandrioli S, *et al.*: **Mandibular Condyle Osteoma.** *Ann Maxillofac Surg.* 2019; **9**(2): 434–438.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Andrade NN, Gandhewar TM, Kapoor P, *et al.*: **Osteochondroma of the mandibular condyle - Report of an atypical case and the importance of computed tomography.** *J Oral Biol Craniofac Res.* 2014; **4**(3): 208–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Kubota Y, Takenoshita Y, Takamori K, *et al.*: **Levandowski Panographic analysis in the diagnosis of hyperplasia of the coronoid process.** *Br J Oral Maxillofac Surg.* 1999; **37**(5): 409–411.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Nolte JW, Karssemakers LH, Grootendorst DC, *et al.*: **Panoramic imaging is not suitable for quantitative evaluation, classification, and follow up in unilateral condylar hyperplasia.** *Br J Oral Maxillofac Surg.* 2015; **53**(5): 446–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Mutoh Y, Ohashi Y, Uchiyama N, *et al.*: **Three-dimensional analysis of condylar hyperplasia with computed tomography.** *J Craniomaxillofac Surg.* 1991; **19**(2): 49–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. López DF, Botero JR, Muñoz JM, *et al.*: **Mandibular and temporomandibular morphologic characteristics of patients with suspected unilateral condylar hyperplasia: a CT study.** *Dental Press J Orthod.* 2020; **25**(2): 61–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Avinash KR, Rajagopal KV, Ramakrishnaiah RH, *et al.*: **Computed tomographic features of mandibular osteochondroma. case report.** *Dentomaxillofac Radiol.* 2007; **36**(7): 434–436.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. Alyamani A, Abuzinada S: **Management of patients with condylar hyperplasia: A diverse experience with 18 patients.** *Ann Maxillofac Surg.* 2012; **2**(1): 17–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Pripatnanont P, Vittayakittipong P, Markmanee U, *et al.*: **The use of SPECT to evaluate growth cessation of the mandible in unilateral condylar hyperplasia.** *Int J Oral Maxillofac Surg.* 2005; **34**(4): 364–368.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Saridin CP, Gilijamse M, Kuik DJ, *et al.*: **Evaluation of temporomandibular function after high partial condylectomy because of unilateral condylar hyperactivity.** *J Oral Maxillofac Surg.* 2010; **68**(5): 1094–1099.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Yang Z, Reed T, Longino BH: **Bone Scintigraphy SPECT/CT Evaluation of Mandibular Condylar Hyperplasia.** *J Nucl Med Technol.* 2016; **44**(1): 49–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Sembronio S, Tel A, Costa F, *et al.*: **An Updated Protocol for the Treatment of Condylar Hyperplasia: Computer-Guided Proportional Condylectomy.** *J Oral Maxillofac Surg.* 2019; **77**(7): 1457–1465.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 02 February 2021

<https://doi.org/10.5256/f1000research.51577.r78061>

© 2021 Castelino R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Renita Lorina Castelino

Department of Oral Medicine and Radiology, AB Shetty Memorial Institute of Dental Sciences, Mangalore, Karnataka, India

It is a well written manuscript covering all the required aspects for diagnosing a case of condylar hyperplasia. The authors have provided a well documented case substantiated with appropriate images and radiographs. This will help students and practitioners to comprehend the clinical and radiographic features of condylar hyperplasia and its progression.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Oral Medicine, Oral Radiology, Forensic odontology, Innovations in dentistry, 3D imaging and printing, Lasers.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 29 January 2021

<https://doi.org/10.5256/f1000research.51577.r78064>

© 2021 Battepati P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Prashant M. Battepati

Department of Preventive Dental Sciences, University of Taif, Taif, Saudi Arabia

The manuscript provides sufficient details regarding the specified case. Though the conclusion or the findings of the case report does not add to the available information, the case report as a whole can prove as a source of information for the practitioners looking for information related to such specific cases.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pediatric dentistry, Preventive Dentistry

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research