RESEARCH ARTICLE

REVISED Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey [version 2; peer review: 2 approved]

Dhan Shrestha¹, Suman Prasad Adhikari², Namrata Rawal²,
Pravash Budhathoki³, Subashchandra Pokharel⁴, Yuvraj Adhikari⁴,
Pooja Rokaya⁴, Udit Raut⁴

¹Department of Emergency Medicine, Mangalbare Hospital, Uurlabari, 1, 56600, Nepal

²Department of Neuro-psychiatry, Nepalese Army Institute of Health Sciences, Shree Birendra Hospital, Kathmandu, 3, 44600, Nepal

³Department of Emergency Medicine, Dr Iwamura Memorial Hospital, Bhaktapur, 3, 44800, Nepal

⁴Medical student, Nepalese Army Institute of Health Sciences, Kathmandu, 3, 44600, Nepal

v2 First published: 28 Jun 2021, 10:505
<https://doi.org/10.12688/f1000research.53904.1>
Latest published: 21 Jul 2021, 10:505
<https://doi.org/10.12688/f1000research.53904.2>

Abstract

Background

Good sleep quality is associated with a diverse range of positive outcomes such as better health, less daytime sleepiness, well-being and proper psychological functioning. Sleep deprivation or poor sleep quality leads to many metabolic, endocrine, and immune changes. Many studies have shown changes in sleep schedule along with worsening of sleep quality during the COVID-19 pandemic.

Methods

This cross-sectional study was conducted among students of a medical college in Kathmandu, Nepal from January 13, 2021 to February 15, 2021. A stratified random sampling technique was used. Data were collected using the Pittsburgh Sleep Quality Index (PSQI). Questionnaires that were completely filled were included and analyzed using STATA vs. 15.

Results

168(n=190) medical students filled out the questionnaires with a response rate of 88.42%. Around 30% (n=51) of respondents had poor sleep quality (PSQI total score of > 5) with an average PSQI score of 4.24±2.19. Unadjusted logistic regression analysis showed significantly higher odds of poor sleep quality among females (OR, 2.25; CI, 1.14-4.43) compared with male and the relation persists even after adjusting with age and year in medical school (aOR, 2.81; CI, 1.35-5.86). Adjusting with age and gender, 4th-year MBBS students had 82% lesser odds of having poor sleep quality compared to 2nd-year MBBS students (aOR, 0.18; CI, 0.04-0.76). Our study is suggestive of poor sleep quality common among medical students.

Open Peer Review

Approval Status

	1	2
version 2 (revision) 21 Jul 2021		
version 1 28 Jun 2021	view	view
1. Roshan Acharya , Cape Fear Valley Medical Center, Fayetteville, USA 2. Samata Nepal , Lumbini Medical College Teaching Hospital, Palpa, Nepal		
Any reports and responses or comments on the article can be found at the end of the article.		

Conclusions

More than a quarter of medical students have a poor sleep quality as per our study. So, education and awareness should be raised among medical students regarding the detrimental effects of poor quality sleep on daily activities, physical and mental well-being, and the overall quality of life.

Keywords

COVID-19, Medical students, Nepal, Sleep, Sleep Deprivation

Corresponding author: Dhan Shrestha (medhan75@gmail.com)

Author roles: **Shrestha D:** Conceptualization, Formal Analysis, Methodology, Software, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Adhikari SP:** Conceptualization, Methodology, Project Administration, Supervision, Validation, Writing – Review & Editing; **Rawal N:** Conceptualization, Methodology, Supervision, Validation, Writing – Review & Editing; **Budhathoki P:** Conceptualization, Data Curation, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Pokharel S:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Adhikari Y:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Rokaya P:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Raut U:** Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Shrestha D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Shrestha D, Adhikari SP, Rawal N *et al.* **Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey [version 2; peer review: 2 approved]** F1000Research 2021, 10:505 <https://doi.org/10.12688/f1000research.53904.2>

First published: 28 Jun 2021, 10:505 <https://doi.org/10.12688/f1000research.53904.1>

REVISED Amendments from Version 1

We have edited our manuscript as suggested by the reviewers. In the abstract section, we rectified the minor punctuation and grammatical errors that were present when we first submitted the manuscript. Similarly, we paraphrased a couple of sentences towards the end of abstract. We corrected the tenses of few sentences in the introduction section of the main text. Few redundant words and a sentence were removed and few words were added where necessary to add more meaning to the content. In the method section as well, redundant words and sentences were removed, rearrangement of words was necessary in some sentences and was done. Tenses, verb and voice (active/passive) of few sentences were modified as suggested by the reviewers. Similarly, minor punctuation and grammatical errors were corrected in the result and discussion sections along with paraphrasing of a sentence in the discussion section.

Any further responses from the reviewers can be found at the end of the article

1. Introduction

The novel coronavirus (COVID-19) that first appeared in Wuhan has been spreading around the world.¹ The World Health Organization declared it a pandemic in March 2020.² The pandemic has brought not only fear of infection & death but also unbearable psychological pressure and disturbances.^{3,4} Good sleep quality is associated with a diverse range of positive outcomes such as better health, less daytime sleepiness, well-being, and proper psychological functioning.⁵ Sleep deprivation or poor sleep quality leads to many metabolic, endocrine, and immune changes.⁶ So, the importance of good quality sleep cannot be stressed enough.

Many studies have shown changes in sleep schedule along with worsening of sleep quality during the lockdown.⁷ Findings regarding sleep duration and sleep quality are mixed during the COVID-19 pandemic.^{7–13} Most studies examining differences in sleep quality have shown poor sleep quality during the pandemic.^{12–16} Studies in different parts of the world showed that sleep quality has been poorer during lockdown period relative to the pre-lockdown period.^{7,10–12,14} When compared to pre-lockdown period, there has been a shift to a later bedtime and waking time, with a reduction in nighttime sleep and an increase in daytime napping during lockdown period. Also, there is evidence of greater sleep latency and poorer sleep efficiency during the pandemic.^{13,17} Especially, undergraduate medical students have poor sleep quality, internet addiction, and depression.^{18–20}

Sufficient studies have not been done in Nepal regarding the sleep quality of medical students during the COVID-19 pandemic. A study on Nepalese college students has shown the poor sleep quality among the majority of students.²¹ Higher prevalence of poor sleep quality among medical students as compared to other non-medical students and the general population has also been reported and several factors including medical students' attitudes, knowledge of sleep, and academic demands have been identified as the causative factors.²²

This study aims to assess the impact of this enormous change caused by the COVID-19 pandemic on the sleep quality of medical students of a medical college in Nepal during the COVID-19 pandemic.

2. Methods

2.1 Study design and settings

This is a cross-sectional study done among the undergraduate medical students (first to the fourth year) of Nepalese Army Institute of Health Sciences (NAIHS), Kathmandu, Nepal from January 13, 2021, to February 15, 2021. The Institutional Review Committee (IRC) of Nepalese Army Institute of Health Sciences approved the study (Reference number: 374). At the time of data collection, all the participants were informed about the study and its objectives. Consent was taken from the participants by incorporating the consent form in the questionnaire itself. So, all the participants are understood to have given consent (details of questionnaire and consent form attached as **extended data**).

The study participants were not recruited as front-liners to tackle the COVID-19 pandemic during or before the study period. All of them were restricted in their homes during the lockdown period. None of the participants had been infected by COVID-19 up until the study period. Data was collected from the participants after receiving their consent, via Google forms sent out by email explaining the objective of the study in the form itself. The participation was voluntary and anonymity was assured to the participants. The participants didn't receive any incentives.

2.2 Study sample

The sample size for the study was calculated using the Cochran formula. Total students from 1st to 4th year in NAIHS are 423. The calculated sample size was 169. Considering and adding 10% as a non-response rate, the final sample size was 186. The questionnaire was emailed to 190 participants, among which only 168 responded giving a response rate of 88.42%. Detail of sample size calculation is as follows:

Sampling method: Stratified random sampling

$$\begin{aligned}\text{Sample size: } n &= Z^2 * p * q / e^2 \\ &= (1.96)^2 * 0.76 * (1 - 0.76) / 0.05^2 \\ &= 280.28 \\ &= 280\end{aligned}$$

where,

n = calculated sample size

Z = 1.96 at 95% Confidence Interval

p = prevalence of poor sleep quality taken from previous study (76%)

q = $1 - p$

e = Margin of error (5%)

Total students from 1st to 4th year in NAIHS-COM (N) = 423

$$\begin{aligned}\text{Adjusted sample size } (n') &= n / [1 + \{(n - 1)/N\}] \\ &= 280 / [1 + \{(280 - 1)/423\}] \\ &= 168.72 \\ &= 169\end{aligned}$$

Considering and adding 10% as a non-response rate, our final sample size = 186.

Participants were selected using stratified random sampling in such a way that every student from first to fourth year had an equal chance of being selected. Firstly, a name list of students of the first to the fourth year was obtained from the administration of the institute, and each student was assigned a particular random number. An equal proportion of the male and female students from each year were selected for the study. Since the total number of students in each year was different, the total number of participants was different in each year. Then the study participants were selected randomly using the computer random number generator maintaining an equal proportion of students from each year and an equal proportion of males and females in each year [(first year: 45; female = 15, male = 30); (second year: 45; female = 13, male = 32); (third year: 50; female = 17, male = 33); (fourth year: 50; female = 15, male = 35)].

We emailed the questionnaire to the participants as everyone was at their homes because of the COVID-19 pandemic imposed lockdown.

2.3 Study instrument

We used the standard and validated Pittsburgh Sleep Quality Index (PSQI), which was developed by researchers at the University of Pittsburgh in 1988 AD. The questionnaire included baseline variables like age, sex, academic year, and questions addressing participants' sleep habits and quality i.e. PSQI. The PSQI assesses the sleep quality during the previous month and contains 19 self-rated questions that yield seven components: subjective sleep quality, sleep latency, sleep duration, sleep efficiency, sleep disturbance, use of sleeping medications and daytime dysfunction. Each component was assigned a score that ranges from zero to three, yielding a PSQI score in a range that goes from 0 to 21. A total score of 0 to 4 is considered as normal sleep quality; whereas, scores greater than 4 are categorized as poor sleep quality.²³ The questionnaire was typed in the Google form and was sent via email to the randomly selected study participants.

2.4 Statistical methods

Data collected from students through the Google forms was extracted to Google sheets, cleaned in Excel, and then imported and analyzed using STATA software v.15. Simple descriptive analysis was performed to see the response for every PSQI variable. Then calculation performed following PSQI form administration instructions (attached in **the extended data**). Logistic regression analysis was done taking PSQI score-based category as the dependent variable and age, gender, and years in medical school as the independent variable. Poor sleep quality was the outcome of interest so logistic regression analysis was run for the occurrence of poor sleep quality to good sleep quality. Firstly, binary logistic regression analysis was run across the sleep quality to gender, age, and year in medical school to estimate the unadjusted odds ratio (OR). Then multiple logistic regression ran to estimate adjusted OR across the sleep quality to gender, age, and year in medical school. For logistic regression purposes, PSQI score-based sleep quality was labeled as zero for good sleep quality and one for poor sleep quality, and odds of occurrence of poor quality of sleep to good sleep quality estimated.

3. Results

Among the mailed 190 individuals from 1st to 4th year of a medical school, only 168 filled the Google form making the response rate of 88.42%. The majority (n = 108, 64.29%) were male with a mean age of 21.57 ± 1.52 years. A majority (n = 110, 65.48%) of students were staying single while the rest used to share their room. The average sleep hour in the last month was $7:27:20.09 \pm 1:25:49.79$ hour and sleep latency was 24.92 ± 25.97 minutes. 30.36% (n = 51) of respondents were having poor sleep quality (PSQI total score of > five) with an average PSQI score of 4.24 ± 2.19 (**Table 1**).

Specific questions on trouble sleeping in last month showed 24(14.3%) were having trouble getting sleep within 30 minutes three or for more times a week. Similarly, 12(7.1%) mentioned they wake up in the middle of the night or early morning for three or more times a week. Majority i.e. 44% did not get up to use the bathroom during the past month. Likewise, only 1.2% and 1.8% had trouble sleeping because they could not breathe comfortably and coughed or snored loudly respectively for three or more times a week. When questioned if they had difficulty sleeping because of too cold or too hot feeling three or more times a week, 4.8% and 2.4% responded positively. The majority, 36.9%, and 75.6% did not have trouble sleeping because of bad dreams and of pain respectively during the last month. 98.8% of the participants did not have to take medicine to help him/her sleep during the last month. However, 1.2% had trouble staying awake while driving, eating meals, or engaging in social activities during the past month. Among those who had a roommate or bed partner, 6.9%, 1.7%, and 3.4% snored loudly, took long pause between breaths while asleep, and twitched or jerked legs while asleep respectively for three or more times a week during the past month whereas none of them had episodes of disorientation or confusion during sleep (**Table 2**).

Table 1. Participant's basic details and sleep quality.

Variables		Frequency	Percent
Gender	Female	60	35.71
	Male	108	64.29
Age: Mean $\pm$ SD = 21.57 ± 1.52 (Median: 22, IQR: 20-23)			
Medical school class	1st year, MBBS	39	23.21
	2nd year, MBBS	38	22.62
	3rd year, MBBS	47	27.98
	4th year, MBBS	44	26.19
Do you have a bed partner or roommate?	No bed partner or roommate	110	65.48
	Partner in the same bed	3	1.79
	Partner in the same room, but not the same bed	52	30.95
	Partner/roommate in other room	3	1.79
During the past month, hours of actual sleep (Mean $\pm$ SD): $7:27:20.09 \pm 1:25:49.79$ (Median: 7:30:00.00)			
During the past month, sleep latency (in minutes) (Mean $\pm$ SD): 24.92 ± 25.97 (Median: 15.5)			
PSQI total score (Mean $\pm$ SD): 4.24 ± 2.19 (Median:4)			
PSQI category (sum score < 5 or > 5)	Good sleep quality	117	69.64
	Poor sleep quality	51	30.36

Table 2. Factors affecting sleep in last month at the time of response to the survey.

PSQI scale based questionnaires	Less than once a week	Not during the past month	Once or twice a week	Three or more times a week	Total
5. During the past month, how often have you had trouble sleeping because of you...					
a. Cannot get to sleep within 30 minutes	29(17.3%)	77(45.8%)	38(22.6%)	24(14.3%)	168(100.0%)
b. Wake up in the middle of the night or early morning	48(28.6%)	75(44.6%)	33(19.6%)	12(7.1%)	168(100.0%)
c. Have to get up to use the bath-room	36(21.4%)	74(44.0%)	47(28.0%)	11(6.5%)	168(100.0%)
d. Cannot breathe comfortably	19(11.3%)	141(83.9%)	6(3.6%)	2(1.2%)	168(100.0%)
e. Cough or snore loudly	20(11.9%)	139(82.7%)	6(3.6%)	3(1.8%)	168(100.0%)
f. Feel too cold	41(24.4%)	95(56.5%)	24(14.3%)	8(4.8%)	168(100.0%)
g. Feel too hot	27(16.1%)	127(75.6%)	10(6.0%)	4(2.4%)	168(100.0%)
h. Had bad dreams	60(35.7%)	62(36.9%)	31(18.5%)	15(8.9%)	168(100.0%)
i. Have pain	26(15.5%)	127(75.6%)	10(6.0%)	5(3.0%)	168(100.0%)
7. During the past month, how often have you taken medicine to help you sleep (prescribed or "over the counter")?	1(0.6%)	166(98.8%)	1(0.6%)	-	168(100.0%)
8. During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in the social activity?	18(10.7%)	130(77.4%)	18(10.7%)	2(1.2%)	168(100.0%)
10. If you have a roommate or bed partner, ask him/her how often in the past month you have had...					
a. Loud snoring	9(5.5%)	42(72.4%)	3(5.2%)	4(6.9%)	58(100.0%)
b. Long pauses between breaths while asleep	5(8.6%)	50(86.2%)	2(3.4%)	1(1.7%)	58(100.0%)
c. Legs twitching or jerking while you sleep	8(13.8%)	41(70.7%)	7(12.1%)	2(3.4%)	58(100.0%)
d. Episodes of disorientation or confusion during sleep	4(6.9%)	52(89.7%)	2(3.4%)	-	58(100.0%)

Among 168 respondents, 111(66.1%) responded to have a fairly good sleep quality when asked to rate their overall sleep quality in the last month (Figure 1).

Fifty-eight (34.5%) responded no problem at all when asked for a problem to keep up enough enthusiasm to get things done during the past month, while 27 (16.1%) responded a very big problem (Figure 2).

Unadjusted logistic regression analysis showed significantly higher odds of poor sleep quality among females (OR, 2.25; CI, 1.14-4.43) comparing to male and the relation persisted even adjusting with age and year in medical school (aOR, 2.81; CI, 1.35-5.86) (Table 3). There was no significant difference in the sleep quality across years in medical school while running an unadjusted logistic regression analysis. But, adjusting with age and gender 4th-year MBBS students had 82% lesser odds of having a poor quality of sleep compared with 2nd-year MBBS students (aOR, 0.18; CI, 0.04-0.76) (Table 3).

4. Discussion

The COVID-19 pandemic in 2020 has imposed a substantial effect on psychological, health, and social issues all over the world. Medical students, like everyone else, have also suffered from these issues. Medical education is based more on a practical approach rather than just a theoretical core. Therefore, classes shifting from the classroom to online platforms have, without any doubt, affected medical education more than anything else. However, the impact of the pandemic on the psychological well-being and sleep quality of medical students has not been assessed and addressed to a good extent in

Figure 1. Response on “During the past month, how would you rate your sleep quality overall?”.

Figure 2. Response on “During the past month, how much of a problem has it been for you to keep up enough enthusiasm to get things done?”.

Nepal. Therefore, this study uses the PSQI to determine the sleep quality among the medical students of a medical college in Kathmandu, Nepal during the COVID-19 pandemic.

Our study showed that 30.36% of medical students had poor sleep quality (PSQI global score >5) but a study done just before the pandemic in the Nepalese medical students showed that 44.23% of students had poor sleep quality.²⁴ A study in India during the COVID-19 pandemic had 34.6% of medical students with poor sleep quality which was in line with the result of our study.²⁵ Previous studies from Pakistan on medical students in 2013 and Nepal on undergraduate students in 2015 showed 39.5% and 35.4% poor quality sleepers respectively.^{19,26} Our findings were in contrast to a study in China which showed a high prevalence of sleep disorders among adolescents and young adult students due to the stress and anxiety caused by the pandemic.²⁷ Previous studies in the Nepalese general population and health care workers found a low level of psychological distress due to the COVID-19 pandemic in Nepal which may explain better sleep quality among medical students in Nepal.^{28,29}

One of the positive findings of this study is that none of the participants had to take medication to help them sleep whereas 10.2% of Saudi Arabian physicians used sleeping pills once or twice a week during the COVID-19 pandemic.³⁰ 14.3% of our study participants had trouble falling asleep within 30 minutes thrice or more in a week, which was less compared to 24.7% in Saudi Arabian physicians during the COVID-19 pandemic.³⁰ Our study showed significantly higher odds of

Table 3. Binary and multiple logistic regression taking age, gender, and years in medical school for PSQI score.

Variables		Unadjusted			Adjusted		
		OR	[95% Conf. Interval]	p-value	OR	[95% Conf. Interval]	p-value
Years in Medical school	1st year, MBBS®						
	2nd year, MBBS	1.04	.4043604	2.674842	.7708897	.2731356	2.175736
	3rd year, MBBS	1.133333	.4640643	2.767816	.5756672	.174416	1.900014
	4th year, MBBS	.4444445	.1610749	1.226329	.1766567	.0412005	.7574558
Age		.9777632	.7871435	1.214545	1.361479	.9609949	1.928861
Gender	Male®						
	Female	2.252747	1.144979	4.432282	2.811422	1.348437	5.86167

Note ®: Reference category, 95% Confidence interval, and 5% standard error considered with p-value or <0.05 as a level of significance. * Significant result. PSQI score based category for sleep was assigned as 0/1 [good quality of sleep (0) to poor quality of sleep (1)].

poor quality sleep among female students (OR, 2.25; CI, 1.14-4.43). A study on Pakistani medical students in 2013 also showed that females had poorer quality sleep i.e. 44% as compared to 32.8% male poor sleepers.²⁶ Similarly, Goweda RA *et al.* also found that sleep disorders were more common among female medical students based on a study done in Saudi Arabia during the COVID-19 pandemic.³¹ Increased incidence of poor sleep quality among females is justifiable as sleep disorder symptoms are usually greater in women as compared to men.^{28,32} So, more focus should be shed upon female students when programs are formulated to improve the sleep quality among medical students. Sleep quality among students across years in medical school was not significantly different while running unadjusted logistic regression analysis in the case of our study. Studies from Brazil and Saudi Arabia assessed the sleep quality of first and second-year medical students to be poor as compared to other years.^{30,33} We found that the fourth-year medical students were less likely to have a poor sleep quality compared to second-year students which may be due to greater academic experience, and exposures during higher academic classes. In contrast, a previous study in China found that senior high school students had greater sleep problems due to increased academic burden and difficulty compared to junior school students.²⁷

Though our study shows less number of poor quality sleepers as compared to other studies discussed,^{19,24,25} 30.36% is still a big number. We cannot stay still and be satisfied based on the result of this study and the concerned authorities should try their best on bringing this number down to as minimum as possible. Because, poor quality sleep, without a doubt, will have a long-term impact on the mental well-being of the students. And no country would want future doctors who are not mentally healthy.

The number of students with poor sleep quality might decrease after the fear and restrictions because of COVID-19 subsides and life will return to normality. Alternatively, the number might go up, as the students return to their hectic schedules of classes, clinical postings, and a lot of studying. Since the scenario post-COVID is still unpredictable, frequent monitoring of sleep health and habits in medical students for raising awareness about sleep quality and problems should be planned and implemented.

Our study is not without limitations. We used a self-reporting questionnaire. Therefore, information bias is a major risk for this study. Our study was conducted among the students of a single institution. Therefore, the results might not be extrapolated to all the medical colleges of Nepal. Likewise, recall bias and subjectivity bias could have also affected the result of our study. Despite the limitations and biases, this study will surely provide a reference for further researches in this particular field.

5. Conclusions

Poor sleep quality is prevalent in 30.36% of medical students of a medical college in Nepal during the COVID-19 pandemic. Further studies have to be done to delve deeper into the sleep habits and problems of the medical students to determine the risk factors and causation of poor sleep quality. Furthermore, awareness should be raised about the importance of proper amount and sleep quality as well as detrimental effects of poor quality sleep on daily activities, physical and mental well-being, and the overall quality of life.

Data availability

Underlying data

Figshare. Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey. DOI: <https://doi.org/10.6084/m9.figshare.14770182.v2>.³⁴

This project contains the following underlying data:

- We used the standard and validated Pittsburgh Sleep Quality Index (PSQI), which was developed by researchers at the University of Pittsburgh in 1988 AD. The questionnaire included baseline variables like age, sex, academic year, and questions addressing participants' sleep habits and quality i.e. PSQI. The PSQI assesses the sleep quality during the previous month and contains 19 self-rated questions that yield seven components: subjective sleep quality, sleep latency, sleep duration, sleep efficiency and sleep disturbance, and daytime dysfunction. Each component is to be assigned a score that ranges from zero to three, yielding a PSQI score in a range that goes from 0 to 21. A total score of 0 to 4 is considered as normal sleep quality; whereas, scores greater than 4 are categorized as poor sleep quality.
- Data collected from students through the Google forms were extracted to Google sheets, cleaned in Excel, and then imported and analyzed using STATA 15. Simple descriptive analysis was performed to see the response for every PSQI variable. Then calculation performed following PSQI form administration instructions.

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC BY 4.0 Public domain dedication).

Extended data

Figshare. Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey. DOI: <https://doi.org/10.6084/m9.figshare.14770182.v2>.³⁴

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC BY 4.0 Public domain dedication).

Authors' contributions

DBS, SPA, NR, PB, SP, YA, PR, and UR contributed to the concept and design, methodology, and data collection. DBS contributed to the analysis, and interpretation of data. DBS, PB, SP, YA, PR, and UR contributed to the literature search, and initial manuscript drafting. SPA, NR involved in the revision and intellectual interpretation of the manuscript.

All authors were involved in drafting and revising the manuscript and approved the final version.

Acknowledgments

We would like to thank all the participants in this research for their voluntary participation.

References

- WHO | Novel Coronavirus – China: **WHO [Internet]**. 2020 [cited 2021 May 18].
[Reference Source](#)
- WHO Director-General's opening remarks at the media briefing on COVID-19-11 March 2020 [Internet]**. [cited 2021 May 18].
[Reference Source](#)
- Cao W, Fang Z, Hou G, *et al.*: **The psychological impact of the COVID-19 epidemic on college students in China.** *Psychiatry Res.* 2020 May 1; **287**: 112934.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Luo M, Guo L, Yu M, *et al.*: **The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public – A systematic review and meta-analysis.** *Psychiatry Res.* Elsevier Ireland Ltd; 2020; **291**: p. 113190.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harvey AG, Stinson K, Whitaker KL, *et al.*: **The subjective meaning of sleep quality: A comparison of individuals with and without insomnia.** *Sleep.* 2008 Mar 1 [cited 2021 May 18]; **31**(3): 383–393.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Reference Source](#)
- Aldabal L: **Metabolic, Endocrine, and Immune Consequences of Sleep Deprivation.** *Open Respir Med J.* 2011 Jun 30; **5**(1): 31–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cellini N, Canale N, Mioni G, *et al.*: **Changes in sleep pattern, sense of time and digital media use during COVID-19 lockdown in Italy.** *J Sleep Res.* 2020 Aug 1 [cited 2021 May 18]; **29**(4): e13074.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- yu LL, Wang J, yong O-y X, *et al.*: **The immediate impact of the 2019 novel coronavirus (COVID-19) outbreak on subjective sleep status.** *Sleep Med.* 2021 Jan; **77**: 348–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wright KP, Linton SK, Withrow D, *et al.*: **Sleep in university students prior to and during COVID-19 Stay-at-Home orders.** *Curr Biol.* CellPress; 2020; **30**: p. R797–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Coiro MJ, Asraf K, Tzischinsky O, *et al.*: **Sleep quality and COVID-19-related stress in relation to mental health symptoms among Israeli and U.S. adults.** *Sleep Heal.* 2021 Apr; **7**(2): 127–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blume C, Schmidt MH, Cajochen C: **Effects of the COVID-19 lockdown on human sleep and rest-activity rhythms.** *Curr Biol.* CellPress; 2020; **30**: p. R795–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Marelli S, Castelnuovo A, Somma A, *et al.*: **Impact of COVID-19 lockdown on sleep quality in university students and administration staff.** *J Neurol.* 2021 Jan; **268**(1): 8–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gupta R, Grover S, Basu A, *et al.*: **Changes in sleep pattern and sleep quality during COVID-19 lockdown.** *Indian J Psychiatry.* 2020 Jul; **62**(4): 370–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Casagrande M, Favieri F, Tambelli R, *et al.*: **The enemy who sealed the world: effects quarantine due to the COVID-19 on sleep quality, anxiety, and psychological distress in the Italian population.** *Sleep Med.* 2020 Nov; **75**: 12–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cellini N, Conte F, De Rosa O, *et al.*: **Changes in sleep timing and subjective sleep quality during the COVID-19 lockdown in Italy and Belgium: age, gender and working status as modulating factors.** *Sleep Med.* 2021 Jan; **77**: 112–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gao C, Scullin MK: **Sleep health early in the coronavirus disease 2019 (COVID-19) outbreak in the United States: integrating longitudinal, cross-sectional, and retrospective recall data.** *Sleep Med.* 2020 Sep; **73**: 1–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Benham G: **Stress and sleep in college students prior to and during the COVID-19 pandemic.** *Stress Heal.* 2020.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Attal BA, Bezdan M, Abdulqader A: **Sleep Quality and Its Correlates among Yemeni Medical Students: A Cross-Sectional Study.** *Sleep Disord.* 2021 Jan; **2021**: 1–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bhandari PM, Neupane D, Rijal S, *et al.*: **Sleep quality, internet addiction and depressive symptoms among undergraduate students in Nepal.** *BMC Psychiatry.* 2017 Mar 21 [cited 2021 May 18]; **17**(1): 1–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Reference Source](#)
- Aylie NS, Mekonen MA, Mekuria RM: **The psychological impacts of COVID-19 pandemic among university students in bench-sheko zone, South-West Ethiopia: A community-based cross-sectional study.** *Psychol Res Behav Manag.* 2020 Sep 30 [cited 2021 May 18]; **13**: 813–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

21. Khadka R, Bista S, Baskota S, *et al.*: **Sleep Quality among College Students in Kathmandu Valley, Nepal.** *Nepal Med J.* 2019 Dec; **2**(2): 1–4.
[Publisher Full Text](#)
22. Azad MC, Fraser K, Rumana N, *et al.*: **Sleep disturbances among medical students: A global perspective.** *J Clin Sleep Med.* American Academy of Sleep Medicine; 2015; **11**: p. 69–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Buysse DJ, Reynolds CF, Monk TH, *et al.*: **The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research.** *Psychiatry Res.* 1989 May 1; **28**(2): 193–213.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Sundas N, Ghimire S, Bhusal S, *et al.*: **Sleep quality among medical students of a tertiary care hospital: A descriptive cross-sectional study.** *J Nepal Med Assoc.* 2020 Feb 29 [cited 2021 May 18]; **58**(222): 76–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Reference Source](#)
25. Saraswathi I, Saikarthik J, Kumar KS, *et al.*: **Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: A prospective longitudinal study.** *PeerJ.* 2020 Oct 16 [cited 2021 May 18]; **8**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Surani AA, Zahid S, Surani A, *et al.*: **Sleep quality among medical students of Karachi.** *Pak. J Pak Med Assoc.* 2015 Apr; **65**(4): 380–2.
[PubMed Abstract](#)
27. Zhou SJ, Wang LL, Yang R, *et al.*: **Sleep problems among Chinese adolescents and young adults during the coronavirus-2019 pandemic.** *Sleep Med.* 2020 Oct 1 [cited 2021 Feb 20]; **74**: 39–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Shrestha DB, Thapa BB, Katuwal N, *et al.*: **Psychological distress in Nepalese residents during COVID-19 pandemic: A community level survey.** *BMC Psychiatry.* BioMed Central Ltd; 2020 [cited 2021 Feb 20]; **20**.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Kafle K, Shrestha DB, Baniya A, *et al.*: **Psychological distress among health service providers during COVID-19 pandemic in Nepal.** *PLoS One.* 2021 Feb 1 [cited 2021 May 18]; **16**(2 February): e0246784.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Alnofaiey YH, Alshehri HA, Alosaimi MM, *et al.*: **Sleep disturbances among physicians during COVID-19 pandemic.** *BMC Res Notes.* 2020 Dec 1 [cited 2021 May 18]; **13**(1): 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Goweda RA, Hassan-Hussein A, Alqahtani MA, *et al.*: **Prevalence of sleep disorders among medical students of Umm Al-Qura University, Makkah, Kingdom of Saudi Arabia.** *J Public Health Res.* 2020 [cited 2021 May 18]; **9**(S1): 45–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Klink M, Quan SF: **Prevalence of reported sleep disturbances in a general adult population and their relationship to obstructive airways diseases.** *Chest.* 1987 Apr 1 [cited 2021 May 18]; **91**(4): 540–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
33. de C CC, de Oliveira FK, Pizzamiglio DS, *et al.*: **Qualidade de sono em estudantes de medicina: Comparação das diferentes fases do curso.** *J Bras Pneumol.* 2017 Jul 1 [cited 2021 May 18]; **43**(4): 285–9.
[Publisher Full Text](#)
34. Shrestha D, Adhikari SP, Rawal N: **Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey.** *figshare.* Online resource. 2021.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 06 July 2021

<https://doi.org/10.5256/f1000research.57334.r88342>

© 2021 Nepal S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Samata Nepal

Department of Community Medicine, Lumbini Medical College Teaching Hospital, Palpa, Nepal

Abstract:

- Please use punctuation wherever required, e.g. "well-being and..." instead of "well-being, and..."
- "from January 13, 2021 to February..." instead of "from January 13, 2021, to February..."
- The sentence starting with a number is not appropriate in the Results section, second sentence. Please consider "Around 30% (n=51) of respondents..." or "Thirty percent (n=51) of respondents..."

Introduction:

- Paragraph 2, line 5: "Compared to the pre-lockdown...", this sentence needs to be paraphrased. Consider "When compared to pre-lockdown period, there was a later shift to bedtime and waking time, which reduced sleep during the night and increased daytime napping".
- Paragraph 4, line 1: The first sentence is not needed, please consider "This study aims..."

Methods:

- Paragraph 1, line 2: Consider "January 13, 2021 to February 15".
- Paragraph 3, line 3: "The questionnaire was...", please remove this sentence as the same sentence is repeated in the next paragraph.
- The last line of 2.2 is not required. "We emailed the..." has been mentioned already.
- 2.3 Study instrument, line 5: Please consider "component is assigned score..."
- 2.3 Study instrument, line 7: Please consider "The questionnaire was typed in the Google

form and...". The use of passive format is better.

- Please mention clearly if the questionnaire was emailed or the link of the Google form was emailed. Please consider using uniformity throughout the Methods section.
- 2.4 Statistical methods: line 1: "Google forms was extracted...".
- 2.4 Statistical methods, line 5: Please consider "Logistic regression analysis was done...". The use of passive format is better.
- Repeated use of the word "ran" in 2.4 can be avoided, e.g. "logistic regression analysis was run...", "multiple logistic regression ran...".
- Please consider uniformity in writing either "good/poor quality sleep" or "good/poor quality of sleep".

Results:

- Table 1: The table with the variable "Do you have a bed partner or roommate?" - the last row has a sub-variable: " Partner/roommate in another room". Please clarify this option.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public health

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 01 July 2021

<https://doi.org/10.5256/f1000research.57334.r88340>

© 2021 Acharya R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Roshan Acharya

Department of Internal Medicine, Cape Fear Valley Medical Center, Fayetteville, NC, USA

This study addresses one of the important problems prevalent among medical students. It has been conducted following the proper research methods. The title of the study is appropriate. The manuscript seems to be technically sound. The statistical analysis has been done appropriately. All the data underlying the findings are also made fully available. The conclusion includes the results, and recommendations are given accordingly. The paper seems to be well-written despite a few minor errors.

My comments regarding the paper are as follows:

Major correction: None

Minor corrections:

Abstract:

- *"Unadjusted logistic regression analysis showed significantly higher odds of poor quality of sleep among females (OR, 2.25; CI, 1.14-4.43) comparing to male and the relation persists even adjusting with age and year in medical school (aOR, 2.81; CI, 1.35-5.86)"*

Please put a full stop(.) after this sentence.

- *"So, education and awareness should be aroused among medical students regarding the detrimental effects of poor quality sleep on daily activities, physical and mental well-being, and the overall quality of life."*

In this sentence, the replacement of the word 'aroused' with 'raised' would be better.

Introduction:

- *"Studies in different parts of the world showed that sleep quality was poorer during the lockdown period relative to the pre-lockdown period. 7, 10-12, 14 Compared to the pre-lockdown period, there was a shift to a later bedtime and waking time, with a reduction in nighttime sleep and an increase in daytime napping during the lockdown period. Also, there was evidence of greater sleep latency and poorer sleep efficiency during the pandemic."*

In the second paragraph, you have used past tense. I think it would be better to use present tense instead.

Methods:

- *"The stratified random sampling method was used."*

This statement has been mentioned thrice in the methods section itself, once in **Study design and**

settings, twice in **Study sample**. Please mention it only once, wherever you think is suitable.

Study sample:

- *"Detail of sample size calculation is as follow:"*

Here 'as follows' is the correct one, please correct it.

Study instrument:

- *"Each component is to be assigned a scored that ranges from zero to three, yielding a PSQI score in a range that goes from 0 to 21."*

In this sentence, correct 'assigned a scored' to 'assigned a score'

Discussion:

- *"Our study showed that 30.36% of medical students had poor quality of sleep (PSQI global score >5) but a study has done just before the pandemic in the Nepalese medical students showed that 44.23% of students had poor sleep quality."*

Here, two points; First, please write 'study done' in place of 'study has done'.

You have discussed and compared the findings of the 2nd and the 4th year, but not compared them to other years. Also, there is no discussion on why the score of the 3rd year is the highest. Please address this as well.

Second: PSQI change it to 'equal or greater than 5'. Please see my last point for further clarification.

Conclusion:

- *" Furthermore, awareness should be aroused about the importance of proper amount and quality of sleep as well as detrimental effects of poor quality sleep on daily activities, physical and mental well-being, and the overall quality of life."*

In this sentence, the replacement of the word 'aroused' with 'raised' would be better.

Table 1:

In the methodology it was mentioned that: A total score of 0 to 4 is considered as normal sleep quality; whereas, scores greater than 4 are categorized as poor sleep quality. But in Table 1, it is otherwise: *PSQI category (sum score < 5 or > 5)*. It should be corrected in Table 1 for uniformity.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Clinical research

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research