SYSTEMATIC REVIEW

The effectiveness of EMR implementation regarding reducing documentation errors and waiting time for patients in outpatient clinics: a systematic review [version 1; peer review: 1 approved, 1 approved with reservations]

Salem Albagmi

Department of Health Information Management, Prince Sultan Military College of Health Sciences, P.O. Box 33048, Dammam, 31448, Saudi Arabia

V1 First published: 29 Jun 2021, 10:514
<https://doi.org/10.12688/f1000research.45039.1>
 Latest published: 11 Oct 2021, 10:514
<https://doi.org/10.12688/f1000research.45039.2>

Abstract

Background: Electronic medical records (EMRs) refer to the digital copies of paper notes prepared in the physician's office, outpatient clinics and other departments in health care institutes. EMRs are considered to be significant and preferable to paper records because they allow providers to keep accurate track of patient data and monitoring over time, thus reducing errors, and enhance overall health care quality. The aim of this systematic review was to highlight the significance of EMRs and the effectiveness of implementation regarding reducing documentation errors and waiting time for patients in outpatient clinics.

Methods: PubMed, Central, Ovid, Scopus, Science Direct, Elsevier, Cochrane, WHO website and the McMaster University Health Evidence website from 2005-2020 were searched to identify studies that investigated the association between the EMR implementation and documentation error and waiting time for patients. A reviewer screened identified citations and extracted data according to the PRISMA guidelines and data was synthesized in a narrative manner.

Results: After full text examination of the articles selected for this literature review, the major themes of relevance that were identified in the context of reducing documentation errors and waiting time for patients in outpatient clinic include: reduction of medical errors because of fewer documentation errors resulting from EMR implementation and reduction of waiting time for patients due to overall improvement of system workflow after use of EMRs.

Conclusion: In summary of the reviewed evidence from published material, the implementation of an EMR system in any outpatient setting appears to reduce documentation errors (medication dose errors, issues of prescription errors). It was also seen that in many settings, waiting time for patients in outpatient clinics was reduced

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
11 Oct 2021	view	view
version 1		
29 Jun 2021	view	view

1. **Clair M. Sullivan**, The University of Queensland, Brisbane, Australia

2. **Shankar Srinivasan** , Rutgers University, Newark, USA

Any reports and responses or comments on the article can be found at the end of the article.

with EMR use, while in other settings it was not possible to determine if any significant improvement was seen in this aspect after EMR implementation.

Keywords

EMR, medical documentation, documentation errors

This article is included in the **Health Services** gateway.

Corresponding author: Salem Albagmi (salbagmi@psmchs.edu.sa)

Author roles: Albagmi S: Conceptualization, Data Curation, Investigation, Methodology, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Albagmi S. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Albagmi S. **The effectiveness of EMR implementation regarding reducing documentation errors and waiting time for patients in outpatient clinics: a systematic review [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 10:514 <https://doi.org/10.12688/f1000research.45039.1>

First published: 29 Jun 2021, 10:514 <https://doi.org/10.12688/f1000research.45039.1>

Introduction

The general definition of electronic medical record (EMR) systems is an electronic record of health-associated data of a person which can be made, collected, organized and consulted by authorized doctors as well as health care staff in a health care institution. EMRs usually refer to digital copies of the paper notes made in the doctor's office, outpatient clinics and other health care institutions (Hyman *et al.*, 2012). EMRs consist of the doctor's official notes and other data or information documented by and for the doctors within that office, clinic, or health care institution (Manca, 2015). EMRs are generally utilized by the providers for diagnostic purposes and treatment (George and Bernstein, 2009). EMRs are far more important than real paper records since they allow the providers to keep accurate track of patient data over time, help in identifying patients who have to be scheduled for preventive care or vital screenings, allow consistent monitoring of patients, and enhance the overall provided health care quality (DesRoches *et al.*, 2008; Donsa *et al.*, 2016; Cloete, 2015).

The EMR system once implemented has the acumen to give significant advantages to doctors and outpatient clinical health centers as well as the patients and the overall system of health care institutes (Agrawal and Wu, 2009). The EMR system has been shown to smooth-out the workflow pathways while enhancing the overall quality of patient care as well as patient safety (DesRoches *et al.*, 2008; Franklin and Puaar, 2019). The benefits provided by the implementation of an EMR system in a health care facility are many and include the following:

- 1) Doctors and other health care staff like the nursing staff have quick and instant access to patient medical data or records like previous diagnoses, known drug reactions as well as allergies, laboratory results, and currently prescribed drugs (DesRoches *et al.*, 2008; Amato *et al.*, 2017).
- 2) It is possible for the outpatient clinic staff to gain access to new and past test results from other departments or even other providers in clinics that use multiple care services (Manca, 2015).
- 3) Computerized provider order entry (CPOE) is an easier and quicker way of entering and sending medical, laboratorial, radiological and pharmacy prescriptions than paper documents, thus reducing the probability of errors.
- 4) EMRs are linked to provision of computerized decision-support systems which are accurate in preventing adverse drug interactions and enhancing the overall use of best clinical practices in outpatient settings (Kavanagh, 2017).
- 5) Use of EMRs ensures that there is safe, private electronic communication between the health care providers and patients (Kavanagh, 2017).
- 6) The patient is also able to access their personal health records, disease management strategies as well as other needed health-related information sources which was not always possible with paper records (Noraziani *et al.*, 2013).
- 7) Implementation of an EMR system also computerizes the patient administration systems like their scheduling protocols (Noraziani *et al.*, 2013).
- 8) EMRs provide standardized and streamlined methods of storing electronic data and allow timely reporting that in turn improves patient safety as well as disease surveillance programs (Radley *et al.*, 2013; Noraziani *et al.*, 2013).

Medication errors and increased waiting time for patients due to manual documentation on paper have been found to be significantly associated with compromised health care quality (Keers *et al.*, 2013). There should be a system that can be used to minimize errors and improve the quality of life of patients. Literature is available discussing the magnificent contribution of EMRs and they have been found to improve the health care quality (Wali *et al.*, 2020; Lin *et al.*, 2020). However, limited studies are available discussing the variables that we have considered in our study as the outcome variable. Therefore, we conducted this systematic review to particularly focus on the reduction in documentation errors and patient waiting time in the outpatient clinic setting by the implementation of EMRs and provide an evidence-based infrastructure to healthcare institutions for better services and facilities.

Methods Reporting

The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines have been used in this literature review since this is an already validated methodology for choosing the final articles to be used from the results obtained within published literature relevant to this research topic (Albagmi, 2020). The PRISMA guidelines, as well as being the format for conducting and reporting systematic literature reviews, were selected due to the fact that the overall

conceptual ideas and the main aims of the research topic of this project are well suited to the PRISMA guidelines. In addition, in order to carry out an accurate or valid literature review for the purposes of summarizing the various aspects of a chosen public health topic, reviews have been documented as being of more relevance when using already validated methodologies like the PRISMA guidelines (Moher *et al.*, 2009; Page *et al.*, 2021).

Search strategy and eligibility criteria

The search strategy was performed in a manner that the methodology can be repeated and employed (see Box 1). A systematic search of the literature published between 2005 and 2020 was performed on PubMed, Central, Ovid, Scopus, Science Direct, Elsevier, Cochrane using free keywords. The reference lists of relevant articles (including primary and secondary results), and grey literature (including PROSPERO and reports of relevant stakeholder organizations - WHO website and the McMaster University Health Evidence website) were screened. The search for relevant articles to be included in the literature review was done after first choosing the appropriate key words and key concepts pertaining to the main research goals of this literature review.

The formulated keywords and concepts were: “efficiency or impact which EMR implementation has had in terms of reducing documentation errors in outpatient clinics” and “efficiency or impact which EMR implementation has had in terms of reducing the waiting time for patients in outpatient clinics.” In accordance with chosen search terms, the study search made use of the keywords as well as the phrases in combination with the Boolean operators “AND” and “OR” in the following manner for search in relevant databases: “efficiency or impact which EMR implementation has had in terms of reducing documentation errors in outpatient clinics” and/or “efficiency or impact which EMR implementation has had in terms of reducing the waiting time for patients in outpatient clinics” (Cronin, Ryan and Coughlan, 2008; Timmins & McCabe, 2005; Wakefield, 2014).

Each of the chosen databases and websites were used for the search keeping in mind the inclusion and exclusion criteria to find relevant published articles. The inclusion/exclusion criteria used were as follows.

Inclusion criteria:

1. All disease and health settings.
2. Articles in the English language.

Box 1. Search strategy (PubMed).

Search: **efficiency or impact which EMR implementation, outpatient clinics**

"efficiencies"[All Fields] OR "efficiency"[MeSH Terms] OR "efficiency"[All Fields] OR "efficiencies"[All Fields] OR "efficient"[All Fields] OR "efficiently"[All Fields] OR "efficients"[All Fields] OR ("impact"[All Fields] OR "impactful"[All Fields] OR "impacting"[All Fields] OR "impacts"[All Fields] OR "tooth, impacted"[MeSH Terms] OR ("tooth"[All Fields] AND "impacted"[All Fields]) OR "impacted tooth"[All Fields] OR "impacted"[All Fields]) AND ("empir musical rev"[Journal] OR "emr"[All Fields]) AND ("implementability"[All Fields] OR "implementable"[All Fields] OR "implementation"[All Fields] OR "implementation s"[All Fields] OR "implementational"[All Fields] OR "implementations"[All Fields] OR "implementer"[All Fields] OR "implementers"[All Fields] OR "implementation"[All Fields]) AND ("ambulatory care facilities"[MeSH Terms] OR ("ambulatory"[All Fields] AND "care"[All Fields] AND "facilities"[All Fields]) OR "ambulatory care facilities"[All Fields] OR ("outpatient"[All Fields] AND "clinics"[All Fields]) OR "outpatient clinics"[All Fields])

Translations

efficiency: "efficiencies"[All Fields] OR "efficiency"[MeSH Terms] OR "efficiency"[All Fields] OR "efficiencies"[All Fields] OR "efficient"[All Fields] OR "efficiently"[All Fields] OR "efficients"[All Fields]

impact: "impact"[All Fields] OR "impactful"[All Fields] OR "impacting"[All Fields] OR "impacts"[All Fields] OR "tooth, impacted"[MeSH Terms] OR ("tooth"[All Fields] AND "impacted"[All Fields]) OR "impacted tooth"[All Fields] OR "impacted"[All Fields]

EMR: "Empir Musical Rev"[Journal: _jid101513485] OR "emr"[All Fields]

implementation,: "implementability"[All Fields] OR "implementable"[All Fields] OR "implementation"[All Fields] OR "implementation's"[All Fields] OR "implementational"[All Fields] OR "implementations"[All Fields] OR "implementer"[All Fields] OR "implementers"[All Fields] OR "implementation"[All Fields]

outpatient clinics: "ambulatory care facilities"[MeSH Terms] OR ("ambulatory"[All Fields] AND "care"[All Fields] AND "facilities"[All Fields]) OR "ambulatory care facilities"[All Fields] OR ("outpatient"[All Fields] AND "clinics"[All Fields]) OR "outpatient clinics"[All Fields]

3. Peer-reviewed articles published 2005-2020.
4. Studies that reported the impact of EMR in reducing documentation error and patient's waiting time in outpatient clinic settings, either in segregation or as one of the reporting outcomes in a multicomponent studies, were included.

Exclusion criteria:

1. Reminder services for appointment and other healthcare activities were not included.
2. Conference proceeding and abstracts were also excluded.

Study selection and data collection

Studies were independently assessed by a single reviewer (SA) and were reported using the PRISMA flow diagram (Figure 1). Initial screening of studies was based on the information mentioned in the titles and abstracts. Full-paper screening was conducted by the same independent reviewer and extracted in to a standardized structure. A form was created, focused on the detail study characteristics. Primarily, the reviewer collected the data in the form and subsequently it was managed in a master Microsoft Excel sheet (2019) and then tabulated separately for each study. Study investigators were not approached to confirm the data as the included studies were well-defined and outcome was clearly mentioned.

Data items

The data collected for each study included the name of first author, year of publication, setting characteristics, study duration, participants and, outcomes.

Primary outcomes

Primary outcomes were any measure related to effectiveness of EMR in:

1. Reducing documentation error,
2. Reducing patient waiting time.

Figure 1. The PRISMA flowchart showing search and selection methodology, PRISMA 2020 flow diagram.

Table 1. Definition of variables.

Variable	Definition
Electronic medical records (EMRs)	EMRs usually refer to digital copies of the paper notes made in the doctor's office, outpatient clinics and other health care institutions
EMR implementation	Use of a electronic software for the documentation by replacing the existing manual documentation method on paper.
Study designs	Type of the study design that was used to conduct the research for example observational studies such as cross sectional, case control studies, cohort studies and interventional studies such as randomized controlled trials. Also the type of reporting the research study that is primary (original articles) and secondary researches (review articles).
Outcome variables	
Impact of EMR on Documentation error	The absence of or decreased in the number of documentation error
Impact of EMR on Patient's waiting time	Time required to complete the documentation is decreased while patient is waiting. Hence the overall patient's waiting time is decreased.

Definitions of variables that were sought out are described in [Table 1](#).

Data analysis and risk of bias assessment

This was a systematic review and hence did not require any statistical analysis. Data was synthesized in a narrative manner. Study quality and bias assessment was done by using a hybrid of the Downs and Black scale and the Newcastle-Ottawa Scale that includes 11 items ([Downs et al., 1998](#); [Wells et al., 2017](#)) under four themes of reporting, external validity, internal validity for bias and internal validity for confounding to capture the observational and experimental study designs

Results

Search results

A search of the above-mentioned chosen databases and sources and the results obtained are shown in [Figure 1](#) (with the use of the PRISMA flowchart for the search strategy for the research question). This literature search made use of many relevant and varied sources so as to obtain the most accurate articles.

After the search was done, a total of 93 articles were found from the databases and other sources. After examining for and removing the duplicates, 30 articles were left. Through the application of the above-mentioned inclusion or exclusion criteria, nine articles were excluded. This resulted in a remainder of 21 articles which were further examined for relevance to the aims of this study's topic. Following this, nine articles were further excluded for not having relevance in determining the effectiveness of EMR implementation in terms of decreasing or reducing documentation errors and waiting time for patients in outpatient clinics. The remaining 12 articles were utilized for this literature review on the basis of their being relevant to the main aims of this research study (see [Figure 1](#) for the PRISMA flowchart) ([Moher et al., 2009](#); [Page et al., 2021](#)).

Selected study findings and data

The findings and data of the articles chosen for inclusion in this literature review are as shown in [Table 2](#).

Risk of bias assessment

The results of the risk of bias assessment are presented in [Figure 2](#). For question number 6, not all studies recruited participants; out of 12, three studies recruited participants and other studies evaluated databases.

Key themes identified in the data findings

The review of the selected articles revealed two key themes in the context of how effective the implementation of the EMR system is in reducing documentation errors and waiting time for patients in outpatient clinics, namely: there is a reduction in medical errors because of fewer documentation errors and there is some degree of reduction of waiting time for patients due to overall improvement of system workflow in some settings, after use of EMRs. The two themes tend to overlap and are interlinked. Thus, each aspect has an impact on the other factors and may ultimately improve care quality for patients attending outpatient visits.

Table 2. The findings and data of the articles chosen for inclusion in this literature review.

#	First author	Year of publication	Study design	Risk of bias	Outcome
First group of studies: Reduction of medical errors because of fewer documentation errors resulting from electronic medical record (EMR) implementation					
1	Priestman, W	2018	Retrospective literature review	Documentation error is studied as one of the variables in this study	312 potential articles were found in initial search, and 117 were used in the review. The findings show that in general, EMR implementation is related to enhancements in documentation and thus leads to decreased prescribing errors. (Priestman <i>et al.</i> , 2018).
2	Manca, P.	2015	Retrospective review	It was a short review and sufficient data was not included.	<p>In summary the findings of this study show that EMRs enhance quality of care, patient outcomes, and safety by reducing documentation errors through better management, reduction in medication errors, reduction in unnecessary testing and better communication between the primary care providers, patients, and other providers linked in care provision.</p> <ul style="list-style-type: none"> • EMRs have improved efficiencies in workflow and waiting times for patients by decreasing the time needed to pull charts, improved access to full patient data, help in managing prescriptions, improving scheduling of patient appointments, and provision of remote access to patients' charts. • EMRs recorded point-of-care data which informed and improved practice through quality improvement programs and use of practice-level interventions for reducing documentation error and improving patient safety (Manca, 2015).
3	DesRoches, C. M	2008	Survey study design	It was survey, may caused response bias and conducted on only ambulatory care setting.	<p>The implementation of EMRs in ambulatory or outpatient clinics by doctors in the USA was found to have a positive effect in reducing documentation errors, which helped in improving many aspects of overall provided patient care by enhancing these factors:</p> <ul style="list-style-type: none"> -Quality of clinical decisions and quality of faster communication with other providers -Quality of communication with patients -Timely access to medical records -Avoiding medication errors in documentation (DesRoches <i>et al.</i>, 2008).
4	George, J	2009	Retrospective literature review	Conducted on only prenatal network.	This study documented that use of EMRs within a prenatal network resulted in a decrease of documentation-related errors and corresponding risks (George and Bernstein, 2009).

Table 2. Continued

#	First author	Year of publication	Study design	Risk of bias	Outcome
5	Agrawal, A	2009	Pilot study design that evaluated the implementation of an EMR system as an intervention	Categorized discrepancy as a severity score according to potential cure and harm was not evaluated.	The findings and evaluation data documented that use of the EMR system coordinated documentation records in a multidisciplinary process based in both outpatient departments and inpatient wards. This resulted in reduced medication errors on admission. This suggests that EMR systems can be a vital tool in improving patient safety (Agrawal and Wu, 2009).
6	Radley, D. C	2013	Systematic literature review	It was a review and every study had the different definition of medication error.	The findings of this study showed that implementation of EMR resulted in medication error reduction in many outpatient clinical settings through corresponding reduction of documentation errors in the electronic prescribing done using computerized provider order entry systems (Radley et al., 2013).
7	Schwartzberg, D	2015	Retrospective literature review	Short period, missing data	The EMR system was implemented with the aim of reducing and eliminating iatrogenic injury caused by avoidable documentation errors in the hospital's paper-order medication entries. The findings show that the EMRs were useful in reducing errors (Schwartzberg et al., 2015).
Second group of studies: Reduction of waiting time for patients due to overall improvement of system workflow after use of EMRs					
8	Jabour, A. M	2020	Observational study design	Data was not stratified according to patient's condition and demographics.	The results showed no significant difference in the amount of time spent by patients in the reception area ($P=0.26$), in the waiting area ($P=0.57$), consultation time ($P=0.08$), and at the pharmacy ($P=0.28$) between the electronic health record and paper based groups. However, there was a significant difference ($P<0.001$) in the amount of time spent on all tasks between the primary healthcare centres located in metropolitan and rural areas. The longitudinal observation also showed reduction in the registration time (from 5.5 [SD 3.5] min to 0.9 [SD 0.5] min), which could be attributed to the introduction of a Web-based booking system (Jabour, 2020).
9	AlSarheed, A. H.	2016	Survey study design	Response bias	The study results suggest that software improvements and enhancement of the EMR system in a Saudi hospital's outpatient clinics reduced the waiting time for their patients (AlSarheed, 2016).

Table 2. *Continued*

#	First author	Year of publication	Study design	Risk of bias	Outcome
10	Cho, K. W	2017	Application of theoretical analysis	Only 03 hospitals were examined and waiting time varied with hospital size.	The authors made use of digital data collected from outpatients' reception times and consultation finish times* for calculating the arrival and service rates, respectively (through application of the queuing theory for analysis of outpatients' waiting times). The findings of this study verified that the implementation of EMR contributed to the enhancement of patient services by reducing outpatients' waiting time, or by increasing efficiency (Cho <i>et al.</i> , 2017).
11	Vahdat, V	2018	Quantitative study design utilizing a discrete-event simulation model	Conducted only in dermatology clinic.	The findings suggest that small changes to processes like addition of a few minutes for extra documentation time within the exam room (due to use of EMRs) resulted in significant delays in the timeliness of patient care (Vahdat <i>et al.</i> , 2018).
12	Noraziani, K	2013	Retrospective literature review	Lack of interoperability	One of the findings of this review is that use of EMRs in outpatient clinics may save time as it reduces patient waiting times through faster, easier workflow and organizational workflow efficiency (Noraziani <i>et al.</i> , 2013).

Figure 2. Quality assessment by Downs and Black scale and the Newcastle-Ottawa Scale.

Discussion

The main objectives of this study were to do a literature review to identify the effectiveness of EMR implementation in regard to reducing documentation errors and waiting time for patients in outpatient clinics. The data collected and the emerging themes indicate that there is a decrease in medical errors as a direct result of fewer documentation errors in EMR use (Priestman *et al.*, 2018; DesRoches *et al.*, 2008; Manca, 2015; Radley *et al.*, 2013) and there is a decrease in waiting time for patients because of overall improvement of system workflow in some settings, after use of EMRs (Jabour, 2020; AlSarheed, 2016; Cho *et al.*, 2017; Noraziani *et al.*, 2013; Jin, Sivakumar and Lim, 2013), although a contrasting increase was observed in some outpatient studies (Vahdat *et al.*, 2018).

The implementation of EMRs in outpatient or ambulatory clinics by providers in the US in one study showed a beneficial effect in reducing documentation errors, which in turn enhanced several factors of overall patient care. Improvements were seen in the quality of clinical decisions, quick communication between providers as well as with patients, timely access to all patient medical records in one place and prevention of medication errors in documentation (DesRoches *et al.*, 2008). Many studies have shown that implementation of an EMR permits doctors to avoid documentation errors since there is improved access to comprehensive patient histories such as clinical data (Ruano *et al.*, 2016). This in turn reduces waiting time for the patients as doctors spend less time looking for lab reports and other patient data.

The EMR implementation thus provides advantages like fast remote access to patient history, enhanced laboratory result access, medication error alerts, and reminders for scheduling preventive care (Manca, 2015; Radley *et al.*, 2013). It was also observed that implementation of various forms of the EMR systems can be a vital tool in improving patient safety in outpatient and inpatient settings (Agrawal and Wu, 2009). As a point of comparison, it was seen that researchers studied and compared the impact of errors in paper-based and computerized EMR diabetes management with decision support in hospitalized patients with type 2 diabetes (Donsa *et al.*, 2016; Vishwanath, Singh and Winkelstein, 2010). This could be applicable in outpatient clinics too. The results from one study suggest that software improvements and enhancement of the EMR system in a Saudi hospital's outpatient clinics reduced the waiting time for their patients (Al Sarheed, 2016).

However, some studies have shown contrasting results to those showing improvements in waiting times of outpatient clinic visitors. A study designed to quantify the downstream effect on patient wait times as well as the total duration of stay because of tiny increments in encounter times resulting from implementation of a new EMR system indicated that use of EMRs added extra documentation time in outpatient clinics and overall led to significant delays in the waiting times for patients (Vahdat *et al.*, 2018).

Limitations

This review has some limitations. First, due to the strict exclusion and inclusion criteria, the studies that were relevant according to the one of the outcomes but not to the main question were not included. Conference proceedings were excluded that might have contained findings of worth. Second, there was a lack of data within studies regarding setting, type of study and population. In addition, a large number of studies were found on documentation errors as compared to patient waiting time.

Conclusion

In summary, EMR implementation appears to enhance documentation, screening performance as well as prescription error prevention. The findings presented in this literature review will be of value to outpatient clinics and units that are looking to implement EMR systems. The waiting time for outpatient clinic visits can also be reduced by EMR use as there is implementation of new tools and functionality like the Web-based booking system, which can reduce the waiting time as well as duration of patients' visits (Jabour, 2020; Noraziani *et al.*, 2013). However, this needs to be further validated in future studies as some studies actually indicate that implementation of an EMR system led to increased waiting times and delays in care provision in outpatient clinics.

Data availability

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Reporting guidelines

Figshare: PRISMA checklist for "The effectiveness of EMR implementation regarding reducing documentation errors and waiting time for patients in outpatient clinics: a systematic review" <https://doi.org/10.6084/m9.figshare.14791356.v1> (Albagmi, 2021).

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

References

- Adurota O: *Implementation of Electronic Health Records in an Outpatient Clinic in Maryland*. Doctoral dissertation, Brandman University; 2019.
- Agrawal A, Wu WY: **Reducing medication errors and improving systems reliability using an electronic medication reconciliation system.** *Jt Comm J Qual Patient Saf.* 2009; **35**(2): 106–114.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Albagmi S: **May PRISMA_2020_checklist_45039.pdf.** *figshare.* Figure. 2021.
[Publisher Full Text](#)
- AlSarheed AH: *The Impact of Enhancing Outpatient Clinic Management Software in Reducing Waiting Time in Saudi Hospitals*. Doctoral dissertation, Prince Sultan University; 2016. Retrieved on 20th September, 2020.
[Reference Source](#)
- Amato MG, Salazar A, Hickman TTT, *et al.*: **Computerized prescriber order entry-related patient safety reports: analysis of 2522 medication errors.** *J Am Med Inform Assoc.* 2017; **24**(2): 316–322.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cho KW, Kim SM, Chae YM, *et al.*: **Application of queuing theory to the analysis of changes in outpatients' waiting times in hospitals introducing EMR.** *Health Inform Res.* 2017; **23**(1): 35–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cloete L: **Reducing medication errors in nursing practice.** *Cancer Nursing Practice.* 2015; **14**(1).
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cronin P, Ryan F, Coughlan M: **Undertaking a literature review: a step-by-step approach.** *Br J Nurs.* 2008; **17**(1): 38–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Downs SH, Black N: **The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions.** *J Epidemiol Community Health.* 1998; **52**(6): 377–384.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DesRoches CM, Campbell EG, Rao SR, *et al.*: **Electronic health records in ambulatory care—a national survey of physicians.** *N Engl J Med.* 2008; **359**(1): 50–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Donsa K, Beck P, Höll B, *et al.*: **Impact of errors in paper-based and computerized diabetes management with decision support for hospitalized patients with type 2 diabetes. A post-hoc analysis of a before and after study.** *Int J Med Inform.* 2016; **90**: 58.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Franklin BD, Puaar S: **What is the impact of introducing inpatient electronic prescribing on prescribing errors? A naturalistic stepped wedge study in an English teaching hospital.** *Health Informatics J.* 2019; **1460458219833112**.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hyman D, Laire M, Redmond D, *et al.*: **The use of patient pictures and verification screens to reduce computerized provider order entry errors.** *Pediatrics.* 2012; **130**(1): e211–e219.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jabour AM: **The Impact of Electronic Health Records on the Duration of Patients' Visits: Time and Motion Study.** *JMIR Med Inform.* 2020; **8**(2): e16502.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jin X, Sivakumar AI, Lim SY: **A simulation based analysis on reducing patient waiting time for consultation in an outpatient eye clinic.** *2013 Winter Simulations Conference (WSC).* IEEE; 2013; (pp. 2192–2203).
[Publisher Full Text](#)
- Kavanagh C: **Medication governance: preventing errors and promoting patient safety.** *Br J Nurs.* 2017; **26**(3): 159–165.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Keers RN, Williams SD, Cooke J, *et al.*: **Causes of medication administration errors in hospitals: a systematic review of quantitative and qualitative evidence.** *Drug Saf.* 2013; **36**(11): 1045–1067.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lin HL, Wu DC, Cheng SM, *et al.*: **Association between Electronic Medical Records and Healthcare Quality.** *Medicine.* 2020; **99**(31).
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Manca DP: **Do electronic medical records improve quality of care? Yes.** *Can Fam Physician.* 2015; **61**(10): 846.

[PubMed Abstract](#) | [Free Full Text](#)

Moher D, Liberati A, Tetzlaff J, et al.: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *Ann Intern Med.* 2009; **151**(4): 264–269.

[PubMed Abstract](#) | [Publisher Full Text](#)

Noraziani K, Nurul'Ain A, Azhim MZ, et al.: **An overview of electronic medical record implementation in healthcare system: Lesson to learn.** *World Applied Sciences J.* 2013; **25**(2): 323–332.

[Publisher Full Text](#)

Page MJ, Moher D, Bossuyt PM, et al.: **PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews.** *bmj.* 2021; 372.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Priestman W, Sridharan S, Vigne H, et al.: **What to expect from electronic patient record system implementation: lessons learned from published evidence.** *J Innov Health Inform.* 2018; **25**(2): 92–104.

[PubMed Abstract](#) | [Publisher Full Text](#)

Radley DC, Wasserman MR, Olsho LE, et al.: **Reduction in medication errors in hospitals due to adoption of computerized provider order entry systems.** *J Am Med Inform Assoc.* 2013; **20**(3): 470–476.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Ruano M, Villamañán E, Pérez E, et al.: **New technologies as a strategy to decrease medication errors: how do they affect adults and children differently?** *World J Pediatr.* 2016; **12**(1): 28–34.

[PubMed Abstract](#) | [Publisher Full Text](#)

Schwartzberg D, Ivanovic S, Patel S, et al.: **We thought we would be perfect: medication errors before and after the initiation of**

Computerized Physician Order Entry. *J Surg Res.* 2015; **198**(1): 108–114.

[PubMed Abstract](#) | [Publisher Full Text](#)

Singh H, Graber ML, Kissam SM, et al.: **System-related interventions to reduce diagnostic errors: a narrative review.** *BMJ Qual Saf.* 2012; **21**(2): 160–170.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Timmins F, McCabe C: **How to conduct an effective literature search.** *Nurs Stand.* 2005; **20**(11): 41–47.

[PubMed Abstract](#) | [Publisher Full Text](#)

Vahdat V, Griffin JA, Stahl JE, et al.: **Analysis of the effects of EHR implementation on timeliness of care in a dermatology clinic: a simulation study.** *J Am Med Inform Assoc.* 2018; **25**(7): 827–832.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vishwanath A, Singh SR, Winkelstein P: **The impact of electronic medical record systems on outpatient workflows: a longitudinal evaluation of its workflow effects.** *Int J Med Inform.* 2010; **79**(11): 778–791.

[PubMed Abstract](#) | [Publisher Full Text](#)

Wakefield A: **Searching and critiquing the research literature.** *Nurs Stand.* 2014; **28**(39): 49–57.

[PubMed Abstract](#) | [Publisher Full Text](#)

Wali RM, Alqahtani RM, Alharazi SK, et al.: **Patient satisfaction with the implementation of electronic medical Records in the Western Region, Saudi Arabia, 2018.** *BMC Fam Pract.* 2020; **21**(1): 1–6.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wells GA, Shea B, O'Connell DA, et al.: **The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses.** 2000.

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 03 August 2021

<https://doi.org/10.5256/f1000research.48111.r89947>

© 2021 Srinivasan S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shankar Srinivasan

School of Health Professions, Rutgers University, Newark, NJ, USA

The manuscript describes the systematic review of published literature on the effectiveness of electronic health records in reducing documentation caused errors and also waiting time for ambulatory settings. The duration chosen is quite appropriate (2005 to 2020), given that the widespread implementation (at least in the US) was from 2009 onwards. The author has done due diligence with the selection of the manuscripts for the review by including (and specifying) the appropriate inclusion and exclusion criteria so that another researcher can reproduce the study accordingly. The analyses, results and conclusions drawn are robust and well written in responding specifically to the outcomes in question and also the results for those two outcomes.

The author, however, hasn't specified or explained why they chose those two outcomes and not others, such as prescription contraindication caused issues, quality of point of care, and so on, which are also included in much literature on EHR/EMR effectiveness.

Secondly, the manuscript would have been better enhanced with some recommendations for future implementations of EHR/EMR.

Lastly, is there a difference in the types of ambulatory settings (surgical, physical therapy, urgent care)? These would have enhanced the quality and usefulness of the paper but they do not detract from the overall quality and rigor of the work reported in the existing manuscript and the way it was conducted.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Health Outcomes Research, Health Data Analytics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 17 Sep 2021

Salem Albagmi

Comment 1:

The author, however, hasn't specified or explained why they chose those two outcomes and not others, such as prescription contraindication caused issues, quality of point of care, and so on, which are also included in much literature on EHR/EMR effectiveness.

- **Page No. 5, Line No. 97-107, Actual Content**
- **Removed/Edited/Replaced Content:** Rationale for selecting both variables in the introduction section

Comment 2:

Secondly, the manuscript would have been better enhanced with some recommendations for future implementations of EHR/EMR.

- **Page No. 19, Line No. 295-310, Actual Content**
- **Removed/Edited/Replaced Content:** Future perspective of EHR is added

Comment 3:

Lastly, is there a difference in the types of ambulatory settings (surgical, physical therapy, urgent care)? These would have enhanced the quality and usefulness of the paper but they do not detract from the overall quality and rigor of the work reported in the existing manuscript and the way it was conducted.

- **Page No. 17, Line No. 269-274, Actual Content**
- **Removed/Edited/Replaced Content:** Discussion related to EMR/EHR system is added in the context of types

Competing Interests: Non-Financial Competing Interests

Reviewer Report 19 July 2021

<https://doi.org/10.5256/f1000research.48111.r88680>

© 2021 Sullivan C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Clair M. Sullivan

Centre for Health Services Research, Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia

Thank you for this paper.

A few overarching comments:

1. This paper is very doctor centric. Although it is true that EMRs contain doctor's notes, they actually allow integration of notes from all members of a multidisciplinary team. Please review the paper to include this.
2. The two outcome variables chosen, reduced waiting time and reduced documentation errors, are both important; but I am curious as to why these two were selected out of the 100s of available impacts? Is there a logic behind this? If so, please outline as it appears a little random currently.
3. The terms ambulatory care and outpatients can be confusing. Are you referring to specialised ambulatory care services? Or primary care? These settings are vastly different and so it is difficult to compare a primary care provider in general practice to a specialised neurosurgery outpatients department.
4. Please review grammar and syntax as this appears a little clumsy at times e.g avoid calling EMRs magnificent

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Partly

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 Sep 2021

Salem Albagmi

Comment 1:

This paper is very doctor centric. Although it is true that EMRs contain doctor's notes, they actually allow integration of notes from all members of a multidisciplinary team. Please review the paper to include this.

- **Page No. 3, Line No. 62-66, Actual Content**
- **Removed/ Edited/ Replaced Content:** The following aspect is added in the introduction section

Comment 2:

The two outcome variables chosen, reduced waiting time and reduced documentation errors, are both important; but I am curious as to why these two were selected out of the 100s of available impacts? Is there a logic behind this? If so, please outline as it appears a little random currently.

- **Page No. 5, Line No .97-107, Actual Content**
- **Removed/ Edited/ Replaced Content:** Rationale for selecting both variables in the introduction section

Comment 3:

The terms ambulatory care and outpatients can be confusing. Are you referring to specialized ambulatory care services? Or primary care? These settings are vastly different and so it is difficult to compare a primary care provider in general practice to a specialized neurosurgery outpatients department.

- **Removed/ Edited/ Replaced Content:** Not catered - the paper is written and discussed in the context of specialized neurosurgery outpatients department

Competing Interests: Non-Financial Competing Interests

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research