Check for updates

METHOD ARTICLE

REVISED Modeling the impact of early interventions on the transmission dynamics of coronavirus infection [version 2; peer review: 2 approved]

Christopher Saaha Bornaa ¹, Baba Seidu ², Yakubu Ibrahim Seini³

¹Department of Mathematics and ICT Education, School of Science, Mathematics and Technology Education, CK Tedam University of Technology and Applied Sciences, Navrongo, UER, +233, Ghana

²Department of Mathematics, Faculty of Mathematical sciences, CK Tedam University of Technology and Applied Sciences,, Navrongo, UER, +233, Ghana

³Department of Engineering, School of Engineering, University for Development Studies,, Tamale,, UER, +233, Ghana

V2 First published: 30 Jun 2021, 10:518
<https://doi.org/10.12688/f1000research.54268.1>

Latest published: 17 Aug 2021, 10:518
<https://doi.org/10.12688/f1000research.54268.2>

Abstract

A deterministic model is proposed to describe the transmission dynamics of coronavirus infection with early interventions. Epidemiological studies have employed modeling to unravel knowledge that transformed the lives of families, communities, nations and the entire globe. The study established the stability of both disease free and endemic equilibria. Stability occurs when the reproduction number, R_0 , is less than unity for both disease free and endemic equilibrium points. The global stability of the disease-free equilibrium point of the model is established whenever the basic reproduction number R_0 is less than or equal to unity. The reproduction number is also shown to be directly related to the transmission probability (β), rate at which latently infected individuals join the infected class (δ) and rate of recruitment (Λ). It is inversely related to natural death rate (μ), rate of early treatment (τ_1), rate of hospitalization of infected individuals (θ) and Covid-induced death rate (σ). The analytical results established are confirmed by numerical simulation of the model.

Keywords

Coronavirus, stability, simulation, reproduction number, interventions, transmission

Open Peer Review

Approval Status

	1	2
version 2 (revision) 17 Aug 2021	 view	
version 1 30 Jun 2021	 view	 view

1. **Mehmet Yavuz** , Necmettin Erbakan University, Konya, Turkey
University of Exeter, Cornwall, UK
2. **Ebenezer Bonyah** , Akenaten Appiah-Menka University of Skills Training and Entrepreneurial Development, Kumasi, Ghana

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the **Pathogens** gateway.

This article is included in the **Emerging Diseases and Outbreaks** gateway.

This article is included in the **Coronavirus** collection.

Corresponding author: Christopher Saaha Bornaa (cbornaa@cktutas.edu.gh)

Author roles: Bornaa CS: Conceptualization, Formal Analysis; Seidu B: Methodology, Software; Seini YI: Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Bornaa CS *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bornaa CS, Seidu B and Seini YI. **Modeling the impact of early interventions on the transmission dynamics of coronavirus infection [version 2; peer review: 2 approved]** F1000Research 2021, 10:518 <https://doi.org/10.12688/f1000research.54268.2>

First published: 30 Jun 2021, 10:518 <https://doi.org/10.12688/f1000research.54268.1>

REVISED Amendments from Version 1

The introduction of the manuscript has been updated to reflect the current scheme of COVID-19 infection.

The appropriate punctuation marks are also inserted to improve upon the flow of the language.

A column on the source of the data has been added to table 1 and how parameter estimation is done for clarity as requested by the reviewer.

A more practical interpretation of the results is included under the discussion as requested by the reviewer.

An appropriate reference is also added.

Any further responses from the reviewers can be found at the end of the article

Introduction

Accurate detection of the underlying agent that causes a disease is very important. Globally, many infections remain undetected and therefore untreated, causing complications and major human and economic consequences. Managing diseases in our modern world requires strategic interventions. Several of these interventions have been explored in the case of the COVID-19 pandemic ranging from the use of personal protective equipment (PPEs), lockdowns, social distancing, etc. Such interventions are implemented at particular points during the infection cycle. These are before the infection, early stages of the infection (asymptomatic) and later stages of the infection (symptomatic). Before the infection, individuals may be immunized as a control measure. Immunization is a cost-effective and highly successful health intervention used to manage emerging infectious diseases such as coronavirus. This is a process where a person, through vaccination, becomes protected against an emerging disease. Early treatment of the disease can also be a good strategy in managing infectious diseases. It is therefore important to identify infected persons early enough to put them on treatment. The emergence of COVID-19 is a worry to many in the world. This virus has come with three sub-categories known as alpha, beta and gamma and an introduction of the fourth subgroup called delta coronaviruses [Naik et al., 2020]. Early treatment of SARS-CoV-2 would speed up the recovery process, reduce the likelihood of developing severe outcomes and reduce demand on health care systems [Kim et al., 2020; Strain et al., 2005]. Strain et al. [2005] studied the effect of early treatment of primary infection on cellular reservoir clearance of HIV-1 and noted a positive effect. The question arises of whether this will be the case for coronavirus infections. Individuals should also be encouraged to go for voluntary testing as this will increase case detection, thereby reducing the number of secondary infections [Kim et al., 2020]. In every infection, the exposure of an individual to the source of infection is followed by an incubation period which varies from one infectious disease to the other. For example, the incubation period for coronavirus disease 2019 (Covid-19) is 14 days. During the incubation period the infected individual may be asymptomatic and unaware of their infection. This is followed by the onset of symptoms and varied health behaviours of the infected individual. Some researchers support the notion that antiviral therapy hastens viral clearance in asymptomatic infection [Strain et al., 2005; Hu et al., 2020] thereby halting the progression to symptomatic infection and the subsequent impact on the health system and the society at large.

Isolating persons who have been exposed to a coronavirus disease in order to prevent transmission has been a long established public health strategy [Kim et al., 2020]. This is done either in isolation homes or at hospitals in severe cases. Treatment interventions are offered in such circumstances.

This paper looks at the effect of early interventions on the transmission dynamics of coronavirus infection by employing mathematical modelling. Several mathematical models have been used as an engine to unravel knowledge in many epidemiological studies [Asamoah et al., 2017, 2020a, 2020b; Bornaa et al., 2017, 2020; Seidu et al., 2020; Agosto et al., 2015; Ivorra Benjamin and Ramos, 2015]. Most of these studies have yielded exceptional results that transformed the lives of families, communities, nations and the entire globe. Mathematical models have also been used to study pest and worm infestation in agriculture and aquaculture. Mathematical modeling is used in economics to observe, understand, and make predictions about human economic behavior.

Model formulation

Consider $N(t)$ as the size of the total population at time t . The population is then subdivided into five classes: Susceptible, $S(t)$, the latently infected (Exposed) $E(t)$, the clinically symptomatic (Infected), $I(t)$, the hospitalized in a facility (Hospitalized) $H(t)$ and Recovered $R(t)$ Classes. Thus, the population is given as

$$N(t) = S(t) + E(t) + I(t) + H(t) + R(t).$$

People are recruited into the susceptible population through birth and immigration at rate Λ and leave by having active contact with the viral source and being infected at rate β . These individuals progress to the latently infected class $E(t)$. The susceptible population also decreases by natural death at the rate μ . Thus;

$$\frac{dS}{dt} = \Lambda - (\beta I + \mu) S.$$

The population of the latently infected (exposed) class decreases whenever an exposed individual begins to show clinical symptoms and is moved into the infected class $I(t)$ at rate δ . This may be due to lack of intervention or intervention failure. It also decreases through natural and disease induced deaths, and recovery from early intervention (treatment) at rates μ , σ and τ_1 respectively. Thus;

$$\frac{dE}{dt} = \beta S I - (\tau_1 + \delta + \sigma + \mu) E.$$

The population of clinically infected individuals decreases due to hospitalization, natural death and disease induced death at rates θ , μ and σ respectively. Thus;

$$\frac{dI}{dt} = \delta E - (\theta + \sigma + \mu) I.$$

The population of the individuals that are hospitalized increases whenever the clinically infected individuals are taken to the hospital at rate θ , then decreases due to natural death, disease induced death and recovery at rate τ_2 . Thus;

$$\frac{dH}{dt} = \theta I - (\tau_2 + \sigma + \mu) H.$$

The population of the recovered individuals increases whenever there is recovery from the exposed class due to early treatment, and also from hospitalized individuals due to recovery after treatment at rates τ_1 and τ_2 respectively, and decreases by natural death. For the period under consideration, the recovered is assumed to have permanent immunity.

$$\frac{dR}{dt} = \tau_1 E + \tau_2 H - \mu R.$$

The description of the dynamics of the disease infection is depicted by following set of equations.

$$\left. \begin{aligned} \frac{dS}{dt} &= \Lambda - (\beta I + \mu) S \\ \frac{dE}{dt} &= \beta S I - (\tau_1 + \delta + \sigma + \mu) E \\ \frac{dI}{dt} &= \delta E - (\theta + \sigma + \mu) I \\ \frac{dH}{dt} &= \theta I - (\tau_2 + \sigma + \mu) H \\ \frac{dR}{dt} &= \tau_1 E + \tau_2 H - \mu R \\ (S(0), E(0), I(0), H(0), R(0)) &\in \mathbb{R}_{\geq 0}^5 \end{aligned} \right\} \quad (1)$$

For purposes of analysis, let $k_1 = \tau_1 + \delta + \sigma + \mu$, $k_2 = \theta + \sigma + \mu$, and $k_3 = \tau_2 + \sigma + \mu$.

We begin to discuss some basic properties of model (1) in the next section. This includes the positivity and boundedness, equilibrium states, local and global stability of the equilibria and sensitivity analysis. Numerical simulation is also employed to confirm the analytical results. Finally, the results are discussed and are conclusions drawn.

Qualitative properties of the model

Positivity and boundedness

Lemma 0.1. *If only all solutions of model (1) start in Ω , they remain in it for all $t \geq 0$.*

Also, for the model (1), the region $\Omega = \{(S, E, I, H, R) \in \mathbb{R}_{\geq 0}^5 \mid N \leq \frac{\Lambda}{\mu}\}$ is a positively invariant set.

Proof. Define $\zeta(x) = \{x(t) = 0 \text{ and } (S, E, I, H, R) \in \mathbb{R}_{\geq 0}^5\}$, $\forall x \in \{S, E, I, H, R\}$.

The following therefore is obtained from model (1);

$$\begin{cases} \left. \frac{dS}{dt} \right|_{\zeta(S)} = \Lambda > 0 \\ \left. \frac{dE}{dt} \right|_{\zeta(E)} = \beta I S > 0 \\ \left. \frac{dI}{dt} \right|_{\zeta(I)} = \delta E > 0 \\ \left. \frac{dH}{dt} \right|_{\zeta(H)} = \theta I > 0 \\ \left. \frac{dR}{dt} \right|_{\zeta(R)} = (\tau_1 E + \tau_2 H) > 0 \end{cases}$$

As stated in Lemma 2 of Yang et al. [1996], a solution of model (1) is always such that $(S(t), E(t), I(t), H(t), R(t)) \in \mathbb{R}_{\geq 0}^5$. The first part of the lemma is therefore proved.

We have

$$\frac{dN}{dt} = \Lambda - N\mu - (I + H + E)\sigma \leq \Lambda - \mu N$$

from $N = S + E + I + H + R$. Thus, $N(t) \leq N(0)e^{-\mu t} + \frac{\Lambda}{\mu}(1 - e^{-\mu t})$.

Therefore if $0 \leq N(0) \leq \frac{\Lambda}{\mu}$ then $\limsup_{t \rightarrow \infty} N(t) \leq \frac{\Lambda}{\mu}$.

Thus, all solutions of the model remain in Ω whenever they start in it. The second part of the lemma is also proved and hence the whole Lemma is proved.

We therefore conclude that model (1) is mathematically and epidemiologically well-posed within Ω [Hethcote, 2000].

Equilibrium points of the model

There are basically two equilibria for model (1); the disease-free, $\mathcal{E}_0$ and endemic, $\mathcal{E}^*$ equilibria. The solution of the system

$$\frac{dS}{dt} = \frac{dE}{dt} = \frac{dI}{dt} = \frac{dH}{dt} = \frac{dR}{dt} = 0$$

gives the equilibrium points of the model.

Assume that an equilibrium point of the model is typically $(S^*, E^*, I^*, H^*, R^*)$, then model (1) gives;

$$\left. \begin{aligned} S^* &= \frac{\Lambda}{\beta I + \mu}, \\ E^* &= \frac{\beta \Lambda I}{(\beta I + \mu)k_1}, \\ I^* &= \frac{\delta \beta \Lambda - \mu k_1 k_2}{\beta} \text{ or } 0 \\ H^* &= \frac{\theta I}{k_3}, \\ R^* &= \frac{(\tau_1 \beta \Lambda k_3 + \tau_2 \theta (\beta I + \mu) k_1) I}{(\beta I + \mu) k_1 \mu}. \end{aligned} \right\} \quad (2)$$

Theorem 0.1. *The model (1) has unique endemic and disease free equilibria.*

Proof. The solution of 2 for $I^* = 0$ is $\mathcal{E}_0 = (S_0, 0, 0, 0, 0)$, where $S_0 = \frac{\Lambda}{\mu}$. When $I^* = \frac{\delta \beta \Lambda - \mu k_1 k_2}{\beta}$ is substituted into (2), the endemic equilibrium is obtained, thus proved.

Now, the basic reproduction number is obtained by employing the next-generation matrix method of [Zhisheng and Van den Driessche \[2012\]](#). Thus;

$$\mathcal{R}_0 = \frac{\beta \Lambda \delta}{\mu k_1 k_2}$$

Local stability of equilibria

The stability of an equilibrium point is determined by considering all the roots of the characteristic equation of the Jacobian Matrix of model (1). If all the roots are negative then the equilibrium point is said to be locally asymptotically stable. The Jacobian Matrix $\mathcal{J}$ of the model is given as

$$\mathcal{J} = \begin{bmatrix} -\beta I - \mu & 0 & -\beta S & 0 & 0 \\ \beta I & -k_1 & \beta S & 0 & 0 \\ 0 & \delta & -k_2 & 0 & 0 \\ 0 & 0 & \theta & -k_3 & 0 \\ 0 & \tau_1 & 0 & \tau_2 & -\mu \end{bmatrix},$$

Local stability of disease-free equilibrium point ($\mathcal{E}_0$)

It can be shown that at $\mathcal{E}$, $-\mu$ and $-k_3$ are roots of the characteristic equation of the Jacobian Matrix $\mathcal{J}$. Clearly all these roots are negative. The roots that remain must therefore satisfy [equation 3](#) at $\mathcal{E}_0$. This is the characteristic polynomial of the Jacobian Matrix $\mathcal{J}_{\mathcal{E}}$,

where

$$\mathcal{J}_{\mathcal{E}} = \begin{bmatrix} -k_1 & \frac{\beta \Lambda}{\mu} \\ \delta & -k_2 \end{bmatrix}$$

The roots of [equation 3](#) are the eigenvalues of $\mathcal{J}_{\mathcal{E}}$.

$$\Phi_2 x^2 + \Phi_1 x + \Phi_0 = 0 \quad (3)$$

where

$$\left. \begin{aligned} \Phi_2 &= 1, \\ \Phi_1 &= k_2 + k_1 \\ \Phi_0 &= (1 - R_0)k_1 k_2 \end{aligned} \right\}$$

Whenever $\mathcal{R}_0 < 1$, all the coefficients Φ_i , $i = 0, 1, 2$, are positive. This suggests lemma 2 by using Descartes rule of signs.

Lemma 0.2. *The $\mathcal{E}_0$ is locally asymptotically stable whenever $\mathcal{R}_0 < 1$ and unstable whenever $\mathcal{R}_0 > 1$.*

Local stability of endemic equilibrium point ($\mathcal{E}^*$)

Again, the roots that remain must, after establishing that $-\mu$ and $-k_3$ are roots of the characteristic equation of the Jacobian Matrix $\mathcal{J}$, satisfy [equation 4](#). This is the characteristic polynomial of the Jacobian Matrix $\mathcal{J}_{\mathcal{E}^*}$ at $\mathcal{E}^*$, where

$$\mathcal{J}_{\mathcal{E}^*} = \begin{bmatrix} -i\beta - \mu & 0 & -\beta S \\ i\beta & -k_1 & \beta S \\ 0 & \delta & -k_2 \end{bmatrix}$$

The roots of [equation 4](#) are the eigenvalues of $\mathcal{J}_{\mathcal{E}^*}$.

$$\Pi_3 x^3 + \Pi_2 x^2 + \Pi_1 x + \Pi_0 = 0 \quad (4)$$

where

$$\left. \begin{aligned} \Pi_3 &= 1, \\ \Pi_2 &= k_2 + k_1 + \beta I + \mu \\ \Pi_1 &= (1 - R_0)k_2k_1 + k_2\beta I + k_2\mu + k_1\beta I + k_1\mu \\ \Pi_0 &= (1 - R_0)k_2k_1\mu + k_2k_1\beta I \end{aligned} \right\}$$

Whenever $\mathcal{R}_0 \leq 1$, all the coefficients Π_i , $i = 0, 1, 2, 3$ are positive. This also suggests lemma 3 by using Descartes rule of signs.

Lemma 0.3. *The $\mathcal{E}^*$ is locally asymptotically stable whenever $\mathcal{R}_0 \leq 1$ and unstable whenever $\mathcal{R}_0 > 1$.*

Global stability of disease-free equilibrium point ($\mathcal{E}_0$)

The technique of [Castillo-Chavez et al. \[2002\]](#) is employed to study the global stability of $\mathcal{E}_0$. If we let $Y = (S, R)$ and $X = (S_1, Q, I, H)$, model (1) can now be re-written as

$$\left. \begin{aligned} \frac{dY}{dt} &= \mathcal{F}(Y, X) \\ \frac{dX}{dt} &= \mathcal{G}(Y, X) \end{aligned} \right\} \quad (5)$$

where

$$\mathcal{F} = \begin{bmatrix} \Lambda - \beta SI - \mu S \\ \tau_1 E + \tau_2 H - \mu R \end{bmatrix}$$

and

$$\mathcal{G} = \begin{bmatrix} -k_1 & \beta S & 0 \\ \delta & -k_2 & 0 \\ 0 & \theta & -k_3 \end{bmatrix}.$$

According to [Castillo-Chavez et al. \[2002\]](#), the following conditions establish the global stability of $\mathcal{E}_0$.

$$\begin{aligned} H1 : & \text{For } \left. \frac{dY}{dt} \right|_{X=0}, Y^* \text{ is globally asymptotically stable} \\ H2 : & \mathcal{G}(Y, X) = \mathcal{L}Z - \hat{\mathcal{G}}(Y, X), \hat{\mathcal{G}}(Y, X) \geq 0 \text{ for } (Y, X) \in \mathcal{D}, \end{aligned} \quad (6)$$

where $\mathcal{L} = D_X \mathcal{G}(Y^*, 0)$ is the Jacobian of $\mathcal{G}(Y, X)$ with respect to X at $\mathcal{E}_0$.

If the reduced system of (1) given by

$$\left. \frac{dY}{dt} \right|_{X=0} = \begin{bmatrix} \Lambda - \mu S \\ -\mu R \end{bmatrix}$$

is considered, then disease-free equilibrium point $Y^* = \mathcal{E}_0$ is clearly a globally asymptotically stable point of the reduced system. Also, let $\mathcal{G}(Y, X) = \mathcal{L}X - \hat{\mathcal{G}}(Y, X)$, where

$$\mathcal{L} = D_X \mathcal{G}(Y^*, 0) = \begin{bmatrix} -k_1 & \beta S & 0 \\ \delta & -k_2 & 0 \\ 0 & \theta & -k_3 \end{bmatrix}$$

and

$$\hat{\mathcal{G}}(Y, X) = \lambda N \begin{bmatrix} \beta SI - k_1 E \\ \delta E - k_2 I \\ \theta I - k_3 H \end{bmatrix}$$

Substituting $\mathcal{E}_0$ into $\hat{G}(Y, X)$ gives zero components and condition H_2 is satisfied. Whenever $\mathcal{R}_0 \leq 1$ the $\mathcal{E}_0$ is globally asymptotically stable hence establishing the following results.

Lemma 0.4. *The disease-free equilibrium point $\mathcal{E}_0$ of the model is globally asymptotically stable whenever $\mathcal{R}_0 \leq 1$.*

Global stability of endemic equilibrium point ($\mathcal{E}^*$)

In this section, we try to find out whether or not the endemic equilibrium is globally stable using Lyapunov functions technique.

Theorem 0.2. *The endemic equilibrium of model (1) is globally asymptotically stable whenever $\mathcal{R}_0 > 1$.*

Proof. Consider the Lyapunov function

$$\begin{aligned} V = & \left(S - S^* - S^* \ln \frac{S}{S^*} \right) + \left(E - E^* - E^* \ln \frac{E}{E^*} \right) \\ & + \left(I - I^* - I^* \ln \frac{I}{I^*} \right) + \left(H - H^* - H^* \ln \frac{H}{H^*} \right) \\ & + \left(R - R^* - R^* \ln \frac{R}{R^*} \right). \end{aligned}$$

If we take the derivative of V we obtain

$$\begin{aligned} \frac{dV}{dt} = & \left(\frac{S - S^*}{S} \right) \frac{dS}{dt} + \left(\frac{E - E^*}{E} \right) \frac{dE}{dt} \\ & + \left(\frac{I - I^*}{I} \right) \frac{dI}{dt} + \left(\frac{H - H^*}{H} \right) \frac{dH}{dt} \\ & + \left(\frac{R - R^*}{R} \right) \frac{dR}{dt}, \\ = & \left(\frac{S - S^*}{S} \right) [\Lambda + -(\beta I + \mu) S] \\ & + \left(\frac{E - E^*}{E} \right) [\beta S I - (\tau_1 + \delta + \sigma + \mu) E] \\ & + \left(\frac{I - I^*}{I} \right) [\delta E - (\theta + \sigma + \mu) I] \\ & + \left(\frac{H - H^*}{H} \right) [\theta I - (\tau_2 + \sigma + \mu) H] \\ & + \left(\frac{R - R^*}{R} \right) [\tau_1 E + \tau_2 H - \mu R]. \end{aligned} \tag{7}$$

Table 1. Descriptions and baseline values of model parameters.

Par.	Description	Average value/ day	Source
μ	Natural death rate	0.01	Asamoah et al (2020)
Λ	Rate of recruitment	20	estimated
τ_1	Rate of early treatment	0.5	estimated
β	Transmission probability	0.002	estimated
σ	COVID-19 induced death rate	0.02	estimated
θ	Rate of hospitalization of infected persons	0.3	estimated
τ_2	Recovery rate of treated infected individuals	0.2	estimated
δ	Rate at which latently infected individuals join the infected class	0.0012	estimated

Table 2. Sensitivity Indexes.

Parameters	β	Λ	δ	μ
Sensitivity Index ($\gamma_{x_i}^{\mathcal{R}_0}$)	1.0000	1.0000	0.9977	-1.0491
Parameters	τ_1	θ	σ	
Sensitivity Index ($\gamma_{x_i}^{\mathcal{R}_0}$)	-0.9413	-0.9091	-0.0983

Figure 1. Simulation results illustrating the local stability of $\mathcal{E}_0$ for $\mathcal{R}_0 < 1$. **Figure 2. Plot for sensitivity of parameters of $\mathcal{R}_0$.**

(a) plot of τ_1 and θ against $\mathcal{R}_0$ (b) plot of β and δ against $\mathcal{R}_0$ **Figure 3.** Solution curves depicting the impact of τ_1 , θ , β and δ on $\mathcal{R}_0$.

(a) Susceptible class

(b) Exposed class

(c) Infected class

(d) Hospitalized class

(e) Recovered class

Figure 4. Solution curves depicting the impact of τ_1 on each of the classes.

After some algebraic manipulation $\frac{dV}{dt} = \mathbf{A} - \mathbf{B}$, where

$$\mathbf{A} = \frac{(S_0 E - S E_0) I \beta}{E} + \frac{\delta E (I - I_0)}{I} + \frac{\theta I (H - H_0)}{H} + \frac{\tau_1 E (R - R_0)}{R} + \frac{\tau_2 H (R - R_0)}{R},$$

and

$$\mathbf{B} = \mu (S - S_0) + k_1 (E - E_0) + k_2 (I - I_0) + k_3 (H - H_0) + \mu (R - R_0).$$

(a) Susceptible class

(b) Exposed class

(c) Infected class

(d) Hospitalized class

(e) Recovered class

Figure 5. Solution curves depicting the impact of δ on each of the classes varied from initial value of 0.0012 at 0.5 intervals.

Therefore whenever $\mathbf{A} < \mathbf{B} \Rightarrow \frac{dV}{dt} < 0$, and also $\mathbf{A} - \mathbf{B} = 0 \Rightarrow \frac{dV}{dt} = 0$, this is possible if $S = S^*, E = E^*, I = I^*, H = H^*, R = R^*$. Therefore, the largest compact invariant set for model (1) is $\{\mathcal{E}^*\}$. Hence the global asymptotic stability of $\mathcal{E}^*$ is established in the positive region $\mathbb{R}_{\geq 0}^5$ if $\mathbf{A} < \mathbf{B}$ for $\mathcal{R}_0 > 1$ as in Lyapunov-LaSalle's stability theorem [LaSalle, 1968],

Sensitivity analysis of the reproduction number ($\mathcal{R}_0$)

Parameters play a very important role in the dynamical behaviour of models; therefore the study of the impact of changes in the values of these parameters cannot be underestimated [Seidu et al., 2020; Bormaa et al., 2021; Chitnis et al.,

Figure 6. Solution curves depicting the impact of θ on each of the classes varied from initial value of 0.3 at 0.5 intervals.

2008]. The influential parameters of the model are considered by beginning to value the changes that occur in their values. Sensitivity analysis is therefore employed to help identify such influential parameters of the model. The normalized forward sensitivity technique is employed to study the sensitivity of $\mathcal{R}_0$ to model parameters.

Definition: Let model output $\mathcal{R}_0$ be differentiable dependent on model parameter x . The normalized forward sensitivity index of $\mathcal{R}_0$ with respect to x is defined by

Figure 7. Solution curves depicting the impact of β on each of the classes varied from initial value of 0.002 at 0.5 intervals.

$$\gamma_{x_i}^{\mathcal{R}_0} = \frac{\partial \mathcal{R}_0}{\partial x_i} \times \frac{x_i}{\mathcal{R}_0} \bigg|_{(x_1, x_2, \dots, x_n)} \quad (8)$$

Parameters estimation

We estimated the model parameters from fitting different combinations of parameters in model (1) to the actual number of Covid cases in Ghana provided by the Ghana Health services from January, 2021 to June, 2021. The estimation of parameters was then carried out using the least squares method which minimises summation of the square errors given by $\sum (M(t,p) - N_{\text{real}})^2$ subject to model (1), where N_{real} is the real reported Covid data, and $M(t,p)$ denotes the solution of the model corresponding to the number of Covid cases over time t with the set of estimated parameters, denoted by p .

The model parameter values, except the natural death rate, in Table 1 are then used as baseline values for the purposes of the sensitivity test and numerical simulation only. The local sensitivity indexes are therefore recorded in Table 2 taking $\mathcal{R}_0$ as a model output.

Numerical simulation

Considering the baseline parameter values in Table 1, numerical experiments are performed, with initial values of $S = 0.411e + 5$, $E = 0.493e + 2$, $I = 0.243e + 2$, $H = 0.318e + 2$, $R = 0.97e + 3$, to verify the analytical results established. The simulation is carried out using matlab version 1.0 for MathWorks R2017a release.

Simulation to illustrate the local stability of the disease free equilibrium of model (1) is run for $\mathcal{R}_0 < 1$.

Figure 2 depicts the bar graph of the sensitivity of $\mathcal{R}_0$ to marginal changes in each of the parameters that constitute $\mathcal{R}_0$.

To illustrate the impact of early treatment on the dynamics of coronavirus infection, we now present the early treatment parameter τ_1 , which is varied (from initial value of 0.5 at 0.5 intervals) and observe its impact on the spread of coronavirus infection, with all other parameters kept constant (see Figure 4a, b, c, d and e).

Each parameter under consideration is varied (from initial value at 0.5 intervals) to observe its impact on the spread of Covid-19 infections, with all other parameters kept constant (see Figure 5a, b, c, d and e for δ ; Figure 6a, b, c, d and e for θ ; Figure 7a, b, c, d and e for β).

Discussions, conclusions and recommendations

A deterministic model is proposed to describe the transmission dynamics of coronavirus infection with early interventions. The study describes the basic qualitative properties of the model and carried out numerical simulations to confirm the qualitative results. The model is shown to have unique disease-free and endemic equilibria (see equation 2, theorem 1 and the proof). The disease-free equilibrium is locally asymptotically stable when $\mathcal{R}_0 < 1$ (see Figure 1). The endemic equilibrium is also locally asymptotically stable when $\mathcal{R}_0 \leq 1$. The global stability of the disease-free and endemic equilibria are respectively established under the conditions; $\mathcal{R}_0 \leq 1$ and $\mathcal{R}_0 > 1$.

Analysis of the responsiveness of the model parameters to marginal changes is also carried out to identify the most important factors to consider in the fight against the spread of the disease. The results show, in order of importance, that a marginal change in any of the following parameters will affect directly or inversely the reproduction number thereby affecting the transmission dynamics of the disease. These parameters are: the death rate μ , transmission probability β , recruitment rate Λ , rate at which exposed individuals join the infected class δ , early treatment rate τ_1 , rate of hospitalization of infected individuals θ and coronavirus induced death rate σ (see Table 2).

We observe from Table 2 that Λ , β and δ are directly proportional to $\mathcal{R}_0$ in order of importance. In other words, an increase (decrease) in any of these parameters will lead to an increase (decrease) in $\mathcal{R}_0$ as confirmed by Figure 3b on β and δ . It therefore suggests, from the result, that all attempts should be made to reduce the transmission probability and the rate at which latently infected individuals join the infected class. The parameters that are however inversely proportional to $\mathcal{R}_0$ are μ , θ , τ_1 and σ . An increase (decrease) in any of these parameters will lead to a decrease (increase) in $\mathcal{R}_0$ as confirmed by Figure 3a on τ_1 and θ . This is also reflected in Figure 4 (a, b, c, d and e). Increasing early treatment reduces the infected population (see Figure 4b, c and d) and increases the non-infected population (see Figure 4a and e). This is also supported by Kim et al. [2020] and Strain et al. [2005] when they carried out independent studies into early treatment of infectious diseases.

Many health care facilities are overstretched in terms of space, equipment and personnel because of the pandemic. This challenge can be handled by either providing more space, equipment and personnel, or by increasing the recovery rate of

treated infected individuals. This study has confirmed that an increase in recovery rate reduces the hospitalized individuals and thereby reduces the pressure on health care facilities to take in more coronavirus and other patients (see [Figure 6d](#)).

It is assumed that the recovered gained permanent immunity because they develop antibodies that react and fight against initial infections. We also observe increases in the rate at which the exposed joins the infected (symptomatic) class (δ). When this happens the symptomatic are moved into hospitals for treatment and subsequently recovery. The effect is that the infected and recovered populations are increased but the susceptible population is reduced because of the immunity gained by the recovered (see [Figure 5a, b, c, d and e](#)).

It is also shown that as transmission probability increases the infected increase and the susceptible decrease (see [Figure 7a, b, c and d](#)).

It is recommended that:

- early treatment of coronavirus should be considered. This can be effectively carried out when proper surveillance is enforced to identify asymptomatic individuals who are mostly at their early stage of infection. People should also be encouraged to use immune boosters to reduce the rate at which the exposed joins the infected while early treatment is enforced.
- efforts should be made to reduce the rate of infection by insisting on people observing all the coronavirus protocols announced by World Health Organisation and other health institutions.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Author contributions

- Christopher Saaha Borna: Conceptualization and algebraic analysis
- Baba Seidu: Numerical analysis
- Yakubu Ibrahim Seini: Proof reading, supervision and advisory role

References

- Agusto FB, Teboh-Ewungkem MI, Gumel AB: **Mathematical assessment of the effect of traditional beliefs and customs on the transmission dynamics of the 2014 ebola outbreaks.** *BMC Med.* 2015; **13**, 12. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Asamoah JKK, Oduro FT, Bonyah E, *et al.*: **Modelling of rabies transmission dynamics using optimal control analysis.** *J Applied Mathematics.* 2017; **2017**. [Publisher Full Text](#)
- Asamoah JKK, Borna CS, Seidu B, *et al.*: **Mathematical analysis of the effects of controls on transmission dynamics of sars -cov-2.** *Alexandria Engineering J.* 2020a. [Reference Source](#)
- Asamoah JKK, Owusu M, Oduro JF, *et al.*: **Global stability and cost-effectiveness analysis of covid-19 considering the impact of the environment: using data from ghana.** *Chaos, Solitons and Fractals.* 2020b; page 110103. [Publisher Full Text](#)
- Baloba EB, Seidu B, Borna CS: **Mathematical analysis of the effects of controls on the transmission dynamics of anthrax in both animal and human populations.** *Computational Mathematical Methods Med.* 2020; **02** 2020. [Publisher Full Text](#)
- Borna CS, Seini YI, Seidu B: **Modelling zoonotic diseases with treatment in both human and animal populations.** *Commun. Math. Biol. Neurosci.* 2017; **2017**. [Publisher Full Text](#)
- Borna CS, Seidu B, Daabo MI: **Mathematical analysis of rabies infection.** *J Applied Mathematics.* 2020; **17**: 2020. [Publisher Full Text](#)
- Borna CS, Seidu B, Makinde OD: **Mathematical analysis of the impact of vaccination and poor sanitation on the dynamics of poliomyelitis.** *Int J Nonlinear Sciences Numerical Simulation.* 2021; **70**. [Publisher Full Text](#)
- Castillo-Chavez C, Blower S, Driessche P, *et al.*: **Mathematical approaches for emerging and reemerging infectious diseases: models, methods, and theory.** Springer; 2002.
- Chitnis N, Hymanand J, Cushing J: **Determining important parameters in the spread of malaria through the sensitivity analysis of a mathematical model.** *Bull. Math. Biol.* 2008; **70**: 1272–1296. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hethcote HW: **The mathematics of infectious diseases.** *SIAM review.* 2000; **42**(4): 599–653.
- Hu Z, Song C, Xu C, *et al.*: **Clinical characteristics of 24 asymptomatic infections with covid-19 screened among close contacts in nanjing, china.** *Sci China life Sci.* 2020; **63**: 706–711. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [Reference Source](#)
- Ivorra Benjamin ND, Ramos AM: **A mathematical model to predict the risk of human diseases spread between countries—validation and application to the 2014–2015 ebola virus disease epidemic.** *Bull Math*

Biol. 2015; **77**, 9.

[PubMed Abstract](#) | [Publisher Full Text](#)

Kim P, Cushing JM, Xu C, *et al.*: **Therapy for early covid-19-a critical need.** *JAMA*. 2020.

[PubMed Abstract](#) | [Publisher Full Text](#)

LaSalle J: **Stability theory for ordinary differential equations.** *J Differential Equations*. 1968; **4**.

[Publisher Full Text](#)

Naik PA, Yavuz M, Qureshi S, *et al.*: **Stuart Townley Modeling and analysis of COVID-19 epidemics with treatment in fractional derivatives using real data from Pakistan.** *Eur Phys J Plus*. 2020; **135**: 795. [Publisher Full Text](#)

Seidu B, Bornaa CS, Makinde OD: **An ebola model with hyper-susceptibility.** *Chaos, Solitons and Fractals*. 2020; **138**: 109938.

[Publisher Full Text](#)

Strain CM, Little SJ, *et al.*: **Effect of treatment, during primary infection, on establishment and clearance of cellular reservoirs of hiv-1.** *J Infect Dis*. 2005; **191**.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yang X, Chen L, Chen J: **Permanence and positive periodic solution for the single-species nonautonomous delay diffusive models.** *Computers Mathematics Applications*. 1996; **32**: 116–312.

Zhisheng S, Van den Driessche P: **Global stability of infectious disease models using lyapunov functions.** *SIAM J Applied Mathematics*. 2012; **73**. [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 01 September 2021

<https://doi.org/10.5256/f1000research.76442.r92124>

© 2021 Yavuz M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mehmet Yavuz

¹ Department of Mathematics and Computer Sciences, Faculty of Science, Necmettin Erbakan University, Konya, Turkey

² Department of Mathematics, College of Engineering, Mathematics and Physical Sciences, University of Exeter, Cornwall, TR10, UK

The revised version of the paper has been properly prepared. It can be indexed as it is.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mathematical modelling in epidemiology, fractional calculus, applied mathematics, optimal control, bifurcation and chaos.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 25 August 2021

<https://doi.org/10.5256/f1000research.57739.r89506>

© 2021 Bonyah E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ebenezer Bonyah

Department of Mathematics Education, Akenten Appiah-Menka University of Skills Training and

Entrepreneurial Development, Kumasi, Ghana

- The authors appreciate that there are many mathematical models available on COVID-19.
- The authors must provide the knowledge gap that this work is filling.
- The authors must present anything new on the model formulated.
- The authors must also provide a reason for using this numerical scheme for this study.
- For the conclusions, I suggest for this to be in the past tense.

Is the rationale for developing the new method (or application) clearly explained?

Yes

Is the description of the method technically sound?

Yes

Are sufficient details provided to allow replication of the method development and its use by others?

Partly

If any results are presented, are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions about the method and its performance adequately supported by the findings presented in the article?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mathematical Biology, Mathematical Modelling and Mathematics Education

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 26 July 2021

<https://doi.org/10.5256/f1000research.57739.r88778>

© 2021 Yavuz M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mehmet Yavuz

¹ Department of Mathematics and Computer Sciences, Faculty of Science, Necmettin Erbakan University, Konya, Turkey

² Department of Mathematics, College of Engineering, Mathematics and Physical Sciences, University of Exeter, Cornwall, TR10, UK

1. Authors should put proper punctuation at the end of all equations.
2. In numerical simulations, the authors need to explain where they have taken the parameter values from. Are they from the literature or estimated ones?
3. What makes the proposed methods and COVID-19 model suitable for this unique task? What new development to the proposed model have the authors added (compared to the existing approaches)? These points should be clarified.
4. What is the novelty of your work? There are some similar papers that have been investigated in epidemiology, especially COVID-19, although there are some minor differences in the structure of models. Please state it clearly.
5. Have you employed any assumptions in your COVID-19 model and problem formulation? Please explain briefly.
6. More physical interpretations should be given. Please provide corresponding explanations of the figures in terms of their physical meanings and pointing out the novelty of the paper. How do figures support your scheme?
7. The authors should improve the introduction by including the recent development within the frame of the mentioned COVID-19 models. Also, their mathematical investigations and numerical simulations with the help of recently published papers should be considered by comparing their current model. Sensitivity analysis of the Reproduction number can be discussed deeply. In this connection, I suggest some interesting results:
 - Modeling and analysis of COVID-19 epidemics with treatment in fractional derivatives using real data from Pakistan¹.
 - A New Mathematical Modeling of the COVID-19 Pandemic Including the Vaccination Campaign. *Open Journal of Modelling and Simulation* ².
 - New Approaches to the Fractional Dynamics of Schistosomiasis Disease Model. *Physica A: Statistical Mechanics and its Applications* ³.

References

1. Naik PA, Yavuz M, Qureshi S, Zu J, et al.: Modeling and analysis of COVID-19 epidemics with treatment in fractional derivatives using real data from Pakistan. *Eur Phys J Plus*. 2020; **135** (10): 795 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Yavuz M, Coşar F, Günay F, Özdemir F: A New Mathematical Modeling of the COVID-19 Pandemic Including the Vaccination Campaign. *Open Journal of Modelling and Simulation*. 2021; **09** (03): 299-321 [Publisher Full Text](#)
3. Yavuz M, Bonyah E: New approaches to the fractional dynamics of schistosomiasis disease

model. *Physica A: Statistical Mechanics and its Applications*. 2019; **525**: 373-393 [Publisher Full Text](#)

Is the rationale for developing the new method (or application) clearly explained?

Partly

Is the description of the method technically sound?

Yes

Are sufficient details provided to allow replication of the method development and its use by others?

Yes

If any results are presented, are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions about the method and its performance adequately supported by the findings presented in the article?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mathematical modelling in epidemiology, fractional calculus, applied mathematics, optimal control, bifurcation and chaos.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research