RESEARCH ARTICLE

Parental coronavirus disease vaccine hesitancy for children in Bangladesh: a cross-sectional study [version 1; peer review: 2 approved, 1 approved with reservations]

Mohammad Ali ^{1,2}, Sohel Ahmed³, Atia Sharmin Bonna⁴, Abu-sufian Sarkar ⁵, Md. Ariful Islam⁶, Tania Akter Urmi⁷, Tasnuva Samarukh Proma⁸

¹Department of Physiotherapy and Rehabilitation, Uttara Adhunik Medical College, Dhaka, 1230, Bangladesh

²Hasna Hena Pain Physiotherapy and Public Health Research Center, Uttara Model Town, Dhaka, 1230, Bangladesh

³Department of Physiotherapy, Mount Adora Hospital, Sylhet, 3100, Bangladesh

⁴Save the Children, Dhaka, 1212, Bangladesh

⁵Bashundhara Kings Football Club, Dhaka, 1229, Bangladesh

⁶Zaman Mordan Hospital, Sherpur, 2100, Bangladesh

⁷Jatio protibondhi seba o sahajjo kendro, Gopalganj, 8100, Bangladesh

⁸Advanced Physiotherapy & Rehab Solution, Women's Children's & General Hospital, Dhaka, 1209, Bangladesh

V1 First published: 25 Jan 2022, 11:90
<https://doi.org/10.12688/f1000research.76181.1>

Latest published: 02 Mar 2022, 11:90
<https://doi.org/10.12688/f1000research.76181.2>

Abstract

Background: Coronavirus disease 2019 (COVID-19) requires mass immunization to control the symptoms and global spread of severe acute respiratory syndrome coronavirus 2. Data from developed countries reported a high prevalence of parental COVID-19 vaccine hesitancy. However, parental vaccine hesitancy data in countries with low and middle income are scarce. The goal of this study was to assess the prevalence of parental vaccine hesitancy and identify subgroups with higher odds of vaccine hesitancy.

Methods: A cross-sectional study was conducted on parents of children aged <18 years from October 10, 2021, to October 31, 2021. Parents aged ≥18 years underwent face-to-face interviews in randomly selected places in Bangladesh using a vaccine hesitancy questionnaire. Predictors were identified using binary logistic regression analysis.

Results: Data from 2633 eligible parents were analyzed. Overall, 42.8% reported COVID-19 vaccine hesitancy for their youngest child. The final model suggested that children's age, parent's age, religion, occupation, monthly household income, permanent address, living location, the status of tobacco use, adherence with the regular government vaccination programs (other than COVID-19), perception about COVID-19 vaccine efficacy among Bangladeshi children, self-vaccination intention, report about family members' illness or death from COVID-19, and perceived COVID-19 threat were the independent predictors of parental vaccine hesitancy.

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)			
02 Mar 2022			
version 1			
25 Jan 2022			

1. **Shakil Ahmed** , ICDDR, Dhaka, Bangladesh

2. **F. M. Moinuddin**, Mayo Clinic, Rochester, USA

3. **Manzur Kader** , Karolinska Institutet, Stockholm, Sweden

Any reports and responses or comments on the article can be found at the end of the article.

Conclusions: Vaccine hesitation varied based on sociodemographic characteristics, religion, behavior, and perceived COVID-19 threat. Therefore, interventions focused on addressing vaccine hesitancy among subgroups are warranted.

Keywords

Bangladesh, COVID-19, developing countries, parents, pediatrics, vaccine hesitancy.

This article is included in the **Sociology of Health** gateway.

This article is included in the **Emerging Diseases and Outbreaks** gateway.

This article is included in the **Sociology of Vaccines** collection.

Corresponding author: Mohammad Ali (alibup2018@gmail.com)

Author roles: **Ali M:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Ahmed S:** Investigation, Validation, Writing – Review & Editing; **Bonna AS:** Investigation, Validation, Writing – Review & Editing; **Sarkar As:** Investigation, Validation, Writing – Review & Editing; **Islam MA:** Investigation, Validation, Writing – Review & Editing; **Urmi TA:** Investigation, Validation, Writing – Review & Editing; **Proma TS:** Investigation, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2022 Ali M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ali M, Ahmed S, Bonna AS *et al.* **Parental coronavirus disease vaccine hesitancy for children in Bangladesh: a cross-sectional study [version 1; peer review: 2 approved, 1 approved with reservations]** F1000Research 2022, 11:90 <https://doi.org/10.12688/f1000research.76181.1>

First published: 25 Jan 2022, 11:90 <https://doi.org/10.12688/f1000research.76181.1>

Introduction

Mass immunization against coronavirus disease 2019 (COVID-19) is one of the gold standard measures to control the spread of symptomatic severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and end the global pandemic.¹ Many countries have targeted vaccinating at least 70% of their total population to achieve herd immunity, including individuals aged ≤ 18 years.^{2,3} However, vaccine hesitancy, which is defined as a delay in acceptance or refusal of vaccines despite the availability of vaccination services,⁴ is a significant threat to the smooth uptake of vaccinations worldwide.⁵

Since December 2019, >250 million COVID-19 cases have been identified globally, and >50 million people have died of this disease, with a significantly high prevalence in older adults.⁶ However, the infection rate among children and adolescents is not negligible, and they can carry and spread the virus.⁷ Furthermore, unvaccinated populations are supposedly suitable hosts for new variants.⁸

Some recent data indicate that a few countries, for example, the USA are unlikely to reach the 70% target for herd immunity; however, vaccinating 22% of the American population, that is, the pediatric population, would boost community protection against COVID-19.³ Nonetheless, more than one in three parents in the USA were vaccine-hesitant for their children.³ In China, this rate was 52.5%.⁹ Along with sociodemographic variabilities and religious beliefs, overall vaccine hesitancy has also varied by political theology, people's perceived pandemic threat, or the socioeconomic status of the target population.^{10–12} Additionally, revealing adverse effects, the vaccine's effectiveness in children, and research availability among the child's age groups may play a crucial role when parents decide to vaccinate their children. Furthermore, a survey revealed that along with potential immediate adverse effects of the vaccines, the possible long-term harmful effects were a growing concern for parents.¹³

The triumph of immunization programs, among others, relies on the vaccination of a wide proportion of pediatric and adult populations in low- and middle-income countries where variants of concern, such as the SARS-CoV-2 B.1.617.2 Delta variant, have been detected.^{14,15} In Bangladesh, by November 2021, only 18% of the whole population was fully vaccinated against this disease.¹⁶ However, approximately 35% of Bangladeshi are aged <18 years.¹⁷ To achieve herd immunity, this young cohort should be included in the mass vaccination program. Thus, the Bangladesh government planned to vaccinate students aged between 12 and 17 years. Vaccination among the young student cohort started in cities, including Dhaka, to a limited extent from November 1, 2021.¹⁸

There is limited information about vaccine hesitancy among parents of children aged <18 years. In Bangladesh, a previous study revealed that 32% of the study population aged >18 years refused to be vaccinated against COVID-19.¹⁰ We hypothesized that parental vaccine hesitancy would not match the rate in the adult population. This study sought to (1) conduct a nationally representative assessment of parental vaccine hesitancy and (2) identify subgroups of parents with higher odds of vaccine hesitancy.

Methods

Ethics statements

The Institutional Review Board of Uttara Adhunik Medical College and Hospital approved this study (Approval number: UAMC-IRB-2021/09). Written informed consent both for participation and publication of data was obtained from all participants.

Study design and participants

This cross-sectional study was conducted in Bangladesh from October 10, 2021, to October 31, 2021. A margin of 2% error, a confidence level of 95%, and a response distribution of 50% were used to calculate the sample size to target fathers/mothers of 80 million children and obtain a minimum sample size of 2401 participants.^{19,20} Approximately 3000 parents aged ≥ 18 years with children aged <18 years who permanently live in Bangladesh were conveniently invited to participate in an individualized interview session using a previously employed vaccine hesitancy questionnaire.^{10,21,35} We received data from 2703 parents, as a result of a 10% refusal rate. However, 36 parents did not answer all the questions, thus were excluded. We also excluded 34 data points for the contradicting answers to the question. These exclusions meant that, only 2633 respondents were included in the final analysis.³⁵

Study questionnaire

In the first part of the questionnaire, participants were queried regarding parental vaccine hesitancy and perceived COVID-19 threat. First, parents were asked about the likelihood of vaccinating their youngest children. Parental vaccine hesitancy, the dependent variable and a vital outcome of the study, was measured using the question, "If a vaccine that would be effective against coronavirus disease among children was available, how likely are you to get your children vaccinated?" (response options: very likely, somewhat likely, not likely, and definitely not). Second, participants were

asked two questions regarding the perceived COVID-19 threat: (1) “How likely is it that your children or a family member could get infected with coronavirus in the next year?” (response options: very likely, somewhat likely, not likely, and definitely not). (2) “How concerned are you that your children or a family member could get infected with coronavirus in the next year?” (response options: very concerned, concerned, slightly concerned, and not concerned at all).

The second part of the questionnaire included a wide array of sociodemographic questions for both children and parents. A set of structured questions assessed the child's health (healthy/disabled), age, and sex. Information on parents' sex, age, religion, current marital status, education, employment status, monthly household income (Bangladeshi taka), permanent address, and region of residence (north, south, and central zones in Bangladesh, including Dhaka), current residence type (own/rented/others), family type (nuclear or extended, number of children, current tobacco use status, religious practice habits, and political affiliation was collected. Additionally, parents were asked some additional COVID-19 vaccine-related questions: “Do you think the COVID-19 vaccine will be effective among Bangladeshi children?” (response options: no, yes, or skeptical), “Have you received or plan to receive the COVID-19 vaccine,” “Did you or your family member(s) test positive for COVID-19,” and “Have you lost any of your family member(s) to COVID-19?” The last three questions received dichotomous (yes or no) answers.

Sampling technique and data collection

Data was collected from all eight divisions of Bangladesh. A dual-stage cluster sampling technique was used to include potential samples for this study. We randomly chose marketplaces, shopping malls, waiting rooms of large hospitals, diagnostic centers, bus and rail stations, and residences and processed them as a cluster in the first stage. To obtain data from the parents of a child with a disability, we also visited randomly selected centers for disabled children. The list of given data collection sites was collected from the divisions' websites. In the second stage, we chose participants conveniently. Data from only the father or mother of a child were taken from a family with children to avoid repeating data.

Eight teams of two persons were created. A team member read the questions aloud to the interviewees individually. Acceptable options were asked and recorded in the question paper. Subsequently, the answers were checked and confirmed by the second team member. The coinvestigator reviewed the data collection sheets for completeness, accuracy, and internal consistency and secured them with the principal investigator. Individual face-to-face interviews were conducted to ensure the participants' privacy. All participants were informed of the voluntary nature of their participation, and the interviews were conducted in Bangla.

Statistical analysis

The crucial outcome of this study was vaccine hesitancy. We dichotomized the four responses to the vaccine hesitancy question as either a positive (very likely and somewhat likely) or a negative (not likely and definitely not) attitude towards the COVID-19 vaccine. The Fisher exact test was used for two nominal variables, and chi-square test was used for more than two nominal variables to assess vaccine hesitancy rates and draw comparisons between the groups. Binary logistic regression analyses were performed to identify the predictors of parental COVID-19 vaccine hesitancy and compute adjusted odds ratios (AORs) with a 95% confidence interval (CI). Factors significantly associated with vaccine hesitancy in the descriptive analysis were included in the regression model. A goodness-of-fit test for the adjusted logistic regression model was performed using the Hosmer-Lemeshow test. The significance level was set at $p < 0.05$, and SPSS (version 22.0; IBM Corp; RRID: SCR_002865) was used to perform all data analyses.

Results

Parents and children's characteristics

Overall, 2633 parents aged 34.97 ± 7.87 years (mean $\pm$ standard deviation) were included in the analysis, with 52.8% (1390) being women. In total, 396 (15%) parents of children with a physical disability were included. Among the children, 1372 (52.1%) were boys, and 1206 (45.8%) were in the 0–4-year-old group. Most parents (653, 24.8%) were in the 31–35-year-old group. Overall, 2358 (89.4%) parents were Muslim, 1791 (68%) were a nuclear family member, 1075 (40.8%) had two children, 1022 (38.8%) had a low education level, 756 (28.7%) were homemakers, and 833 (31.6%) had a low-middle household income. Among all participants, 1528 (58%) were from the village, 1323 (50%) were living in the central zone, including Dhaka, 1695 (64.4%) were tobacco non-users, 1797 (68.2%) were regular religious practitioners, and 1032 (39.2%) were politically neutral respondents. A total of 177 (6.3%) parents did not adhere to the regular government vaccination programs other than COVID-19, and 1458 (55.4%) remained skeptical about COVID-19 vaccine effectiveness for Bangladeshi children. Furthermore, 722 (27.4%) parents were either not vaccinated or did not receive the COVID-19 vaccine; however, 752 (28.6%) parents reported that they or their family members tested positive for COVID-19, and 151 (5.7%) lost a loved one to COVID-19. Details of the responses to the questions regarding the likelihood of children or family members' infection by COVID-19 and the level of concern about children or family members' disease in the next year are shown in [Table 1](#).

Table 1. Descriptive analysis: Sociodemographic characteristics, COVID-19 threat, and parental vaccine hesitancy.

Variables	Total sample n (%)	Likelihood of vaccinating children		P value
		Not likely/definitely not n (%)	Very likely/somewhat likely n (%)	
All participants	2633 (100)	1126 (42.8)	1507 (57.2)	N/A
<i>Children's health</i>				0.507
Healthy	2237 (85)	957 (42.8)	1280 (57.2)	
Disabled	396 (15)	169 (42.7)	227 (57.3)	
<i>Children's age group</i>				0.000
0–4	1206 (45.8)	649 (53.8)	557 (46.2)	
5–9	870 (33)	344 (39.5)	526 (60.5)	
10–14	354 (13.4)	98 (27.7)	256 (72.3)	
15–<18	203 (7.7)	35 (17.2)	168 (82.8)	
<i>Children's sex</i>				0.000
Male	1372 (52.1)	537 (39.1)	835 (60.9)	
Female	1261 (47.9)	589 (46.7)	672 (53.3)	
<i>Parents' age group</i>				0.000
18–25	268 (10.2)	146 (54.5)	122 (45.5)	
26–30	604 (22.9)	329 (54.5)	275 (45.5)	
31–35	653 (24.8)	288 (43.8)	367 (56.2)	
36–40	563 (21.4)	223 (39.6)	340 (60.4)	
41–45	285 (10.8)	86 (30.2)	199 (69.8)	
46–50	162 (6.2)	37 (22.8)	125 (77.2)	
≥ 51	98 (3.7)	19 (19.4)	79 (80.6)	
<i>Parents' sex</i>				0.237
Female	1390 (52.8)	604 (43.5)	786 (52.2)	
Male	1243 (47.2)	522 (42)	721 (58)	
<i>Marital status</i>				0.438
Married	2527 (96)	1082 (42.8)	1445 (57.2)	
Divorced or widowed	106 (4)	44 (41.5)	62 (58.5)	
<i>Religion</i>				0.000
Muslim	2358 (89.4)	1069 (45.4)	1285 (54.6)	
Hindu	258 (9.8)	56 (21.7)	202 (78.3)	
Buddhist	6 (.2)	0 (0)	6 (100)	
Christian	15 (.6)	1 (6.7)	14 (93.3)	
<i>Type of family</i>				0.167
Extended family	842 (32)	372 (44.2)	470 (55.8)	
Nuclear family	1791 (68)	754 (42.1)	1037 (57.9)	
<i>Number of children</i>				0.993
One	924 (35.1)	395 (42.7)	529 (57.3)	
Two	1075 (40.8)	461 (42.9)	614 (57.1)	
Three or more	634 (24.1)	270 (42.6)	364 (57.4)	

Table 1. *Continued*

Variables	Total sample n (%)	Likelihood of vaccinating children		P value
		Not likely/definitely not n (%)	Very likely/somewhat likely n (%)	
Educational qualification				0.000
≤ High school	1022 (38.8)	473 (46.3)	549 (53.7)	
College education	594 (22.6)	316 (53.2)	278 (46.8)	
Graduate	608 (23.1)	236 (38.8)	372 (61.2)	
Post-graduate	409 (15.5)	101 (24.7)	308 (75.3)	
Occupation				0.000
Service	677 (25.7)	248 (36.6)	429 (63.4)	
Business	472 (17.9)	198 (41.9)	274 (58.1)	
Unemployed	179 (6.8)	132 (73.7)	47 (26.3)	
Student	56 (2.1)	34 (30.7)	22 (39.3)	
Home maker	756 (28.7)	364 (48.1)	392 (51.9)	
Healthcare	216 (8.2)	64 (29.6)	152 (70.4)	
Daily labor	277 (10.5)	86 (31)	191 (69)	
Monthly household income (₹)				0.000
<₹15 000	799 (30.3)	401 (50.2)	398 (49.8)	
₹ 15000–30000	833 (31.6)	409 (49.1)	424 (50.9)	
₹ 31000–45000	433 (16.4)	150 (34.6)	283 (65.4)	
>₹ 45000	568 (21.6)	166 (29.2)	402 (70.8)	
Current residence type				0.030
Own	1436 (54.5)	646 (45)	790 (55)	
Rented	1075 (40.8)	427 (39.7)	648 (60.3)	
Others	122 (4.6)	53 (43.4)	69 (56.6)	
Permanent address				0.000
Village	1528 (58)	687 (45)	814 (55)	
Semi-urban	535 (20.3)	247 (45.2)	288 (53.8)	
City	570 (21.6)	192 (33.7)	378 (66.3)	
Current living location				0.000
Central zone	1323 (50.2)	542 (41.0)	781 (59.0)	
North zone	921 (35)	472 (51.2)	449 (48.8)	
South zone	389 (14.8)	112 (28.8)	277 (71.2)	
Present tobacco user				0.000
No	1695 (64.4)	679 (40.1)	1016 (59.9)	
Yes	938 (35.6)	447 (47.7)	491 (52.3)	
Regular religious practice				0.099
No	836 (31.8)	377 (45.1)	459 (54.9)	
Yes	1797 (68.2)	749 (41.7)	1048 (58.3)	
Political affiliation				0.000
Ruling party	779 (30.3)	283 (35.8)	516 (64.6)	
Opposition	296 (11.2)	175 (59.1)	121 (40.9)	
Neutral	1032 (39.2)	488 (47.3)	544 (52.7)	
Prefer not to say	506 (19.2)	180 (35.6)	326 (64.4)	

Table 1. Continued

Variables	Total sample n (%)	Likelihood of vaccinating children		P value
		Not likely/definitely not n (%)	Very likely/somewhat likely n (%)	
Vaccinated/plan to vaccinate children under regular (other than COVID-19) govt. vaccination programs				0.000
No	177 (6.7)	103 (58.2)	74 (41.8)	
Yes	2456 (93.3)	1023 (41.7)	1433 (58.3)	
Do you think the COVID-19 vaccine will be effective in Bangladeshi children?				0.000
No	167 (6.3)	151 (90.4)	16 (9.6)	
Yes	1008 (38.3)	54 (5.4)	954 (94.6)	
Skeptical	1458 (55.4)	921 (63.2)	537 (36.8)	
Have you taken or plan to take the COVID-19 vaccine?				0.000
No	722 (27.4)	653 (88)	87 (12)	
Yes	1911 (72.6)	491 (25.7)	1420 (74.3)	
Have you or your family member(s) tested positive for COVID-19?				0.000
No	1881 (71.4)	947 (50.3)	934 (49.7)	
Yes	752 (28.6)	179 (23.8)	573 (76.2)	
Have you lost any of your family member(s) to COVID-19?				0.000
No	2482 (94.3)	1105 (44.5)	1377 (55.5)	
Yes	151 (5.7)	21 (13.9)	130 (86.1)	
Perceived likelihood of children or family members' infection in the next year				0.000
Very likely	345 (13.1)	51 (14.8)	294 (85.2)	
Somewhat likely	1678 (63.7)	665 (39.6)	1013 (60.4)	
Not likely	451 (17.1)	296 (65.6)	155 (34.4)	
Definitely not	159 (6)	114 (71.7)	45 (28.3)	
Level of concern about children or family members' infection in the next year				0.000
Very concerned	386 (14.7)	72 (18.7)	314 (81.3)	
Concerned	1020 (38.7)	384 (37.6)	636 (62.4)	
Slightly concerned	673 (25.6)	303 (45)	370 (55)	
Not concerned at all	554 (21)	367 (66.2)	187 (33.8)	

Results of descriptive analysis

Overall, 42.8% of parents reported hesitancy towards the COVID-19 vaccine for their youngest child. Closer analysis revealed that 26.05% of parents were very likely, 31.18% were somewhat likely, and 36.31% were not likely to vaccinate their child. While only 6.46% were definitely not vaccinating their child against COVID-19 (Figure 1). The incidence of vaccine hesitancy was significantly high among the parents of 0–4-year-old children (53.8%; $p=0.000$), parents of girls (46.7%; $p=0.000$), young parents (54.5%; $p=0.000$), Muslims (45.4%; $p=0.000$), parents who received college education (53.2%; $p=0.000$), unemployed parents (73.7%; $p=0.000$), parents with a household income of $< \text{BD}15,000$ (50.2%; $p=0.000$), those who lived in their own house (45%; $p=0.030$), came from a village (45%; $p=0.000$), lived in the north zone (51.2%; $p=0.000$), tobacco users (47.7%; $p=0.000$), and parents politically affiliated with opposition parties (59.1%; $p=0.000$). Similarly, participants who did not vaccinate or will not vaccinate their child with regular vaccines (other than COVID-19) available under government programs (58.2%; $p=0.000$), those who did not believe in the effectiveness of the COVID-19 vaccine for Bangladeshi children (90%; $p=0.000$), and those who did not or will not receive the COVID-19 vaccine for themselves (88%; $p=0.000$) showed high vaccine hesitancy. Parents who were not likely to believe that their children or a family member could be infected with COVID-19 in the next year (71.7%; $p=0.000$) and those not concerned about their children or a family member getting COVID-19 in the next year (66.2%; $p=0.000$) showed high levels vaccine hesitancy (Table 1).

Figure 1. Likelihood of COVID-19 vaccine acceptance/refusal by Bangladeshi parents for children aged <18.

Results of regression analysis

Subgroups with statistically significant higher odds of vaccine hesitancy were found to be parents of children aged 0–4 years (AOR=5.87, 95% CI=2.91–11.85; $p=0.000$), parents aged 26–30 years (AOR=2.73, 95% CI=1.04–7.16; $p=0.035$), Muslims (AOR=24.27, 95% CI=2.36–248.74; $p=0.007$), unemployed parents (AOR=2.94, 95% CI=1.35–6.41; $p=0.007$), parents with a household income of <₹15 000 (AOR=1.49, 95% CI=0.962–1.84; $p=0.009$), those from a semi-urban area (AOR=1.61, 95% CI=1.09–2.38; $p=0.016$), lived in the north zone (AOR=3.71, 95% CI=2.37–5.82; $p=0.000$), did not vaccinate or will not vaccinate their child with regular vaccines (other than COVID-19) available under government programs (AOR=1.93, 95% CI=1.19–3.14; $p=0.007$), did not believe in the effectiveness of COVID-19 vaccine for Bangladeshi children (AOR=5.80, 95% CI=3.12–10.78; $p=0.000$), and did not or will not receive the COVID-19 vaccine for themselves (AOR=10.15, 95% CI=7.16–14.39; $p=0.000$).

Contrarily, participants who were non-tobacco users (AOR=0.71, 95% CI=0.53–0.96; $p=0.025$), appeared to be very likely to believe that their children or a family member could be infected with COVID-19 in the next year (AOR=0.21, 95% CI=0.07–0.44; $p=0.000$), and very concerned about their children or a family member contracting COVID-19 in the next year (AOR=0.34, 95% CI=0.21–0.58; $p=0.000$) had significantly lower odds of COVID-19 vaccine hesitancy (Table 2).

Discussion

This nationally representative comprehensive study found a significant prevalence of COVID-19 vaccine hesitancy among parents in Bangladesh for their children. There were substantial differences in COVID-19 vaccine hesitancy according to sociodemographic factors and perceived COVID-19 threat among parents. The logistic regression model revealed that the children's age and parents' age, religion, occupation, monthly household income, permanent address, current living location, tobacco use, adherence to the regular government pediatric vaccination programs (other than COVID-19), perception about COVID-19 vaccine effectiveness for children in Bangladesh, self-vaccine hesitancy, loss of a family member due to COVID-19, and perceived COVID-19 threat can all be used independently to predict parental vaccine hesitancy for children aged <18 years.

To the best of our knowledge, this is the first study to examine parental COVID-19 vaccine hesitancy in Bangladesh. Furthermore, very limited data are available for parental vaccine hesitancy in Indian subcontinental countries. Thus, there is limited information about the previous hesitancy rate in this region. However, the rate observed in this study (42.8%) is

Table 2. Binary logistic regression: predictors of parental vaccine hesitancy in study participants.

Variables	Adjusted OR	Standard error	95% CI		P-value
<i>Children's age group (year)</i>					
0–4	5.876	0.358	2.914	11.850	0.000
5–10	2.845	0.348	1.438	5.631	0.003
11–14	1.101	0.359	0.545	2.225	0.789
15–<18	References				
<i>Children's sex</i>					
Male	0.918	0.127	0.716	1.176	0.498
Female	Reference				
<i>Parents' age group</i>					
18–25	1.513	0.522	0.544	4.205	0.427
26–30	2.732	0.492	1.042	7.165	0.041
31–35	2.755	0.481	1.073	7.075	0.035
36–40	2.737	0.476	1.077	6.958	0.034
41–45	2.524	0.487	0.972	6.552	0.057
46–50	1.101	0.522	0.396	3.064	0.853
≥ 51	Reference				
<i>Religion</i>					
Muslim	24.277	1.187	2.369	248.740	0.007
Hindu	18.704	1.206	1.758	198.977	0.015
Others	Reference				
<i>Educational qualification</i>					
≤ High school	1.002	0.279	0.579	1.731	0.995
College education	1.292	0.246	0.798	2.093	0.297
Graduate	1.007	0.223	0.651	1.560	0.974
Postgraduate	Reference				
<i>Occupation</i>					
Service	2.327	0.303	1.285	4.213	0.005
Business	1.463	0.310	0.797	2.686	0.219
Unemployed	2.943	0.398	1.350	6.415	0.007
Student	4.389	0.480	1.714	11.242	0.002
Home maker	2.105	0.281	1.214	3.650	0.008
Healthcare	2.775	0.370	1.344	5.728	0.006
Daily labor	Reference				
<i>Monthly household income (₹)</i>					
<₹15 000	1.499	0.266	0.962	1.840	0.009
₹ 15000–30000	0.826	0.210	0.547	1.248	0.364
₹ 31000–45000	0.559	0.216	0.366	0.854	0.007
>₹ 45000	Reference				
<i>Current residence type</i>					
Own	1.202	0.310	0.655	2.205	0.552
Rented	0.984	0.311	0.535	1.810	0.959
Others	Reference				

Table 2. *Continued*

Variables	Adjusted OR	Standard error	95% CI		P-value
<i>Permanent address</i>					
Village	1.259	0.177	0.891	1.780	0.192
Semi-urban	1.614	0.198	1.095	2.381	0.016
City	Reference				
<i>Current living location</i>					
Central zone including Dhaka	3.112	0.213	2.049	4.727	0.000
North zone	3.716	0.230	2.370	5.827	0.000
South zone	Reference				
<i>Present tobacco user</i>					
No	0.716	0.149	0.535	0.959	0.025
Yes	Reference				
<i>Political affiliation</i>					
Ruling party	1.001	0.188	0.692	1.448	0.997
Opposition	1.310	0.261	0.785	2.188	0.301
Neutral	0.971	0.176	0.687	1.371	0.865
I prefer not to say	Reference				
<i>Vaccinated/plan to vaccinate children under regular (other than COVID-19) govt. vaccination programs</i>					
No	1.937	0.247	1.193	3.144	0.007
Yes	Reference				
<i>Do you think the COVID-19 vaccine will be effective for Bangladeshi children</i>					
No	5.805	0.316	3.124	10.786	0.000
Yes	0.052	0.171	0.037	0.073	0.000
Skeptical	Reference				
<i>Have you taken or plan to take the COVID-19 vaccine</i>					
No	10.152	0.178	7.161	14.392	0.000
Yes	Reference				
<i>Have you or your family member(s) tested positive for COVID-19</i>					
No	1.320	0.164	0.956	1.822	0.091
Yes	Reference				
<i>Have you lost any of your family member(s) to COVID-19</i>					
No	2.502	0.337	1.293	4.839	0.006
Yes	Reference				
<i>Perceived likelihood of children or family members' infection in the next year</i>					
Very likely	0.206	0.383	0.097	0.437	0.000
Somewhat likely	0.413	0.330	0.216	0.788	0.007
Not likely	0.687	0.329	0.360	1.309	.253
Definitely not	Reference				
<i>Level of concern about children or family members' infection in the next year</i>					
Very concerned	0.345	0.267	0.205	0.583	0.000
Concerned	0.502	0.207	0.334	0.754	0.001
Slightly concerned	0.593	0.202	0.399	0.881	0.010
Not concerned at all	Reference				

similar to that found in the USA (42%)²¹ and slightly lower than that found in China (52%).⁹ Contrarily, the observed rate was significantly higher than that in Brazil, Malaysia, and Saudi Arabia.^{8,22,23} The high parental vaccine hesitancy rate in Bangladesh induces further concern when vaccinating an optimum number of the subcontinental population.

Health behavior theory is centrally influenced by disease risk perception. Herein, a strong association was found between perceived COVID-19 threat and parental vaccine hesitancy. Parents who thought their children or family members were not likely to be at risk of contracting COVID-19 were highly hesitant toward vaccinate their children. Similarly, parents who were not concerned about children or family members' infections were hesitant. Furthermore, vaccine hesitancy was significantly higher among those who did not believe or remain skeptical about COVID-19 vaccine efficacy among Bangladeshi children than those who did. These findings were consistent with the results of previous studies that measured COVID-19 vaccine hesitancy among the adult population.^{10,24,25} Ignorance, belief in conspiracy theories, and even denial of the existence of COVID-19 may influence one's thought of self-vaccination or vaccinating a child.²⁶ Therefore, further studies are warranted to understand the in-depth association between COVID-19 threat and vaccine hesitancy among different population groups.

Parents who reported unemployment, an education level lower than or equal to high school, a household income of <₳15 000-₳30 000, and those from the village or semi-urban area were significantly more vaccine-hesitant. Similarly, a previous study found high vaccine hesitancy among unemployed, low education level, and lower-income groups of parents in several high- and middle-income countries.^{3,8,22,27} Furthermore, our previous study on the adult Bangladeshi population found a similar scenario.¹⁰ Global research and studies conducted in the USA and Saudi Arabia among the general population reported identical results.²⁸⁻³⁰

A previous study found a higher prevalence of vaccine hesitancy among younger parents and parents of children aged between 0 and 4 years,²¹ and our analysis yielded similar results. However, unlike a previous study in Malaysia (a multi-ethnic country),²³ we found high vaccine hesitancy among Muslim parents compared to non-Muslim parents in Bangladesh. Hence, more studies are warranted to understand the influence of religion on the decisions regarding vaccine acceptance and rejection.

Our study found a significantly high prevalence of parental vaccine hesitancy among tobacco users. A previous study also found a similar result, citing the association between unhealthy life practices and vaccine hesitancy among tobacco users.¹⁰ Interestingly, vaccine hesitancy among parents living in the north zone of Bangladesh has also been shown to be high. This is likely because the north zone of Bangladesh is a tobacco-producing area with a high poverty level. Therefore, this information may explain the high prevalence of vaccine hesitancy among parents who are tobacco users and living in the north zone of Bangladesh.

Additionally, this study found a strong association between self-vaccination intention and vaccination decision for children. The incidence of parental vaccine hesitancy was 10 times higher among parents who did not receive or will not receive the COVID-19 vaccine for themselves than among those who did and will receive the COVID-19 vaccine. Furthermore, parents who did not get their children vaccinated with regular vaccines other than COVID-19 were also highly hesitant toward the COVID-19 vaccine for their children, indicating stubborn vaccine hesitancy among groups of people. Therefore, special advocacy targeting these groups is recommended when including them in the vaccination program to eradicate vaccine-preventable diseases. Conversely, we found high vaccination willingness among parents who reported that their family member(s) had either tested positive for or died of COVID-19. The harmful effect of COVID-19 may encourage patients to make a favorable decision about vaccinating their children when it is available.

Strengths and limitations

This is the first study to reveal the rate of parental vaccine hesitancy for children in Bangladesh. This study included parents' data from all eight divisions of Bangladesh by randomly selected data collection sites; 52% women, 10% Hindu parents, and 1% other non-Muslim parents participated, providing a good representation of the population. Additionally, we conducted anonymous face-to-face interviews to reduce social desirability bias, minimize non-response, and maximize the quality of collected data. Data from parents of children with disabilities have also increased the generalizability of our findings.

Nevertheless, this study has several limitations. Previous studies have found that vaccine hesitancy is complicated, time and location vary, and adherence-specific matter depends on the perceived behavioral nature of the community.³¹⁻³³ We conducted this study when the COVID-19 detection rate in the community was significantly lower than it has been, which may have influenced the perceived threat of the disease and the vaccine hesitancy rate. This study did not measure social and traditional media influences, which may have confounded the results.³⁴ Lastly, our questionnaire did not include questions specific to attitudes, beliefs, or mistrust about the vaccine.

Conclusions

To ensure the optimum coverage of vaccines, the government, public health officials, and advocates should be prepared to address parental vaccine hesitancy to reach their target and establish programs to improve childhood COVID-19 vaccine literacy among parents. Availability of safety and efficacy data for COVID-19 vaccines for children in social and traditional media, community and healthcare centers, and mosques/temples would likely positively impact community dwellers' attitudes toward childhood COVID-19 vaccination. Engaging community and religious leaders, family physicians, and trustworthy relatives should accelerate advocacy programs to reduce parental vaccine hesitancy for their children in the community.

Data availability

Underlying data

Open Science Framework: Parental coronavirus disease vaccine hesitancy for children in Bangladesh: a cross-sectional study. <https://doi.org/10.17605/OSF.IO/43G5M>³⁵

The project contains the following underlying data:

- Parental Vac Hesitancy F1000.sav (raw data from questionnaires)

Extended data

Open Science Framework: Parental coronavirus disease vaccine hesitancy for children in Bangladesh: a cross-sectional study. <https://doi.org/10.17605/OSF.IO/43G5M>³⁵

The project contains the following extended data:

- Parental Vac Questionnaire.docx

Reporting guidelines

Open Science Framework: Parental coronavirus disease vaccine hesitancy for children in Bangladesh: a cross-sectional study. <https://doi.org/10.17605/OSF.IO/43G5M>³⁵

This project contains the following reporting guidelines checklist:

- STROBE_checklist_Par_Vac_Hesi.docx

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

The authors would like to thank the participants for providing the information used to conduct the study. Also, the authors would like to thank Zarin Tasnim and Umme Salma Khan for assisting with data collection.

References

1. Hodgson SH, Mansatta K, Mallett G, *et al.*: **What defines an efficacious COVID-19 vaccine? A review of the challenges assessing the clinical efficacy of vaccines against SARS-CoV-2.** *Lancet Infect. Dis.* 2021 Feb 1 [cited 2022 Jan 13]; **21**(2): e26–e35. [PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
2. Randolph HE, Barreiro LB: **Herd Immunity: Understanding COVID-19.** *Immunity.* 2020 [cited 2021 Oct 28]; **52**: 737–741. Cell Press. [PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
3. Alfieri NL, Kusma JD, Heard-Garris N, *et al.*: **Parental COVID-19 vaccine hesitancy for children: vulnerability in an urban hotspot.** *BMC Public Health.* 2021 Dec 13 [cited 2021 Oct 28]; **21**(1): 1662. [PubMed Abstract](#) | [Publisher Full Text](#)
4. MacDonald NE: **Vaccine hesitancy: Definition, scope and determinants.** *Vaccine.* 2015 Aug 14 [cited 2021 Jan 16]; **33**(34): 4161–4164. [Publisher Full Text](#) | [Reference Source](#)
5. Cascini F, Pantovic A, Al-Ajlouni Y, *et al.*: **Attitudes, acceptance and hesitancy among the general population worldwide to receive the COVID-19 vaccines and their contributing factors: A systematic review.** *iScience.* 2021 Oct [cited 2021 Oct 29]; **40**: 101113. [PubMed Abstract](#) | [Publisher Full Text](#)

6. World Health Organization: *Weekly Operational Update on COVID-19*. World Heal Organ; 2021; (February): 1–10.
[Reference Source](#)
7. Kostoff RN, Calina D, Kanduc D, *et al.*: **Why are we vaccinating children against COVID-19?** *Toxicol. Rep.* 2021 Jan 1; **8**: 1665–1684.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Bagateli LE, Saeki EY, Fadda M, *et al.*: **COVID-19 Vaccine Hesitancy among Parents of Children and Adolescents Living in Brazil.** *Vaccine.* 2021 Sep 30 [cited 2021 Oct 5]; **9**(10): 1115.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
9. Zhang M-X, Lin X-Q, Chen Y, *et al.*: **Determinants of parental hesitancy to vaccinate their children against COVID-19 in China.** *Expert Rev. Vaccines.* 2021 Aug 23 [cited 2021 Oct 5]; **20**: 1339–1349.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Ali M, Hossain A: **What is the extent of COVID-19 vaccine hesitancy in Bangladesh? A cross-sectional rapid national survey.** *BMJ Open.* 2021 Aug 24 [cited 2021 Aug 24]; **11**(8): e050303.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
11. Jamison AM, Quinn SC, Freimuth VS: **"You don't trust a government vaccine": Narratives of institutional trust and influenza vaccination among African American and white adults.** *Soc. Sci. Med.* 2019 Jan [cited 2021 Nov 4]; **221**: 87–94.
[Publisher Full Text](#) | [Reference Source](#)
12. Kempe A, Saville AW, Albertin C, *et al.*: **Parental Hesitancy About Routine Childhood and Influenza Vaccinations: A National Survey.** *Pediatrics.* 2020 Jul 1 [cited 2021 Nov 4]; **146**(1): e20193852.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
13. Hamel L, Lopes L, Kearney A, *et al.*: **KFF COVID-19 vaccine monitor: Parents and the pandemic.** *KFF.* 2021 [cited 2021 Nov 4].
[Reference Source](#)
14. Faria NR, Mellan TA, Whittaker C, *et al.*: **Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil.** *Science (80-).* 2021 May 21 [cited 2021 Nov 4]; **372**(6544): 815–821.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Mlcochova P, Kemp SA, Dhar MS, *et al.*: **SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion.** *Nature.* 2021 Nov 4 [cited 2021 Nov 4]; **599**(7883): 114–119.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
16. Mathieu Edouard RH: **Coronavirus (COVID-19) Vaccinations - Statistics and Research - Our World in Data.** *Nat. Hum. Behav.* 2021 [cited 2021 Nov 4]; 5–7.
[Reference Source](#)
17. United Nations Population Fund: *Handbook of Disease Burdens and Quality of Life Measures*. New York, NY: Springer New York; 2010 [cited 2021 Nov 7]; 4344–4344.
[Reference Source](#)
18. Children 12 and above to get jabs | The Daily Star; [cited 2021 Nov 7].
[Reference Source](#)
19. Taherdoost H: **Determining sample size; How to calculate survey sample size.** *Int J Econ Manag Syst.* 2017; **2**(2): 237–239.
[Reference Source](#)
20. James E, Joe W, Chadwick C: **Organizational research: Determining appropriate sample size in survey research.** *Inf. Technol. Learn. Perform. J.* 2001; **19**(1): 43–50.
21. Szilagyi PG, Shah MD, Delgado JR, *et al.*: **Parents' Intentions and Perceptions About COVID-19 Vaccination for Their Children: Results From a National Survey.** *Pediatrics.* 2021 Oct; **148**(4): e2021052335.
[Publisher Full Text](#) | [PubMed Abstract](#) | [Reference Source](#)
22. Aldakhil H, Albedah N, Alturaiki N, *et al.*: **Vaccine hesitancy towards childhood immunizations as a predictor of mothers' intention to vaccinate their children against COVID-19 in Saudi Arabia.** *J. Infect. Public Health.* 2021 Oct 1 [cited 2021 Nov 6]; **14**(10): 1497–1504.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
23. Mohd Azizi FS, Kew Y, Moy FM: **Vaccine hesitancy among parents in a multi-ethnic country, Malaysia.** *Vaccine.* 2017 May 19 [cited 2021 Oct 6]; **35**(22): 2955–2961.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
24. Caserotti M, Girardi P, Rubaltelli E, *et al.*: **Associations of COVID-19 risk perception with vaccine hesitancy over time for Italian residents.** *Soc. Sci. Med.* 2021 Mar 1 [cited 2021 Jun 17]; **272**: 113688.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Kucukkarapinar M, Karadag F, Budakoglu I, *et al.*: **COVID-19 Vaccine Hesitancy and Its Relationship With Illness Risk Perceptions, Affect, Worry, and Public Trust: An Online Serial Cross-Sectional Survey From Turkey.** *Psychiatry Clin Psychopharmacol.* 2021 Apr 12; **31**(1): 98–109.
[Publisher Full Text](#) | [Reference Source](#)
26. Bertin P, Nera K, Delouée S: **Conspiracy Beliefs, Rejection of Vaccination, and Support for hydroxychloroquine: A Conceptual Replication-Extension in the COVID-19 Pandemic Context.** *Front. Psychol.* 2020 Sep 18; **0**: 2471.
27. Teasdale CA, Borrell LN, Shen Y, *et al.*: **Parental plans to vaccinate children for COVID-19 in New York city.** *Vaccine.* 2021 Aug; **39**(36): 5082–5086.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Lazarus JV, Ratzan SC, Palayew A, *et al.*: **A global survey of potential acceptance of a COVID-19 vaccine.** *Nat. Med.* 2020 Oct 20 [cited 2021 Jan 16]; **27**: 225–228.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Al-Mohaithef M, Padhi BK: **Determinants of COVID-19 Vaccine Acceptance in Saudi Arabia: A Web-Based National Survey.** *J. Multidiscip. Healthc.* 2020 Nov 20 [cited 2021 Feb 5]; **13**: 1657–1663.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
30. Malik AA, McFadden SM, Elharake J, *et al.*: **Determinants of COVID-19 vaccine acceptance in the US.** *EClinicalMedicine.* 2020 Sep [cited 2021 Jan 14]; **26**: 100495.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Xiao X, Wong RM: **Vaccine hesitancy and perceived behavioral control: A meta-analysis.** *Vaccine.* 2020 Jul 14 [cited 2021 Feb 5]; **38**(33): 5131–5138.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
32. Larson HJ, Jarrett C, Eckersberger E, *et al.*: **Understanding vaccine hesitancy around vaccines and vaccination from a global perspective: A systematic review of published literature, 2007–2012.** *Vaccine.* 2014; **32**: 2150–2159. Elsevier BV.
[Publisher Full Text](#)
33. Wilson K, Nguyen HH, Brehaut H: **Acceptance of a pandemic influenza vaccine: a systematic review of surveys of the general public.** *Infect Drug Resist.* 2011 Oct 20 [cited 2021 Feb 5]; **4**(1): 197.
[Publisher Full Text](#) | [Reference Source](#)
34. Reno C, Maietti E, Di Valerio Z, *et al.*: **Vaccine Hesitancy towards COVID-19 Vaccination: Investigating the Role of Information Sources through a Mediation Analysis.** *Infect Dis Rep.* 2021 Aug 12 [cited 2021 Nov 7]; **13**(3): 712–723.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
35. Ali M: **Parental Coronavirus Disease Vaccine Hesitancy for Children in Bangladesh: A Cross-sectional Study.** *OSF.* 2022 January 14.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ? ✓ ✓

Version 1

Reviewer Report 11 February 2022

<https://doi.org/10.5256/f1000research.80146.r120978>

© 2022 Kader M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Manzur Kader

Post doctoral researcher, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

Abstract

- In methods, the Authors say "Predictors were identified using binary logistic regression analysis. Rephrase it like "Factors associated with vaccine hesitancy were identified using binary logistic regression analysis". Because identifying predictors require a stronger study design than the cross-sectional study.
- Replace the word "face-to-face interviews" with an interview-administered questionnaire. Because "face-to-face interviews" is commonly used in a qualitative study
- Write the name of the questionnaire used to assess vaccine hesitancy

Results

Results should be more specific:

- children's age (specify the age?), parent's age (specify the age?, religion (specify?), occupation (specify what occupation?), monthly household income (what level?), permanent address, living location (zone? Specify)
- Write "associated factors with parental vaccine hesitancy", instead of "the independent predictors of parental vaccine hesitancy"

Introduction

- Authors provide some data of prevalence of vaccine hesitancy, for example, the USA, and China (high-resourced countries)." Provide some similar examples from low-resourced countries if available as the study setting.
- Provide a clear justification of vaccine hesitancy, why this might happen? In relation to fear, safety, the effectiveness of vaccines? It's not clear the setting of reference 13, specify it.

- Authors say, “along with sociodemographic variabilities and religious beliefs....” Specify the sociodemographic variabilities.

Methods

- “completeness, accuracy, and internal consistency and secured them” Did you use any statistical test to assess such internal validity or reliability of the questionnaire? For example, to assess internal consistency requires Cronbach's alpha. Or did you assess it just by visual perception? specify.
- Was the vaccine hesitancy questionnaire validated in a similar population like in Bangladesh? write some more about the questionnaire.

Results

- Write higher secondary, instead of “College education”. Write BDT instead of “<”

Discussion

- Write a statement if the results agree with the hypothesis?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public health, Global Health, Epidemiology,

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 27 Feb 2022

Mohammad Ali, Uttara Adhunik Medical College, Dhaka, Bangladesh

Dear reviewer, thank you very much for your precise suggestions. We have revised our manuscript in light of your directions.

Abstract:**Method**

1. We relaced the word predictor by associated factors.
2. Most of the studies used a web-based data collection method in the pandemic time. However, we use the in-person interviews (face-to-face) method to get high-quality data.
3. The question was named as vaccine hesitancy questionnaire.

Result

1. & 2. We rewrote the result section.

Main text:**Introduction:**

1. Data regarding parental vaccine hesitancy in low and middle-income countries were scarce; thus, we could not add more references.
2. We amended the sentence.
3. Example added.

Method:

1. We can not extract Cronbach's alfa value for a questionnaire consisting of only one.
2. This questionnaire was used for a vaccine hesitancy study among adults in Bangladesh (Ali & Hossain, 2021).

Result:

1. The symbol is well accepted and used by the British Medical Journal.

Discussion:

1. The statement has been added in the discussion section.

References:

Ali, M., & Hossain, A. (2021). What is the extent of COVID-19 vaccine hesitancy in Bangladesh? A cross-sectional rapid national survey. *BMJ Open*, 11(8), e050303. <https://doi.org/10.1136/bmjopen-2021-050303>

Competing Interests: None

Reviewer Report 09 February 2022

<https://doi.org/10.5256/f1000research.80146.r121413>

© 2022 Moinuddin F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

F. M. Moinuddin

Mayo Clinic Neuro-Informatics Laboratory, Mayo Clinic, Rochester, MN, USA

The paper presented intends to show the prevalence of parental vaccine hesitancy in Bangladesh. They found that vaccine hesitation varied based on sociodemographic characteristics, religion, behavior, and perceived COVID-19 threat. The paper may be useful for the policymaker to identify the obstacle in the mass vaccination program. The paper is well written and well-planned. The methodology is acceptable. However, the following comments are recommended:

1. The authors should change $p=0.000$ to <0.001 for better understanding in Table 1.
2. Please add a zero before a decimal point in Table 1.
3. The COVID vaccine is recommended for students aged between 12 and 17 years in Bangladesh. However, the study participant who had children below 12 years is more than 78% of the total participants. The author should mention the justification for including this group in their discussion.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Methodology and structure

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 27 Feb 2022

Mohammad Ali, Uttara Adhunik Medical College, Dhaka, Bangladesh

Dear reviewer, thank you very much for your review and insight. We have revised our paper

according to your suggestions.

1. We replaced 0.000 by <0.001.
2. In table 1, zero has been added before the decimal point.
3. You are right; we get most data for children under 12. However, the vaccine is being approved for all the pediatric population (0-<18 years). This study will be useful when the COVID-19 vaccine is approved for children under 12 years.

Competing Interests: None

Reviewer Report 08 February 2022

<https://doi.org/10.5256/f1000research.80146.r120976>

© 2022 Ahmed S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shakil Ahmed

Maternal and Child Health Division, ICDDR, Dhaka, Bangladesh

Abstract:

- The abstract requires a more precise statement in each section, i.e., Background, Methods, Results, and Conclusion.
- The Methods are not clear. As the first line of the “Methods” indicates, the parents of children <18 years were interviewed between October 10 – October 31, 2021. However, the following line points out that the parents aged ≥18 years underwent face-to-face interviews. At first glance, it is not easy to understand the study population of this study.
- The key independent predictors of parental vaccine hesitancy should contain more numbers, i.e., odds ratio (OR) and 95% CI’s upper and lower values. Otherwise, it will not be easy for the readers to grasp the primary findings of this study just by reading the abstract.

Introduction:

- It would be more standard if the author would use the word “more than/greater than/similar terms” instead of using the sign “>” in the introduction.
- There are some grammatical errors in the introduction section (especially in the articles). Please correct.
- It’s not clear why vaccine hesitancy research is essential for Bangladesh. A clear justification is required with proper reference and evidence.

- Why are the parents aged ≥ 18 years important? Is there any evidence?

Methods:

Study design and participants:

- Please clarify what do you mean by Parents aged ≥ 18 years? Whether it is the father, aged ≥ 18 years or the mother ≥ 18 years or both? What did you do if one of the parents was aged < 18 years?
- More clear inclusion and exclusion criteria should be mentioned in the manuscript.
- What language did you use in the vaccine hesitance questionnaire to collect the data?
- It would be great for the general readers if the authors could add the sample size formula in the manuscript
- Did the authors validate the vaccine hesitancy questionnaire for the Bangladeshi population as they mentioned they were using a previously employed vaccine hesitancy questionnaire? Is there any cultural validation evidence?

Study questionnaire:

- It would be great if the authors could attach the vaccine hesitancy questionnaire as supplementary documents for the ease of the readers both English and Bengali (if any), which they have used to collect data
- The authors did not add any funding information. Please add.

Sampling technique and data collection:

How did the authors choose the marketplaces, shopping malls, waiting rooms of large hospitals, diagnostic centres, bus and rail stations randomly? Did they acquire any list? If yes, then from where did you acquire the list? Please provide evidence (if there's any)

Statistical analysis:

- How do you dichotomize the four responses to vaccine hesitancy? Is there any evidence or previous research which have done the same? Please provide the proper evidence.
- Age, Number of children, Educational qualification, and monthly household income are ordinal variables. What analysis did you perform for these variables to compare with the vaccine hesitancy?
- Why did you consider all the variables in the logistic regression model? Please explain.

Results:

- As you have collected data from the eight divisions of Bangladesh, it would be logical if the author could show the difference in divisional vaccine hesitancy of Bangladesh. Then it would be easier for the policymaker to emphasize the efforts where it is more required

Parents and children characteristics (Table 1):

- Please use the term "lost a family member" instead of using "lost a loved one."

- Please do not refer the readers as – “Details of the responses to the questions regarding the likelihood of children or family members' infection by COVID-19 and the level of concern about children or family members' disease in the next year are shown”. The author should describe the findings.

Results of descriptive analysis (Table 1):

- The author should not use the content of the outcome variable as the independent variables and use it in the analysis, such as – “Perceived likelihood of children or family members' infection in the next year”, “Level of concern about children or family members' infection in the next year”. Please correct.
- For a better understanding, please mention the meaning of the sign “□” as “Bangladeshi Taka/BDT/Taka/similar term” in the footnote of the table or the manuscript.

Results of regression analysis (Table 2):

- Please mention the comparison group in the table-2's header. Otherwise, it is hard to interpret for the readers.
- The author should not use the content of the outcome variable as the independent variables and use it in the analysis, such as – “Perceived likelihood of children or family members' infection in the next year”, “Level of concern about children or family members' infection in the next year”. Please correct.
- The religion variable's result might be overestimated. The author considered the lowest number of population group “Other”, as the reference group. This variable should be categorized into “Muslim and Non-muslim”.

Strengths and Limitations

- What did the author mean by 1%, non-muslim parents? Here, the groups should be mentioned as written in Table 1.

Conclusions

- What are the recommendations for the policymakers, and how do the policymakers should address the recommendations are not clear. Please elaborate.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Sexual and reproductive health, Maternal health, Child health, Non-communicable diseases, Pandemics, COVID-19, Gender research

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 27 Feb 2022

Mohammad Ali, Uttara Adhunik Medical College, Dhaka, Bangladesh

Dear reviewer, thank you very much for your time and insight. We have amended our manuscript based on your directions. Please find the point-by-point solution to the problems.

Abstract:

1. We have amended the abstract according to your suggestion.
2. The correction has been made in the method section. We replaced < by below where appropriate.
3. The result section has been rewritten. However, we can not add AOR and CI due to word restrictions of the abstract.

Main text:**Introduction:**

1. We replaced symbols with words where applicable.
2. Please be assured that this manuscript is grammatical-error-free; however, we rechecked and made amendments where needed.
3. We amended the justification of the study.
4. The age of marriage in Bangladesh is 18 years and above, thus, we have taken data only from adult parents.

Methods:**Study design**

1. We exclude parents <18 years. We clarified inclusion and exclusion criteria.
2. We use the Bangla language, clearly mentioned in the method section.
3. The sample size formula has been described clearly; please find the formula with appropriate references.
4. This is a questionnaire consisting of one question. British Medical Journal has published data using this questionnaire in Bangladesh (Ali & Hossain, 2021).

Questionnaire

1. Please find the questionnaire in the supplementary section.
2. Funding information has been added; find the information at the upper end of the article.

Sampling technique

1. We clearly described that the list had been collected from the division's website.

Statistical analysis

1. Yes, we followed previously published studies when categorising the vaccine intention (Ali & Hossain, 2021; Khubchandani et al., 2021).
2. We used descriptive analysis for categorical variables.
3. The final model has included significant variables found in the descriptive analysis. Find the information in the statistical analysis section.

Results

1. Our aim was not to compare the hesitancy considering division; rather, we compared the zones of Bangladesh. Future studies should compare the prevalence of COVID-19 vaccine hesitancy according to the divisions.
2. We replaced the loved one with a family member. Thanks for the suggestion.
3. To reduce the words, we refer a few to the table.
4. We use perceived threat as the independent variable, which is one of our core findings. This variable has been used in previous studies as an independent variable.
5. Symbol of Bangladeshi taka is well accepted any used previously by British Medical Journal. We kept the symbol.
6. Mentioned.
7. We used the variable as it is at the core of vaccine hesitancy. We discussed the implications in the discussion section.
8. We have shown the diversity of religions in the study that clearly reflects the general Bangladeshi population. However, when we take this variable in regression, we combined Christian and Buddhists (total 21 samples).
9. We corrected the error. Thank you for your meticulous observation.

Conclusion

1. We rewrote the conclusion section.

Finally, we would like to thank you again for your time and dedication when reviewing the paper. Your suggestion made the paper improve than before.

References:

Ali, M., & Hossain, A. (2021). What is the extent of COVID-19 vaccine hesitancy in Bangladesh? A cross-sectional rapid national survey. *BMJ Open*, 11(8), e050303. <https://doi.org/10.1136/bmjopen-2021-050303>

Khubchandani, J., Sharma, S., Price, J. H., Wiblishauser, M. J., Sharma, M., & Webb, F. J. (2021). COVID-19 Vaccination Hesitancy in the United States: A Rapid National Assessment. *Journal of Community Health*, 1–8. <https://doi.org/10.1007/s10900-020-00958-x>

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research