Check for updates

RESEARCH ARTICLE

Generalized assessment of the impact of Parkinson's disease on functional condition [version 1; peer review: 2 not approved]

Olena Myalovitska, Iryna Lobanova

Department of Neurology, National Medical University, Kiev, 01601, Ukraine

V1 First published: 03 May 2013, 2:121
<https://doi.org/10.12688/f1000research.2-121.v1>
 Latest published: 03 May 2013, 2:121
<https://doi.org/10.12688/f1000research.2-121.v1>

Abstract

Individuals who develop Parkinson's disease are confronted not only with the physical manifestations of the disorder, but also with the psychosocial issues that impact on quality of life.

The aim of the current work was to assess the level of psychosocial and physical functioning impairment in 95 patients with Parkinson's disease (48 men and 47 women aged 65–79, average age 70.3 ± 0.5 , stages of Parkinson's disease according to Hoehn and Yahr scale - 2–2.5). This research was carried out with the help of "Functional Limitation Profile" and "Sickness Impact Profile - 68" tests.

According to the "Functional Limitation Profile" test, the patients with a diagnosis of Parkinson's disease most commonly showed functional condition impairment in 3 categories: "walk" (92% of cases), "social interaction" (85% of cases) and "clarity of mind" (79% of cases). Less frequent condition impairments were changes in the categories "body care and motion" (72% of cases), "emotions" (65% of cases) and "work" (56% of cases). The least impaired were the categories "communication" (22% of cases), "leisure and entertainment" (24% of cases) and "food" (19% of cases). According to the "Sickness Impact Profile - 68" test, patients with Parkinson's disease showed functional condition impairment most often in the categories "mobility control", "social behavior" and "degree of mobility". Patients showed impairment less frequently in the categories "somatic autonomy", "psychic autonomy and communication". The least impaired category was "emotional stability". The highest percent in relation to the maximum possible point was shown in the category "mobility control", and the lowest in the category "emotional stability".

Thus, this study demonstrates significant impact of the disease on the functional condition of these patients. Functional limits, connected with the disease appearance, are most often present in the motion and social sphere of patients' life and less frequently in the emotional sphere.

Open Peer Review

Approval Status **XX**

	1	2
version 1	X	X
03 May 2013	view	view

1. **Alberto Albanese**, IRCCS C Mondino Institute of Neurology Foundation, Milan, Italy
2. **Iracema Leroi**, University of Manchester, Manchester, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Iryna Lobanova (Lobanova_Iryna@ukr.net)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2013 Myalovitska O and Lobanova I. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Myalovitska O and Lobanova I. **Generalized assessment of the impact of Parkinson's disease on functional condition [version 1; peer review: 2 not approved]** F1000Research 2013, 2:121 <https://doi.org/10.12688/f1000research.2-121.v1>

First published: 03 May 2013, 2:121 <https://doi.org/10.12688/f1000research.2-121.v1>

Introduction

Parkinson's disease is a chronic degenerative neurological disorder that adversely affects an individual's motor functioning, which leads to disability, and impacts on their quality of life. Individuals who develop Parkinson's disease are confronted not only with the physical manifestations of the disorder, but with the psychosocial issues that also impact on quality of life. Psychosocial aspects of Parkinson's disease may present as subtle changes with progression of the disease^{1,2,3}

The purpose of the present research was to assess the level of psycho-social and physical functioning impairment in patients with Parkinson's disease.

Methods

95 patients with Parkinson's disease have been examined (48 men and 47 women aged 65–79, average age 70, 3 ± 0.5). An interview and complete neurologic examination, including the Hoehn and Yahr scale⁴, were performed on the same day. Stages of Parkinson's disease according to Hoehn and Yahr scale were 2–2.5.

Having the aim to assess the impact of the disease on the everyday life of patients to the fullest extent, their interviewing included the use of the tests "Functional Limitation Profile" and "Sickness Impact Profile–68"^{7–15}.

While carrying out "Functional Limitation Profile" test, the change of patient's behavior in cases of Parkinson's disease was assessed in 12 categories of life activity: "walk", "body care and motion", "movement", "household activity", "leisure and entertainment", "social interaction", "emotions", "clarity of mind", "sleep and rest", "food", "communication", "work".

While carrying out "Sickness Impact Profile–68" test, functional condition of patients was assessed in 6 categories of life activity. "Somatic autonomy" category reflected the extent of help that the patient requires when performing everyday activities (dressing, standing, walking, eating etc). "Mobility control" category characterized the degree of control of motion functions, including walking and actions with hands. "Psychic autonomy and communication" category described behavior, connected with mental functions and verbal communication. "Mobility degree" category included the ways the disease limits household and professional activity. "Social Behavior" category reflects social sphere of activity. "Emotional stability" category reflects disease impact on the emotional sphere.

Results

While carrying out "Functional Limitation Profile" test, all patients showed functional conditions changes. Impairments in the category "walk" were present in 87 patients (92% of cases). Most patients noted a slowing down in walking speed (92%), shortening of the distance they could walk without rest (79%) and the need for outside assistance during walking (44%).

Changes in the category "body care and motion" were observed in 69 patients (72% of cases). Impairments of this category were connected with slowing of movements—bradykinesia (85%), patients' reduced ability to maintain balance (72%) and the need for outside assistance (43%).

Limitations in the category "movement" were present in 42 patients (44% of cases) and were manifested in the inability to leave the house (15%) or room (7%), the need for support at a certain time of the day (22%).

In the category "household activity" changes were observed in 40 patients (42% of cases), and consisted of reducing the amount of housework (80%), increasing the time needed for rest (63%), and inability to perform hard work (63%).

In the category "leisure and entertainment" impairments were present in 23 patients (24% of cases). The patients showed decreased period of time spent on leisure (65%) and sports (52%).

Changes in the category "social interaction" were observed in 81 patients (85% of cases). Patients rarely participated in social activities (93%), and isolated themselves from the work team (51%).

Changes in the category "emotion" were noticed in 62 patients (65% of cases). Patients demonstrated depressed mood background (65%), talked about the future with hopelessness (85%).

Impairments in the category "clarity of mind" were noticed in 75 patients (79% of cases). Patients showed impairment of memory (85%) and concentration (85%), slowing of thought processes (80%).

In the category "sleep and rest", patients showed sleepiness in the daytime and insomnia at night (41%) and required additional time for rest because of constant fatigue (80%). Changes in this category were noted in 19 patients (38% of cases).

In the category "food", impairments were observed in 18 patients (19% of cases). The most common impairments included decreased appetite (55%), and need for outside assistance when eating (17%).

Changes in the category "communication" were observed in 21 patients (22% of cases). Patients had problems with writing (93%), and difficulties in communication because of speech dysfunction (85%).

Impairments in the category "work" were present in 53 patients (56% of cases). Some of the patients did not work at all (57%), whilst some performed some of the work at home (19%), or did less work than usual (85%), worked fewer hours (80%) performed only light work (42%), or produced lower-quality work (42%). The frequency of functional limitations is given in [Table 1](#) and [Figure 1](#).

Figure 1 The structure of functional limitations in patients with Parkinson's disease according to the "Functional Limitation Profile" test data.

Table 1 The frequency of functional limitations cases in patients with acute disseminated encephalomyelitis according to the "Functional Limitation Profile" test data.

Behavior categories	Number of patients	%
Walk	87	92
Social interaction	81	85
Clarity of mind	75	79
Body care and motion	69	72
Emotions	62	65
Work	53	56
Movement	42	44
Household activity	40	42
Sleep and rest	39	41
Leisure and entertainments	23	24
Communication	21	22
Food	18	19

The frequency of functional limitation cases in patients with Parkinson's disease according to "Sickness Impact profile-68" test and their assessment in points are given in [Table 2](#), [3](#) and [Figure 2](#), [3](#).

Conclusion

The analysis of investigation of patients with Parkinson's disease (duration 3–5 years) according to the tests "Functional Limitation Profile" and "Sickness Impact Profile-68" showed significant changes of their functional condition connected with the occurrence of the disease.

Table 2 The frequency of functional limitation cases in patients with Parkinson's disease according to "Sickness Impact profile-68" test data.

Behavior categories	Number of patients	%
Social Behavior	82	86
Mobility Control	78	50
Degree of mobility	72	34
Somatic Autonomy	67	71
Psychic autonomy and communication	55	58
Emotional stability	35	37

The patients with Parkinson's disease according to the test "Functional Limitation Profile" showed functional condition impairment most often in 3 categories: "walk", "social interaction" and "clarity of mind", while changes in the categories "body care and motion", "emotion" and "work" were less frequent. The least impaired were the categories "communication" and "leisure and entertainment".

The patients with Parkinson's disease diagnosis according to the test "Sickness Impact Profile-68" showed functional condition impairment most often in the categories "mobility control", "social behavior" and "degree of mobility", less frequently in the categories "somatic autonomy" and "psychic autonomy and communication". The least impaired category was "emotional stability". The highest percent in relation to the maximum possible point was observed in the category "mobility control", and the lowest in the category "emotional stability".

Table 3 Assessment (in points) of disease impact on functional condition of patients according to “Sickness Impact profile–68” test data.

Behavior categories	Average point	Total point	Maximum possible point	% in relation to the maximum possible point
Mobility control	6.6±0.4	278	540	51
Social behavior	5.9±0.4	70	270	26
Psychic autonomy and communication	4±0.3	99	450	22
Degree of mobility	5.3±0.2	189	540	28
Somatic autonomy	3.6±0.2	123	495	25
Emotional stability	2.3±0.5	82	765	11

Figure 2 The structure of functional limitations in patients with Parkinson's disease according to the “Sickness Impact Profile–68” test data.

Figure 3 Assessment in points of functional limitations in patients with Parkinson's disease according to the “Sickness Impact Profile–68” test data (% in relation to the maximum possible point).

Thus, according to the results of two tests, the presence of the disease affects the motion and social sphere of life in the patients with Parkinson's disease the most and their emotional state the least.

For the most accurate assessment of the disease impact on the functional condition of the patients diagnosed with Parkinson's disease, a secondary investigation (depending on the rate of the disease progression) should be carried out. Such investigation of disease dynamics enables detection of possible changes in the categories of life that are most affected by the disease occurrence. Assessment of the quality of life of patients at different stages of the disease with the help of Hoehn and

different stages of the disease with the help of Hoehn and Yahr Scale is a very perspective direction of future studies.

Author contributions

O. Myalovitska and Lobanova conceived the study. Lobanova designed the experiments and carried out the research. All authors prepared the first draft of the manuscript and were involved in the revision of the graft manuscript and have agreed to the final content.

Competing interests

No relevant competing interests were disclosed.

References

1. Bergner M, Bobbitt RA, Carter WB, *et al*: The Sickness impact profile: development and final revision of a health status measure. *Med Care*. 1981; **19** (8): 787–805.
2. Bergner M, Bobbitt RA, Kressel S, *et al*: The sickness impact profile: conceptual formulation and methodology for the development of a health status measure. *Int J Health Serv*. 1976; **6** (3): 393–415.
3. Bergner M, Bobbitt RA, Pollard WE, *et al*: The sickness impact profile: validation of a health status measure. *Med Care*. 1976; **14** (1): 57–67.
4. de Bruin AF, Diederiks JP, de Witte LP, *et al*: Assessing the responsiveness of a functional status measure: the Sickness Impact Profile versus the SIP68. *J Clin Epidemiol*. 1997; **50** (5): 529–40.
5. Nanda U, McLendon PM, Andresen EM, *et al*: The SIP68: an abbreviated sickness impact profile for disability outcomes research. *Qual Life Res*. 2003; **12** (5): 583–595.
6. Hoehn MM, Yahr MD: Parkinsonism: onset, progression and mortality. *Neurology*. 1967; **17**: 427–442.
7. Chrischilles EA, Rubenstein LM, Voelker MD, *et al*: Linking clinical variables to health-related quality of life in Parkinson's disease. *Parkinsonism Relat Disord*. 2002; **8** (3): 199–209.
8. Findley L, Eichhorn T, Janca F: Factors impacting on quality of life in Parkinson's disease: results from an international survey. *Mov Disord*. 2002; **17** (1): 60–67.
9. Hanna KK, Cronin-Golomb A: Impact of Anxiety on Quality of Life in Parkinson's Disease. *Parkinsons Dis*. 2012; **2012**: 640707.
10. Kuopio AM, Marttila RJ, Helenius H, *et al*: The quality of life in Parkinson's disease. *Mov Disord*. 2000; **15** (2): 216–223.
11. Schrag A: Quality of life and depression in Parkinson's disease. *J Neurol Sci*. 2006; **248**: 151–157.
12. Schrag A, Jahanshahi M, Quinn N: How does Parkinson's disease affect quality of life? A comparison with quality of life in the general population. *Mov Disord*. 2000; **15** (6): 1112–1118.
13. Schrag A, Jahanshahi M, Quinn N: What contributes to quality of life in patients with Parkinson's disease? *J Neurol Neurosurg Psychiatry*. 2000; **69**: 308–12.
14. Schenkman M, Ellis T, Christiansen C, *et al*: Profile of Functional Limitations and Task Performance Among People With Early- and Middle-Stage Parkinson Disease. *Phys Ther*. 2011; **91** (9): 1339–1354.
15. Williams SJ: Measuring health status? A review of the Sickness Impact and functional limitations profiles. *Health Care Anal*. 1996; **4** (4): 273–283.

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 26 September 2013

<https://doi.org/10.5256/f1000research.1194.r1909>

© 2013 Leroi I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Iracema Leroi

Institute of Brain, Behaviour and Mental Health, University of Manchester, Manchester, UK

This is an uncontrolled study using assessments tools that do not appear to have been validated in Parkinson's disease. The description of the methodology, particularly the recruitment and testing procedure, is insufficient. The literature on the topic has not been adequately reviewed. The authors have not included a Discussion of their findings. The analysis is simplistic.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 16 Oct 2013

Iryna Lobanova, National Medical University, Kiev, Ukraine

The quality of life was assessed using "Functional Limitation Profile" and "Sickness Impact Profile - 68" tests. These allow for assessment of the physical and psychological health of patients and their social functions, as well as global self-assessment of their health. Investigating using this scale does not require a control group of virtually healthy people because the changes assessed by this scale can be observed only among sick people. We used this methodology of generalized quality of life assessment on patients with Parkinson's disease (which is reflected in the introduction to our work).

Competing Interests: No competing interests were disclosed.

Reviewer Report 13 May 2013

<https://doi.org/10.5256/f1000research.1194.r943>

© 2013 Albanese A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alberto Albanese

IRCCS C Mondino Institute of Neurology Foundation, Milan, Italy

The methodology seems to be poor and is insufficiently described. There is no control group.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 16 Oct 2013

Iryna Lobanova, National Medical University, Kiev, Ukraine

The quality of life was assessed using "Functional Limitation Profile" and "Sickness Impact Profile - 68" tests. These allow for assessment of the physical and psychological health of patients and their social functions, as well as global self-assessment of their health. Investigating using this scale does not require a control group of virtually healthy people because the changes assessed by this scale can be observed only among sick people. We used this methodology of generalized quality of life assessment on patients with Parkinson's disease (which is reflected in the introduction to our work).

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research