Check for updates

CASE REPORT

Case Report: Delayed intra-abdominal bleeding following trans-vaginal ultrasonography guided oocyte retrieval for *in vitro* fertilization in patients at risk for thrombo-embolic events under anticoagulant therapy [version 1; peer review: 3 approved]

Roy Mashiach^{1,2}, David Stockheim^{1,2}, Mati Zolti^{1,2}, Raoul Orvieto^{1,3}

¹Department of Obstetrics and Gynecology, Chaim Sheba Medical Center, Ramat Gan, Israel

²Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

³Faculty of Health Sciences, Ben Gurion University of the Negev, Beer Sheva, Israel

V1 First published: 16 Sep 2013, 2:189
<https://doi.org/10.12688/f1000research.2-189.v1>

Latest published: 18 Oct 2013, 2:189
<https://doi.org/10.12688/f1000research.2-189.v2>

Abstract

We report herein, two cases of massive delayed (2 and 4 days) intra abdominal hemorrhage following ovum pick-up (OPU), in patients at risk for thrombo-embolic events, who concomitantly used therapeutic doses of low molecular weight heparin (LMWH). We discuss the possible mechanisms involved in causing the aforementioned delayed bleeding, and call for re-evaluation of the presently accepted anticoagulant co-treatment regimen. These case reports should direct physicians' attention and keep them alert, while conducting IVF treatment to this subgroup of high risk patients.

Keywords

IVF, ovum pick-up, bleeding, anticoagulant

Open Peer Review

Approval Status

	1	2	3
version 2 (update) 18 Oct 2013	 view	 ↑	
version 1 16 Sep 2013	 view	 view	 view

- Fatma Ferda Verit**, Suleymaniye Maternity, Research and Training Hospital, Istanbul, Turkey
- Pedro Barri**, Institut Universitari Dexeus of the Universitat Autònoma de Barcelona, Barcelona, Spain
- William Ledger**, University of New South Wales, Sydney, Australia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Raoul Orvieto (raoul.orvieto@sheba.health.gov.il)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2013 Mashiach R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Mashiach R, Stockheim D, Zolti M and Orvieto R. **Case Report: Delayed intra-abdominal bleeding following trans-vaginal ultrasonography guided oocyte retrieval for *in vitro* fertilization in patients at risk for thrombo-embolic events under anticoagulant therapy [version 1; peer review: 3 approved]** F1000Research 2013, 2:189
<https://doi.org/10.12688/f1000research.2-189.v1>

First published: 16 Sep 2013, 2:189 <https://doi.org/10.12688/f1000research.2-189.v1>

Introduction

Nowadays, ultrasonographically guided trans-vaginal ovum pick-up (OPU) is considered an accepted and safe method for oocyte retrieval worldwide. Nevertheless, the multiple punctures of the vaginal vault can injure or tear the vaginal mucosa, ovaries, intra-abdominal organs or blood vessels, resulting in mild to severe internal and external bleeding and other complications^{1,2}. Clinically significant blood loss after OPU is actually uncommon³, with a reported incidence of severe intra- or retroperitoneal bleeding varying from 0 to 1.3%².

Intraperitoneal bleeding tends to be severe with acute hemodynamic deterioration, whereas retroperitoneal bleeding usually has a later and more indolent presentation. Intra-abdominal bleeding should be suspected immediately after OPU with the development of signs and symptoms of anemia. While a drop in hemoglobin concentration is an indication for prompt blood transfusion, hemodynamic deterioration dictates surgical intervention with subsequent hemostasis. Most of cases are diagnosed several hours following OPU, with a reported interval between OPU and surgical intervention, ranging from 3 to 18 hours^{4,5}.

In patients with an underlying coagulopathy, the interval between OPU and the appearance of clinical signs and symptoms of severe intra-abdominal bleeding or surgical intervention, was reported to be longer. While, Battaglia *et al.*⁶ reported a 3 hour interval between OPU and surgical intervention in a patient with coagulation factor XI deficiency⁶, intervals of 2 and 10 days, were reported in patients suffering from essential thrombocythemia⁷ and factor VIII deficiency⁸, respectively. Notably, none of the aforementioned patients were treated with anticoagulant medications.

The situation is even more complicated for women with known thrombophilia and/or history of a thrombo-embolic event, who use anticoagulant drugs. The hitherto published research regarding this issue is scarce, with only one study reporting on 24 patients treated concomitantly with anticoagulation therapy, who underwent 68 oocyte retrieval procedures⁹. Since IVF treatment was not associated with any medical complication, such as bleeding or thromboembolic complications, Yinon *et al.* have concluded that among patients considered as high risk for a thromboembolic event, the introduction of low molecular weight heparin (LMWH), as a cycle protective treatment, is safe⁹.

Here, we report two cases of massive delayed intra-abdominal hemorrhage following OPU, in patients at risk for thromboembolic events, who concomitantly used LMWH. These cases aim to direct physicians' attention and keep them alert, while treating this subgroup of at risk patients.

Case report

Two patients were referred to our IVF clinic for egg/embryos collection for surrogacy because of the risk associated with pregnancy. Both were under therapeutic doses of LMWH, which according to a senior haematologist consultant, were co-administered during the controlled ovarian hyperstimulation (COH) for IVF, until 16 h

before the oocyte retrieval and resumed 12 h after the procedure. This study was approved by the institutional review board of Sheba Medical Centre.

Case 1:

A 37-year-old patient with essential thrombocytosis, was diagnosed incidentally following routine examination 7 years ago. One year ago, she developed sinus vein thrombosis, necessitating therapeutic doses of anticoagulant therapy. Controlled ovarian hyperstimulation included the multidose GnRH antagonist protocol, with follicle stimulating hormone (FSH) daily dose of 112.5 to 150 IU. She achieved a peak E2 level of 1538 pmol/l with one leading follicle at each ovary. OPU was reported as uneventful with a retrieval of 3 oocytes. A routine complete blood count (CBC) 2 h following OPU revealed a stable hemoglobin (Hb) concentration, as compared to her baseline level (9.92 g/dL vs 10.52 g/dL, respectively).

Two days following OPU she was re-admitted to our ward, because of severe abdominal and shoulder pain, abdominal bloating and tenesmus. On physical examination her abdomen was swollen and tender, with floating blood as evidenced by transabdominal ultrasonography. She was hospitalized and closely observed hemodynamically. During the next 4 h, her Hb concentration dropped to 7.2 g/dL with clinical deterioration, mandating resuscitation, and urgent exploration.

During exploratory laparoscopy, a massive hemoperitoneum was found (approximately 2.5 L of blood), and a profuse bleeding was observed from a large vein in the posterior aspect of the left infundibulo-pelvic ligament. The vessel was successfully coagulated. The patient was transfused with 4 units of blood, 3 unit of cryoprecipitate, and 2 units of fresh frozen plasma. Her postoperative course was uneventful and she was discharged 4 days later with a hemoglobin level of 9.4 g/dL.

Case 2:

A 32-year-old patient, suffering from Budd–Chiari syndrome as a consequence of a myeloproliferative disorder, necessitating therapeutic doses of anticoagulant therapy. Controlled ovarian hyperstimulation included the long GnRH agonist protocol, with HMG daily dose of 375 to 450 IU. She achieved a peak E2 level of 2349 pmol/l with four leading follicles at both ovaries. OPU was reported as technically uneventful, however no oocytes were retrieved.

Three days following OPU, the patient started to feel unwell and complained of lower abdominal pain and dyspnea with stable Hb concentration (10.43–10.95 g/dL). A day later, she gradually became pale, tachycardiac, with a drop in Hb level (8.84 g/dL) that continued (8.66 g/dL) despite a blood transfusion, mandating urgent exploratory laparoscopy. During laparoscopy, a massive hemoperitoneum was observed (approximately 2.5 L of blood), with an active bleeding from a tear of the right ovarian capsule. The tear was successfully coagulated with an accurate hemostasis. The patient was transfused with 5 units of blood, and 2 units of fresh frozen plasma. Her postoperative course was uneventful and she was discharged a week later with a hemoglobin level of 11.68 g/dL.

Discussion

We report herein, to our knowledge, the first 2 cases of massive delayed intra-abdominal hemorrhage following OPU, in patients at risk for thromboembolic event, who concomitantly used *therapeutic* doses of LMWH. We are not aware of any guidelines that have been published by any professional society, regarding OPU under LMWH. Moreover, since most guidelines relate to patients undergoing surgeries that may apply strict hemostatic measures, which are not available during US guided OPU, our report challenges and calls for re-evaluation of the anticoagulation regimen, provided to this subgroup of at risk patients, during an IVF treatment cycle.

Following OPU, the cessation of bleeding requires the interaction of the injured blood vessel with the circulating platelets and clotting proteins in order to form a stable platelet-fibrin clot. Any abnormalities of these factors may result in clinically significant bleeding. In the present cases, both patients received therapeutic doses of LMWH (enoxaparin, 60mg bid), which were discontinued 16 h prior to OPU and resumed 12 h after the procedure. Moreover, intra abdominal bleeding occurred 2 and 4 days following OPU.

Several plausible mechanisms might be responsible for the aforementioned delayed bleeding: (a) early postoperative use of enoxaparin and unfractionated heparin was reported to be associated with a significant increase in re-exploration for postoperative bleeding, often at a significantly delayed time period after the initial surgery¹⁰. Moreover, this delay was especially common with enoxaparin; (b) while the effect of LMWH and unfractionated heparin on the coagulation cascade is well established, we may wonder whether these substances also affect the fibrin nanostructure, which plays a major role in the mechanical strength and lysis of clots. Yeronahos *et al.*¹¹ provided quantitative evidence showing that both LMWH and unfractionated heparin, independent of their other actions on the coagulation cascade, directly alter the nanostructure of fibrin fibers, contributing to improved fibrinolysis.

We therefore believe that the presently accepted anticoagulant co-treatment regimen, including the dose and timing of administration, should be challenged and undergo a critical re-evaluation. For example, delaying and reducing the first dose of LMWH post OPU, with gradual increments might offer some advantages. However, whether this modification will increase the tendency

toward thrombosis, should be further investigated. Moreover, since intra abdominal bleeding obviously occurs, and may occur several days following OPU, we recommend that the patient be kept in the ward for observation for at least 2–4 days following OPU.

In the presented cases we also observed a tear of the ovarian capsule and an inadvertent puncture of a blood vessel within the infundibulo-pelvic ligament. Both complications might have been prevented by meticulous follicular aspiration using a small diameter aspiration needle (19G), and preferably under the guide of color Doppler ultrasonography¹², respectively.

In conclusion, massive delayed intra-abdominal hemorrhage may occur following OPU, in patients at risk for thromboembolic event, who concomitantly used LMWH. These case-reports should direct physicians' attention and keep them alert, while conducting an IVF treatment to this subgroup of high risk patients.

Author contributions

Roy Mashiach: coordinated and participated in the patients' treatment, contributed to the discussion, assisted in writing the paper and edited it in all its revisions.

David Stockheim: participated in the patients' treatment, contributed to the discussion, assisted in writing the paper and edited it in all its revisions.

Mati Zolti: participated in the patients' treatment, contributed to the discussion, assisted in writing the paper and edited it in all its revisions.

Raoul Orvieto: the principal investigator, designed the study, participated in the patients' treatment, contributed to the discussion, assisted in writing the paper and edited it in all its revisions.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Bennett SJ, Waterstone JJ, Cheng WC, *et al.*: **Complications of transvaginal ultrasound-directed follicle aspiration: a review of 2670 consecutive procedures.** *J Assist Reprod Genet.* 1993; **10**(1): 72–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Orvieto R, Ben Rafael Z: **Bleeding, severe pelvic infection and ectopic pregnancy following ART.** In: *Textbook of Assisted Reproductive Techniques: Volume 2: Clinical Perspectives.* 4th Edition. D.K. Gardner, A. Weissman, C. Howles, Z. Shoam, Eds. Informa Healthcare, London, UK Chapt. 2012; **62**: 374–81.
[Reference Source](#)
- Yih MC, Goldschlag D, Davis OK, *et al.*: **Complete blood counts (CBC) after oocyte retrieval: what is normal?** *Fertil Steril.* 2001; **76**(2): S115–6.
[Publisher Full Text](#)
- Dicker D, Ashkenazi J, Feldberg D, *et al.*: **Severe abdominal complications after transvaginal ultrasonographically guided retrieval of oocytes for in vitro fertilization and embryo transfer.** *Fertil Steril.* 1993; **59**(6): 1313–15.
[PubMed Abstract](#)
- Liberty G, Hyman JH, Eldar-Geva T, *et al.*: **Ovarian hemorrhage after transvaginal ultrasonographically guided oocyte aspiration: a potentially catastrophic and not so rare complication among lean patients with polycystic ovary syndrome.** *Fertil Steril.* 2010; **93**(3): 874–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Battaglia C, Regnani G, Giulini S, *et al.*: **Severe intraabdominal bleeding after transvaginal oocyte retrieval for IVF ET and Coagulation factor XI deficiency: a case report.** *J Assist Reprod Genet.* 2001; **18**(3): 178–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- El-Shawarby SA, Margara RA, Trew GH, *et al.*: **Thrombocytopenia and hemoperitoneum after transvaginal oocyte retrieval for in vitro fertilization.** *Fertil Steril.* 2004; **82**(3): 735–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kart C, Guven S, Aran T, *et al.*: **Life threatening intraabdominal bleeding after oocyte retrieval successfully managed with angiographic embolization.** *Fertil Steril.* 2011; **96**(2): e99–e102.
[PubMed Abstract](#) | [Publisher Full Text](#)

9. Yinon Y, Pauzner R, Dulitzky M, *et al.*: **Safety of IVF under anticoagulant therapy in patients at risk for thrombo-embolic events.** *Reprod Biomed Online*. 2006; **12**(3): 354–3580.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Jones HU, Muhlestein JB, Jones KW, *et al.*: **Early postoperative use of unfractionated heparin or enoxaparin is associated with increased surgical re-exploration for bleeding.** *Ann Thorac Surg*. 2005; **80**(2): 518–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Yeromonahos C, Marlu R, Polack B, *et al.*: **Antithrombin independent effects of heparins on fibrin clot nanostructure.** *Arterioscler Thromb Vasc Biol*. 2012; **32**(5): 1320–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Shalev J, Orvieto R, Meizner I: **Use of color Doppler sonography during follicular aspiration in patients undergoing *in vitro* fertilization may reduce the risk of blood vessel injury.** *Fertil Steril*. 2004; **81**(5): 1408–10.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 30 October 2013

<https://doi.org/10.5256/f1000research.2391.r1846>

© 2013 Ledger W. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

William Ledger

Obstetrics and Gynaecology, University of New South Wales, Sydney, Australia

This article is built around 2 case reports of clinically significant intra-abdominal bleeding in the days following egg collection for the purposes of IVF, in patients receiving therapeutic doses of Heparin. The authors use the case reports as a focus for discussion of the management of patients prescribed this medication prior to IVF.

The reports are provisionally novel to warrant publication. The use of English is somewhat idiosyncratic and requires minor review. The text preceding the first case report includes the suggestion that both eggs and embryos may be collected for surrogacy. Only eggs can be collected, and this should be amended.

The most important part of the article is in the discussion. The authors suggest a review of anticoagulant co-treatment in the light of their cases. I do not think that their experience would be sufficient risk of thrombotic complications. This problem is mentioned but should be further highlighted by the authors. I would also suggest they further emphasize the use of a smaller gage aspiration needle for egg collection in these higher risk cases.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 03 Nov 2013

Raoul Orvieto, Chaim Sheba Medical Center, Ramat Gan, Israel

We would like to thank Prof. Ledger for his valuable comments.

Eggs were collected and fertilized, and the embryos were frozen for future FET cycles to the surrogate. The issue of the thrombotic risk and the suggestion of a smaller gage needle

both appear in the 2nd version.

Competing Interests: No competing interests were disclosed.

Reviewer Report 09 October 2013

<https://doi.org/10.5256/f1000research.2391.r1844>

© 2013 Barri P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pedro Barri

Department of Obstetrics, Gynaecology and Reproduction, Institut Universitari Dexeus of the Universitat Autònoma de Barcelona, Barcelona, Spain

This is an observational case-report on two cases of delayed intra-abdominal bleeding following OPU for IVF in patients under anticoagulant therapy.

This article is acceptable, but some changes should be included:

1. There is no information either on the type of needles used for aspiration, or on the aspiration pressure.
2. Why does a young patient (32 yrs) have a negative OPU?
3. How many needle punctures have been performed?
4. There is no information on the outcomes of the oocytes retrieved in case 1.
5. The discussion section is well structured, but authors should present a final recommendation on daily doses and the duration of LMWH.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 10 Oct 2013

Raoul Orvieto, Chaim Sheba Medical Center, Ramat Gan, Israel

We would like to thank Prof. Barri for his valuable comments.

1. The follicles were aspirated by experienced physicians using a fine (19 G) needle (William

- A. Cook Australia PTY LTD, Australia) and 140 mmHg suction pressure.
2. The negative OPU was unexplained (empty follicle?).
3. Punctures were started in the ovary with the follicles closest to the vagina, and each follicle was sequentially punctured without reinserting the needle through the vaginal wall.
4. In case 1, OPU was reported as uneventful with a retrieval of 3 oocytes, of which 2 were fertilized and cryopreserved.
5. As we already mentioned and further clarify: "We therefore believe that the presently accepted anticoagulant co-treatment regimen, including the dose and timing of administration, should be challenged and undergo a critical re-evaluation. For example, delaying [24 h after the procedure] and reducing the first dose [half] of LMWH post OPU, with gradual increments [after 2 days of half a daily dose] might offer some advantages. However, whether this modification will increase the tendency toward thrombosis, should be further investigated [by multidisciplinary experts]. Moreover, since intra abdominal bleeding obviously occurs, and may occur several days following OPU, we recommend that the patient be kept in the ward for observation for at least 2–4 days following OPU."

Competing Interests: No competing interests

Reviewer Report 02 October 2013

<https://doi.org/10.5256/f1000research.2391.r1843>

© 2013 Verit F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fatma Ferda Verit

Department of Obstetrics and Gynecology, Suleymaniye Maternity, Research and Training Hospital, Istanbul, Turkey

I think this article can be accepted without any revisions. The paper quality is good and it adds new knowledge to the literature.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research