OPINION ARTICLE

Ultimate drivers of native biodiversity change in agricultural systems [version 1; peer review: 2 approved]

David A Norton¹, Nick Reid², Laura Young¹

¹School of Forestry, University of Canterbury, Christchurch, 8140, New Zealand

²Environmental Management, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia

V1 First published: 14 Oct 2013, 2:214
<https://doi.org/10.12688/f1000research.2-214.v1>
Latest published: 14 Oct 2013, 2:214
<https://doi.org/10.12688/f1000research.2-214.v1>

Abstract

The ability to address land degradation and biodiversity loss while maintaining the production of plant and animal products is a key global challenge. Biodiversity decline as a result of vegetation clearance, cultivation, grazing, pesticide and herbicide application, and plantation establishment, amongst other factors, has been widely documented in agricultural ecosystems. In this paper we identify six ultimate drivers that underlie these proximate factors and hence determine what native biodiversity occurs in modern agricultural landscapes; (1) historical legacies; (2) environmental change; (3) economy; (4) social values and awareness; (5) technology and knowledge; and (6) policy and regulation. While historical legacies and environmental change affect native biodiversity directly, all six indirectly affect biodiversity by influencing the decisions that land managers make about the way they use their land and water resources. Understanding these drivers is essential in developing strategies for sustaining native biodiversity in agricultural landscapes into the future.

This article is included in the **Agriculture, Food and Nutrition** gateway.

This article is included in the **Ecology and Global Change** gateway.

Open Peer Review

Approval Status

	1	2
version 1
14 Oct 2013	view	view

1. **Bruce Clarkson**, University of Waikato, Hamilton, New Zealand

2. **Rachel Standish**, University of Western Australia, Crawley, Australia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: David A Norton (david.norton@canterbury.ac.nz)

Competing interests: No competing interests were disclosed.

Grant information: This research was supported by a grant provided to DN by Landcare Research (1213-27-006B).
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2013 Norton DA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Norton DA, Reid N and Young L. **Ultimate drivers of native biodiversity change in agricultural systems [version 1; peer review: 2 approved]** F1000Research 2013, 2:214 <https://doi.org/10.12688/f1000research.2-214.v1>

First published: 14 Oct 2013, 2:214 <https://doi.org/10.12688/f1000research.2-214.v1>

Introduction

Global land use and land cover have changed markedly over the past two centuries as a result of rapid population growth and the increasing demand for food and fibre products. As much as 75% of the Earth's ice-free surface has been directly modified by human activities, mainly through urbanisation, timber harvesting, cultivation, and livestock grazing¹. These changes have resulted in land and water degradation and consequential species extinctions. Ongoing human population growth (perhaps to 9.3 billion by 2050²), including a massive increase in the size of the world's middle class³ and the concomitant increase in the consumption of meat, dairy products and luxury goods derived from animals (e.g. leather and wool), will see the demand for agricultural products increase further. This increase in demand is likely to be met through further losses of natural areas and the transformation of low-production systems to high-input farming and grazing systems. The ongoing expansion and intensification of agriculture will put more pressure on native biodiversity and the ecosystem services that biodiversity provides to humanity^{4,5}. Sound environmental management of agriculture has never been more important for human well-being, the maintenance of global life-support systems, and the survival of planetary biodiversity.

Our concern in this paper is for the native biodiversity that evolved in an area rather than the exotic diversity that is associated with agriculture and came to that place from elsewhere⁶. Exotic biodiversity forms the basis of most agricultural systems worldwide, including crop and livestock species, as well as the many other species that have benefitted from human activities. Maintaining the genetic diversity of exotic species used in agriculture is critical, but is not our focus. Not all changes due to agricultural management will be negative for native biodiversity: some species will prosper from changes, and remnants of some native communities will persist in a matrix of agricultural development⁷. As a general rule however, remnant native taxa tend to be generalists and agricultural development usually results in the homogenisation of native biodiversity⁸.

Degradation of agricultural lands and the inevitable loss of associated native biodiversity within these landscapes is one of the main consequences of the expansion in the extent and intensity of agricultural production at all scales, from local to global⁹. Addressing land degradation and biodiversity loss while maintaining the production of plant and animal products is a key global challenge. Understanding the pressures or drivers that lead to environmental degradation and biodiversity loss is important in meeting this challenge¹⁰. Indeed, it is not possible to restore degraded systems without understanding these drivers and addressing them¹¹. The importance of anthropogenic drivers of change for predicting current and future ecosystem condition was strongly emphasised in the Millennium Ecosystem Assessment^{9,12}. Drivers are any natural or human-induced factors that directly or indirectly cause ecosystem change and biodiversity loss and include both the proximate drivers that result in actual change on the ground and the ultimate drivers that underlie these. Our focus in this article on the ultimate drivers of native biodiversity loss is motivated not by any intent to undermine the obvious importance of proximate causes of change, such as vegetation clearance, cultivation, grazing, pesticide and fertiliser application or plantation establishment, but by a desire to

better understand the underlying or ultimate influences leading to day-to-day management decisions by land managers that directly affect native biodiversity.

In this article we outline a framework for considering ultimate drivers of biodiversity change in agricultural landscapes under six headings (Figure 1): (1) historical legacies; (2) environmental change; (3) economy; (4) social values and awareness; (5) technology and knowledge; and (6) policy and regulation. While environmental change and historical legacies directly affect native biodiversity, all six indirectly affect biodiversity by influencing the decisions that land managers make about the way they use land and water resources. These land management decisions are critical and have a range of flow-on effects for biodiversity. We use examples from Australia and New Zealand to illustrate these underlying drivers of biodiversity change, but similar examples can be found in and applied to any agricultural system worldwide.

Ultimate drivers

The current distribution of biodiversity in agricultural ecosystems reflects the interactions between the environment (climate, land-form, soils, etc.), and the effects of past human activities (habitat creation, modification and loss, introduction of agricultural species, commensals, pesticide and fertiliser application, etc.⁷). Legacies from these historical influences on biodiversity, together with a range of new, primarily anthropogenic, drivers will shape future biodiversity patterns directly through their influence on biodiversity (e.g. land clearance or invasive species), but most often indirectly through their influence on land and water management practices (e.g. the intensity of management inputs), which in turn have direct effects on biodiversity⁹. The various drivers interact across multiple spatial and temporal scales¹² and can work synergistically. It is often synergistic effects that are of most concern for biodiversity conservation^{13–15}. Without a clear understanding of ultimate drivers of land-use (and hence biodiversity) change, it is not possible to identify and implement appropriate strategies for biodiversity conservation, strategies that are urgently required if biodiversity decline is to be reversed over the majority of the Earth's surface^{16–18}.

The way that land managers respond to these drivers in terms of their farm management practices will be critical for native biodiversity, and the outcomes could either be positive or negative depending on the particular response taken. Unfortunately to date, most of the outcomes for native biodiversity have been negative. Developing a good understanding of these drivers and the ways they influence biodiversity is essential if we are to have the ability to influence the way that agricultural landscapes are managed in order to obtain better outcomes for native biodiversity. We now review each of these drivers.

Historical legacies

Historical legacies include the many events that have occurred in the past but have an ongoing influence on both land management practices and biodiversity today. Two types of historical legacy are particularly important: effects of past land management activities on soils (e.g. erosion, salinization, compaction, pesticide application, acidification, nutrient inputs) and ongoing adjustments of remnant biotas to historical habitat destruction and fragmentation.

Figure 1. Relationship between different drivers of change and native biodiversity. While all drivers indirectly affect biodiversity through their effect on land-use practices (thin black lines), only environmental change and historical legacies directly affect biodiversity (thick black lines). Interactions (dotted lines) also occur amongst the different drivers.

Both of these directly affect biodiversity, while changes in soil biogeochemistry also influence farm management practices and hence biodiversity indirectly.

In some agricultural landscapes, periods of intensive land use (e.g. irrigation or fertiliser application) have resulted in fundamental changes in ecosystem attributes, especially in relation to soil properties (e.g. through salinization¹⁹). These modifications have pushed ecosystems across thresholds of change that may take decades, centuries or millennia to reverse²⁰, with their effects influencing both land management practices and biodiversity for the foreseeable future. A good example of an historical legacy having a long-term effect on biodiversity is the influence of phosphorus fertiliser and exotic legume seed inputs in tableland pastures in south-eastern Australia²¹. Elevated soil P and N levels, coupled with clearance

of native timber, resulted in epidemic numbers of native scarabs, particularly Christmas beetles (*Anoplognathus* spp., Scarabaeidae). These beetles, in concert with other defoliating insects, have caused widespread dieback of pasture eucalypts, and continue to hamper establishment of eucalypts in intensively developed pasture land^{21,22}. Elevated soil P and N are good for grass growth and the loss of tree cover has resulted in increased stocking rates because of the greater area of pasture. However, the loss of shade and shelter increases sheep mortality in extreme weather and the loss of woodland and scattered trees has caused reductions in native birds, mammals (microbats and arboreal marsupials) and lizards^{23,24}.

Remnant biotas are still adjusting to the effects of fragmentation and the extinction debt associated with past and current habitat clearance^{25,26}. For many species, particularly those within forested

habitats, the altered environmental conditions that occur as a result of fragmentation reduce the quality of the habitat as edge effects alter forest remnant interior microclimates (e.g. making them warmer, windier and dryer²⁷). Both fragmentation itself and management practices on adjacent agricultural land can alter the disturbances that naturally occurred in these systems (e.g. fire or flooding regimes) or introduce novel disturbances such as grazing, chemical drift and weed invasion, which affect the remnant biota²⁸. Habitat loss also reduces the number of remaining individuals of a species, increasing their vulnerability to disturbance and the likelihood of local extinction, especially as isolation reduces the chance of recolonisation²⁹. At the same time, altering species distributions, both of natives and exotics, means that new species are becoming established in agricultural landscapes causing further changes to remnant biotas, the full impacts of which are yet to be felt³⁰ (see next section).

Environmental change

While climate change and species invasion are only two of a range of environmental changes occurring globally (others include increased nitrogen deposition, land use change and intensification, and CO₂ enrichment), they exert perhaps the strongest influence on both agriculture and biodiversity (we see land-use change as one of the proximate causes of biodiversity change). Climate change and species invasion can both directly affect biodiversity, but also have strong effects on land management practices with flow-on effects on biodiversity.

While the long-term consequences of climate change will be shifts in average rainfall and temperature patterns, it is the changes in the frequency and intensity of extreme events such as droughts, floods, wind and snow storms or frosts that are likely to be most significant for both farm production and biodiversity in the short

to medium term^{31,32}. Native biodiversity will be affected directly through changes in species distributions³³, including invasive species, and altering interactions between species³⁴. Biodiversity will also be affected indirectly through changing land-use patterns as farmers change their management practices to cope with climate change³⁵. While the potential direct effects of climate change on biodiversity are generally well appreciated, the indirect effects are less well understood.

One example of a possible indirect effect of climate change will be through intensification of farm management practices as farmers seek to buffer themselves against the vagaries of unpredictable weather. Irrigation can be used to guarantee grass and crop growth as summer droughts become more frequent, while exotic grasses and legumes can be used to increase productivity of native dryland grasslands to better buffer these systems³⁶. For example, a succession of below-average annual rainfall years in the first decade of the 21st century in the Mackenzie Basin in New Zealand's eastern South Island (Figure 2) has increased pressure on farmers to establish irrigation or plant legume crops such as lucerne as a buffer against the effects of drought. Intensification such as this has a number of flow-on effects for biodiversity, both as a result of habitat loss as remnant vegetation is cleared to accommodate more intensive farming systems^{37,38} and through flow-on effects to other parts of the ecosystem (e.g. eutrophication of waterways as a result of increased fertiliser inputs and irrigation, or altered river flows as a result of abstraction and regulation³⁹).

Invasive species are another important part of environmental change and have the potential to be a key driver of the future condition of biodiversity. In particular, a vast pool of introduced species,

Figure 2. 1927–2009 annual rainfall, Lake Tekapo (Station H40041), Mackenzie Basin, New Zealand (44.002 S, 170.441 E). Data (downloaded from www.cliflo.niwa.co.nz) are expressed as a proportional deviance from the 1927–2009 average value.

especially plants, occur in gardens or as individuals in the wild but do not yet have an obvious impact on either agriculture or native biodiversity⁴⁰. These species may never become a problem, but equally, should environmental conditions or land management practices change, or if they simply cross an abundance threshold, then they could have major impacts in the future. It is difficult to predict when or why a species will start to rapidly expand, but it is clear that for many species this might not occur for years or even decades after the species is first naturalised⁴¹. An example of this is the European daisy, mouse-ear hawkweed (*Hieracium pilosella*, Asteraceae), which was present in New Zealand as early as 1878 (it most likely arrived as a contaminant in grass seed) but wasn't recognised as a problem until the 1970s when it started to rapidly expand across a wide range of sites in the eastern South Island (Figure 3). This species has become a major pest in the rain-shadow mountains of this region where it has invaded grasslands, significantly reducing both livestock feed and biodiversity values through outcompeting other plant species for water and nutrients⁴².

Economy

The economy (from local to global) is perhaps the biggest driver of agricultural decision making, and hence of change in native biodiversity. Except in subsistence systems, farmers are exposed to the vagaries of international markets including the price of farm inputs (especially fuel and fertiliser), changing consumer preferences and demands, the availability of substitutes (e.g. synthetics for natural fibres) and cheap imports, and the effects of national and global economies (e.g. cycles of recessions and booms) on key farm costs such as mortgage interest rates⁴³. Reduced farm incomes also flow on through taxation to the funding available for conservation initiatives at a regional and national government level. While the state of the global economy does not have a direct effect on biodiversity, it has two main consequences that can have marked flow-on effects for biodiversity through effects on farm management: (1) changes in the type of farming operation that is undertaken in response to economic conditions (e.g. the shift towards dairy farming in New Zealand in response to high global dairy prices), and (2) changes in the profitability of the farming operation itself, thus affecting both what the farmer can afford to do (e.g. weed or pest control) and indirectly what government can afford to do in terms of biodiversity conservation (e.g. through financial incentives to farmers to undertake conservation work) through lower taxes.

If prices squeeze profit margins^{44,45}, farmers might seek to increase economies of scale in their operations, which might affect biodiversity, for example, through an increase in the size of individual paddocks with a resultant loss of remnant areas of biodiversity (as already occurs when large pivot irrigators are installed³⁷). Alternatively farmers might be forced to scale back their operations and especially the use of inputs such as fertiliser, which could be good for biodiversity; this occurred in New Zealand when widespread farm subsidies were removed in the 1980s⁴⁶. Another conundrum that can occur is that declines in farm returns can be good for biodiversity as farmers reduce their investment in on-farm improvements enabling native plants and animals to reoccupy areas they had been excluded from by agricultural practices. However, good financial returns could be either positive or negative for biodiversity: positive in that farmers have more disposable income to spend on

non-essential activities such as biodiversity conservation, but negative in that in other situations farmers might choose to intensify their management and increase productivity to take greater advantage of higher returns.

One of the key causes of increased agricultural production over the last 50 years has been the increased use of fertiliser, especially nitrogen-based fertilisers⁴. Fertiliser production is dependent on both supply (especially for fertilisers based on mining naturally occurring deposits, such as phosphorus) and the energy costs of fertiliser production. Spikes in global fertiliser prices in the mid-1970s and late-2000s led to reductions in fertiliser application in New Zealand as farmers had to trade off productivity gains against the increased cost of fertiliser. Cheap fertiliser enables farmers to intensify their management with obvious impacts on biodiversity, while high fertiliser prices generally mean that fertiliser use declines. It is clear that the supply of both fertiliser and fuel is not finite⁴⁷, but the consequences for biodiversity of rapidly increasing prices of both are unclear. Technology may well provide alternatives to traditional fertilisers or alternative ways to retain key nutrients within agricultural systems might be developed. For example, various types of holistic management aim to reduce or even eliminate the use of inputs such as fertilisers by adopting alternative grazing systems and through the increased use of other nutrient sources such as N-fixing trees. Holistic management (e.g. <http://holisticmanagement.org/>) and regenerative agriculture (e.g. <http://regenag.com/web/>) are two examples of these types of approaches. Both are likely to be positive for biodiversity as they encourage a more environmentally sympathetic approach to land management and are attractive to farmers who already have a personal empathy for the environment. High fertiliser prices might force farmers to effectively 'abandon' some parts of their properties or alternatively farmers might choose to increase their stocking rates across their whole property to effectively 'mine' the resources present in order to maintain a certain level of profit in the face of rising prices.

Social values and awareness

The way we value biodiversity and agriculture has a fundamentally important influence on what happens to native biodiversity in agricultural landscapes. This is particularly evident with respect to the way in which society's valuation of agricultural production versus biodiversity conservation has changed through time. As our values and awareness has changed, so too have the ways that we approach land management and hence its impacts on native biodiversity.

When European settlers first colonised Australia and New Zealand, elements of the native biodiversity were seen as a hindrance to 'good' land management⁴⁸. The new settlers struggled to understand the local environment while land management approaches were biased by the European agricultural tradition and 'what worked best'. Few people at that time believed that indigenous ecosystems, especially timbered areas, held any value beyond that of the timber and grass they supported, or the land on which they grew. In fact woody vegetation and wetlands were widely seen as a limitation to 'progress'. Settlers actively sought to impose a European mantle across the land by clearing timber, draining wetlands, planting Northern Hemisphere species, and introducing livestock⁴⁸. Government policy at the time in both Australia and New Zealand actively

Figure 3. Rapid expansion of *Hieracium pilosella* in short tussock grasslands, Tara Hills Station, eastern South Island, New Zealand. (Peter Espie, unpublished data with permission).

promoted this through the setting of land clearance targets that needed to be met before land ownership could be gained (e.g. the Robertson Land Acts in the 1870s in New South Wales). Links between forest cover and soil and water values were not appreciated, while stock numbers in rangelands reached densities never matched subsequently as flocks of sheep in particular exploited the available forage⁴⁹.

Values and awareness have changed over time, both in Australia where aridity and frequent droughts made it difficult to impose the traditional European agricultural model over most of the continent, and in New Zealand where it was realised that a lack of trees resulted in severe soil erosion and loss of the productive potential of the land that farmers were seeking to utilise⁵⁰. More recently the importance of wetlands as filters to reduce the impacts of agricultural pollutants has also been recognised⁵¹ (Figure 4). Soil and water conservation became necessary and new land management practices were developed to cope with the profound changes that resulted from European-style farming. Contour banking of sloping croplands in Australia⁵² and tree planting with exotic species in New Zealand were undertaken to reduce soil erosion⁵⁰. While the importance of setting aside natural areas for biodiversity conservation has been long recognised (e.g. the national park and reserve systems in Australia and New Zealand date back to the late nineteenth century), the intrinsic value of native biodiversity in agricultural landscapes above and beyond its utilitarian value for grazing and timber, and especially its role in providing key ecosystem services (e.g. pollination) has only been recognised in recent decades. Policy and regulation in this area still lags behind the preservation policies that underpin public conservation lands and a shift of emphasis is

urgently needed within land management, research and policy to address this imbalance⁵³.

Technology and knowledge

Technology and knowledge are also important drivers of change in agricultural systems. The availability and affordability of new technology has revolutionised agriculture over the last 50 years. Technological advances in plant breeding and genetic engineering, the use of new herbicides and pesticides, the development of larger, more powerful machinery (tractors, irrigators, etc.), and the application of precision agriculture have all enabled farmers to produce more food and fibre at lower cost, and to move into environments that previously were not able to be farmed (e.g. the heavy soils on alluvial floodplains that early post-war machinery could not cultivate; Figure 5). The development of new genotypes through plant breeding programmes and the incorporation of genetic engineering technology to increase disease resistance and productivity and facilitate weed control has also allowed farmers to increase output of food and fibre production^{4,54}.

Technology is both positive and negative for biodiversity: positive in that many of the new technological advances have enabled farmers to better target their management interventions, thus sparing areas with higher biodiversity values (e.g. through the use of GPS to guide application of fertiliser or herbicide) and in being able to increase productivity in one part of the farm and spare or rest other areas and avoiding adjacent natural areas such as wetlands⁵⁵. Technological advances have also enabled the application of more environmentally sustainable management practices such as direct

Figure 4. Riparian fencing and recent restoration plantings, Westland, New Zealand. (Photo: David Norton with permission).

drilling as opposed to ploughing⁵⁶, especially where native pastures can be direct drilled with production species without affecting the native biodiversity present (called pasture-cropping⁵⁷). However, the advent of new technology has also enabled farmers to farm new areas, especially areas that might otherwise have been marginal or too difficult to crop, and to increase the size of individual management units (e.g. paddocks) and farms in order to capitalise on economies of scale, further increasing impacts on native biodiversity through homogenising farmscapes⁸.

Some technological advances have been both positive and negative. For example, direct drilling enhances soil condition by removing the need for cultivation, but the increased use of herbicides and spray drift may affect adjacent native ecosystems. Genetic engineering has allowed farmers to grow crops that are resistant to glyphosate and certain insect herbivores. Genetically modified glyphosate-resistant crops allow over-spraying with herbicides to control weeds in the crop, both with adverse effects due to herbicide drift into adjacent native vegetation and concomitant loss of habitat of native fauna⁵⁸. In addition, increasing use of glyphosate has fostered selection of glyphosate-resistant weeds, which are now widespread across Australian irrigated farming districts⁵⁹. However, the development through genetic engineering of *Bacillus thuringiensis* (Bacillaceae) insect resistance in commercial crops such as rice and cotton has been positive for native biodiversity, as it requires less pesticide application, which may well have benefits for native invertebrates^{60,61}. However, genetic engineering has not yet been able to address abiotic stressors of plant productivity such as drought⁶², although Monsanto have recently developed a genetically modified form of maize including a gene that enables plants to decrease their soil water absorption rate under dry conditions⁶³. Such developments could potentially provide farmers with another tool for improving resilience in the face of climate change, although traditional plant breeding at this stage appears just as effective as genetic modification⁶³.

Knowledge is a key driver of change in agricultural systems, both in relation to farming practices and biodiversity. Knowledge feeds into social values and awareness as well as having a direct impact on the way land is managed and hence on biodiversity. Much of the impact of past agricultural practices on biodiversity has occurred because farmers (and land managers more generally) have not had

the relevant information to enable them to realise the adverse consequences of their activities. The recent recognition of the importance of biodiversity conservation in agricultural landscapes, especially in 'new' countries like Australia and New Zealand, has limited the attention that biologists have given to these areas. Several studies have highlighted the lack of ecological research undertaken in agricultural landscapes^{64,65}. For example, a survey of New Zealand ecological literature published over the period 1968–1997, found that 65% of articles focused exclusively on protected areas while 18% focused exclusively on non-protected areas (mainly agricultural and plantation forest landscapes), despite non-protected areas accounting for nearly 70% of the total land area⁶⁶. While native biodiversity in agricultural landscapes has received more attention in recent years⁷, there are still substantial gaps in our knowledge in terms of just what native biodiversity is present there, how this biodiversity is impacted by agricultural practices, and what ecosystem services this biodiversity provides to agricultural production specifically and society more generally.

Policy and regulation

Policy and regulation develop in part from social value systems, but in many ways are not in tune with current societal values or awareness of issues. This occurs because the politicians and bureaucrats who develop policy are often not willing to tackle current societal issues (as is clearly evidenced by the gun debate in the USA), perhaps because of the pressures associated with short election cycles, and because the vast majority of society simply does not have any idea of the diversity of issues that are regulated and hence is not involved in debates about them. This leaves policy making and regulation in the hands of a very small group of politicians, bureaucrats and lobby groups. In addition, historical legacies of past policy, including the case law associated with legislation, drive ongoing policy development. Notwithstanding this, policy and regulation exert a very powerful influence on land management decision making⁶⁷, with both positive and negative outcomes for biodiversity. Policy influences decision making at every level from international to national and regional levels. Protectionist trade tariffs are one example where international policy can directly influence the types of decisions made by farmers³⁰, for example on the type of farming that is undertaken. The removal of trade barriers is also likely to further facilitate the flow of invasive species as trade increases facilitating these species in reaching new areas through increased dispersal opportunities⁶⁸.

Domestic policy and associated regulation also have a marked impact on a farmer's decision making. Regulation can be particularly poor in mitigating the many small incremental actions of multiple actors. This is illustrated by vegetation clearance rules that are widely used to restrict the amount of native vegetation cleared in Australia and New Zealand ([Supplementary material](#)). However, if every farmer in a region cleared what was legally possible under exemptions associated with such rules, biodiversity would continue to decline. Furthermore, some farmers look at vegetation clearance rules as a cost of business, with the fine or penalty associated with breaking the rule being trivial compared to the potential profits that would result from utilising newly cleared areas. Blunt regulatory instruments are weak at controlling gradual land-use intensification. (e.g. the loss of palatable native plants from fertilised and over-sown

Figure 5. Cultivating heavy cracking clay soils on inland floodplains, New South Wales, Australia. (Photo: Leah MacKinnon with permission).

pastures subjected to increased stocking rates), nor do they deal with the habitat loss resulting from species invasions. Some policy can have unintended positive outcomes for biodiversity. In New Zealand, a change in government in the early 1980s resulted in the removal of a range of farm subsidies. As a result of this, fertiliser price increased locally, even though global fertiliser prices were low, resulting in farmers using less fertiliser and in some cases abandoning less productive land with many of these areas now regenerating back into native forest^{69,70}.

Conclusions

The effects of different drivers on farmers' decision making and hence on land management activities and native biodiversity are complex and often interrelated. The following example illustrates how two of these drivers can affect native biodiversity in a live-stock production system through their influence on farm management practice. Both environmental change and the international market place can place substantial pressure on the profitability of a farm business, especially where financial commitments such as mortgage payments are an issue. In the case of environmental change, this can be through changes in the frequency and intensity of extreme climatic events such as droughts, heatwaves, floods, windstorms or frosts that are significant for farm production^{31,32}. In the case of international markets, factors such as increasing costs of external inputs such as fuel and fertiliser coupled with fluctuating returns for farm products are critical. One response to these types of pressure is to intensify farm management practices in order to buffer the farm business against the vagaries of unpredictable weather or markets. Intensification can involve one or more management actions (the proximate drivers of change), including increasing the carrying capacity of existing pastures (e.g. through cultivation or topdressing with seed and fertiliser), bringing new land into production through native vegetation clearance, or by overgrazing the existing forage base (as a short-term strategy to cope with immediate financial challenge). All of these can result in land degradation and loss of native biodiversity.

Agricultural systems are of course very diverse and each driver will have a different effect depending on the nature of the particular

system including the type of farming system (grazing, cropping, irrigated, rain-fed, with or without woody vegetation, etc.), the productivity of the system (annual yield of grain, milk solids, meat, etc.), and the amount and distribution of native biodiversity present. The potential to intensify farm management for whatever reason is the primary cause of native biodiversity vulnerability³⁰, as intensification almost always results in the loss of native species, both where this involves clearing areas of remnant native vegetation or through transformation of current farmed areas from low-producing to high-producing states. In contrast, native biodiversity in systems that have relatively little potential for intensification, either because they have already been intensively developed (e.g. irrigated arable or dairy farming systems) or because productivity is limiting and cannot be readily enhanced (e.g. rangeland systems lacking water), is likely to be less vulnerable.

An appreciation of the drivers of biodiversity change within agricultural ecosystems is important as it enables us to better plan management activities in order to achieve desired outcomes for native biodiversity. Distinguishing between direct and indirect drivers is particularly important, as it is the indirect or ultimate drivers that are likely to have the greatest influence on future biodiversity given that the primary use of these ecosystems is for the production of food and fibre. The drivers discussed here are ultimately responsible for most impacts on native biodiversity in agricultural systems, both directly but especially through their strong influence on farm management practice. Management in turn dictates the nature and habitat qualities of the production matrix and shifts remnants of native ecosystems into structural and compositional states further from their historical condition⁷¹. These changes are likely to require novel approaches for the long-term management of native biodiversity⁷² as well as an explicit evaluation of the value of this biodiversity, both for itself (intrinsic value) and for the ecosystem services that it provides to farm production and to society more generally.

In order to address the effects of the different drivers of native biodiversity in agricultural landscapes, mechanisms are required to encourage farmers to incorporate management of native species into overall farm management practices. In some cases, all that is required is to facilitate the farmer to continue with their existing management approach. However, more often mechanisms are required to assist farmers to change their management or adopt new practices that are more sympathetic to native biodiversity⁷. In selecting the most appropriate approach for doing this, it is necessary to consider the relative levels of public (external) and private (internal) benefits of a given land management practice⁷³ and to ensure that these are appropriately accounted for. Simply putting in place regulations that limit carrying capacity or vegetation clearance do not necessarily address the underlying (ultimate) reasons driving management decisions. In fact, poorly thought through regulations can actually result in unintended or perverse outcomes (e.g. proposed vegetation clearance rules resulting in a flurry of clearance before regulations are enacted). Therefore, in order to address the indirect effects of the ultimate drivers of biodiversity change considered here, other approaches need to be considered. Such approaches could better align on-farm decision making with positive biodiversity outcomes, for example, by offering financial incentives to farmers to retain native vegetation and provide ecosystem services such as clean water and carbon sequestration^{74,75}.

Author contributions

DN and NR conceived the article, LY researched examples to include, DN prepared the first draft, and DN, NR and LY substantially revised the article for publication and have agreed to the final content.

Competing interests

No competing interests were disclosed.

Grant information

This research was supported by a grant provided to DN by Landcare Research (1213-27-006B).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank Peter Bellingham and Jason Tylianakis for constructive comments on earlier versions of this article.

Supplementary material

Examples of a vegetation clearance rules in three New Zealand district plans (local government land planning statutes).

Waikato District Plan rule 25.43.1

(www.waikatodistrict.govt.nz/CMSFiles/ca/cab18ac7-cba6-49e1-888b-4c9d5efdc1cb.pdf)

Vegetation clearance of indigenous vegetation or habitat of indigenous fauna in a Landscape Policy Area or Conservation Policy Area is a permitted activity if:

(a) it does not exceed 1000m² or 1% of contiguous indigenous vegetation or habitat of indigenous fauna, whichever is the lesser, per contiguous area per site in any 3-year period, and is limited to:

- (i) a building platform for a permitted or approved building, or structure(s) or access, or
- (ii) gathering of plants in accordance with Maori custom and values, or

(b) it does not exceed 3000m² or 3% of contiguous indigenous vegetation or habitat of indigenous fauna, whichever is the lesser, per contiguous area per site in any 3-year period, and is limited to:

- (i) maintaining or reinstating productive pasture and productive forests and maintenance of tracks and fences through the removal of manuka and/or kanuka and/or treeferns that are more than 10m from a water body and less than 15 years old or less than 5m in height and any under-storey under such manuka or kanuka or treeferns growing on land that was previously in productive use.

Hurunui District Plan rule A7.2.1

(www.hurunui.govt.nz/assets/Documents/District%20Plan/2-A-07%20Natural%20Environment.pdf)

(d) Wetlands

There shall be no clearance or cultivation or damage by deposition of material or infilling of indigenous vegetation in any wetland exceeding 1000 square metres in area.

(e) Clearance of indigenous vegetation

(i) No clearance of indigenous vegetation not already significantly modified by any farming practice, other than as provided for in

- Section A2 – Landscape
- Section A7 – Natural Environment
- Section B2 – Coastal Environment
- Section B3 – Hurunui Lakes Area

shall be permitted of greater than 1 hectare over a 5-year period on any separate certificate of title.

Mackenzie District Plan rule 12.1.1g

(www.mackenzie.govt.nz/Site/Documents_and_policy/key_documents/district_plan.aspx)

Short Tussock Grasslands

On each of the individual farm properties existing in the Mackenzie Basin Map as at 1 January 2002 in any continuous period of five years there shall be no clearance including cultivation above the following thresholds of short tussock grasslands, consisting of silver or blue (*Poa* species), or *Elymus solandri*, or fescue tussock where tussocks exceed 15% canopy cover:

- (i) 40 hectares or less – Permitted Activity
- (ii) Greater than 40 hectares – Discretionary Activity.

References

1. Ellis EC, Ramankutty N: **Putting people in the map: anthropogenic biomes of the world.** *Front Ecol Environ.* 2008; **6**: 439–447.
[Publisher Full Text](#)
2. Anon **World Population Prospects: The 2010 Revision, Highlights and Advance Tables.** Department of Economic and Social Affairs, Population Division Working Paper No. ESA/P/WP.220. United Nations, New York. 2011.
[Reference Source](#)
3. **United Nations Development Programme. The Rise of the South: Human Progress in a Diverse World.** Human Development Report 2013. United Nations Development Programme, New York. 2013.
[Reference Source](#)
4. Tilman D, Cassman KG, Matson PA, *et al.*: **Agricultural sustainability and intensive production practices.** *Nature.* 2002; **418**(6898): 671–677.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Lindenmayer D, Cunningham S, Young A: **Land Use Intensification: Effects on Agriculture, Biodiversity and Ecological Processes.** CSIRO Publishing, Australia. 2012.
[Reference Source](#)
6. Crosby AW: **Ecological Imperialism: The Biological Expansion of Europe, 900–1900.** Revised edition. Cambridge University Press, Cambridge. 2004.
[Reference Source](#)
7. Norton D, Reid N: **Nature and Farming: Sustaining Native Biodiversity in Agricultural Landscapes.** CSIRO Publishing, Melbourne. 2013.
[Reference Source](#)
8. Lockwood JL, McKinney ML: **Biotic Homogenisation.** Kluwer Academic, New York. 2001.
[Publisher Full Text](#)
9. **Millennium Ecosystem Assessment Ecosystems and Human Well-being. Synthesis.** Island Press, Washington D.C. 2005.
[Reference Source](#)
10. Mattison EH, Norris K: **Bridging the gaps between agricultural policy, land-use and biodiversity.** *Trends Ecol Evol.* 2008; **23**(11): 610–616.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Hobbs RJ, Norton DA: **Towards a conceptual framework for restoration ecology.** *Restoration Ecology.* 1996; **4**(2): 93–110.
[Publisher Full Text](#)
12. Nelson GC, Bennett E, Berhe AA, *et al.*: **Anthropogenic drivers of ecosystem change: an overview.** *Ecology and Society.* 2006; **11**(2).
[Reference Source](#)
13. Brook BW, Sodhi NS, Bradshaw CJ: **Synergies among extinction drivers under global change.** *Trends Ecol Evol.* 2008; **23**(8): 453–460.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Didham RK, Tylianakis JM, Gemmell NJ, *et al.*: **Interactive effects of habitat modification and species invasion on native species decline.** *Trends Ecol Evol.* 2007; **22**(9): 489–496.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Sala OE, Chapin FS 3rd, Armesto JJ, *et al.*: **Global biodiversity scenarios for the year 2100.** *Science.* 2000; **287**(5459): 1770–1774.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Norris K: **Agriculture and biodiversity conservation: opportunity knocks.** *Conservation Letters.* 2008; **1**(1): 2–11.
[Publisher Full Text](#)
17. Scherr SJ, McNeely JA: **Biodiversity conservation and agricultural sustainability: towards a new paradigm of 'ecoagriculture' landscapes.** *Philos Trans R Soc Lond B Biol Sci.* 2008; **363**(1491): 477–494.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Balmford A, Green R, Phalan B: **What conservationists need to know about farming.** *Proc Biol Sci.* 2012; **279**(1739): 2714–2724.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Cramer VA, Hobbs RJ: **Ecological consequences of altered hydrological regimes in fragmented ecosystems in southern Australia: impacts and possible management responses.** *Austral Ecol.* 2002; **27**(5): 546–564.
[Publisher Full Text](#)
20. Suding KN, Hobbs RJ: **Threshold models in restoration and conservation: a developing framework.** *Trends Ecol Evol.* 2009; **24**(5): 271–279.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Reid N, Landsberg J: **Tree decline in agricultural landscapes: what we stand to lose.** In: Hobbs RJ, Yates CJ, eds. *Temperate Eucalypt Woodlands in Australia: Biology, Conservation, Management and Restoration.* Surrey Beatty & Sons, Chipping Norton, Victoria, 2000; pp 127–166.
[Reference Source](#)
22. Reid N: **Maintaining tree cover health in temperate pastoral landscapes.** In: Barlow T, Thorburn R, eds. *Balancing Conservation and Production in Grassy Landscapes.* Environment Australia, Biodiversity Group, Canberra, 2000; pp 177–187.
[Reference Source](#)
23. Lumsden LF, Bennett AF: **Bats in rural landscapes: a significant but largely unknown faunal component.** In: Barlow T, Thorburn R, eds. *Balancing Conservation and Production in Grassy Landscapes.* Environment Australia, Biodiversity Group, Canberra, 2000; pp 42–50.
[Reference Source](#)
24. Maron M: **Agricultural change and paddock tree loss: implications for an endangered subspecies of red-tailed black-cockatoo.** *Ecological Management and Restoration.* 2005; **6**(3): 206–211.
[Publisher Full Text](#)
25. Schrott GR, With KA, King AW: **On the importance of landscape history for assessing extinction risk.** *Ecological Applications.* 2005; **15**: 493–506.
[Publisher Full Text](#)
26. Tilman D, May RM, Lehman CL, *et al.*: **Habitat destruction and the extinction debt.** *Nature.* 1994; **371**: 65–66.
[Publisher Full Text](#)
27. Murcia C: **Edge effects in fragmented forests: implications for conservation.** *Trends Ecol Evol.* 1995; **10**(2): 58–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Hobbs RJ, Huenneke LF: **Disturbance, diversity, and invasion: implications for conservation.** *Conserv Biol.* 1992; **6**(3): 324–337.
[Publisher Full Text](#)
29. Diamond JM: **The island dilemma: lessons of modern biogeographic studies for the design of natural reserves.** *Biol Conserv.* 1975; **7**(2): 129–146.
[Publisher Full Text](#)
30. Didham RK, Denmead LH, Deakin EL: **Riches to rags: the ecological consequences of land use intensification in New Zealand.** In: Lindenmayer D, Cunningham S, Young A, eds. *Land Use Intensification: Effects on Agriculture, Biodiversity and Ecological Processes.* CSIRO Publishing, Australia, 2012; pp 73–83.
[Reference Source](#)
31. Sillmann J, Roegner E: **Indices for extreme events in projections of anthropogenic climate change.** *Climatic Change.* 2008; **86**(1–2): 83–104.
[Publisher Full Text](#)
32. Smith MD: **The ecological role of climate extremes: current understanding and future prospects.** *Journal of Ecology.* 2011; **99**(3): 651–655.
[Publisher Full Text](#)
33. Walther GR, Post E, Convey P, *et al.*: **Ecological responses to recent climate change.** *Nature.* 2002; **416**(6879): 389–395.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Tylianakis JM, Didham RK, Bascompte J, *et al.*: **Global change and species interactions in terrestrial ecosystems.** *Ecol Lett.* 2008; **11**(12): 1351–1363.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Howden SM, Soussana JF, Tubiello FN, *et al.*: **Adapting agriculture to climate change.** *Proc Natl Acad Sci U S A.* 2007; **104**(50): 19691–19696.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Woodman RF: **Trifolium ambiguum. (Caucasian clover) in montane tussock grasslands, South Island, New Zealand.** *New Zealand Journal of Agricultural Research.* 1999; **42**(3): 207–222.
[Publisher Full Text](#)
37. Maron M, Fitzsimons JA: **Agricultural intensification and loss of matrix habitat over 23 years in the west Wimmera, south-eastern Australia.** *Biological Conservation.* 2007; **135**(4): 587–593.
[Publisher Full Text](#)
38. Weeks ES, Walker S, Dymond JR, *et al.*: **Patterns of past and recent conversion of indigenous grasslands in the South Island, New Zealand.** *New Zealand Journal of Ecology.* 2013; **37**(1): 127–138.
[Reference Source](#)
39. Quinn JM, Stroud MJ: **Water quality and sediment and nutrient export from New Zealand hill-land catchments of contrasting land use.** *New Zealand Journal of Marine and Freshwater Research.* 2002; **36**(2): 409–429.
[Publisher Full Text](#)
40. Lee WG, Williams P, Cameron E: **Plant invasions in urban environments: the key to limiting new weeds in New Zealand.** In: Suckling DM, Stevens PS, eds. *Managing Urban Weeds and Pests. Proceedings of a New Zealand Plant Protection Society Symposium, 9 August, 1999.* The New Zealand Plant Protection Society, Lincoln, 2000; pp 43–58.
41. Mack RN, Simberloff D, Lonsdale WM, *et al.*: **Biotic invasions: causes, epidemiology, global consequences, and control.** *Ecological Applications.* 2000; **10**(3): 689–710.
[Publisher Full Text](#)
42. Espie PR: **Hieracium in New Zealand: Ecology and Management.** AgResearch, Mosgiel, New Zealand. 2001; 66.
[Reference Source](#)
43. Lambin EF, Geist HJ, Lepers E: **Dynamics of land-use and land-cover change in tropical regions.** *Annual Review of Environment and Resources.* 2003; **28**: 205–241.
[Publisher Full Text](#)
44. Gardiner B, Reid N: **Sustainable livestock production.** In: Cottle D, ed. *International Sheep and Wool Handbook.* Nottingham University Press, Nottingham, 2010; pp 445–471.
[Reference Source](#)

45. Reid N, Norton D, Stafford Smith M: **Environmental management**. In, Cottle D, Kahn L, eds. *Beef Cattle Production and Trade*. CSIRO Publishing, Victoria, in press. 2013.
46. MacLeod CJ, Moller H: **Intensification and diversification of New Zealand agriculture since 1960: an evaluation of current indicators of land use change**. *Agriculture, Ecosystems and Environment*. 2006; **115**(1–4): 201–218.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Czúcz B, Gathman JP, McPherson GR: **The impending peak and decline of petroleum production: an underestimated challenge for conservation of ecological integrity**. *Conserv Biol*. 2010; **24**(4): 948–956.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Park G: **Nga Uruora/the Groves of Life: Ecology and History in a New Zealand Landscape**. Victoria University Press, Wellington. 1995.
[Reference Source](#)
49. O'Connor KF: **The implications of past exploitation and current developments to the conservation of South Island tussock grasslands**. *New Zealand Journal of Ecology*. 1982; **5**: 97–107.
[Reference Source](#)
50. Maclaren JP: **Environmental Effects of Planted Forests in New Zealand**. New Zealand Forest Research Institute, Rotorua. 1996.
[Reference Source](#)
51. Kadlec RH, Wallace SD: **Treatment Wetlands, second edition**. CRC Press. Boca Raton, FL 2009.
[Reference Source](#)
52. Freebairn DM, Wockner GH: **A study of soil erosion on vertisols of the eastern Darling Downs, Queensland. I. Effects of surface conditions on soil movement within contour bay catchments**. *Australian Journal of Soil Research*. 1986; **24**(2): 135–158.
[Publisher Full Text](#)
53. Moller H, MacLeod CJ, Rosin C, et al.: **Intensification of New Zealand agriculture: implications for biodiversity**. *New Zealand Journal of Agricultural Research*. 2008; **51**(3): 253–263.
[Publisher Full Text](#)
54. Huffman WE, Evenson RE: **Science for Agriculture: A Long Term Perspective**. Iowa State University Press, Ames IA. 1993.
[Reference Source](#)
55. Tscharrnke T, Batary P, Clough Y, et al.: **Combining biodiversity conservation with agricultural intensification**. In, Lindenmayer D, Cunningham S, Young A, eds. *Land Use Intensification: Effects on Agriculture, Biodiversity and Ecological Processes*. CSIRO Publishing, Australia, 2012; pp 7–15.
[Reference Source](#)
56. Peigné J, Ball BC, Roger-Estrade J, et al.: **Is conservation tillage suitable for organic farming? A review**. *Soil Use and Management*. 2007; **23**(2): 129–144.
[Publisher Full Text](#)
57. Millar GD, Badgery WB: **Pasture cropping: a new approach to integrate crop and livestock farming systems**. *Animal Production Science*. 2009; **49**(10): 777–787.
[Publisher Full Text](#)
58. Cerdeira AL, Duke SO: **The Current Status and Environmental Impacts of Glyphosate-Resistant Crops: A Review**. *J Environ Qual*. 2006; **35**(5): 1633–1658.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Owen MJ, Powles SB: **Glyphosate-resistant rigid ryegrass (*Lolium rigidum*) populations in the Western Australian grain belt**. *Weed Technology*. 2010; **24**(1): 44–49.
[Publisher Full Text](#)
60. Whitehouse MEA, Wilson LJ, Fitt GP: **A comparison of arthropod communities in transgenic Bt and conventional cotton in Australia**. *Environmental Entomology*. 2005; **34**(5): 1224–1241.
[Publisher Full Text](#)
61. Cattaneo MG, Yafuso C, Schmidt C, et al.: **Farm-scale evaluation of the impacts of transgenic cotton on biodiversity, pesticide use and yield**. *Proc Natl Acad Sci U S A*. 2006; **103**(20): 7571–7576.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Sinclair TR: **Challenges in breeding for yield increase for drought**. *Trends Plant Sci*. 2011; **16**(6): 289–293.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Eisenstein M: **Plant breeding: Discovery in a dry spell**. *Nature*. 2013; **501**(7468): S7–S9.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Robertson AI: **The gaps between ecosystem ecology and industrial agriculture**. *Ecosystems*. 2000; **3**(5): 413–418.
[Publisher Full Text](#)
65. Stiling P: **What do ecologists do?** *Bulletin of the Ecological Society of America*. 1994; **75**(2): 116–121.
[Reference Source](#)
66. Norton DA: **Is the amount and focus of ecological research in New Zealand sufficient to sustain indigenous biodiversity on private land?** *New Zealand Journal of Ecology*. 2001; **25**: 77–82.
[Reference Source](#)
67. Haggerty J, Campbell H, Morris C: **Keeping the stress off the sheep? Agricultural intensification, neoliberalism, and 'good' farming in New Zealand**. *Geoforum*. 2009; **40**(5): 767–777.
[Publisher Full Text](#)
68. Hulme PE: **Trade, transport and trouble: managing invasive species pathways in an era of globalization**. *J Appl Ecol*. 2009; **46**(1): 10–18.
[Publisher Full Text](#)
69. Wilson H: **Living in Raoul country: the changing flora and vegetation of Banks Peninsula**. In, Burrows CJ, ed. *Etienne Raoul and Canterbury Botany 1840-1996*. Canterbury Botanical Society, Christchurch, 1998; pp 101–121.
[Reference Source](#)
70. Standish RJ, Sparrow AD, Williams PA, et al.: **A state-and-transition model for the recovery of abandoned farmland in New Zealand**. In, Hobbs RJ, Suding KN, eds. *New Models for Ecosystem Dynamics and Restoration*. Island Press, Washington, 2009; pp 189–205.
[Reference Source](#)
71. Hobbs RJ, Arico S, Aronson J, et al.: **Novel ecosystems: theoretical and management aspects of the new ecological world order**. *Global Ecology and Biogeography*. 2006; **15**(1): 1–7.
[Publisher Full Text](#)
72. Seastedt TR, Hobbs RJ, Suding KN: **Management of novel ecosystems: are novel approaches required?** *Front Ecol Environ*. 2008; **6**(10): 547–553.
[Publisher Full Text](#)
73. Pannell DJ: **Public benefits, private benefits, and policy intervention for land-use change for environmental benefits**. *Land Economics*. 2008; **84**: 225–240.
[Reference Source](#)
74. Morton SR, Stafford Smith DM, Friedel MH, et al.: **The stewardship of arid Australia: ecology and landscape management**. *J Environ Manage*. 1995; **43**(3): 195–217.
[Publisher Full Text](#)
75. Salzman J: **Creating markets for ecosystem services: notes from the field**. *New York University Law Review*. 2005; **80**(6): 870–961.
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 10 February 2014

<https://doi.org/10.5256/f1000research.2650.r3305>

© 2014 Standish R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rachel Standish

School of Plant Biology, University of Western Australia, Crawley, Australia

As the title of this paper suggests, it reviews the ultimate drivers of biodiversity change in agricultural systems using examples from eastern Australia and New Zealand. The authors are well-respected for their contribution to ecological research, and particularly for informing the options for the conservation of biological diversity in these landscapes. Therefore, the review is as I'd expect it to be—authoritative and conceptually sound. In particular, Figure 1 will be a useful framework to prompt discussions among land managers and other stakeholders. An example of a decision informed by the framework would be a valuable addition to the paper.

More generally, however, the review paints a fairly pessimistic view for biodiversity conservation in agricultural landscapes and reading it left me wanting to know more about the strategies mentioned in the final sentence of the abstract. The authors deliberately reviewed the ultimate drivers of biodiversity change, because unlike proximate drivers, these can be managed—but how? For example, what ecologists might do to fill the knowledge gaps? How might the emerging carbon market be developed to benefit biodiversity and farmers in agricultural landscapes of Australia and New Zealand¹? What are some other examples of potential win-win farming enterprises? How might trade-offs among ecosystem services, including both ecological and socio-economic perspectives, affect outcomes for biodiversity in these landscapes? There is some evidence that traditional, as opposed to novel, approaches such as fencing to exclude livestock and unassisted regeneration of abandoned farmlands can help biodiversity conservation in agricultural landscapes². What are the incentives for farmers to adopt such practices?

These are my other, more minor points:

1. Historical legacies are particularly important in the ancient landscapes of south-western Australia³⁴. Land-use legacies combine with seed dispersal and recruitment limitations to form ecological thresholds to recovery^{5 6}. The socio-economic drivers of land-use change are also documented for this region⁷⁸.
2. I agree that land-use intensification generally results in the loss of native species, but what about the options for biodiversity conservation at larger scales if intensification is paired

with land-sparing? Indeed, the current global debate about land-sparing vs sharing would be relevant to mention here.

3. Nested within social values and awareness are the management goals we as a society set for biodiversity conservation. I think this section of the paper could introduce some of the novel approaches mentioned in the discussion. For example, where and when might it be appropriate to consider novel ecosystems as goals for management in place of historic ecosystems?
4. While it is probably true that agricultural landscapes are understudied compared with other landscapes in Australia as well as New Zealand, this point seems to detract from the arguably more valid point that there's decades worth of research from which to base good decisions ⁹.

Thank you for the invitation to review this paper. I enjoyed thinking about it.

References

1. Standish RJ, Hulvey KB: Co-benefits of planting species mixes in carbon projects. *Ecological Management & Restoration*. 2014; **15** (1): 26-29 [Publisher Full Text](#) | [Reference Source](#)
2. Prober SM, Standish RJ, Wiehl G: After the fence: vegetation and topsoil conditions in grazed, fenced, and benchmark eucalypt woodlands of fragmented agricultural landscapes. *Australian Journal of Botany*. 2011; **59** (4): 369-381 [Publisher Full Text](#) | [Reference Source](#)
3. Standish RJ, Cramer VA, Hobbs RJ, Kobryn HT: Legacy of land-use evident in soils of Western Australia's wheatbelt. *Plant and Soil*. 2006; **280** (1-2): 189-207 [Publisher Full Text](#) | [Reference Source](#)
4. Standish RJ, Cramer VA, Hobbs RJ: Land-use legacy and the persistence of invasive *Avena barbata* on abandoned farmland. *Journal of Applied Ecology*. 2008; **45** (6): 1576-1583 [Publisher Full Text](#) | [Reference Source](#)
5. Standish RJ, Cramer VA, Wild SL, Hobbs RJ: Seed dispersal and recruitment limitation are barriers to native recolonization of old-fields in western Australia. *Journal of Applied Ecology*. 2007; **44** (2): 435-445 [Publisher Full Text](#) | [Reference Source](#)
6. Cramer VA, Hobbs RJ, Standish RJ: What's new about old fields? Land abandonment and ecosystem assembly. *Trends Ecol Evol*. 2008; **23** (2): 104-112 [PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
7. Allison HE, Hobbs RJ: Resilience, Adaptive Capacity, and the "Lock-in Trap" of the Western Australian Agriculture Region. *Ecology and Society*. 2004; **9** (1). [Reference Source](#)
8. Smith FP: Who's planting what, where and why - and who's paying?: An analysis of farmland revegetation in the central wheatbelt of Western Australia. *Landscape and Urban Planning*. 2008; **86** (1): 66-78 [Publisher Full Text](#) | [Reference Source](#)
9. Prober SM, Smith FP: Enhancing biodiversity persistence in intensively used agricultural landscapes: A synthesis of 30 years of research in the Western Australian wheatbelt. *Agriculture, Ecosystems & Environment*. 2009; **132** (3-4): 173-191 [Reference Source](#)

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 December 2013

<https://doi.org/10.5256/f1000research.2650.r2330>

© 2013 Clarkson B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bruce Clarkson

Environmental Research Institute, University of Waikato, Hamilton, New Zealand

This is an insightful opinion piece and deserving of publication.

The introduction in particular is succinct and to the point. The key scene setting statement for me is: "As a general rule however, remnant taxa tend to be generalists and agricultural development usually results in homogenisation of native biodiversity".

I have twice in my career said as much in "Conservation at the end of the century: challenges, crises and contradictions"¹ and [Lee, Meurk & Clarkson, 2008](#) - "only a small generalist subset of the indigenous flora and fauna can persist in, or recolonise, intensified agricultural landscapes"².

The statement is critical because it begs the question is it even realistic to think that indigenous biodiversity can ever be adequately represented within essentially intensive productive agricultural lands?

My only criticisms (minor) of the paper are around areas where I would argue for a different or slightly different emphasis. We all have different backgrounds and so depending on our experiences there are likely to be questions of emphasis. I deal with these in turn below:

Page 5: left hand column last sentence regarding good financial returns; the non-essential activities are **perceived** non-essential activities, which if an ecosystems services or natural capital accounting model was in place may not be viewed as non-essential.

Page 5: right hand column last paragraph; the European settler approach was also a pragmatic one, driven (economic driver) by the products required for export to the home country. As the historian WH Oliver has succinctly noted, this agricultural land use was driven by the imperative to survive in a new land by producing trade goods demanded by colonial masters across the sea, and the land has disproportionately suffered as a result. The settlers were just responding to a market signal, and there was no other market?

Page 6: left hand column last sentence; seems that this is one place that the lack of comment policy and regulatory incentives, and the development of self-organised responses to the problem e.g. QE II Trust and Landcare groups, could have been addressed? Additionally the [Lee et al., 2008](#) paper², in response to the [Moller et al., 2008](#) article³, took a view that regulation needed more emphasis than incentive, but even that may still be insufficient to stop an extinction pathway for indigenous biodiversity under agricultural intensification?

Page 7: left column top para; it's not just new technology that enables development of previously marginal lands, but ready access and/or cheaper availability of existing technology e.g. the swamp forests on the Egmont ringplain habitat reduction by modern drain digger vehicles.

Page 7: left column last para; the claim that farmers lacked the relevant information led to an inability to realise adverse consequences of their activities seems too simplistic. What about barriers to adoption, the need for first adopters, lack of awareness, failure to observe nature, ignorance, bloody mindedness, deliberate ignoring of warnings. There are multiple reasons for a lack of realisation?

Page 7: Policy and regulation section takes a very narrow view focusing on vegetation clearance rules by themselves, but without any quantification of how clearance rules may have slowed clearance or how when coupled with incentives e.g. covenanting incentives linked to SNA scheduling to retain has had a positive impact on the Egmont ringplain.

Page 8 Conclusions: I reiterate here the lack of mention of policy and regulatory incentives and the development of self-organised responses to the problem e.g. QE II Trust and Landcare groups. Making statements like "...mechanisms are required to encourage farmers..." or "simply putting in place regulations... do not necessarily address..." or "...other approaches need to be considered" just begs a lot a questions and a single answer by offering financial incentives is not nuanced enough: what types of incentives? Environmental, Biodiversity or Carbon Credits, tax relief? Rates relief? These are positive financial incentives but what about negative financial incentives, such as trading barriers or forms of environmental accreditation etc. Is it other approaches or just a wider diversity of approaches including regulation an positive and negative financial incentives?And who is responsible for doing this - Central Government and Local Government?

And if all of this new approach is put in place, is the prognosis for native biodiversity in agricultural systems: very optimistic, reservedly optimistic or not optimistic?

A valuable contribution; thanks for the opportunity to comment.

References

1. Clarkson B: Conservation at the end of the century: challenges, crises and contradictions. In: New Zealand Botanical Society Conference. *New Zealand Botany at the end of the Millennium*. 17-19th June 1999.
2. Lee WG, Meurk CD, Clarkson BD: Agricultural intensification: whither indigenous biodiversity?. *New Zeal J Agr Res*. 2008; **51** (4): 457-460 [Publisher Full Text](#) | [Reference Source](#)
3. Moller H, MacLeod CJ, Haggerty J, Rosin C, et al.: Intensification of New Zealand agriculture: Implications for biodiversity. *New Zeal J Agr Res*. 2008; **51** (3): 253-263 [Publisher Full Text](#) | [Reference Source](#)

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research