Check for updates

SHORT RESEARCH ARTICLE

REVISED Accumulation of oocytes and/or embryos by vitrification: a new strategy for managing poor responder patients undergoing pre implantation diagnosis [version 2; peer review: 2 approved, 1 approved with reservations]

Alexia Chatziparasidou, Martine Nijs, Martha Moisidou, Oraiopoulou Chara, Christina Ioakeimidou, Christos Pappas, Nicos Christoforidis

Embryolab, Assisted Reproduction Unit, Kalamaria, Thessaloniki, 551 34, Greece

v2 First published: 12 Nov 2013, 2:240
<https://doi.org/10.12688/f1000research.2-240.v1>
 Latest published: 03 Mar 2014, 2:240
<https://doi.org/10.12688/f1000research.2-240.v2>

Abstract

Background: Low (or poor) responder patients are women who require large doses of stimulation medications and produce less than an optimal number of oocytes during IVF cycles. Low responder patients produce few oocytes and embryos, which significantly reduces their chances for success in a preimplantation genetic diagnosis (PGD) cycle. Accumulation of vitrified oocytes or embryos before the actual PGD cycle is a possible strategy that might increase patient's chances for a healthy pregnancy.

Aim of the study: This retrospective study evaluates the efficacy of a PGD program in low responder patients after repeated ovarian stimulation cycles with cumulative vitrification of oocytes and embryos.

Methods: Over a period of 30 months, 13 patients entering the PGD program were identified as poor responders after their first ovarian stimulation. These patients started a PGD cycle for one of the following indications: history of recurrent implantation failure (n=1), cystic fibrosis (n=1), X-linked microtubular myopathy (n=1), recurrent miscarriages (n=5), Duchene muscular dystrophy (n=1), chromosomal translocation (n=1) and high sperm aneuploidy (n=1). After multiple ovarian hormonal stimulations patients had either all mature oocytes (Group A; 3 patients) or all of their day 2 embryos vitrified (group B; 10 patients). Mean total number of oocyte collections per patient was 2.3 (range: 2 - 5 cycles).

Results: In the actual PGD cycle, all vitrified oocytes from group A patients were warmed and underwent intra cytoplasmic sperm injection (ICSI) followed by culture up to day 3. For group B patients all vitrified day 2 embryos were warmed and cultured overnight. On day 3 of culture, all embryos from Group A and B had blastomere biopsy

Open Peer Review

Approval Status

	1	2	3
version 2			
(revision)
03 Mar 2014	view		view

version 1
12 Nov 2013	view	view	

1. **Pedro Barri**, Institut Universitari Dexeus of the Universitat Autònoma de Barcelona, Barcelona, Spain
2. **Joep PM Geraedts**, Maastricht University, Maastricht, The Netherlands
3. **Alex Simon**, Hadassah Medical Center, Jerusalem, Israel

Any reports and responses or comments on the article can be found at the end of the article.

followed by genetic analysis. In group A, 20 embryos were found suitable for biopsy and genetic analysis; at least one healthy embryo was available for transfer for each patient. For group B, 72 embryos in total were available for biopsy and PGD. All patients, except one, had at least one healthy day 5 embryo for transfer (mean number of 2.1 embryos per transfer). Nine patients had a clinical pregnancy; 7 patients delivered a healthy baby.

Conclusion: Low responder patients entering a PGD program might increase their chances for a healthy pregnancy by repeat ovarian stimulation in combination with cumulative oocyte or embryo vitrification.

Corresponding author: Martine Nijs (martine.nijs@telenet.be)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2014 Chatziparasidou A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Chatziparasidou A, Nijs M, Moisidou M *et al.* **Accumulation of oocytes and/or embryos by vitrification: a new strategy for managing poor responder patients undergoing pre implantation diagnosis [version 2; peer review: 2 approved, 1 approved with reservations]** F1000Research 2014, 2:240 <https://doi.org/10.12688/f1000research.2-240.v2>

First published: 12 Nov 2013, 2:240 <https://doi.org/10.12688/f1000research.2-240.v1>

REVISED Amendments from Version 1

Patients recruitment criteria and patients demographic data are now included. More details on embryo biopsy outcomes have been supplied. Discussion section is more developed and discusses comparable high survival rates for oocytes and embryos and development and implantation rates for both fresh and vitrified embryos. Percentages in Table 1 are listed using decimal points.

See referee reports

Introduction

Low responder patients undergoing hormonal stimulation for an IVF or ICSI treatment have a reduced potential to produce an adequate number of oocytes and hence also embryos^{1,2}. Especially for patients seeking a healthy pregnancy through preimplantation genetic diagnosis (PGD), this low production of oocytes and embryo(s) in one cycle will significantly reduce their chances of success. Multiple consecutive ovarian stimulation cycles combined with serial vitrification of oocytes and embryos obtained before the actual PGD could be an option to increase the chances for these patients. Until now, only one successful case report has been presented by Chung *et al.*³ where a normal birth was obtained after serial vitrification of oocytes from 5 consecutive ovarian stimulation cycles for a patient carrying reciprocal translocations.

This retrospective cohort study evaluates the efficacy of a PGD program in low responder patients after repeated ovarian stimulation and accumulation of vitrified oocytes or embryos before genetic analysis, in combination with PGD on embryos obtained from a fresh ICSI cycle.

Methods

Setting and study design

This retrospective cohort study was performed over a 30 month-period (2011–2013) at Embryolab, a private fertility treatment centre in Thessaloniki, Greece.

Cycles and patients studied

During the 30 month period 13 patients of those entering the PGD program showed to be poor responders. Patients were counselled on both options (serial oocyte or embryo vitrification) with clear explanations on pro and cons of each option. Patients selected themselves for serial oocyte or embryo vitrification. PGD patients with more than 6 oocytes or 5 embryos from their first fresh PGD cycle were excluded from the study. Study patients started a PGD cycle for one of the following indications: history of recurrent implantation failure (n=1), cystic fibrosis (n=1), X-linked microtubular myopathy (n=1), recurrent miscarriages (n=5), Duchene muscular dystrophy (n=1), chromosomal translocation (n=1) and high sperm aneuploidy (n=1). Baseline characteristics for patients were mean age of 35,2 years; mean antral follicle count of 7; mean body mass index of 24,6 kg/m² and a mean FSH on day 2 of cycle: 7,43 IU.

Ovarian stimulation of patients

Patient's ovarian stimulation protocol consisted of a standard down-regulation protocol or antagonist protocol⁴. Hormonal stimulation

treatment showed these patients to be poor responders and very few oocytes could be harvested at the time of the first oocyte collection. Following counseling, couples opted for serial vitrification of oocytes (group A) or embryos (group B) from repeat ovarian stimulation cycles. Allocation to either group was based on the outcome of a medical counseling session with the patient. One to two extra hormonal stimulation cycles were initiated to obtain an accumulated minimum of 6 mature oocytes (group A) or alternatively of 5 embryos (group B) for each patient.

IVF Laboratory protocols

Oocyte collection was carried out 36 hours post-hCG administration. Fresh semen samples were prepared by density gradient centrifugation and one wash step (Quinn's Advantage Sperm Washing Medium, Sage). ICSI was performed according to standard procedures⁵. Oocytes were checked for presence of 2 pronuclei 18–22 hours post oocyte collection. Fertilised oocytes were group-cultured in 0.7 ml droplets (Cleavage medium, Sage) and embryo quality was checked daily under a microscope using a standard protocol¹⁰. Oocytes and day 2 embryos were vitrified and warmed using the methods described by Kuwayama *et al.* (Cryotop, Cryotec, http://cryotech-japan.jp/method/warming_Protocol.htm)⁶ and stored in liquid nitrogen.

Pre implantation genetic diagnosis

Embryos were biopsied on day 3 of development. Three different genetic techniques were applied, depending on the indication: fluorescent *in situ* hybridization (FISH)⁷ was used for patients suffering from X-linked microtubular myopathy, Duchene muscular dystrophy, high sperm aneuploidy or recurrent implantation failure; polymerase chain reaction (PCR)⁸ was the technique used for patients at risk for offspring with cystic fibrosis. Array complete genome hybridisation techniques (aCGH)⁹ were applied for patients at risk for recurrent miscarriage or for reciprocal translocations. Biopsied embryos were cultured individually in 50 µl droplets under oil (washed sterile oil, Sage, USA) until day 5 for transfer (Cleavage medium, Sage, USA). Embryo quality was checked daily under the microscope according to a standard protocol¹⁰.

Transfer of embryos

Embryo(s) were transferred under abdominal ultrasound guidance (Logic 400 MD) to the patient in 0.1 ml of medium (Cleavage medium, Sage) using a Wallace- (Smiths or Labotect soft catheter (Genetec). Clinical pregnancy was defined as the presence of a gestational sac with fetal heartbeat by ultrasound imaging at 8–10 weeks after embryo transfer.

Laboratory quality

The IVF laboratory at Embryolab has ISO 9001:2000 accreditation (2007) and has been assessed in accordance to ISO 15189-2007.

Given the retrospective nature and lack of identifiable health data used in the study, no institutional review board approval was needed. Patients signed an informed consent before the start of the treatment.

Results

During the 30 month study period, 13 patients were shown to be poor responders because of failure to produce a sufficient number

of oocytes or embryos to continue their PGD analysis (< 6 mature oocytes or < 5 embryos on day 2). Mean age of the patients was 35.2 years (range: 31–41 years, SD: 3.4). After medical counseling all 13 patients agreed to accumulate their oocytes or embryos by vitrification, and hence underwent repeat hormonal stimulations and oocyte collections (mean: 2.3; range 2–5 stimulations) until a sufficient number was stored (>6 mature oocytes or >5 embryos on day 2). Mean total number of oocyte collections per patient (cycles) was 2.3 (range: 2–5 cycles). Details on laboratory and clinical outcomes are listed in [Table 1](#). On day 3 of culture, a total of 92 embryos were biopsied and diagnosed genetically. In total 40 embryos were diagnosed as normal, 43 as abnormal and for 9 embryos no result was obtained. Mean number of biopsied embryos per patient was 7.2 (+SD: 2.1) and a mean average number of 2.1 embryos per patient were transferred. Eleven supernumerary embryos, diagnosed as being normal, were vitrified post-biopsy. One patient with a history of repeated failure of implantation had no healthy embryos available for transfer. This patient had a total of 5 embryos biopsied (2 from cryostorage and 3 fresh). Twelve out of 13 patients had an embryo transfer of a healthy embryo (92.3%) and 9 patients had a clinical pregnancy (75% clinical pregnancy rate in patients with embryo transfer). In total, 2 patients miscarried and 7 patients delivered a healthy baby (7/12; 58.3% delivery rate). Two twin pregnancies were noted; both patients had delivery of healthy babies. No frozen-thawed embryo transfer was done for any of the patients.

Discussion

Low responder patients undergoing IVF are characterised by a low number of oocytes retrieved because of suboptimal oocyte maturation, poor embryo quality, hormonal stimulation cycle or embryo transfer cancellation². Cobo *et al.*¹¹ demonstrated in a prospective study that accumulation of oocytes by vitrification is a successful strategy for managing low responder patients in 'classical' IVF/ICSI

treatments: delivery and cumulative delivery rates per patient were statistically higher in the low responder group (36.4%) than the low responder fresh group (23.7%). Our study could demonstrate, although on a limited number of patients, that this accumulation strategy can also be applied for a specific patient population, namely patients undergoing PGD for specific genetic diseases. Although we did not compare our outcomes to those of a control group of low responder fresh PGD patients from our center, we could demonstrate that the strategy to accumulate vitrified oocytes or embryos from consecutive hormonal stimulation cycles resulted in a sufficient number of embryos available for genetic diagnosis. As a consequence, a high percentage of patients had transfer of an embryo diagnosed to be negative for the specific genetic test (92.3%). It is evident that in order to accumulate oocytes and embryos by vitrification for the management of low responder patients, an efficient and well-established oocyte vitrification system needs to be in place. Survival rates after warming of these oocytes and embryos need to be optimal (between 80 and 100%); if this is not the case, this approach should not be offered to low responder patients. Our laboratory has high survival rates for oocytes and embryos (up to 100%) and comparable development and implantation rates for both fresh and vitrified embryos (Cryotop and Cryotec vitrification methods^{6,11}) are obtained; hereby confirming outcomes of Rienzi *et al.* (2009)¹² and Ku *et al.* (2012)¹³.

Although the treatment costs can be double or triple compared to one single hormonal stimulation for ICSI with PGD, the total costs of the accumulated cycles are lower because patients have to pay for only one ICSI procedure (in case of accumulation of oocytes) and only one genetic analysis combined with one embryo transfer.

Moreover, this accumulation strategy resulted in higher outcomes (58.3% delivery rate per transfer) as compared to the 24% delivery

Table 1. Clinical and laboratory outcomes for poor responder PGD patients after serial vitrification of oocytes or embryos.

	Group A Vitrification of Oocytes	Group B Vitrification of Embryos
Number of patients with vitrification	3	10
Number of cycles with vitrification	6	18
Total number oocytes/embryos vitrified from repeat cycles	15	44
Survival after warming number (%)	15 (100%)	44 (100%)
Number of oocytes/embryos obtained in ultimate fresh cycle	22	28
Total number of embryos available for PGD on day 3	20	72
Number of patients with transfer of at least 1 healthy embryo	3	9
Mean number of embryos per transfer	2.1	
Number of patients with positive hCG test	9/12 (75.0%)	
Number of patients with healthy delivery	7/12 (58.3%)	

rate per fresh embryo transfer presented by the ESHRE PGD consortium for 2008¹⁴.

This retrospective cohort study demonstrates, although on a limited number of patients, that low responder patients in need of PGD can benefit from serial vitrification of oocytes and/or embryos after repeated ovarian stimulation cycles to improve their chances of a successful pregnancy. Future studies should address the ideal number of vitrified oocytes and/or embryos necessary in order to increase success in low responder patients undergoing PGD.

Author contributions

AC, NC, IC and CP conceived the study. AC, MN and NC designed the research. AC, MM and OC carried out the research. MM and OC provided expertise in vitrification. MN and AC prepared the first draft of the manuscript. NC and IC contributed to the preparation

of the manuscript. All authors were involved in the revision of the draft manuscript and have agreed to the final content.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgements

These data were partially presented in a poster at the 12th International Conference on Preimplantation Genetic Diagnosis, Istanbul, Turkey, 2013.

We are grateful to the staff of Embryolab and EUROGENETICA SA Genetic Laboratories for assisting in all daily aspects of the treatment of these patients.

References

1. Surrey ES, Schoolcraft WB: **Evaluating strategies for improving ovarian response of the poor responder undergoing assisted reproductive techniques.** *Fertil Steril.* 2000; **73**(4): 667–676.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Karande V, Gleicher N: **A rational approach to the management of low responders in in-vitro fertilization.** *Hum Reprod.* 1999; **14**(7): 1744–1748.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Chung JT, Son WY, Zhang XY, *et al.*: **Normal birth following PGD for reciprocal translocation after serial vitrification of oocytes from a poor responder: a case report.** *RBMonline.* 2012; **25**(5): 521–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Olivennes F, Cunha-Filho J, Fanchin R, *et al.*: **The use of GnRH antagonists in ovarian stimulation.** *Hum Reprod Update.* 2002; **8**(3): 279–290.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Palermo G, Joris H, Devroey P, *et al.*: **Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte.** *Lancet.* 1992; **2**(8810): 17–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Kuwayama M, Vajta G, Kato O, *et al.*: **Highly efficient vitrification method for cryopreservation of human oocytes.** *RBMonline.* 2005; **11**(3): 300–308.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Price CM: **Fluorescence in situ hybridization.** *Blood Rev.* 1993; **7**(2): 127–134.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Beverly C, John PL, John JP, *et al.*: **Polymerase Chain Reaction.** *Methods Mol Biol.* 1993; **15**: 1–29.
[Publisher Full Text](#)
9. Wells D, Alfarawati S, Fragouli E: **Use of comprehensive chromosomal screening for embryo assessment: microarrays and CGH.** *Molec Hum Reprod.* 2008; **14**(12): 703–710.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. **Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group of Embryology.** The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum Reprod.* 2011; **26**(6): 1270–1283.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Cobo A, Garrido N, Crespo J, *et al.*: **Accumulation of oocytes: a new strategy for managing low-responder patients.** *RBMonline.* 2012; **24**(4): 424–432.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Rienzi L, Romano S, Albricci L, *et al.*: **Embryo development of fresh 'versus' vitrified metaphase II oocytes after ICSI: a prospective randomized sibling-oocyte study.** *Hum Reprod.* 2010; **25**(1): 66–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Ku PY, Lee RK, Lin SY, *et al.*: **Comparison of the clinical outcomes between fresh blastocyst and vitrified-thawed blastocyst transfer.** *J Assist Reprod Genet.* 2012; **29**(12): 1353–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Goossens V, Traeger-Synodinos J, Coonen E, *et al.*: **ESHRE PGD Consortium data collection XI cycles from January to December 2008 with pregnancy follow-up to October 2009.** *Hum Reprod.* 2012; **27**(7): 1887–1911.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 03 July 2014

<https://doi.org/10.5256/f1000research.3961.r3937>

© 2014 Barri P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pedro Barri

Department of Obstetrics, Institut Universitari Dexeus of the Universitat Autònoma de Barcelona, Barcelona, Spain

I have carefully read the new version of the article "Accumulation of oocytes and/or embryos by vitrification: a new strategy for managing poor responder patients undergoing pre implantation diagnosis".

I thank the authors for having implemented the suggested changes. I feel comfortable with the article like it is now.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 19 May 2014

<https://doi.org/10.5256/f1000research.3961.r4796>

© 2014 Simon A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alex Simon

Department of Obstetrics and Gynecology, Hadassah Medical Center, Jerusalem, Israel

In the current study the authors suggest to accumulate either embryos or oocytes for PGD/PGS in patients who are low responders. This method was found to be successful in its outcome as well

as money saving. Although patients have to repeat ovum pick-up cycles and vitrify either oocytes or embryos, the PGD/PGS process and embryo transfer is performed only once. The revised manuscript has been corrected according to the referees' suggestions and I found it suitable now for publication. Although this is a retrospective study with heterogeneous patients but still small number of patients, the concept that raised by the author is worth presentation.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 30 January 2014

<https://doi.org/10.5256/f1000research.2565.r3264>

© 2014 PM Geraedts J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Joep PM Geraedts

Department of Genetics and Cell Biology, Maastricht University, Maastricht, The Netherlands

The ability to accumulate oocytes or embryos from multiple cycles before PGD is done, could be an interesting development as is suggested in this article. However, the numbers are too small to be conclusive.

I agree with the remarks of the first referee, furthermore I would like to add the following points for revision:

Table 1:

- The numbers of normal and abnormal embryos should be specified.
- What are the definitions of clinical pregnancy rate per patient with transfer and the pregnancy rate per patient with transfer?
- How many embryos were frozen?
- How many FETs (Frozen Embryo Transfers) were done?
- All percentages should be given using decimal points.

In the discussion it is stated that a high percentage of patients had an embryo transfer of healthy embryos. In my opinion this cannot be concluded from the material presented in the manuscript. In a number of cases FISH was used on day 3 embryos, which means that not PGS, not PGD was done. This screening can only give the results for the chromosomes included in the FISH analysis, while all other chromosomes can still be aneuploid. Furthermore it is known that mosaicism can complicate analysis at day 3.

Finally the information given with respect to the costs is not detailed enough. The prices of an ICSI cycle and the PGD analysis should be given in order to come to conclusions about the financial aspects of this approach.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 10 Feb 2014

Martine Nijs, Assisted Reproduction Unit, Kalamaria, Thessaloniki, Greece

We thank Prof. Dr. Geraedts for his questions, comments and suggestions on our manuscript. We have listed our replies below. They will be implemented in in the final version of the manuscript.

- a. In total 40 embryos were diagnosed as normal, 43 as abnormal and for 9 embryos no result was obtained.
- b. Clinical pregnancy is defined as the presence of a gestational sac with foetal heartbeat by ultrasound at 8-10 weeks after embryo transfer.
- c. Eleven supernumerary embryos, diagnosed as being normal, were frozen post-embryo transfer.
- d. No Frozen embryo transfer was done for these patients.
- e. All percentages will be listed using decimal points.

The discussion will be rephrased and will state the following: 'As a consequence, a high percentage of patients had transfer of an embryo diagnosed to be negative for the specific genetic test (92,3%).'

Our cost calculation of the treatments was based on the following:

- Traditional strategy involves repeat hormonal stimulation, repeat oocyte collection, repeat ICSI and repeat genetic diagnosis tests, each step associated with low chance for embryo transfer because of low number of oocytes collected.
- Cost of the accumulation strategy includes repeat hormonal stimulation, repeat oocyte collection, repeat vitrification and storage, one warming cycle, one ICSI and one genetic diagnosis test with very good chance for an embryo transfer.

Competing Interests: I am the co-author of the paper and have no competing interest.

Reviewer Report 20 November 2013

<https://doi.org/10.5256/f1000research.2565.r2412>

© 2013 Barri P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pedro Barri

Department of Obstetrics, Institut Universitari Dexeus of the Universitat Autònoma de Barcelona, Barcelona, Spain

This is an interesting paper that addresses an important issue, but the sample size is small (13 patients) and there is relevant information lacking. I believe that this article should have the following recommendations taken into consideration:

1. Provide information concerning patients' baseline characteristics (as well as age, AFC and hormonal profile).
2. Could the authors specify how allocation was done? (Please explain "based on the outcome of a medical counselling session with the patient".)
3. In Methods, Ovarian stimulation of patients, in the phrase "to obtain an accumulated minimum of 6 oocytes" the word "mature" should be added after oocytes.
4. How was the cut-off of 6 mature oocytes established? Taking into account the average rates of fertilization and development to D2 embryos, isn't this cut-off too low? (It is true that with this cut-off the obtained results are good but the sample size is small...)
5. In Results, the phrase "until a sufficient number was stored (< 6 mature oocytes or < 5 embryos on day 2)": the < should be a >.
6. One patient did not have any healthy embryos for transfer: could you please explain how many embryos were biopsied in this patient?
7. The total number of embryos available for PGD is given for each group, but could you provide the mean number of biopsied embryos $\pm$ SD per patient?
8. If multiple pregnancy rate is zero it should be specified better, if it is not zero the rate should be given.
9. I agree with the conclusion of the study but, in order to firmly state that vitrification for accumulation purposes in PGD cycles increases the chances of success, the next 2 points should be taken into account:
 - Was the rate of development to day 3 embryos the same in fresh and vitrified and warmed cycles?
 - It could be interesting to analyze the euploidy rate between embryos coming from fresh oocytes vs. vitrified + warmed oocytes; the same for fresh vs. vitrified + warmed embryos.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 09 Dec 2013

Martine Nijs, Assisted Reproduction Unit, Kalamaria, Thessaloniki, Greece

We would like to thank Dr. Pedro Barri for his useful comments concerning our study on the evaluation of the efficacy of a PGD program in low responder patients after repeated ovarian stimulation and accumulation of vitrified oocytes or embryos. Indeed our sample size was small, but still indicative for the usefulness of vitrification as a tool for poor responders in a PGD/PGS program. Hence we opted to describe our observations in a 'short' research article.

1. The following baseline characteristics will be included in the revised article: Mean Age: 35.2 years; Mean AFC:7; Mean BMI: 24.6; Mean D2 FSH: 7.43.
2. This retrospective cohort study was performed over a 30 month-period (2011–2013). Patients were counseled on both options (serial oocyte or embryo vitrification) with clear explanations on the pro and cons of each option. Patients selected themselves for serial oocyte or embryo vitrification.
3. 'Mature' will be added to the specific sentence.
4. Our study population consisted of patients with a low number of eggs retrieved and embryos produced. Our cut off was the minimum number that was possible to be obtained by this patient population. A higher cut off would require additional stimulation cycles which was not an option, as it required more repetitive cycles and a higher treatment cost.
5. The < will be changed to >.
6. This patient had 2 embryos from her first cycle and decided to vitrify them in order to proceed to another stimulated cycle. The aim was to increase the number of the available embryos for biopsy, and hence increase the number of having at least one healthy embryo to transfer after the screening. The next stimulation cycle resulted in 3 fresh embryos. In total 5 embryos were biopsied (2 thawed and 3 fresh). None of the embryos tested was genetically healthy and the transfer was cancelled.
7. The mean number of biopsied embryos per patient: 7.2 (+SD: 2.1).
8. Out of the 9 pregnancies obtained, two twin pregnancies were noted; both patients had delivery of healthy babies.
9. According to our in house data there is no difference in development or implantation

rate of fresh versus vitrified embryos, hereby confirming results of [Rienzi *et al.* \(2009\)](#) and [Ku *et al.* \(2012\)](#). This observation can be included in the discussion part. Unfortunately, we do not have in house data on the euploidy status of vitrified oocytes. [Forman *et al.* \(2012\)](#) however, did not observe an increase in aneuploidy rates after vitrification and warming of embryos.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research