Check for updates

RESEARCH NOTE

Understanding the association between chromosomally integrated human herpesvirus 6 and HIV disease: a cross-sectional study [version 1; peer review: 2 approved, 1 approved with reservations]

Mundeep K. Kainth¹, Susan G. Fisher^{2,3}, Diana Fernandez², Amneris Luque⁴, Caroline B. Hall^{1,3,4+}, Anh Thi Hoang¹, Anisha Lashkari¹, Alexandra Peck¹, Lubaba Hasan¹, Mary T. Caserta¹

¹Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Rochester Medical Center, Rochester, NY, USA

²Department of Public Health Sciences, University of Rochester Medical Center, Rochester, NY, USA

³Department of Pharmacy, Temple University School of Medicine, Philadelphia, PA, USA

⁴Division of Infectious Diseases, Department of Medicine, University of Rochester Medical Center, Rochester, NY, USA

+ Deceased author

V1 First published: 09 Dec 2013, 2:269
<https://doi.org/10.12688/f1000research.2-269.v1>
 Latest published: 01 Jun 2017, 2:269
<https://doi.org/10.12688/f1000research.2-269.v2>

Abstract

We conducted a cross-sectional investigation to identify evidence of a potential modifying effect of chromosomally integrated human herpes virus 6 (ciHHV-6) on human immunodeficiency virus (HIV) disease progression and/or severity. ciHHV-6 was identified by detecting HHV-6 DNA in hair follicle specimens of 439 subjects. There was no statistically significant relationship between the presence of ciHHV-6 and HIV disease progression to acquired immunodeficiency syndrome. However, after adjusting for use of antiretroviral therapy, all subjects with ciHHV-6 had low severity HIV disease; these findings were not statistically significant. A multi-center study with a larger sample size will be needed to more precisely determine if there is an association between ciHHV-6 and low HIV disease severity.

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 01 Jun 2017			
version 1 09 Dec 2013	view	view	view
1. Susana N. Asin , Dartmouth Medical School, Hanover, NH, USA 2. Mario Clerici , University of Milano, Milano, Italy 3. David Camerini , University of California, Irvine, CA, USA			
Any reports and responses or comments on the article can be found at the end of the article.			

Corresponding author: Mundeep K. Kainth (Mundeep_Kainth@urmc.rochester.edu)

Competing interests: Mary T. Caserta has received funding from the HHV-6 Foundation for unrelated projects. No other competing interests were disclosed.

Grant information: Mundeep K. Kainth received a Strong Children's Research Center Bradford Fellowship and a Developmental Center for AIDS Research pilot award. She was supported by NIH/NIAID 2 T32 AI 007464-16 during her fellowship training.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2013 Kainth MK *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Kainth MK, Fisher SG, Fernandez D *et al.* **Understanding the association between chromosomally integrated human herpesvirus 6 and HIV disease: a cross-sectional study [version 1; peer review: 2 approved, 1 approved with reservations]** F1000Research 2013, 2:269 <https://doi.org/10.12688/f1000research.2-269.v1>

First published: 09 Dec 2013, 2:269 <https://doi.org/10.12688/f1000research.2-269.v1>

Editorial note:

30th May 2017. Dataset 1 has been removed from this article, as it was found to contain identifying information. A new version of this article has been published with an updated version of this dataset, with the identifying data removed. The article type has also been updated to "Research Note" to reflect a change in our article type naming conventions.

Introduction

The relationship between chromosomally integrated human herpesvirus 6 (ciHHV-6) and the progression of human immunodeficiency virus (HIV) infection in humans has, to date, never been examined.

HHV-6 causes ubiquitous human infection with approximately 100% of the population seropositive by 3 years of age. The virus infects lymphocytes with a predominant tropism for CD4+ T cells¹. In individuals with ciHHV-6, the entire viral genome is present in every nucleated cell of the body and transmitted in a Mendelian fashion^{2,3}. The presence of HHV-6 DNA in hair follicle DNA specimens is a marker for ciHHV-6⁴. Published data have demonstrated the prevalence of ciHHV-6 to be approximately 0.85%⁵. If ciHHV-6 is associated with HIV infection, then this could uncover new possibilities for understanding HIV disease severity and progression.

Numerous studies have attempted to unravel the relationship between HHV-6 and HIV. Initially, the identification of HHV-6 (not ciHHV-6) DNA in 100% of saliva samples from HIV-infected adults with high CD4+ T-cell counts suggested a protective effect of HHV-6 on the progression of HIV^{6,7}. HHV-6 produces a functional chemokine, U83A, which binds C-C chemokine receptor 5 (CCR5) and blocks HIV infection of susceptible cells *in vitro* providing a plausible mechanism for HHV-6 inhibition of HIV⁸.

Alternatively, *in vitro* activation of the HIV long terminal repeat (LTR) sequences by HHV-6 has been described suggesting that HHV-6 may promote HIV disease progression⁹. More recent data from animal models have linked infection with HHV-6 and the production of RANTES, the ligand for CCR5, with the acceleration of HIV disease via increased HIV virulence^{10,11}. Clinical studies have found reactivation or persistence of HHV-6 antigen and IgM antibody in patients with progression from asymptomatic HIV to AIDS¹².

After acquiring HIV infection, approximately 80% of adults gradually develop low CD4+ T-cell counts over 5–10 years in the peripheral blood with a high HIV viral burden, and are otherwise referred to as 'typical progressors' (TP). A small proportion of individuals develop low CD4+ T-cell counts and high viral load within two years of initial infection; these individuals are identified as 'rapid progressors' (RP). The mechanisms surrounding these differences are unclear^{13,14}. Additionally, despite prior investigations, there is also no clear understanding of the role of HHV-6 in the pathophysiology of HIV disease.

We conducted an exploratory investigation to estimate the prevalence of ciHHV-6 in an HIV-infected population and to determine

if there is a potential modifying effect of ciHHV-6 on HIV disease progression or severity.

Materials and methods**Study design**

This cross-sectional study included patients seen at the Infectious Diseases Clinic at Strong Memorial Hospital in Rochester, NY. The inclusion criterion limited the population to subjects with HIV infection over the age of 18. The exclusion criteria were lack of visible hair and inability to consent. Enrollers were available four days a week, 3–6 hours a day to approach eligible subjects.

Consent and ethical considerations

Written informed consent for publication of their clinical details and/or clinical images was obtained from the patient/parent/guardian/ relative of the patient. This study protocol was reviewed and approved by the Research Subject Review Board at the University of Rochester (RSRB00032054).

Data collection

Approximately one-half of the patients' clinical data was obtained through the HIV Data Registry supported by the Developmental Center for AIDS Research (D-CFAR). The remainder of the patient data were obtained from electronic medical records with the use of a standardized chart abstraction tool (see data file) that included HIV diagnoses dates, date of AIDS defining illnesses, and viral load/CD4+ T cell counts. Demographic data such as race/ethnicity and sex/gender were provided by subject self identification and included in grant agency reports. Hair follicles were collected by enrollers by pulling 2–3 strands of scalp hair with gloved hands.

Based on established standards, the definition of RP HIV included patients with CD4+ T cell counts of less than 200 or an AIDS-defining illness within 2 years of diagnosis of HIV infection. Subjects who had confirmed HIV infection did not meet criteria for RP were placed in a TP group. Diagnosis of HIV infection was identified by a positive ELISA/Western blot test or indicated by a provider in chart documentation based on patient history. Testing for HIV during the study period was performed by antibody testing (Vitros ECI) at Strong Memorial Hospital. Confirmation by Western blot (INNO-LIA HIV I/II Score) was performed at the Mayo Clinic in Rochester, MN.

We also compared HIV disease severity in ciHHV-6 positive and negative patients because initial disease progression from HIV to AIDS may not necessarily correlate with current disease severity in subjects on antiretroviral treatment (ART). Subjects were considered to have severe disease if CD4+ T cell counts were less than 200 cells/uL and viral load was greater than 10,000 copies/mL. If CD4+ T cell counts were greater than 200 cells/uL, and viral load was less than 10,000 copies/mL, subjects were considered to have non-severe disease. All subjects with any remaining combination of viral load and CD4+ T cell counts were considered to have 'moderately' severe disease.

HHV-6 testing

Hair samples were chosen to identify HIV positive subjects with ciHHV-6 and to exclude subjects with acquired HHV-6 infection.

Hair follicle samples were digested with proteinase K followed by DNA extraction using QIAamp DNA mini kit (Valencia, CA). DNA samples were tested by a nested qualitative PCR amplifying the HHV-6 U38 DNA polymerase gene as previously described¹⁵. In order to verify the presence of ciHHV-6, qualitative PCR was repeated on all of the positive samples followed by quantitative real time PCR for the HHV-6 U4 gene as described by Zhen *et al.*¹⁶ with modifications developed in our laboratory¹⁷.

Data analysis

All data was analyzed with SAS (v9.2). Prevalence rates of ciHHV-6 were compared between the entire cohort of HIV-infected subjects and the prevalence of ciHHV-6 in the general population⁵. Next, the prevalence of ciHHV-6 was compared between RP and TP subjects to determine if there was a significant difference of ciHHV-6 between these two groups. We also determined whether the presence of ciHHV-6 was associated with markers of HIV disease severity ascertained at the time of enrollment.

Student's t-test was used to compare age and the chi-square test for gender, ethnicity, and race. An analysis of subjects positive for ciHHV-6 and markers of HIV disease progression was performed comparing prevalence rates between the RP and TP groups and the three severity groups using Fisher's exact test. An association was considered to be statistically significant at $p < 0.05$.

Results

Cohort

The clinic population available for this study included 1035 HIV-infected patients, 714 men and 321 women between 18 and 79 years of age (African American – 418 (40%); White – 453 (44%); Hispanic – 155 (7%); other – 9 (<0.01%)). Approximately 115 patients per week were invited to join this study during routine clinic visits from October 2010 to April 2012 and 463 subjects were enrolled. From this group, 1 subject withdrew and 9 were erroneously enrolled twice. 4 subjects with unobtainable hair follicle DNA were excluded. Clinical data on 215 subjects were obtained from the University of Rochester D-CFAR database. For the remaining 248 subjects, electronic medical records were reviewed. When discrepancies between HIV and AIDS diagnosis dates occurred, the earliest date identified on the chart was selected. HIV diagnosis date was unavailable for 10 subjects, with a final number of 439 subjects with complete data available for analysis.

Approximately 1/3 of the total cohort of subjects identified themselves as female, 2/3 male, and less than 1% were transgender. Reported racial distributions for enrolled subjects were: White 205 (47%), Black-African American 171 (39%), White-Black 5 (1%), unknown or other 54 (12.3%), Asian 1, and Hawaiian Pacific-Islander 1 (Table 1). Fifty-eight (13.2%) subjects reported their ethnicity as Hispanic. Age ranged from 18 to 74 years with a mean age of 46.7 years and a median of 48 years.

ciHHV-6 results

Eight ciHHV-6 positive samples out of a total of 439 hair follicle samples were identified by initial qualitative PCR testing. Upon

re-testing, only three samples were positive. Real-time quantitative PCR analysis verified 4 positive samples (Table 2). Due to the higher specificity of the assay, the quantitative PCR data were used in the remaining study analyses.

Prevalence of ciHHV-6 in study population

The calculated prevalence estimate of ciHHV-6 in the cohort was 0.91% with a 95% CI of [0.37–2.31%]. Of the four positive ciHHV-6 subjects, 2 were female and 2 were male; 2 were white and 2 were black.

Table 1. Demographic data by presence or absence of CI-HHV 6^a detected by quantitative PCR.

	Positive		Negative		Total	
Gender	n	%	n	%	n	%
Male	2	0.5	287	65.4	289	65.8
Female	2	0.5	145	33.0	147	33.5
Transgender	0	0	3	0.7	3	0.7
Race	n	%	n	%	n	%
Asian	0	0	1	0.2	1	0.2
Black/AA ^b	2	0.5	169	38.5	171	39.0
Other	0	0	8	1.8	8	1.8
Unknown	0	0	48	11.0	48	11.0
White	2	0.5	203	46.2	205	46.7
White-Black	0	0	5	1.1	5	1.1
White-Haw/PI ^c	0	0	1	0.2	1	0.2
Ethnicity	n	%	n	%	n	%
Hispanic	0	0	58	13.2	58	13.2
Non-Hispanic	4	0.9	347	79.0	351	80.0
Unknown	0	0	30	6.8	30	6.8

^aCI-HHV6 (Chromosomally Integrated HHV6)

^bAA (African-American)

^cHaw/PI (Hawaiian/Pacific-Islander)

Table 2. Quantitative PCR results from hair follicle specimens for ciHHV-6^a positive subjects.

Patient ID number	Log ₁₀ gec per µg of DNA ^b
165	7.93
254	5.61
258	8.05
369	7.87

^aciHHV-6 (chromosomally integrated human herpes virus 6)

^bValues >5 Log₁₀ genome equivalent copies (gec) per µg of DNA are considered to be positive for high viral loads of HHV-6 DNA indicating chromosomal integration

HIV disease progression and ciHHV-6

Of 340 (77%) TP subjects, 3 (0.68%) were positive for ciHHV-6. Ninety-nine (23%) subjects were classified as having RP HIV disease based on the time to AIDS diagnosis after the date of HIV infection. Out of these 99 subjects, one (0.23%) was positive for ciHHV-6. Due to the low ciHHV-6 prevalence rate in the study population, the difference between ciHHV-6 among RP and TP is not statistically significant ($p < 0.42$).

HIV disease severity and ciHHV-6

Patients were categorized into three severity groups based on their CD4+ T cell count and HIV viral loads as described above. In order to control for antiretroviral therapy (ART) use, 9 individuals who were ART naïve were excluded. 332 subjects (84%) had non-severe disease, 42 (11%) had moderately severe disease, and 19 (5%) had severe disease. All of the subjects with ciHHV-6 had non-severe HIV disease. Given the small numbers, this finding did not reach statistical significance ($p < 0.51$).

Data on the presence of chromosomally integrated human herpes virus 6 (ciHHV-6) and human immunodeficiency virus (HIV) disease progression and severity

2 Data Files

<http://dx.doi.org/10.6084/m9.figshare.863785>

Discussion

This pilot study is the first to identify the presence of ciHHV-6 in an HIV-infected population. Because ciHHV-6 is inherited in a Mendelian fashion, our data confirmed that the prevalence rate of ciHHV-6 in our total HIV-infected cohort was similar to the published prevalence rate in otherwise healthy populations. We hypothesized that ciHHV-6 would be associated with rapid HIV disease progression or markers of disease severity. However, we did not identify a significant association between HIV disease progression and ciHHV-6 status. All four subjects with ciHHV-6 had non-severe disease; yet this was also not significant due to the small numbers of individuals with ciHHV-6 identified.

The strengths of the study include that a large number of HIV-infected subjects were available for study and the ease of hair sample collection. The testing was appealing to subjects and allowed efficient sample storage. The major limitation of this study was the low prevalence of ciHHV-6 in the HIV-infected cohort. Additionally, we were not always able to identify dates for HIV infection and AIDS

diagnosis to establish clinical disease progression. We attempted to overcome differential misclassification by evaluating CD4+ T cell counts and viral loads at the time of sample collection to establish a subject's HIV disease severity. This last method revealed that all 4 subjects with ciHHV-6 were in the low disease severity group, but due to the low ciHHV-6 prevalence, we were unable to conduct further statistical analyses.

While earlier *in vitro* studies have suggested a protective effect of HHV-6 infection on HIV disease, our study is the first clinical investigation to identify a possible protective effect of ciHHV-6 on HIV disease severity. However, due to a lack of statistical significance of the association between ciHHV-6 and HIV disease severity, more data will need to be collected to assess this relationship.

Author contributions

MTC conceived the study. CBH designed the experiments. MKK, ATH, AL, AP and LH carried out the research. SGF and IDF contributed to the design of the study and provided expertise in statistics. MKK, SGF, IDF and MTC prepared the first draft of the manuscript. All surviving authors were involved in the revision of the draft manuscript and have agreed to the final content.

Competing interests

Mary T. Caserta has received funding from the HHV-6 Foundation for unrelated projects. No other competing interests were disclosed.

Grant information

Mundee K. Kainth received a Strong Children's Research Center Bradford Fellowship and a Developmental Center for AIDS Research pilot award. She was supported by NIH/NIAID 2 T32 AI 007464-16 during her fellowship training.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We are grateful for the assistance of the staff at the Strong Memorial Hospital AIDS Center, including David Clinton, Phyllis Mulvaney and Linda Plano, for their enthusiastic recruitment efforts as well as the research staff at the AIDS Clinical Trials Unit, especially Carol Greisberger and Nurhan Calisir. In memoriam, we dedicate this paper to Caroline Breese Hall, MD, our mentor, educator and friend.

References

1. Salahuddin SZ, Ablashi DV, Markham PD, *et al.*: **Isolation of a new virus, HBLV, in patients with lymphoproliferative disorders.** *Science*. 1986; 234(4776): 596–601. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Tanaka-Taya K, Sashihara J, Kurahashi H, *et al.*: **Human herpesvirus 6 (HHV-6) is transmitted from parent to child in an integrated form and characterization of cases with chromosomally integrated HHV-6 DNA.** *J Med Virol*. 2004; 73(3): 465–473. [PubMed Abstract](#) | [Publisher Full Text](#)
3. Ward KN, Thiruchelvam AD, Couto-Parada X: **Unexpected occasional persistence of high levels of HHV-6 DNA in sera: detection of variants A and B.** *J Med Virol*. 2005; 76(4): 563–570. [PubMed Abstract](#) | [Publisher Full Text](#)
4. Ward KN, Leong HN, Nacheva EP, *et al.*: **Human herpesvirus 6 chromosomal integration in immunocompetent patients results in high levels of viral DNA in blood, sera, and hair follicles.** *J Clin Microbiol*. 2006; 44(4): 1571–1574. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

5. Pellett PE, Ablashi DV, Ambros PF, *et al.*: **Chromosomally integrated human herpesvirus 6: questions and answers.** *Rev Med Virol.* 2012; **22**(3): 144–155.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Fairfax MR, Schacker T, Cone RW, *et al.*: **Human herpesvirus 6 DNA in blood cells of human immunodeficiency virus-infected men: correlation of high levels with high CD4 cell counts.** *J Infect Dis.* 1994; **169**(6): 1342–1345.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Fabio G, Knight SN, Kidd IM, *et al.*: **Prospective study of human herpesvirus 6, human herpesvirus 7, and cytomegalovirus infections in human immunodeficiency virus-positive patients.** *J Clin Microbiol.* 1997; **35**(10): 2657–2659.
[PubMed Abstract](#) | [Free Full Text](#)
8. Catusse J, Parry CM, Dewin DR, *et al.*: **Inhibition of HIV-1 infection by viral chemokine U83A via high-affinity CCR5 interactions that block human chemokine-induced leukocyte chemotaxis and receptor internalization.** *Blood.* 2007; **109**(9): 3633–3639.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Ensoli B, Lusso P, Schachter F, *et al.*: **Human herpes virus-6 increases HIV-1 expression in co-infected T cells via nuclear factors binding to the HIV-1 enhancer.** *EMBO J.* 1989; **8**(10): 3019–3027.
[PubMed Abstract](#) | [Free Full Text](#)
10. Lusso P, Crowley RW, Malnati MS, *et al.*: **Human herpesvirus 6A accelerates AIDS progression in macaques.** *Proc Natl Acad Sci U S A.* 2007; **104**(12): 5067–5072.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Biancotto A, Grivel JC, Lisco A, *et al.*: **Evolution of SIV toward RANTES resistance in macaques rapidly progressing to AIDS upon coinfection with HHV-6A.** *Retrovirology.* 2009; **6**: 61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Ablashi DV, Marsh S, Kaplan M, *et al.*: **HHV-6 infection in HIV-infected asymptomatic and AIDS patients.** *Intervirology.* 1998; **41**(1): 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Fontaine J, Coutlée F, Tremblay C, *et al.*: **HIV infection affects blood myeloid dendritic cells after successful therapy and despite nonprogressing clinical disease.** *J Infect Dis.* 2009; **199**(7): 1007–1018.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Ling B, Veazey RS, Hart M, *et al.*: **Early restoration of mucosal CD4 memory CCR5 T cells in the gut of SIV-infected rhesus predicts long term non-progression.** *AIDS.* 2007; **21**(18): 2377–2385.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Hall CB, Long CE, Schnabel KC, *et al.*: **Human herpesvirus-6 infection in children. A prospective study of complications and reactivation.** *N Engl J Med.* 1994; **331**(7): 432–438.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Zhen Z, Bradel-Tretheway B, Sumagin S, *et al.*: **The human herpesvirus 6 G protein-coupled receptor homolog U51 positively regulates virus replication and enhances cell-cell fusion *in vitro*.** *J Virol.* 2005; **79**(18): 11914–11924.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Caserta MT, Hall CB, Schnabel K, *et al.*: **Human herpesvirus (HHV)-6 and HHV-7 infections in pregnant women.** *J Infect Dis.* 2007; **196**(9): 1296–1303.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 09 January 2014

<https://doi.org/10.5256/f1000research.2563.r2695>

© 2014 Camerini D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

David Camerini

Department of Molecular Biology and Biochemistry, University of California, Irvine, CA, USA

This interesting, well-conducted study investigated the relationship between chromosomally integrated human herpes virus six (ciHHV-6) on HIV disease severity and progression. Individuals with ciHHV-6 constitute just under 1% of the general population and a similar proportion of the 439 HIV positive individuals analyzed in this study. Due to the small sample size, only 4 ciHHV-1, HIV+ individuals were found and no significant conclusions could be drawn regarding any association with HIV disease. Nevertheless, this study is thought provoking and may provide impetus for a larger study that could shed light on this potential viral interaction *in vivo*.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 07 January 2014

<https://doi.org/10.5256/f1000research.2563.r3026>

© 2014 Clerici M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mario Clerici

Department of Biomedical Sciences and Technology, University of Milano, Milano, 20157, Italy

Because of the very low ciHHV-6 positivity in the analyzed HIV-infected population, the author's suggestion that the presence of ciHHV-6 associates with HIV disease severity seems to be too strong and not (yet) justified.

In particular, the abstract is misleading: the authors should add the number of ciHHV-6 positive subjects, and delete/modify the sentence about the severity of disease and ciHHV-6.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 07 January 2014

<https://doi.org/10.5256/f1000research.2563.r2694>

© 2014 Asin S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Susana N. Asin

Department of Microbiology and Immunology, Dartmouth Medical School, Hanover, NH, USA

This study addresses an interesting question: the potential impact of chromosomally integrated Human Herpes Virus 6 (HHV6) on HIV-1 disease progression. The rationale underlying the proposed hypothesis is weak. The authors do not suggest potential mechanisms underlying their hypothesis, which may provide a more provoking discussion. It is appropriate to anticipate that immunosuppression will lead to HHV6 reactivation. Since HHV6 shares tropism for CD4⁺ T cells with HIV-1, HHV6 could accelerate disease progression by impairing the immune response to HIV-1. Based on observation demonstrating that HHV6 replication accelerates progression to AIDS in macaques, I am curious to understand why the authors hypothesize that chromosomally integrated HHV6 could impact HIV-1 disease progression specially without evaluating HHV6 replication/reactivation.

It would have been appropriate to define how the authors evaluated chromosomally integrated DNA. Are the primers used specific for HHV6 or do these primers cross react with other HHV family members? Only describing the target gene will allow the reader to define the potential confounding effects of additional endogenous herpes viruses.

The manuscript is well written and the authors do not draw overambitious conclusions. The fact that only one individual in the rapid progression group had ciHHV6 support the author's conclusion to conduct a multi-center study with a larger sample size. Based on higher HHV6 prevalence rates described in another groups of HIV-1 infected individuals I would be curious to know whether the authors evaluated HHV6 antibodies in this group and how do they explained that, despite high prevalence, the percentage of integrated HHV6 DNA is so low. The pathogenesis of HHV6 in HIV-1 infected individuals should be further investigated.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research