Check for updates

REVIEW

REVISED Intrathecal Drug Delivery (ITDD) systems for cancer pain [version 4; peer review: 2 approved, 1 approved with reservations]

Gaurav Bhatia, Mary E Lau, Katharine M Koury, Padma Gulur

Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital, Boston MA, 02114, USA

v4 **First published:** 28 Mar 2013, 2:96
<https://doi.org/10.12688/f1000research.2-96.v1>
Second version: 11 Jun 2013, 2:96
<https://doi.org/10.12688/f1000research.2-96.v2>
Third version: 17 Jul 2013, 2:96
<https://doi.org/10.12688/f1000research.2-96.v3>
Latest published: 28 Jul 2014, 2:96
<https://doi.org/10.12688/f1000research.2-96.v4>

Abstract

Intrathecal drug delivery is an effective pain management option for patients with chronic and cancer pain. The delivery of drugs into the intrathecal space provides superior analgesia with smaller doses of analgesics to minimize side effects while significantly improving quality of life. This article aims to provide a general overview of the use of intrathecal drug delivery to manage pain, dosing recommendations, potential risks and complications, and growing trends in the field.

Keywords

Intrathecal drug delivery, chronic pain, cancer, analgesic, opioids

Open Peer Review

Approval Status ✓ ? ✓

	1	2	3
version 4 (revision) 28 Jul 2014			
version 3 (update) 17 Jul 2013	✓ ↑		✓ ↑
version 2 (update) 11 Jun 2013	✓ view ↑		? view ↑
version 1 28 Mar 2013	? view	? view	

1. **Michael Erdek**, John Hopkins University, Baltimore, MD, USA
2. **Arun Bhaskar**, The Christie NHS Foundation Trust, Manchester, UK
3. **Marcin Chwistek** , Fox Chase Cancer Center, Philadelphia, PA, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Padma Gulur (pgulur@partners.org)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2014 Bhatia G *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bhatia G, Lau ME, Koury KM and Gulur P. **Intrathecal Drug Delivery (ITDD) systems for cancer pain [version 4; peer review: 2 approved, 1 approved with reservations]** F1000Research 2014, 2:96 <https://doi.org/10.12688/f1000research.2-96.v4>

First published: 28 Mar 2013, 2:96 <https://doi.org/10.12688/f1000research.2-96.v1>

REVISED Changes from Version 3

The order of the authors on the article has been changed from Gaurav Bhatia, Mary E Lau, Padma Gulur, Katharine M Koury to Gaurav Bhatia, Mary E Lau, Katharine M Koury, Padma Gulur at the request of the authors.

See referee reports

Introduction

Intrathecal drug delivery (ITDD or IDD) is a method which provides analgesia by a direct continuous infusion of medication into the cerebrospinal fluid (CSF). The rationale is based on the concept that direct delivery into the CSF within the intrathecal space avoids crossing the blood-brain barrier and allows delivery of smaller doses than oral, intravenous, transdermal or epidural drug administration¹. The spinal cord is a key anatomical site for pain processing and more specifically the dorsal horn of the spinal cord within the intrathecal space. There are various receptor targets within the spinal column that can be utilized in relieving pain. These receptors include opioid (mu, kappa, and delta), GABA, alpha-2, dopaminergic, NMDA as well as sodium and calcium channels². The advantage of delivering the medication directly into the CSF in smaller dosages to target the specific receptors is that it may reduce the systemic side effects and the direct delivery to the site of action allows a more rapid and effective response. As a popular alternative for patients experiencing cancer or chronic pain, intrathecal analgesia has become a therapeutic option for patients who have exhausted all other treatment avenues as well as patients experiencing side effects from their current treatment options. The system uses a small pump that is surgically placed under the skin of the abdomen and delivers medication through a tunneled catheter directly into the intrathecal space³. The pump is refilled periodically through a subcutaneous access port in an office setting utilizing a sterile technique. The pump can be programmed to be a fixed-rate, constant flow device or a variable rate pump with the option of boluses via patient controlled analgesia programming using a wireless radiofrequency transmitter given to the patient¹. In contrast to more conservative oral or intravenous delivery methods, ITDD is becoming increasingly popular for its efficacy in managing cancer and chronic pain. This review will specifically target ITDD uses for patients with cancer-related pain although its use has also been very popular in patients with chronic non-cancer pain.

Cancer pain: epidemiology

The incidence of deaths due to cancer has been estimated to be 6.6 million people globally each year⁴. Pain is often considered as the

most feared symptom amongst individuals living with cancer and it can occur at any point during the course of the illness. Pain is often also the first sign of malignancy⁴. Pain is also often associated with the treatment of cancer and it can present itself as both acute and chronic pain. Furthermore, advancement of the disease process can lead to pain or make the existing pain worse. It is difficult to estimate the prevalence of cancer pain because of a lack of standardization in definitions of pain and in the measures used to assess it⁵. The other factor that makes it challenging in estimating the prevalence is the heterogeneity of nociceptive and neuropathic pain conditions⁶. About 50% of patients will report pain as a symptom at the time of cancer diagnosis and early in the course of disease⁶. Its prevalence increases to 75% or more at advanced stages. A strategy should be devised to evaluate the prevalence of pain in patients with cancer by separating the pain related to the disease process, to its treatment, or to unrelated causes.

Prevalence of pain related to cancer

According to a systematic review of the past 40 years by van den Beuken *et al.* published in the *Annals of Oncology* in 2007⁴, the range of reported prevalence of pain is highest for the following tumors:

- Head and neck (67–91%)
- Prostate (56–94%)
- Uterine (30–90%)
- Genitourinary (58–90%)
- Breast (40–89%)
- Pancreatic (72–85%)

Treatment options

The various treatment options for cancer pain generally depend on the location, character, duration, its origin, and the success or failure of previous and ongoing treatments. Treatment options for cancer-related pain include oral non-steroidal anti-inflammatory drugs, oral opioid analgesics, neuropathic pain medications, parenteral opioids, peripheral nerve blocks, neurolysis such as celiac plexus blocks, local anesthetic injections, spinal cord stimulation, continuous epidural analgesia, and intrathecal pumps. In most cases, multiple forms of treatment are used for the most effective treatment combination. Opiates such as morphine, dilaudid, fentanyl, and

Table 1. Conversion ratios of morphine. All other drugs are converted to Morphine Equivalent Dose (MED) to allow equianalgesic conversions.

Route of administration	Conversion ratio
Oral	300
Intravenous	100
Epidural	10
Intrathecal	1

sufentanil are most commonly administered via the intrathecal drug delivery system for cancer-related and non-malignant pain⁷. There is increasing evidence that intrathecal opioids are superior to oral delivery in malignant pain, especially when narcotic dosage is limited by its side effects¹. It is believed that this advantage results from delivering extremely small doses on target receptors in the spinal column⁸.

There are other agents also available to treat neuropathic pain, spasticity related pain, sympathetic pain, and visceral pain (see [Table 2](#) for examples). In addition, ITDD systems can also be used to deliver chemotherapy agents such as floxuridine and methorexate for the treatment of primary or metastatic cancer. [Table 2](#) below lists the most commonly targeted receptors and the medications used for each receptor.

One of the most important clinical decisions a provider has to make is what agent to use in the intrathecal pump and in some cases

multiple forms of treatment are used for the most effective treatment combination. In this review, we looked at the recommendations of The Polyanalgesic Consensus Conference (PACC) 2012 which lists the algorithms for intrathecal therapies in both neuropathic and nociceptive pain ([Table 3](#) and [Table 4](#)). The algorithm lists the medications that are arranged in a hierarchy on the basis of evidence of efficacy and safety. The approaches are listed from Line 1 (first line approach) to Line 5 (a more advanced approach if previous approaches are unsuccessful).

Non-malignant pain and other uses

A retrospective study¹⁰ involving 120 patients with non-malignant pain syndromes found that the patients benefited from the intrathecal opiate therapy and the average pain reduction was 67.4% after 6 months from implantation. Over ninety percent of the patients were satisfied with the therapy and 81% reported an improvement in their quality of life.

Table 2. Receptor targets for pain medications².

Receptors targeted	Intrathecal drug delivery system medication(s)
Opioid (mu, delta, kappa)	Morphine, hydromorphone, fentanyl, sufentanil, methadone
GABA	Baclofen, midazolam
Sodium channel receptors	Local anesthetics such as bupivacaine, levobupivacaine, ropivacaine
NMDA	Ketamine, methadone
Calcium channel receptors	Ziconotide
Alpha-2 receptors	Clonidine, dexmedetomidine
Other agents (rarely used)	Adenosine, gabapentin, ketorolac, neostigmine, octreotide

Note: The only medications that are FDA approved for ITDD are: Morphine Sulfate, Baclofen, Ziconotide, Floxuridine, and Methotrexate⁹.

Table 3. 2012 polyanalgesic algorithm for intrathecal (IT) therapies in neuropathic pain⁷.

Line 1	Morphine	Ziconotide	Morphine + bupivacaine
Line 2	Hydromorphone	Hydromorphone + bupivacaine OR hydromorphone + clonidine	Morphine + clonidine
Line 3	Clonidine	Ziconotide + opioid	Fentanyl Fentanyl + bupivacaine OR fentanyl + clonidine
Line 4	Opioid + clonidine + bupivacaine	Bupivacaine + clonidine	
Line 5	Baclofen		

Note: Line 6-Experimental agents: Gabapentin, Octreotide, Conceptide, Neostigmine, Adenosine, XEN 2174, AM 336, XEN, ZGX 160 (ongoing experiments in animal models).

Table 4. 2012 polyanalgesic algorithm for intrathecal (IT) therapies in nociceptive pain⁷.

Line 1	Morphine	Hydromorphone	Ziconotide	Fentanyl
Line 2	Morphine + bupivacaine	Ziconotide + opioid	Hydromorphone + bupivacaine	Fentanyl + bupivacaine
Line 3	Opioid + clonidine			Sufentanil
Line 4	Opioid + clonidine + bupivacaine		Sufentanil + bupivacaine or clonidine	
Line 5	Sufentanil + bupivacaine + clonidine			

Note: Line 6-Experimental agents: Gabapentin, Octreotide, Conceptide, Neostigmine, Adenosine, XEN 2174, AM 336, XEN, ZGX 160 (ongoing experiments in animal models).

Besides opiates, other medications such as baclofen (GABA agonist) for spasticity related pain, ziconotide (calcium channel blocker) for severe chronic neuropathic pain, bupivacaine (amide local anesthetic), clonidine (alpha-2 agonist), and ketamine (NMDA antagonist) have also been used in various instances and in specific patient groups. In addition, ITDD systems can also be used to deliver chemotherapy agents such as floxuridine and methorexate for the treatment of primary or metastatic cancer⁹.

One of the most common uses of the ITDD system is the delivery of baclofen for spasticity related pain³. Baclofen is a GABA-A receptor agonist which can have a prominent effect on the motor tone via direct hyperpolarization of the motor horn cells. It is most often utilized in patients suffering from spasticity, cerebral palsy, amyotrophic lateral sclerosis, Stiff-Person syndrome, and those suffering from brain and spinal cord injuries⁷.

Intrathecal drug delivery systems

One option, intrathecal analgesia, has become a therapeutic option for patients who have exhausted all other treatment avenues as well as patients experiencing side effects from their current treatment options³. In intrathecal drug delivery or ITDD, medication is delivered directly into the CSF in the intrathecal space (Figure 1); therefore, a substantially smaller dose of medication is required as compared to the oral or intravenous route. With lesser dosage of the medication and a more direct route to the pain processing center in the spinal cord, the patient generally experiences superior analgesia with fewer side effects such as nausea, pruritis, erythema, flushing, constipation, etc. However, intrathecal delivery of medication also poses significant risks such as respiratory depression, infection, bleeding, epidural/spinal hematomas, spinal cord injury during initial catheter placement, wound dehiscence, pocket seromas, pump malfunction, and costs.

There have been various publications and consensus guidelines in regards to a screening trial before proceeding with pump implantation^{2,3,7}. Previously, trialing was considered to be the standard of care and critical in evaluation of a patient's response and potential side effects from a certain agent. However, more recently, the PACC 2012 expert panel has called it debatable specifically in cancer-pain patients⁷. The issues of opioid-induced hyperalgesia (OIH) or disease progression is difficult to assess from a single-shot trial or even with a brief 72–96 hour infusion. A trial may also lead to underestimation of the failure rate and potential side effects. Still, the rationale for performing trials is based on the intuitive basis that it best mimics the system that would eventually be implanted⁷. Generally speaking, there is currently no solid base to either refute or adopt preimplantation trials but it is generally recommended although not required². The British Pain Society expert panel suggested that trials should always be performed before the implantation of an intrathecal drug delivery system⁷.

There is a spectrum of intrathecal system options that range from a percutaneous catheter/external pump to a totally implanted system. Choice is dependent on a number of factors including, but not limited to, life expectancy, cost and availability of professional expertise and patient's wishes and comfort level.

Figure 1. Image of the spine illustrating placement of the catheter into the intrathecal space which is connected to an implanted pump containing the drug reservoir. Reprinted with the permission of Medtronic, Inc.© 2010.

Percutaneous

In general, percutaneous refers to a medical procedure in which a catheter or a medical device is introduced into the body via needle-puncture of the skin. This approach is favored because the catheter is easy to place (fluoroscopy can be used to confirm the position of the catheter and to maneuver it into the desired location) and is suitable for patients with limited life expectancy. The catheter is then attached to an external pump which delivers the medication directly into the intrathecal space. However, it is generally considered only a temporary option due to limitations such as frequent monitoring for infection, catheter migration, and patient's immobility. The risks from this procedure are similar to performing a spinal anesthetic which includes bleeding, infection, headache, spinal cord damage, etc.

Fully implanted

Fully implanted drug delivery systems are suitable for long-term use. Mobility and functional activity are not particularly adversely affected by these systems. The implantation is performed by a skilled health care provider in an operating room under monitored anesthesia care with local anesthetic infiltration or general anesthesia. The patients require specialized care with a full multi professional infrastructure including regular follow-up appointments for pump refills. The pump can be programmed to be a fixed-rate, constant flow device or a variable rate pump with the option of boluses via patient controlled analgesia programming using a wireless radiofrequency transmitter given to the patient. Fixed rate delivery systems are less expensive than variable rate delivery systems but lack flexibility of medication delivery based on the need of a patient. These systems have a larger reservoir volume which allows for longer intervals between refills. In cases of suspected or actual medication overdose or implant malfunction, the pump's drug reservoir has to be emptied and the device is interrogated by a designated health care professional.

ITDD devices

There are two systems available for implantable intrathecal devices—those that are programmable and those that are fixed-rate. In addition, some systems come with a bolus option that gives patients more control in managing their treatment. The needs of the patient as well as the suitability of the system for use with selected drugs need to be considered when deciding between the options.

Programmable devices

Programmable devices such as Medtronic's SynchroMed system deliver a specific amount of medication intermittently (Figure 2 and Figure 3). These systems are preferred by providers because they allow drug dosages to be changed without invasive intervention as the disease progresses and/or the patient builds tolerance to the medication. In cases of suspected or actual medication overdose or implant malfunction, the pump can be interrogated and deactivated without having to empty the drug reservoir. However, since it is battery driven, the life of this pump is usually 5–8 years and a surgical revision is required to implant a new pump. In addition, regular attendance for refilling is required because the drug reservoir tends to be smaller in volume than those found in fixed-rate devices.

Fixed-rate devices

Constant flow systems such as Codman's 3000 delivers a continuous amount of medication into the intrathecal space. These devices are favorable because they are generally less expensive, do not require a battery to operate and possess a larger drug reservoir that allows the drug to be delivered for a longer period of time before requiring a refill. However, constant rate devices are disadvantageous because they don't give the flexibility to change the programming or a patient controlled bolus option for additional analgesia.

Devices with bolus option

The Medtronic SynchroMed II infusion system comes with a Patient Therapy Manager (PTM) which is a trademark of the Medtronic Corporation¹¹. It is a remote hand-held device which communicates with the implanted pump and allows the patient to self administer boluses based on the medication limits which are pre-programmed by the clinical provider. This system gives patients the ability to

Figure 2. Image of the SynchroMed II Pump with specifications. Pump is programmable to deliver a specific amount of medication at different times and can be increased or decreased depending on the individuals needs. Reprinted with the permission of Medtronic, Inc.© 2009.

Figure 3. X-ray of the SynchroMed II Pump. The pump has 20 and 40 cc reservoirs to extend the time between refills. Drug is refilled via the reservoir fill port. Reprinted with the permission of Medtronic, Inc.© 2010.

activate a bolus option when adequate pain relief is not achieved. The clinical care provider has the ability to set the number of boluses a patient may receive in a 24 hour period. There is a lockout system in place with the programmer that does not allow the patient to self activate the bolus option more than what it is allowed based on the settings created by the clinical provider. This allows for flexibility for the patients who have more intimate needs to control their pain based on certain body positions, movement, time of the day, and/or progression of their underlying disease. There are several safety options built in to the device such as the maximum number of boluses the patient may self administer in a 24 hour period, maximum dosage of the total drug in a 24 hour period, lockout in-between bolus activation, and the duration of time during which a bolus is administered.

Placement¹²

The placement of ITDD consists of two parts: 1) placement of the catheter in the intrathecal space surrounding the spinal cord and 2) placement of the pump/reservoir in the abdomen/flank region. The pump is implanted in the lower abdomen and is attached to the intrathecal catheter which is tunneled under the skin from the

patient's back towards the abdomen. The procedure requires two surgical incisions: 1) Lower back where the catheter is inserted into the intrathecal space and is secured to the fascia. 2) The lower abdominal wall to create a pocket for the pump. The procedure generally takes 3 to 4 hours from start to finish.

Patient selection

Patient selection is an important consideration. Patients who are good candidates for ITDD therapy include those that have significant side effects from oral or IV opioids which hinders titration of these agents to adequate analgesia. Another criterion includes patients who do not achieve adequate analgesia in spite of significant doses of oral opioids. These pumps are implanted by pain physicians and neurosurgeons. When implanted by pain physicians, they continue to manage the pump as well. Neurosurgeons primarily implant the pump which they then send back to pain specialists or palliative specialists to manage. It has been shown that ITDD for pain management can actually help patients on cancer therapies such as chemotherapy or radiation¹³. This is because ITDD has less adverse effects than other pain treatments, so patients can endure more aggressive levels of chemotherapy or radiation.

Preoperative preparation

Prior to the procedure, a thorough evaluation is performed by the anesthesiologist to ensure patient's ability to undergo surgery and anesthesia. Physical exam, blood tests, electrocardiogram, and chest x-rays may be requested by the provider. Current systemic use of analgesics is discontinued as well as anticoagulants and antiplatelet therapies. The position of the pump is proposed and agreed upon by the patient and provider keeping patients' comfort, activity, need for future procedures and rehabilitation matters in consideration. The choice of medication is dependent on the success of past and present medication choices and acceptable side effects. This is usually predetermined from either previous efficacy of a particular medication or during a trial period.

Operating room procedure

Implantation of an ITDD system is a minor surgical procedure that is carried out in the operating room using aseptic precautions, including skin preparation, sterile draping, and the use of full surgical attire. The patient is positioned in a lateral decubitus position with the patient's side for the pump pocket nondependent¹². Real time fluoroscopy is used to identify the ideal location for placement of the catheter via a needle provided by the device manufacturer. A small skin incision is made in the middle of the back to expose the fascia over the bony arch (lamina) of the vertebra and the catheter is placed in the subarachnoid, or intrathecal space and then anchored into the fascia to prevent migration.

Once the catheter is in place and secure, a tunneling device is used to pass the catheter under the skin from the spine to the abdomen where the pump will be implanted. A 4–6 inch skin incision is made in the side of the abdomen below the waistline. The surgeon then creates a pocket for the pump between the skin and muscle layers and the catheter is attached to the pump. Once the connector is attached to the pump, the pump is correctly positioned under the skin and secured to the thick fascia layer overlying the stomach muscles using suture loops. A space inside the pump called the reservoir is

filled with the desired medication and the pump is programmed. The incisions in the back and abdomen are closed with sutures or staples and a dressing is applied. The patient is taken to the recovery area and is monitored per the standard requirements of the institution before being discharged.

Post-operative

Often antibiotics are prescribed to prevent complications and a one week follow-up is recommended. There may be temporary restriction on daily activities to allow for healing of the wound and pump to settle. Once healed, no special care is necessary.

Dosing and titration

As mentioned previously, ITDD delivers medicine directly into the cerebrospinal fluid, bypassing the route that oral medication takes through the body. Generally 1/100th–300th of the amount of medication is used with the pump delivery as compared with the amount when taken orally. A survey conducted among a small group of physicians highly-experienced in ITDD revealed that morphine was the more frequently used and prescribed choice of medication, both alone and in combination with other drugs¹⁴. The concentrations and daily dosages administered of medications varied from patient to patient. Providers often adjusted the dosages of drugs or drug combinations by a fixed percentage, between 10% and 20%, whether increasing or decreasing the dosage. An average of five to eight adjustments in dosage was completed before a different drug or drug combination was introduced in cases of inadequate analgesia¹³.

Efficacy

Studies have suggested that ITDD is effective in providing pain relief to 60–80% of patients experiencing chronic malignant pain⁷. Long-term intrathecal application of opioid medications in cases of cancer pain is substantially more effective than its systemic application. Intrathecally applied opioids exert a strong analgesic effect via spinal receptors, without significantly influencing motor, sensory, and sympathetic reflexes. It has been shown to improve patients' overall mood and reduces the incidence of systemic side effects. As a result, patients have reported an improvement in quality of life and the ability to participate more fully in daily activities. Furthermore, a randomized controlled study illuminates the efficacy of ITDD over comprehensive medical management in treating refractory cancer pain¹³. Patients receiving ITDD not only had better pain control and relief, but also had improved survival, a reduction in drug toxicity, and fewer drug side effects¹³.

Complications

As with any surgical procedure, serious complications may arise post-implant that require immediate attention and perhaps the removal of the device.

Drug related complications

A retrospective study¹⁵ showed that the most common complication was patients' adverse reaction to a drug. Pharmacological complications were the most common immediately post-implant and generally subsided as treatment continued. Serious complications include respiratory depression/arrest, anaphylaxis, and meningitis with introduction of contaminated solution. It was also found that certain opioids (the drugs of choice) were more prone to complications

than others. For example, the incidence of tip catheter granulomas is higher with morphine and hydromorphone as compared to fentanyl which has been upgraded to a first-line drug in the algorithm due to that specific advantage⁷. **Table 5** below lists the most common drug related adverse reactions and it is compiled based on the data presented in the PACC 2012.

Device related complications

Catheter and pump related malfunctions are one of the main sources of complications. In fact, one study¹⁵ found catheter related complications as the most common cause of repeat surgery. Despite suturing the catheter to the underlying fascia, coiling of the tube and leakage at the connection site remains a problem. A study reviewing complications from long-term intrathecal drug therapy found that the annual rate for complications requiring surgical procedure was 10.5%, with 35% being pump related and 65% catheter related². Errors in pump programming have also been reported and resulted in incorrect flow rates. Device related complications have led to infections (meningitis), pockets abscesses, bleeding, pain and discomfort, and blood or fluid in the pump's pocket. **Table 6** lists the most common device related side effects.

Patient related complications

Although rare, patient induced complications generally involve infections at implant location. Immune cells form a mass, known as a granuloma, around the catheter tip to wall off foreign substances. However, the incidence of infections in all cases reviewed during a study was 0.7% per year¹⁶, with all infections appearing within the first three months post-implant. Factors that increased the potential risk of complication included psychological problems, obstructive sleep apnea, immunosuppression, smoking, diabetes active infections, bleeding disorders, and concurrent anticoagulation therapy^{2,7}. A data analysis found that one successful measure to further decrease the chance of infection was adherence to guidelines and recommendations for surgical site infections¹⁷.

Compounding of medications and its considerations

The FDA and the United States Pharmacopoeia have issued standards on compounded sterile products that have clinical, legal, and practical significance^{7,9}. These standards apply to compounding of solutions by various routes, including intrathecal administration. There are strict

provisions in place for considerations such as: training of personnel, a segregated sterile compounding area, air quality of the compounding area, certification and calibration of equipment, a cleaning and disinfection program, and a quality assurance program⁷. Any violation of any of the provisions can result in a civil and criminal law suit in addition to the suspension of the privileges to dispense compounded medications. An outbreak of fungal meningitis resulted in 2012 when several lots of contaminated methylprednisolone vials were used for epidural steroid injections in various states which were obtained from a compounding pharmacy. Shortly following the outbreak, the device manufacturer Medtronic released an "Urgent Medical Device Safety Notification" in November 2012 discouraging the use of unapproved drugs with its delivery system⁹.

Cost effectiveness

With estimates of pain care treatment costs exceeding \$1 billion annually in the United States, cost and efficacy are important factors in clinical decision making. Although the initial cost of intrathecal drug delivery is substantially more, maintenance costs over time are significantly lower than conventional routes of administration. Cost analyses by the group concluded that intrathecal delivery is the most cost-effective route of opioid administration for patients who require long-term management of cancer ($\geq 3-6$ months) or nonmalignant pain ($\geq 11-22$ months)¹⁸.

An analysis by Kumar *et al.* showed that intrathecal therapy is a cost-effective method of treating chronic nonmalignant pain caused by failed back syndrome and that the break-even point occurs at 28 months, after which conventional pain therapy is the more expensive treatment option¹⁹. A separate analysis by Scott Guillemette from Ingenix Consulting on patients suffering from failed back surgery syndrome concluded with similar results. Despite the higher upfront costs, patients utilizing intrathecal drug delivery returned to 'normal living' more quickly than those with conventional therapy and the break-even point between the two points occurred between months 19 and 20²⁰.

Conclusion

The treatment of pain poses an ongoing challenge to the health-care field and better prevention, assessment and treatment of pain is needed. Intrathecal drug delivery can become a critical component

Table 5. Drug related adverse reactions to medications delivered via ITDD².

Adverse reaction	Associated medications
Peripheral edema	Opioids
Hormonal changes	Opioids
Respiratory depression/somnolence	Opioids, benzodiazepines, local anesthetics, baclofen
Granuloma	Opioids (except fentanyl)
Hyperalgesia/tolerance/withdrawal	Opioids, baclofen
Immune suppression	Opioids
Psychosis, suicidality, hallucinations, confusion	Ziconotide, clonidine, baclofen
Urinary retention, weakness, hypotension	Opioids, local anesthetics
Demyelination, necrotizing lesions	Ketamine, dexmedetomidine

Table 6. Device related side effects².

Device related side effects:
Infection/meningitis
Post-dural puncture headaches/CSF leak
Catheter tip granuloma formation
IT catheter and pump malfunctions
Pocket site seroma formation, bleeding, pain and discomfort

of this transformation in treatment when more conservative forms of pain management have failed to provide adequate pain relief. In comparison to more conservative delivery methods such as drugs administered orally or intravenously, ITDD is more effective because the medication is directly introduced into the subarachnoid space⁶. The spinal cord is a key anatomical site for pain processing and more specifically the dorsal horn of the spinal cord within the intrathecal space^{5,8}. The advantage of delivering the medication directly into the CSF in smaller dosages is that it may reduce the systemic side effects and the direct delivery to the site of action allows a more rapid and effective response. As a popular alternative for patients experiencing cancer or chronic pain, intrathecal analgesia has become a therapeutic option for patients who have exhausted all other treatment avenues

as well as patients experiencing side effects from their current treatment options. However, there are limitations such as adequate patient selection, ability to tolerate the procedure, and costs. There are also potential complications related to the device, drug, and procedure that may require hospitalization and immediate removal/revision of the device. Despite numerous studies involving existing and novel drugs, only limited data is available to address the issue of medication safety, efficacy, stability and compatibility with the intrathecal drug delivery system. As this option becomes more popular with patients and physicians, future research should focus on cross-site studies that would provide a more accurate outlook on this type of delivery.

Author contributions

Guarav Bhatia edited the manuscript to its present state. Mary E Lau formulated the initial draft of the article. Padma Gulur conceptualized the article, and both Padma Gulur and Katharine M Koury edited the manuscript to its present state. All authors agreed to the publication of the final manuscript.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Cohen SP, Dragovich A: **Intrathecal analgesia.** *Anesthesiol Clin.* 2007; **25**(4): 863–882.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stearns L, Boortz-Marx R, Du Pen S, *et al.*: **Intrathecal drug delivery for the management of cancer pain: a multidisciplinary consensus of best clinical practices.** *J Support Oncol.* 2005; **3**(6): 399–408.
[PubMed Abstract](#)
- Smith HS, Deer TR, Staats PS, *et al.*: **Intrathecal Drug Delivery.** *Pain Physician.* 2008; **11**(2 Suppl): S89–S104.
[PubMed Abstract](#)
- van den Beuken-van Everdingen MH, de Rijke JM, Kessels AG, *et al.*: **Prevalence of pain in patients with cancer: a systematic review of the past 40 years.** *Ann Oncol.* 2007; **18**(9): 1437–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vranken JH, van der Vegt MH, Kal JE, *et al.*: **Treatment of neuropathic cancer pain with continuous intrathecal administration of S + ketamine.** *Acta Anaesthesiol Scand.* 2004; **48**(2): 249–252.
[PubMed Abstract](#) | [Publisher Full Text](#)
- International Anesthesia Research Society. **Global year against cancer pain: epidemiology of cancer pain.**
[Reference Source](#)
- Deer TR, Prager J, Levy R, *et al.*: **Polyanalgesic Consensus Conference 2012: Recommendations for the management of pain by intrathecal (intraspinal) drug delivery: report of an interdisciplinary expert panel.** *Neuromodulation.* 2012; **15**(5): 436–464; discussion 464–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ahmed S: **Chapter 13: Intrathecal drug delivery for chronic pain management.** *Translational Pain Research.* 2006; **1**: Nova Science Publishers, Inc.
- Medtronic. **Urgent medical device safety notification: use of unapproved drugs with the SynchroMed Implantable Infusion Pump.**
[Reference Source](#)
- Winkelmüller M, Winkelmüller W: **Long-term effects of continuous intrathecal opioid treatment in chronic pain of nonmalignant etiology.** *J Neurosurg.* 1996; **85**(3): 458–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Medtronic. **Personal therapy manager for SynchroMed II physician manual.**
[Reference Source](#)
- Rathmell JP: **Atlas of image-guided intervention in regional anesthesia and medicine.** Second Edition. Lippincott Williams & Wilkins. Philadelphia, PA. 2012.
[Reference Source](#)
- Smith TJ, Staats PS, Deer T, *et al.*: **Randomized clinical trial of an implantable drug delivery system compared with comprehensive medical management for refractory cancer pain: Impact on pain, drug-related toxicity, and survival.** *J Clin Oncol.* 2002; **20**(19): 4040–4049.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hassenbusch SJ, Portenoy RK: **Current practices in intraspinal therapy – A survey of clinical trends and decision making.** *J pain Symptom Manage.* 2000; **20**(2): S4–S11.
[PubMed Abstract](#)
- Krakovsky AA: **Complications associated with intrathecal pump drug delivery: a retrospective evaluation.** *Am J of Pain Med.* 2007; **17**(1): 4–10.
[PubMed Abstract](#)
- Fluckiger B, Knecht H, Grossmann S, *et al.*: **Device-related complications of long-term intrathecal drug therapy via implanted pumps.** *Spinal Cord.* 2008; **46**(9): 639–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Follett KA, Boortz-Marx RL, Drake JM, *et al.*: **Prevention and management of intrathecal drug delivery and spinal cord stimulation system infections.** *Anesthesiology.* 2004; **100**(6): 1582–1594.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mueller-Schwefe G, Hassenbusch SJ, Reig E: **Cost effectiveness of intrathecal therapy for pain.** *Neuromodulation.* 1999; **2**(2): 77–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kumar K, Hunter G, Demeria DD: **Treatment of chronic pain by using intrathecal drug therapy compared with conventional pain therapies: a cost-effectiveness analysis.** *J Neurosurg.* 2002; **97**(4): 803–810.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Guillemette S: **American Academy of Pain Medicine (AAPM) 26th Annual Meeting.** February 3–6, San Antonio, Texas 2010.
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 3

Reviewer Report 27 August 2013

<https://doi.org/10.5256/f1000research.1897.r1577>

© 2013 Chwistek M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Marcin Chwistek

Department of Medical Oncology, Fox Chase Cancer Center, Philadelphia, PA, USA

Competing Interests: I have in the past received payments from Medtronic Spine for participation on their speaker's bureau in reference to balloon kyphoplasty (a division of Medtronic manufactures kyphoplasty kits). I have not received any payments from Medtronic intrathecal pump division.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 09 August 2013

<https://doi.org/10.5256/f1000research.1897.r1428>

© 2013 Erdek M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Michael Erdek

Department of Anesthesiology & Critical Care Med, John Hopkins University, Baltimore, MD, USA

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 2

Reviewer Report 17 June 2013

<https://doi.org/10.5256/f1000research.1520.r1006>

© 2013 Chwistek M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Marcin Chwistek

Department of Medical Oncology, Fox Chase Cancer Center, Philadelphia, PA, USA

This is an informative review (written with the general practitioner in mind) that provides the basic information and rationale for using ITDD for the treatment of cancer pain.

I have one major reservation and a few suggestions:

1. Cancer pain can be managed by a variety of practitioners; in the majority of situations it will fall under the purview of the patient's primary oncologist or a primary care physician, who typically are unfamiliar with ITDDs. It would be very helpful, in my view, to add a paragraph that would outline in more detail when one should start thinking about referring a patient for consideration of ITDD? What criteria does the patient need to meet? E.g. life expectancy? Whom such patients should be referred to? Who will manage the pump after the implantation? How is it going to affect further treatment such as chemotherapy or radiation? What kind of treatment break one can expect? This kind of information is crucial for the treating oncologist. Some of this information is already in the text, however it would be very useful to organize it better and add some more specific and practical information. Perhaps in a form of a table?

A few clarifications may be needed:

Table 2:

- Is Meperidine still used as an opioid in ITDD? Its clinical use, especially in cancer pain, has been basically eliminated.
- NMDA receptor – consider adding Methadone. Methadone is an opioid but also has NMDA activity

Page 7:

- In the US, the commonly used term is "cancer pain", instead of "carcinoma pain"

Marcin Chwistek, MD FAAHPM

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 14 June 2013

<https://doi.org/10.5256/f1000research.1520.r1004>

© 2013 Erdek M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Michael Erdek

Department of Anesthesiology & Critical Care Med, John Hopkins University, Baltimore, MD, USA

I don't see that the question of the difference in thickness between the 20cc and 40cc pumps was addressed.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 22 April 2013

<https://doi.org/10.5256/f1000research.1238.r905>

© 2013 Bhaskar A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Arun Bhaskar

Research Division, The Christie NHS Foundation Trust, Manchester, UK

This article gives a broad overview of the various types of ITDD systems used in the management of complex cancer pain. It also discusses the advantages, the common side-effects and complications, the technique and cost-effectiveness of using ITDD systems. Overall the article is useful in providing an understanding for doctors and nurses not familiar with the intrathecal drug delivery systems.

The article is written at a basic level and although it is useful to beginners and people who are not well versed in ITDD, it does not deliver much to experienced practitioners. I have reservations as I feel that this review article would have been of a higher standard if it was more comprehensive.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response () 10 Jun 2013

Padma Gulur, Anesthesiology, Pain Medicine, UC Irvine Medical Center, USA

Thank you very much for your response. This article was written for general practitioners to gain an understanding of ITDD systems for cancer pain.

Competing Interests: No competing interests were disclosed.

Reviewer Report 10 April 2013

<https://doi.org/10.5256/f1000research.1238.r885>

© 2013 Erdek M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Michael Erdek

Department of Anesthesiology & Critical Care Med, John Hopkins University, Baltimore, MD, USA

Page 3: There can be a need for increasing doses of opioids as tolerance develops.

Page 5: There is a difference in thickness of the 20cc and 40cc pumps.

Page 6: The lamina is not routinely exposed unless laminotomy leads are being placed.

Page 7: Reference [Smith TJ et al. \(2002\) J Clin Oncol.](#) for efficacy.

Page 7: Reference [Follett K et al. \(2004\) Anesthesiology](#) for complications.

Page 7: (Table 5) – Ziconotide is not an opioid.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response () 10 Jun 2013

Padma Gulur, Anesthesiology, Pain Medicine, UC Irvine Medical Center, USA

Thank you very much for your response.

Page 3: There can be a need for increasing doses of opioids as tolerance develops. We

understand and agree with your comment; however, we are unsure as to where you feel this statement should to be added within the article.

Page 5: There is a difference in thickness of the 20cc and 40cc pumps. We understand and agree with your comment; however, we are unsure as to where you feel this statement fits in the article.

Page 6: The lamina is not routinely exposed unless laminotomy leads are being placed.

6.2 Operating room procedure - We have changed the sentence to say that the incision is made to expose the fascia over the bony arch (lamina).

Page 7: Reference Smith TJ et al. (2002) J Clin Oncol. for efficacy. A discussion of this article has been added into the efficacy section to show that ITDD is a better system of pain management than comprehensive medical management for patients with cancer pain.

Page 7: Reference Follett K et al. (2004) Anesthesiology for complications. After reviewing this article, it has been added into our article that in order to further decrease the chances of infection, there must be an adherence to guidelines and recommendations regarding surgical site infections.

Page 7: (Table 5) – Ziconotide is not an opioid. Ziconotide has been taken out of the chart.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research