RESEARCH ARTICLE

The assessment of CD146-based cell sorting and telomere length analysis for establishing the identity of mesenchymal stem cells in human umbilical cord [version 1; peer review: 1 approved, 1 approved with reservations]

Dimitrios Kouroupis¹, Sarah M. Churchman^{2,3}, Dennis McGonagle^{2,3}, Elena A. Jones^{2,3}

¹Department of Biomedical Research, Foundation for Research and Technology-Hellas, Institute of Molecular Biology and Biotechnology, University of Ioannina, Ioannina, 45110, Greece

²Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, LS97TF, UK

³NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals NHS Trust, Leeds, LS97TF, UK

V1 First published: 11 Jun 2014, 3:126
<https://doi.org/10.12688/f1000research.4260.1>
 Latest published: 27 Aug 2014, 3:126
<https://doi.org/10.12688/f1000research.4260.2>

Abstract

Adult stem cells are characterised by longer telomeres compared to mature cells from the same tissue. In this study, candidate CD146⁺ umbilical cord (UC) mesenchymal stem cells (MSCs) were purified by cell sorting from UC tissue digests and their telomere lengths were measured in comparison to donor-matched CD146-negative fraction. UC tissue fragments were enzymatically treated with collagenase and the cells were used for cell sorting, colony-forming fibroblast (CFU-F) assay or for long-term MSC cultivation. Telomere lengths were measured by qPCR in both culture-expanded MSCs and candidate native UC MSCs. Immunohistochemistry was undertaken to study the topography of CD146⁺ cells.

Culture-expanded UC MSCs had a stable expression of CD73, CD90 and CD105, whereas CD146 declined in later passages which correlated with the shortening of telomeres in the same cultures. In three out of four donors, telomeres in candidate native UC MSCs (CD45⁻CD235α⁻CD31⁻CD146⁺) were longer compared to donor-matched CD146⁻ population (CD45⁻CD235α⁻CD31⁻CD146⁻). The frequency of CD45⁻CD235α⁻CD31⁻CD146⁺ cells measured by flow cytometry was ~1000-fold above that of donor-matched CFU-Fs (means 10.4% and 0.01%, respectively). CD146⁺ cells were also abundant *in situ* having a broad topography including high levels of positivity in muscle areas in addition to vessels.

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
27 Aug 2014		view

version 1
11 Jun 2014	view	view

1. **Armin Attar**, Shiraz University of Medical Sciences, Shiraz, Iran

2. **Simone Pacini**, University of Pisa, Pisa, Italy

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Elena A. Jones (e.jones@leeds.ac.uk)

Competing interests: No competing interests were disclosed.

Grant information: SC is supported by NIHR-Leeds Musculoskeletal Biomedical Research Unit (LMBRU). EJ and DM hold funding from Wellcome Trust/EPSRC through WELMEC, a Centre of Excellence in Medical Engineering, under grant number WT088908/Z/09/Z and are additionally supported by LMBRU.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2014 Kouroupis D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Kouroupis D, Churchman SM, McGonagle D and Jones EA. **The assessment of CD146-based cell sorting and telomere length analysis for establishing the identity of mesenchymal stem cells in human umbilical cord [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2014, **3**:126 <https://doi.org/10.12688/f1000research.4260.1>

First published: 11 Jun 2014, **3**:126 <https://doi.org/10.12688/f1000research.4260.1>

Introduction

Perinatal mesenchymal stem cell (MSC) sources are attracting increasing attention as an alternative to 'gold standard' bone marrow (BM) MSCs. Owing to its universal accessibility, umbilical cord (UC) tissue represents an attractive source of MSCs for cell therapy¹. However, the potency of standard culture-expanded UC MSCs, especially towards osteoblasts, chondrocytes and adipocytes, is lower compared to BM MSCs^{2,3}. This can be explained by the fact that conventional UC MSC cultures arise from diverse clonal populations that possess varying degrees of self-renewal leading to mixed cultures that gradually lose their MSC properties⁴. The therapeutic advantages of minimally-expanded UC MSCs would therefore include a better preservation of their native functionality as well as rapid manufacture and reduced cost². However, no agreement yet exists on the native phenotype of UC MSCs⁵⁻⁷, which is an essential pre-requisite for their isolation and minimal expansion.

Prospective isolation of UC MSCs has been attempted after enzymatic digestion followed by selection for specific cell surface markers using fluorescence-activated cell sorting (FACS) or immunomagnetic bead separation techniques. Previous studies showed that CD45-based negative depletion⁵, neural ganglioside (GD2)-⁷ and CD146-based positive selection⁶⁻¹¹ could be used to select for clonogenic and multipotential MSC fractions. Although CD146 is expressed on pericytes, which have been proposed as a reservoir of tissue specific progenitors and MSCs^{12,13}, it is also present on the surface of CD31⁺ UC endothelial cells (ECs)^{2,6,9}, and therefore cannot be used alone to achieve high-purity UC MSC preparations.

Whilst the majority of studies in this field used combinations of surface markers to isolate putative MSCs from different tissues, followed by their long-term clonal expansion and analysis of multipotentiality^{10,14}, another alternative and faster approach could involve measuring the telomere length of freshly-sorted candidate MSC fractions. As immature cells, MSCs by definition should have longer telomeres compared to mature cell types. This idea was first proposed by Flores *et al.* (2008) who showed that stem cell compartments from hair follicles, intestine, testis, cornea and brain are enriched with cells with the longest telomeres¹⁵.

The aims of this study were to confirm that CD146 could be a good MSC marker in the UC and to purify candidate MSCs from UC tissue digests based on the non-haematopoietic (CD45⁺CD235 α ⁻), non-EC (CD31⁻), CD146⁺ phenotype. We measured telomere lengths in CD146⁺ and donor-matched control CD146⁻ populations using quantitative real-time PCR (qPCR) and examined the topography of CD146⁺ cells *in situ* using immunohistochemistry.

Materials and methods

Patients and cells

UC tissue was collected from the UCs of consenting full-term caesarean section patients (n=8). After delivery, UCs were immediately stored in Dulbecco's phosphate buffered saline (DPBS, #14190-250, Invitrogen, Renfrew, UK) at 4°C. All samples were obtained after written informed consent and the protocols were approved by National Research Ethics Committee (07/Q1205/27). Human primary

skin fibroblasts were obtained from Lonza and ATCC (Lonza, Cambridge, UK and ATCC, Middlesex, UK).

MSC and EC isolation from UC tissue

The whole UC tissue was mechanically dissected in small pieces (~0.2±0.1 g) and washed repeatedly with DPBS followed by enzymatic digestion using 600 U/ml collagenase I (#07902, Stem Cell Technologies, Grenoble, France)² for 6 hours. Released cells were resuspended in 1:50 v/v DPBS, filtered through a 70 µm cell strainer (#352350, BD Biosciences, Oxford, UK), centrifuged (650 g) and counted.

For primary MSC culture, UC tissue digests were seeded in 6-well tissue culture plates (#3516, Corning, Amsterdam, The Netherlands) in non-haematopoietic (NH) expansion medium (#130-091-680, Miltenyi Biotec, Bisley, UK) at a density of 5×10⁴ cells/well. After observing 80% cell confluency [denoted passage (p) 0], adherent cells were trypsinised using 0.5% trypsin/EDTA (#15400-054, Invitrogen) and re-seeded at 5×10⁵/25 cm² flask for further passaging which was performed as 1:1 splits until approximately p17. Cultures were fed twice weekly with half media changes. The number and viability of the cells at each passage were evaluated using Trypan blue staining. For primary EC culture, 5×10⁴ cells were seeded into 6-well tissue culture plates coated with 2 µg/cm² fibronectin (#354008, BD Biosciences) in endothelial basal medium (EBM2, #CC-3162, Lonza). On day 2, adherent cells were washed with DPBS and subsequently fed three times per week with half media changes. ECs were passaged using 1:1 splits until p4.

For all cultures, the number of population doublings (PDs) between passages (starting from initial passage; p0) was calculated according to the following equation: PD= log 2 (Nt/Ni), where Ni and Nt are the initial and terminal cell counts, respectively. PDs before p0 were calculated based on colony-forming unit-fibroblast (CFU-F) potential of cells seeded and the number of cells at p0 according to the equation PD= log 2 (N cells at p0/N seeded CFU-Fs). The CFU-F assay was performed in triplicate at the cell seeding density of 5×10⁴ cells/well and individual colonies (>50 cells) were counted on day 14 after staining with 1% w/v Crystal Violet (#V5265, Sigma, Hertfordshire, UK).

Gene expression analysis of UC MSCs

RNA extraction from culture expanded UC MSCs (p4) and fibroblasts was performed using RNA/DNA/Protein purification kit (#23500, Norgen, Ontario, Canada) according to the manufacturer's instructions and RNA yield was quantified by using NanoDrop 2000 spectrophotometer (Thermo, Essex, UK). Single strand cDNA was synthesised using high-capacity cDNA reverse transcription kit (#4368814, Applied Biosystems, Warrington, UK).

A custom designed 48 gene Taqman low density array (TLDA, Applied Biosystems) contained mesenchymal and endothelial lineage-related transcripts and novel surface receptors that could be used to segregate MSCs from fibroblasts based on previously published data^{16,17}. To perform TLDA, 200 ng cDNA were used per port. The results were obtained using an ABI PRISM 7900HT SDS (Applied

Biosystems). Normalisation of transcript levels relative to reference gene *HPRT* was performed using the formula: $2^{-\Delta Ct}$, $\Delta Ct = Ct \text{ value (selected transcript)} - Ct \text{ value (HPRT)}$.

Immunophenotyping of MSC and EC cultures

Phenotypic characterisation was performed on culture expanded MSCs at different passages and on cultured ECs at p4 using: CD31-FITC (#MCA1738F), CD105-PE (#MCA1557PE), CD90-PE (#MCA90PE) (all from Serotec, Kidlington, UK), CD73-PE (#550257), CD146-PE (#550315) (both from BD Pharmingen, Oxford, UK), and CD271-PE (#130-091-885, Miltenyi Biotec). The isotype controls were IgG1-FITC (#550616, BD Pharmingen) and IgG1-PE (#MCA928PE, Serotec). A total of 2×10^5 cells was stained with 5 μ l FITC- or PE-conjugated antibodies, and dead cells were excluded using 2 μ g/ml propidium iodide (PI, #P1304MP, Invitrogen). Cells were acquired using FACScan equipped with CellQuest software version 3.1 (BD Biosciences) and the proportions of the different fractions were calculated as a percentage of total live cells.

Fluorescence-activated cell sorting of candidate UC MSCs

Cell sorting was performed using a MoFlo cell sorter equipped with SUMMIT software (Beckman Coulter, Buckinghamshire, UK). Following collagenase digestion of UC tissue, 2×10^6 cells were split into two tubes. One tube was stained with 5 μ l of neat CD45-FITC (#F0861), CD235 α -FITC (#F0870) (both from DAKO, Cambridge, UK), CD146-PE (BD Pharmingen) and CD31-APC (#130-092-652, Miltenyi Biotec), whereas the other was stained with 2.5 μ l of neat isotype controls alone. After incubation with relevant antibodies and washes, 2 μ g/ml 7-aminoactinomycin D (7-AAD) (#A1310, Invitrogen) was added to exclude dead cells before sorting into four fractions: haemopoietic cell fraction (HC), CD45 $^+$ CD235 α^+ CD31 $^-$; EC fraction, CD45 $^-$ CD235 α^+ CD31 $^+$; double-negative candidate MSC fraction, CD45 $^-$ CD235 α^+ CD31 $^-$ CD146 $^+$ and non-MSC fraction, CD45 $^-$ CD235 α^+ CD31 $^-$ CD146 $^-$. The latter two subsets were processed for telomere length analysis.

Telomere length measurements

QIAamp DNA Mini kit (#51306, Qiagen, Crawley, UK) was used for gDNA extraction from cultured MSCs and freshly-sorted CD146 $^+$ and CD146 $^-$ subsets. Telomere length measurement by SYBR Green qPCR (#4309155, Invitrogen) involved determining the relative ratio of telomere repeat copy number (T) to a single copy gene (*36B4*) copy number (S): T/S ratio, as previously described^{18,19}. Samples were run in triplicate using 20 ng gDNA.

Immunohistochemistry

Immunohistochemistry was used to characterise UC tissue architecture and investigate CD146 cell topography *in situ* to ascertain whether it exhibited the proposed MSC pericyte distribution. Whole UC tissue cross sections were embedded in OCT mounting media (#361603E, VWR, Leicestershire, UK), snap frozen in liquid nitrogen and stored at -80°C. Cryostat sections (6 μ m) were mounted on superfrost slides (#48311-703, VWR) and dried overnight at 37°C. Immunohistochemistry was performed using DAKO REAL detection system (#K4065, DAKO) according to the manufacturer's

instructions. Primary antibodies included: CD31 (#CBL468, working concentration 1:10), CD146 (#MAB16985, working concentration 1:50) (both from Chemicon, Watford, UK), CD34 (#M716501, working concentration 1:100), CD271 (#M3507, working concentration 1:20) (both from DAKO), CD90 (#MCA90, working concentration 1:100) and alpha smooth muscle actin (α SMA, #MCA5781GA, working concentration 1:20) (both from Serotec). Antibody binding was visualized using DAKO REAL DAB+ chromogen (#K3468, DAKO) and slides were counterstained by mounting in Harris haematoxylin (#HHS128, Sigma). Slides were mounted using DPX (#317616, Sigma) and images were captured using CAMEDIA C-7070 camera (Olympus, Tokyo, Japan).

Statistical analysis

The software used for analysis and statistics was GraphPad (GraphPad software, La Jolla, USA). The gene expression results were analysed with Mann-Whitney test for unpaired samples ($p < 0.05$: high significance, 95% confidence interval). The cell sorting and CFU-F results were analysed by Kruskal-Wallis test with Dunn's correction.

Results

Molecular profile of culture-expanded UC MSCs compared to fibroblasts

We initially aimed to confirm the validity of CD146 as a candidate marker of UC MSCs using cultures established following standard MSC protocols². Mesenchymal tri-potentiality of these cultures was demonstrated in our previous study². We studied the expression of 45 MSC- and fibroblast-related transcripts by TLDA in these cultures and compared their expression levels to those of negative control skin fibroblasts. Eighteen transcripts were selected (Table 1) as they showed interesting differences between UC MSCs and fibroblasts. Moreover, five transcripts showed more than two-fold higher expression in UC MSCs. These included *NANOG* (a pluripotency marker), whose expression in UC cultures suggested their greater level of immaturity, as well as *NGFR/CD271* and *MCAM/CD146*, the markers of native BM MSCs^{20,21}. Fibroblasts showed stronger expression of eight transcripts (Table 1).

Expression of CD146 and CD271 during extended culture of UC MSCs

Having shown *NGFR* and *MCAM* transcript expression in culture-expanded UC MSCs, their surface protein expression at p4 and subsequent passages was next investigated. Phenotypic characterization was performed at four different passages (representing approximately 17, 19, 21, 25 PDs). MSC-specific markers CD90, CD73 and CD105²² were uniformly positive showing a stable expression profile throughout the expansion, whereas the absence of EC marker CD31 showed no contamination with ECs (Figure 1A).

Despite the expression of *NGFR* transcript, CD271 surface protein expression was absent in culture-expanded MSCs at all tested time-points (Figure 1A). CD146 expression declined gradually and correlated with telomere loss in the two cultures tested (Figures 1A and 1B). This effect was not evident for CD90 which remained stable (Figure 1B). When comparing the phenotypic profile of UC MSCs

Table 1. Relative gene expression in UC MSCs compared to skin fibroblasts. Normalisation of transcript levels relative to reference gene *HPRT* was performed using the formula: $2^{-\Delta Ct}$, $\Delta Ct = Ct \text{ value (selected transcript)} - Ct \text{ value (HPRT)}$.

Gene	TaqMan assay ID	Description	UC MSCs	FIBs	UC MSCs to FIBs
NGFR	Hs00182120_m1	nerve growth factor receptor	0.11	0.02	5.5
MCAM	Hs00174838_m1	melanoma cell adhesion molecule	1.1	0.35	3.14
SOX9	Hs00165814_m1	SRY (sex determining region Y)-box 9	0.29	0.12	2.42
HIF1B	Hs01121918_m1	aryl hydrocarbon receptor nuclear translocator	2.96	1.23	2.41
NANOG	Hs02387400_g1	Nanog homeobox	0.06	0.03	2.0
HIF1A	Hs00936371_m1	hypoxia inducible factor 1, α subunit	109.63	58.17	1.88
LRP5	Hs00182031_m1	low density lipoprotein receptor related protein 5	7.08	3.91	1.81
VEGFA	Hs00900058_m1	vascular endothelial growth factor A	7.01	4.69	1.49
SFRP4	Hs00180066_m1	secreted frizzled-related protein 4	0.02	0.03	0.66
CEBPA	Hs00269972_s1	CCAAT/enhancer binding protein (C/EBP)	0.03	0.05	0.6
FRZB	Hs00173503_m1	frizzled-related protein	0.005	0.01	0.5
COL1A2	Hs01028971_m1	collagen, type I, $\alpha 2$	75.61	212.62	0.35
SPARC	Hs00277762_m1	osteonectin	28.34	98.73	0.28
VEGFB	Hs00173634_m1	vascular endothelial growth factor B	9.52	33.48	0.28
BMPER	Hs00403062_m1	BMP binding endothelial regulator	3.06	10.84	0.28
DDR2	Hs00178815_m1	discoidin domain receptor tyrosine kinase 2	4.1	17.59	0.23
FZD4	Hs00201853_m1	frizzled family receptor 4	0.79	3.64	0.22
HIF1AN	Hs00215495_m1	hypoxia inducible factor 1, α subunit inhibitor	3.14	15.21	0.21

Figure 1. MSC marker expression in culture-expanded UC MSCs. **A** - Surface expression levels of MSC markers (n=3 donors, y error bars indicate SD). **B** - Donor variation of CD146 and CD90 markers in early (<20 PDs) and late (>20 PDs) cultures (M1: marker expression). Bottom panels - telomere T/S ratios were directly correlated to the expression levels of CD146 and not CD90 during culture-expansion.

with ECs only CD31 revealed high specificity for ECs; CD73, CD105, CD146 were expressed on both UC MSCs and ECs (data not shown).

In combination with the molecular profile of UC MSCs and the previously-published literature^{6,8-11}, these data indicated that uncultured UC CD146⁺ cells could indeed possess longer telomeres than the remaining CD146⁻ cells.

Telomere length measurement in sorted CD146⁺ and CD146⁻ populations

The cell sorting strategy for these experiments is described in **Figure 2A**; live cells (R1), containing distinct populations of ECs (R2) and HCs (R3) were clearly observed. Double negative cells (R4) were further subdivided into CD146⁺ (R5, candidate UC MSC) and CD146⁻ (R6, non-MSC) subsets and sorted into separate RNA lysis buffers. In general, HCs were most abundant (mean 48% of total live cells, n=8). CD31⁺ ECs represented a mean of 1.4% of live cells.

Within the double-negative cells, the CD146⁻ population was predominant over the CD146⁺ population (mean 81% and 17% respectively, $p < 0.05$; **Figure 2A**). When these frequencies were re-calculated in relation to total live cells, the CD146⁺ and CD146⁻ fractions represented a mean of 10.4% and 37.7%, respectively (**Figure 2B**). This was significantly higher than the frequency of CFU-Fs (as a percentage of total live cells) in the same UC digests (mean 0.01%, **Figure 2B**). When the telomere length of both sorted subsets (CD146⁺ and CD146⁻) were tested, the CD146⁺ subset exhibited higher telomere lengths compared to the CD146⁻ subset in three out of four donors (**Figure 2C**), hence the observed differences failed to reach statistical significance.

The high frequency of CD45⁺CD31⁺CD235α⁺CD146⁺ cells compared to the CFU-F frequency and the lack of consistent and significant enrichment for cells with long telomeres indicated that the CD146⁺ UC fraction most likely contained non-haemopoietic cells with varying degrees of maturity.

CD146 expression in UC tissue *in situ*

Initial haematoxylin staining (**Figure 3A**) revealed the basic structure of the UC tissue where the UC vein and artery could be seen. This was surrounded by the single thickness endothelial area (EA) and highly organised muscular perivascular area (PA). The Wharton's jelly (WJ) matrix could also be seen spanning intra-muscular areas.

Immunohistochemistry was next used to investigate marker expression *in situ* including semi-quantitative assessment in different anatomical areas (**Figures 3B and 3C**). Three optical microscope fields (×200) were evaluated per anatomical area of UC tissue. CD271 was present at low levels in the WJ area only. EC-specific CD31 and CD34 were present in the EA of the vessels. The intracellular marker of smooth muscle cells; αSMA²³ was expressed and had its highest positivity in the PA surrounding vessels. CD90 and CD146 were highly expressed in most UC compartments, including the PA;

Figure 2. Sorting strategy and telomere length measurements in putative native UC MSCs. **A** - Cell sorting strategy: live cells (upper right graph, R1) were identified by 7AAD exclusion method. Three distinct populations (bottom right graph, R2/ECs, R3/HCs, and R4/double-negative) were evident. Gating on double-negative subset (R4) revealed two subsets (bottom left graph), CD146⁺ (R5/candidate MSCs) and CD146⁻ (R6/non MSCs). Cells confined to regions R5 and R6 were sorted and processed for telomere length analysis. **B** - The percentage of CD146⁺ and CD146⁻ populations compared to the percentage of CFU-Fs (n=4 donors, y error bars indicate SD, * $p < 0.05$, *** $p < 0.001$). **C** - Telomere T/S ratios in sorted subsets (n=4 donors).

Figure 3. Tissue architecture of UC vein, artery and surrounding Wharton's jelly. A – Images show endothelial area (EA, indicated by arrows), perivascular area (PA, multiple layers of muscle fibres) and Wharton's jelly area. **(B)** Expression of MSC and EC markers in UC tissue. **(C)** Staining of MSC and EC markers in UC tissue (representative donor and cross sections, original magnification $\times 200$; (+) symbol indicates the expression of a marker and (-) symbol indicates the absence of expression of a marker).

however CD146 but not CD90 showed positivity for the EA area. This was consistent with previously published literature⁹. Representative photomicrographs are shown in **Figure 3C**.

Overall, our immunohistochemistry results revealed the expected topography of CD31⁺ and CD34⁺ ECs, the expression of CD271 in the WJ area, and the preferential topography of CD90 and α SMA in PA. Consistent with previous findings⁹, CD146 was expressed in WJ, PA and EA, with the highest proportion of cells present in the PA. Wide distribution of CD146⁺ cells in all anatomical areas of UC tissue was consistent with the high frequency of CD45⁺CD31⁻CD235⁻CD146⁺ cells evident by flow cytometry. This indicated that UC MSC isolation alone, even after the removal of CD31⁺ ECs, was not sufficient to purify native MSCs from the UC tissue.

Data for CD146-based cell sorting and telomere length in umbilical cord

1 Data File

<http://dx.doi.org/10.5256/f1000research.4260.d28700>

UC MSC and fibroblast cultures were evaluated for their expression levels of selected transcripts characterizing multiple cell fates *in vivo*. Ct values obtained using Taqman qPCR technology are shown in data set 1 (UC=Umbilical cord; FIB=fibroblast cell lines). The flow cytometry phenotype profiles of UC MSCs (cultures tested against a panel of MSC markers) and UC ECs (cultures tested against a panel of EC markers) are shown in data set 2. Telomere lengths of cultured UC MSC1 and MSC2 samples are displayed in data set 3 (telomere lengths calculated as T/S ratio in different passages of UC MCS cultures).

Discussion

The present study assessed the possibility of using CD146-based cell sorting and telomere length analysis for establishing the identity of mesenchymal stem cells in human UC. Previous studies have shown that the UC CD146⁺ subset contained MSCs able to differentiate into osteoblasts, chondrocytes and adipocytes^{8,9}. Other studies have shown that 'true' immature stem cells have much longer telomeres compared to the remaining mature cells^{15,24}. Here we tested whether CD146⁺ UC cells, that were depleted of contaminating haemopoietic and endothelial cells, had longer telomeres compared to the corresponding CD146⁻ population.

Initially we performed gene expression analysis, flow cytometry and telomere length measurements on culture-expanded UC MSCs during their extensive passaging. The mesenchymal tripotentiality of these cultures has been demonstrated in our previous study². In this study, gene expression analysis revealed that *MCAM*/CD146 was expressed at higher levels (>3-fold) in UC MSCs compared to fibroblasts, confirming its potential specificity for MSCs. Whereas common MSC-specific markers CD73, CD105, CD90²² displayed stable expression throughout passaging, CD146 expression declined with the increased number of cell divisions at later passages consistent with a loss of multipotential progenitors⁴. The CD271 surface marker was not expressed at any stage during UC MSC cultivation, consistent with the loss observed in BM MSCs^{16,25,26}. Although CD271 is expressed on uncultured BM MSCs, it was not selective for UC blood MSCs^{21,27}. Therefore, CD271-based MSC isolation from UC tissue was not pursued further. However, the decrease in CD146 during expansion correlated with telomere erosion in the same cultures, supporting the idea that CD146 could mark the most immature cells *in vivo*. Based on the present findings and previous data^{9,11}, CD146 was selected as a candidate positive marker for sorting native MSCs from UC digests. CD45/CD235 α and CD31 were used to exclude native HCs and ECs from the analysis, respectively².

Sorted CD146⁺ and CD146⁻ subsets were next compared with respect to their telomere length. Several previous studies provided initial evidence that UC MSCs express telomerase continuously and hence were able to retain long telomeres^{28,29}. In UC blood MSCs, a distinct SSEA-4⁺CD105⁺MSCA-1⁺CD90⁻ cell population was shown to have longer telomeres than the SSEA-4⁺CD105⁺MSCA-1⁺CD90⁺ subset³⁰. Although evident in three out of four cultures, in our study we observed the lack of significant difference in telomere length between CD146⁺ and CD146⁻ subsets potentially indicating contamination of MSCs with more mature cells in the CD146⁺ subset. The high frequency of CD146⁺ cells (10.4%) was in stark contrast to the CFU-F frequency (~0.01%), which was consistent with previous findings^{5,31}. Furthermore, the immunohistochemistry data confirmed the broad reactivity of CD146 with UC tissue anatomical areas⁹. Altogether, these data provided the first indication that the CD146⁺ population was unlikely to represent pure native UC MSCs.

Alternatively, a lack of significant difference in telomere lengths could be explained by the limited sensitivity of qPCR assay to

measure telomere lengths³². Telomere length assessment in the majority of studies using cultured MSC is based on terminal restriction fragment (TRF) analysis^{33–37}, which has the disadvantage of producing a smear of bands rather than a discrete band/point affecting accurate quantifications. Additionally, the large number of cells required for such analysis precluded its use on sorted UC cells. On the other hand, studies investigating the correlations between telomere lengths derived by qPCR and TRF indicated good levels of correlation ($r > 0.823$)^{20,38}, although qPCR was shown to be limited in its ability to measure the longest telomeres³². This potential technical limitation of qPCR could have affected the accuracy of the telomere analysis in the present study.

Our findings could be further compounded by the large donor-to-donor variation, which was evident in culture expanded UC MSCs (~10% between donor 1 and 2, **Figure 1B**), but also with sorted subsets (**Figure 2C**). One recent study has demonstrated that telomere lengths could be heritable, with the parental age at conception being a factor affecting offsprings' telomere length in leukocytes (LTL)³⁸. Although the mode of LTL inheritance has been suggested to be X-linked³⁹, another study shows a paternal mode of heritability⁴⁰. The parental ages of the UC donors were not known/recorded in this study but would be an interesting subject for the future work. Previous studies have suggested that the magnitude of inter-individual variation in telomere lengths could exceed the variation between cell types within the same individual^{41–43}, therefore telomere length measurement approaches should be further refined before they can be used as a tool to identify MSCs *in vivo*.

To conclude, the broad availability of UC tissue makes it amenable to be used in cell therapy and regenerative medicine interventions. To this end, the present study showed that in the UC, CD146⁺ cells were too numerous to be selective for pure native MSCs and were likely to contain more mature cells. Therefore, additional markers would be needed to isolate MSCs from UC tissue.

Data availability

F1000Research: Dataset 1. Data for CD146-based cell sorting and telomere length in umbilical cord, [10.5256/f1000research.4260.d28700](https://doi.org/10.5256/f1000research.4260.d28700)⁴⁴

Consent

Written informed consent has been obtained from full-term caesarean section patients for the use of clinical samples in the present study.

Author contributions

DK, SMC, DM, EJ conceived the study. DK performed the experimental work. All authors were involved in manuscript preparation and have agreed to the final content.

Competing interests

No competing interests were disclosed.

Grant information

SC is supported by NIHR-Leeds Musculoskeletal Biomedical Research Unit (LMBRU). EJ and DM hold funding from Wellcome Trust/EPSRC through WELMEC, a Centre of Excellence in Medical Engineering, under grant number WT088908/Z/09/Z and are additionally supported by LMBRU.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We would like to thank Mrs. Anne English for her valuable technical contribution in histology and flow cytometry in this project.

References

- Troyer DL, Weiss ML: **Wharton's jelly-derived cells are a primitive stromal cell population.** *Stem Cells.* 2008; **26**(3): 591–599.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kouroupis D, Churchman SM, English A, et al.: **Assessment of umbilical cord tissue as a source of mesenchymal stem cell/endothelial cell mixtures for bone regeneration.** *Regen Med.* 2013; **8**(5): 569–581.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bosch J, Houben AP, Radke TF, et al.: **Distinct differentiation potential of "MSC" derived from cord blood and umbilical cord: are cord-derived cells true mesenchymal stromal cells?** *Stem Cells Dev.* 2012; **21**(11): 1977–1988.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Saragaser R, Hanoun L, Keating A, et al.: **Human mesenchymal stem cells self-renew and differentiate according to a deterministic hierarchy.** *PLoS One.* 2009; **4**(8): e6498.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Saragaser R, Lickorish D, Baksh D, et al.: **Human umbilical cord perivascular (HUCPV) cells: a source of mesenchymal progenitors.** *Stem Cells.* 2005; **23**(2): 220–229.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Baksh D, Yao R, Tuan RS: **Comparison of proliferative and multilineage differentiation potential of human mesenchymal stem cells derived from umbilical cord and bone marrow.** *Stem Cells.* 2007; **25**(6): 1384–1392.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Xu J, Liao W, Gu D, et al.: **Neural ganglioside GD2 identifies a subpopulation of mesenchymal stem cells in umbilical cord.** *Cell Physiol Biochem.* 2009; **23**(4–6): 415–424.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tsang WP, Shu Y, Kwok PL, et al.: **CD146+ human umbilical cord perivascular cells maintain stemness under hypoxia and as a cell source for skeletal regeneration.** *PLoS One.* 2013; **8**(10): e76153.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Schugar RC, Chirieleison SM, Wescoe KE, et al.: **High harvest yield, high expansion, and phenotype stability of CD146 mesenchymal stromal cells from whole primitive human umbilical cord tissue.** *J Biomed Biotechnol.* 2009; **2009**: 789526.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Torin A, Li O, Brune JC, et al.: **CD146 expression on primary nonhematopoietic bone marrow stem cells is correlated with *in situ* localization.** *Blood.* 2011; **117**(19): 5067–5077.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Covas DT, Panepucci RA, Fontes AM, et al.: **Multipotent mesenchymal stromal cells obtained from diverse human tissues share functional properties and gene-expression profile with CD146+ perivascular cells and fibroblasts.** *Exp Hematol.* 2008; **36**(5): 642–654.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Crisan M, Yap S, Casteilla L, et al.: **A perivascular origin for mesenchymal stem cells in multiple human organs.** *Cell Stem Cell.* 2008; **3**(3): 301–313.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bianco P, Robey PG, Simmons PJ: **Mesenchymal stem cells: revisiting history, concepts, and assays.** *Cell Stem Cell.* 2008; **2**(4): 313–319.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zimmerlin L, Donnenberg VS, Pfeiffer ME, et al.: **Stromal vascular progenitors in adult human adipose tissue.** *Cytometry A.* 2010; **77**(1): 22–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Flores I, Canela A, Vera E, et al.: **The longest telomeres: a general signature of adult stem cell compartments.** *Genes Dev.* 2008; **22**(5): 654–667.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Churchman SM, Ponchel F, Boxall SA, et al.: **Transcriptional profile of native CD271+ multipotential stromal cells: Evidence for multiple fates, with prominent osteogenic and Wnt pathway signaling activity.** *Arthritis Rheum.* 2012; **64**(8): 2632–2643.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ishii M, Koike C, Igarashi A, et al.: **Molecular markers distinguish bone marrow mesenchymal stem cells from fibroblasts.** *Biochem Biophys Res Commun.* 2005; **332**(1): 297–303.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jones E, English A, Churchman SM, et al.: **Large-scale extraction and characterization of CD271+ multipotential stromal cells from trabecular bone in health and osteoarthritis: implications for bone regeneration strategies based on uncultured or minimally cultured multipotential stromal cells.** *Arthritis Rheum.* 2010; **62**(7): 1944–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cawthon RM: **Telomere measurement by quantitative PCR.** *Nucleic Acids Res.* 2002; **30**(10): e47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bianco P, Sacchetti B, Riminucci M: **Osteoprogenitors and the hematopoietic microenvironment.** *Best Pract Res Clin Haematol.* 2011; **24**(1): 37–47.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Watson JT, Foo T, Wu J, et al.: **CD271 as a marker for mesenchymal stem cells in bone marrow versus umbilical cord blood.** *Cells Tissues Organs.* 2013; **197**(6): 496–504.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dominici M, Le Blanc K, Mueller I, et al.: **Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement.** *Cytotherapy.* 2006; **8**(4): 315–317.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hinz B, Celetta G, Tomasek JJ, et al.: **Alpha-smooth muscle actin expression upregulates fibroblast contractile activity.** *Mol Biol Cell.* 2001; **12**(9): 2730–2741.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wagner W, Bork S, Horn P, et al.: **Aging and replicative senescence have related effects on human stem and progenitor cells.** *PLoS One.* 2009; **4**(6): e5846.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Quirici N, Soligo D, Bossolasco P, et al.: **Isolation of bone marrow mesenchymal stem cells by anti-nerve growth factor receptor antibodies.** *Exp Hematol.* 2002; **30**(7): 783–791.
[PubMed Abstract](#)
- Jones EA, Kinsey SE, English A, et al.: **Isolation and characterization of bone marrow multipotential mesenchymal progenitor cells.** *Arthritis Rheum.* 2002; **46**(12): 3349–3360.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Attar A, Ghalyanchi Langeroudi A, Vassaghi A, et al.: **Role of CD271 enrichment in the isolation of mesenchymal stromal cells from umbilical cord blood.** *Cell Biol Int.* 2013; **37**(9): 1010–1015.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jo C, Kim OS, Park EY, et al.: **Fetal mesenchymal stem cells derived from human umbilical cord sustain primitive characteristics during extensive expansion.** *Cell Tissue Res.* 2008; **334**(3): 423–433.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Allsopp R, Shimoda J, Easa D, et al.: **Long telomeres in the mature human placenta.** *Placenta.* 2007; **28**(4): 324–327.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pacini S, Carnicelli V, Trombi L, et al.: **Constitutive expression of pluripotency-associated genes in mesodermal progenitor cells (MPCs).** *PLoS One.* 2010; **5**(3): e9861.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lu LL, Liu YJ, Yang SG, et al.: **Isolation and characterization of human umbilical cord mesenchymal stem cells with hematopoiesis-supportive function and other potentials.** *Haematologica.* 2010; **95**(10): 1017–1026.
[PubMed Abstract](#)
- Baird DM, Rowson J, Wynford-Thomas D, et al.: **Extensive allelic variation and ultrashort telomeres in senescent human cells.** *Nat Genet.* 2003; **33**(2): 203–207.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stenderup K, Justesen J, Clausen C, et al.: **Aging is associated with decreased maximal life span and accelerated senescence of bone marrow stromal cells.** *Bone.* 2003; **33**(6): 919–26.
[PubMed Abstract](#) | [Publisher Full Text](#)

34. Baxter MA, Wynn RF, Jowitt SN, *et al.*: **Study of telomere length reveals rapid aging of human marrow stromal cells following *in vitro* expansion.** *Stem Cells*. 2004; **22**(5): 675–682.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Bernardo ME, Zaffaroni N, Novara F, *et al.*: **Human bone marrow-derived mesenchymal stem cells do not undergo transformation after long-term *In vitro* culture and do not exhibit telomere maintenance mechanisms.** *Cancer Res*. 2007; **67**(9): 9142–9149.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. De Bari C, Dell'Accio F, Karystinou A, *et al.*: **A biomarker-based mathematical model to predict bone-forming potency of human synovial and periosteal mesenchymal stem cells.** *Arthritis Rheum*. 2008; **58**(1): 240–250.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Kim J, Kang JW, Park JH, *et al.*: **Biological characterization of long-term cultured human mesenchymal stem cells.** *Arch Pharm Res*. 2009; **32**(1): 117–126.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Aviv A, Hunt SC, Lin J, *et al.*: **Impartial comparative analysis of measurement of leukocyte telomere length/DNA content by Southern blots and qPCR.** *Nucleic Acids Res*. 2011; **39**(20): e134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Nawrot TS, Staessen JA, Gardner JP, *et al.*: **Telomere length and possible link to X chromosome.** *Lancet*. 2004; **363**(9408): 507–510.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Nordfjall K, Svenson U, Norrback KF, *et al.*: **Large-scale parent-child comparison confirms a strong paternal influence on telomere length.** *Eur J Hum Genet*. 2009; **18**(3): 385–389.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Friedrich U, Griesse E, Schwab M, *et al.*: **Telomere length in different tissues of elderly patients.** *Mech Ageing Dev*. 2000; **119**(3): 89–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Takubo K, Izumiyama-Shimomura N, Honma N, *et al.*: **Telomere lengths are characteristic in each human individual.** *Exp Gerontol*. 2002; **37**(4): 523–531.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Gadalla SM, Cawthon R, Giri N, *et al.*: **Telomere length in blood, buccal cells, and fibroblasts from patients with inherited bone marrow failure syndromes.** *Aging (Albany NY)*. 2010; **2**(11): 867–74.
[PubMed Abstract](#) | [Free Full Text](#)
44. Kouroupis D, Churchman SM, McGonagle D, *et al.*: **Data for CD146-based cell sorting and telomere length in umbilical cord.** *F1000Research*.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 23 July 2014

<https://doi.org/10.5256/f1000research.4561.r5441>

© 2014 Pacini S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Simone Pacini

Department of Oncology, Transplants and New Advances in Medicine, University of Pisa, Pisa, Italy

I carefully read the manuscript entitled "*The assessment of CD146-based cell sorting and telomere length analysis for establishing the identity of mesenchymal stem cells in human umbilical cord*" from Kouropis *et al.* The aim of the study is verifying the hypothesis that umbilical cord-derived MSCs reside in the CD146-positive fraction, and could be discriminated by their possible longer telomeres. The authors demonstrated that frequency of CD146-positive cells in the culture-initiating suspension was much higher than the frequency of CFU-F confirming that this population also include non MSC-progenitors. Telomere length analysis revealed non-significant differences between CD146-positive and -negative cells, in three of four samples. Thus, the authors correctly conclude that the CD146-positive population was unlikely representing pure native UC-MSCs, and that a prospective isolation of *ex vivo* MSC necessarily should include more antigens to investigate.

My review will be concentrated on some technical aspects regarding conduction of cell sorting and telomere analysis, which should be clarified. My major comments are as follows:

- From the density plots reported in figure 1A, it is not clear if a morphology gate was applied in the SSC/FSC plot, to exclude events of no-cellular origins. A Gate was applied in the dead-cells exclusion plot and in R1 would be counted no-cellular 7-AAD negative events, which could significantly alter the CD146 positive or negative percentages on total living cells. The authors extensively discussed the percentage of CD146-positive frequency in correlation to CFU-F frequency. Thus, the authors should clarify this aspect, and I strongly suggest clearing the flow cytometry data from the non-cellular events.
- As the CD146-positive and -negative populations are not sufficiently separated in the R4 gate, the sorting gates R5 and R6 should be defined by applying isotypic control. Moreover, I suggest reporting the isotypic control plot in the figure, in order to demonstrate the

absence of events in the R5 region.

- Purity, and possible Yield, of sorted cells should be indicated in the results section, in order to reassure the reader of the quality of the sorting process.
- Telomere length has been described as T/S ratio, reported data are close to the value of 1.0 and two of four have values lower than 1.0. How could it be possible to detect telomere repeat copy numbers lower than the single copy gene (36B4)? According to [Cawthon R.M. \(2002\)](#) T/S ratios should be normalized. PCR efficiency could vary gene to gene, and should be evaluated in order to normalize qPCR data with the efficiency correction factor.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 21 Aug 2014

Dimitrios Kouroupis, Foundation for Research and Technology-Hellas, Institute of Molecular Biology and Biotechnology, University of Ioannina, Ioannina, Greece

- *"From the density plots reported in figure 1A, it is not clear if a morphology gate was applied in the SSC/FSC plot, to exclude events of no-cellular origins. A Gate was applied in the dead-cells exclusion plot and in R1 would be counted no-cellular 7-AAD negative events, which could significantly alter the CD146 positive or negative percentages on total living cells. The authors extensively discussed the percentage of CD146-positive frequency in correlation to CFU-F frequency. Thus, the authors should clarify this aspect, and I strongly suggest clearing the flow cytometry data from the non-cellular events."*

We are very grateful to Dr Pacini for his careful reading of our work and for his insightful comments. We believe that Figure 1A that he refers to is in fact Figure 2A as this pertains to our cell sorting experiments from umbilical cord (UC) tissue digests. We can confirm that we did perform FSC/SSC gating to remove anuclear red cells and cellular debris in all our sorting experiments. It was an oversight that this was omitted in the original submission and we thank the reviewer for highlighting this. We have now modified the relevant section of our Materials and Methods as well as Figure 2A to reflect this (gate R1). Additionally, we have added average percentage of cells within R1 out of all events collected into the appropriate Figure Legend.

- *"As the CD146-positive and -negative populations are not sufficiently separated in the R4 gate, the sorting gates R5 and R6 should be defined by applying isotypic control. Moreover, I suggest reporting the isotypic control plot in the figure, in order to demonstrate the absence of events in the R5 region. Response: We appreciate this comment. In our experiments with UC tissue digests, a distinct CD45-CD235a-CD31-CD146+ population was always evident above the isotype control staining."*

We further acknowledge that our original density plots failed to show that our sorting gates were always set apart to ensure maximal sorting purity. We have made new flow cytometry plots for panel 2A to reflect all of these changes (FSC/SSC gating and isotype control panel).

- *“Purity, and possible Yield, of sorted cells should be indicated in the results section, in order to reassure the reader of the quality of the sorting process.”*

We have now provided information on CD146+ and CD146- fractions' cell yields in Figure 2B, left panel. We acknowledge that we did not make clear in our original submission how many sorting experiments we performed in total, we confirm that this is ten for the sorting experiments. Fraction purities post-sort are always >95% using our cell sorter (see [Churchman et al., 2012](#)).

- *“Telomere length has been described as T/S ratio, reported data are close to the value of 1.0 and two of four have values lower than 1.0. How could it be possible to detect telomere repeat copy numbers lower than the single copy gene (36B4)? According to [Cawthon R.M. \(2002\)](#) T/S ratios should be normalized. PCR efficiency could vary gene to gene, and should be evaluated in order to normalize qPCR data with the efficiency correction factor.”*

We appreciate this comment and agree that it is indeed a valid point that we should clarify further. Our method was based upon that of Cawthon R.M. (2002) as stated above and in the text, however when using unexpanded cells sorted from UC the cell numbers were limited because of the amount of tissue processed and inevitable cell losses caused by multiple centrifugation steps prior to cell sorting. Due to this we elected to standardise our telomere length experiments to 1000 sorted cells per fraction (previously overlooked from methods). We accept that we were working at the lower limit of DNA concentrations (acknowledgement of this is now added as a new sentence in Discussion), but believe that with triplicate replicates and extremely comparable standard curves (prepared using immortal cell lines) on each plate for inter-plate comparisons, our work is technically sound, and that our standardisation and technical control was suitable for comparing the differences between the CD146+ and CD146- populations. We did not normalise our T/S ratios and chose to show the data as a direct comparison of sample pairs and therefore some of these values fell below 1. This was also observed in the Cawthon paper; a 'raw' T/S ratio of 0.69 is what he normalised against to give the lowest value of 1 (Figure legend 5). Values below 1 can also be observed in [Guillot et al. \(2007\)](#). We have added an interesting observation that our umbilical cord T/S ratios fall closer to the adult tissues than the foetal tissues in that study. We now present normalized T/S ratios adjusted according to Cawthon R.M. (2002) in Figure 2C and conversion to telomere length in the text. We have also strengthened this figure with new data from a further 3 sorting experiments (new total n=7) and have performed statistical analysis. The CD146+ fraction possesses telomeres with median value 28bp longer than the CD146- fraction, although this was not significant (Wilcoxon).

Competing Interests: No competing interests were disclosed.

Reviewer Report 04 July 2014

<https://doi.org/10.5256/f1000research.4561.r5083>

© 2014 Attar A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Armin Attar

Shiraz University of Medical Sciences, Shiraz, Iran

In the current study, the investigators have used cell sorting techniques as well as telomere length analysis to assess the identity of UC tissue derived mesenchymal stromal cells. The manuscript is scientifically sound and deserves indexation.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research