RESEARCH ARTICLE

Immunoprecipitation and mass spectrometry identify non-cell autonomous Otx2 homeoprotein in the granular and supragranular layers of mouse visual cortex [version 1; peer review: 2 approved]

Namsuk Kim¹, Dario Acampora^{2,3}, Florent Dingli⁴, Damarys Loew⁴, Antonio Simeone^{2,3}, Alain Prochiantz¹, Ariel A. Di Nardo¹

¹CIRB, CNRS UMR 7241 / INSERM U1050, College de France, 11 place Marcelin Berthelot, 75231 Paris Cedex 05, France

²Institute of Genetics and Biophysics, Via Pietro Castellino 111, 80131 Napoli, Italy

³IRCCS Neuromed, 86077 Pozzilli (IS), Italy

⁴Institut Curie, Centre de Recherche, Laboratoire de Spectrométrie de Masse Protéomique, 75248 Paris Cedex 05, France

V1 First published: 30 Jul 2014, 3:178
<https://doi.org/10.12688/f1000research.4869.1>
Latest published: 30 Jul 2014, 3:178
<https://doi.org/10.12688/f1000research.4869.1>

Abstract

Plasticity in the visual cerebral cortex is regulated by the internalization of Otx2 homeoprotein into parvalbumin neurons in cortical layers II/III and IV. However the *Otx2* locus is not active in these neurons and the protein is imported from external sources, including the choroid plexus. Because Otx1 and Otx2 may have redundant functions, we wanted to verify if part of the staining in parvalbumin neurons corresponds to Otx1 transported from cortical layer V neurons. It is demonstrated here that Otx staining in layer IV cells is maintained in *Otx1*-null mice. The immunoprecipitation of extracts from finely dissected granular and supragranular cortex (layers I-IV) gave immunoblots with a band corresponding to Otx2 and not Otx1. Moreover, high-resolution mass spectrometry analysis after immunoprecipitation identifies two peptides within the Otx2 homeodomain. One of these peptides is specific for Otx2 and is not found in Otx1. These results unambiguously establish that the staining in parvalbumin neurons revealed with the anti-Otx2 antibodies used in our previous studies identifies non-cell autonomous Otx2.

Open Peer Review

Approval Status

1

2

version 1

30 Jul 2014

[view](#) [view](#)

1. **Fred Gage**, Salk Institute for Biological Studies, La Jolla, CA, USA
2. **Paola Bovolenta** , Universidad Autonoma de Madrid, Madrid, Spain

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Alain Prochiantz (alain.prochiantz@college-de-france.fr), Ariel A. Di Nardo (ariel.dinardo@college-de-france.fr)

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Région Ile-de-France, the FRM, GRL Program n°2009-00424, ANR grant BRAINEVER n° 11-BLAN-069467, and ERC Advanced Grant HOMEOSIGN n°339379.

Copyright: © 2014 Kim N *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Kim N, Acampora D, Dingli F *et al.* **Immunoprecipitation and mass spectrometry identify non-cell autonomous Otx2 homeoprotein in the granular and supragranular layers of mouse visual cortex [version 1; peer review: 2 approved]** F1000Research 2014, **3**:178 <https://doi.org/10.12688/f1000research.4869.1>

First published: 30 Jul 2014, **3**:178 <https://doi.org/10.12688/f1000research.4869.1>

Introduction

Neural circuits generated during embryonic development are remodeled by environmental inputs during periods of heightened plasticity in postnatal development¹. These critical periods are limited to specific windows of time that are different for each sensory system. In the visual system, the primary visual cortex is subjected to a critical period for ocular dominance plasticity during which connections from a weak or absent eye can be permanently overtaken by the strong eye². The integrated action between inhibitory and excitatory circuits determines critical period onset, with a major role being played by the maturation of fast-spiking parvalbumin (FSPV) interneurons³. We have shown that Otx2 homeoprotein helps determine critical period timing by signaling FSPV cells in mice⁴. Conditional knock-down in heterozygous floxed mice just prior to normal critical period timing is sufficient to delay onset, while cortical infusion of recombinant Otx2 protein accelerates both onset and closure⁴.

Remarkably, Otx2 protein in the cortex is non-cell autonomous. The *Otx2* locus is silent, as shown by PCR, *in situ* hybridization and *Otx2*^{+/GFP} mice, while Otx2 protein is detectable by immunohistochemistry and immunoblot^{4,5}. We have shown that cortical infusion of recombinant Otx2 protein results in specific uptake by FSPV cells, while injection in the retina results in its transport along the visual pathway and into these same cells⁴. Blocking extracellular Otx2 through infusion of antibodies or specific peptides reduces uptake of endogenous Otx2 in FSPV cells^{4,6}. Furthermore, we recently showed that the choroid plexus expresses Otx2 and secretes it into the cerebrospinal fluid⁵. Conditional knockdown of *Otx2* expression in the choroid plexus results in reduced cortical levels of Otx2 protein. Needless to say, all of these approaches that alter Otx2 protein levels in the visual cortex have resulted in changes in cortical plasticity timing⁷.

An outstanding question is whether cortical Otx1 homeoprotein also plays a role in the critical period. Indeed, Otx1 is expressed in the cerebral cortex during development and continues to be expressed by layer V neurons in the adult⁸. It is thus possible that it is secreted by layer V cells and transferred into above granular and supragranular layers where Otx2 protein is detected. Unfortunately, most antibodies (commercial and academic) for Otx1 and Otx2 are pan-Otx thereby ruling out immunohistochemical approaches for conclusive evidence. Since genetic manipulation of the *Otx2* locus has resulted in reduced protein in layer IV visual cortex⁷, we sought confirmation whether Otx1 is also present in these layers in the adult by using *Otx1* knockout mice and proteomic approaches.

Methods

Animals

All experiments were conducted in accordance with European Union Council Directives (86/609/EEC) and conform to Directive 2010/63/EU of the European Parliament. This study falls under project #00704.02 authorized by the French Ministry of Research. Adult male C57Bl/6J mice (Janvier) aged 2 to 3 months were used throughout this study and euthanized by cervical dislocation. *Otx1* knockout mice, in which *Otx1* is replaced with *lacZ* gene, have been previously described¹⁴. Adult (~3 months) male *Otx1*^{LacZ/LacZ} mice were fixed by intracardiac perfusion.

Quantitative RT-PCR

Total RNA from frozen tissue samples were extracted by using RNeasy Mini kits (Qiagen) and were reverse transcribed (~400 ng) with Superscript II and oligo-dT primers (Invitrogen). For real-time PCR, samples were analyzed with a LightCycler Instrument (Roche) and SYBR green (Sigma) detection with the following primers:

Otx2-forward ATTCACGTTTCATGACCAACAG;

Otx2-reverse ATTGACTCCGTATGAGCGGTAT;

Otx1-forward GAACCTTCCTTCTCCGAAATCT;

Otx1-reverse GATCTTCACATCGGACAAATCA;

GAPDH-forward TGACGTGCCGCTGGAGAAAC;

GAPDH-reverse CCGGCATCGAAGGTGGAAGAG.

The experiment was performed in duplicate. For calculating fold expression, gene-to-GAPDH ratios were determined by using the 2^{-ΔΔCt} method referenced to Otx1 expression in the lateral geniculate nucleus (LGN).

Immunoprecipitation and immunoblots

Dissected tissue was lysed in 100 mM Tris-HCl, pH 7.4, 10 mM EDTA, 150 mM NaCl, 1% Triton X-100, and protease inhibitor (Roche) then triturated (22G and 26G needles). Lysates were centrifuged at 16,000 × g for 10 min at 4°C and supernatants were incubated either directly with antibodies or with antibody-coupled Dynabeads (Life Technologies) at 4°C for 16 h. For samples with uncoupled antibodies, Protein A Dynabeads (Life Technologies) were added and incubated at 4°C for 1 h. Proteins were eluted with 2X SDS sample buffer after five washes with lysis buffer. Antibodies were rabbit polyclonal IgG (Abcam ab27478) and anti-Otx2 (rabbit polyclonal, Abcam ab21990) and were coupled to Dynabeads according to manufacturer instructions (Life Technologies). For immunoblots, samples were separated on NuPAGE 4%–12% Bis-Tris precast gels (Invitrogen) and transferred onto PVDF membrane. Membranes were incubated overnight at 4°C with anti-Otx2 (rabbit polyclonal, 1/1,000, Abcam ab21990) and then with HRP-coupled anti-rabbit IgG (1/2,000, GE Healthcare NA934) 1 h at RT.

Immunohistochemistry

For immunostaining, 50 μm floating sections were incubated with anti-Otx2 (rat polyclonal, 1/200, in-house) and biotinylated-WFA (1/100, Sigma L1516) in TBS, 1% Triton-X, 0.2% Tween-20, 10% fetal calf serum, overnight at 4°C. Sections were extensively washed at RT, incubated 2 h at RT with anti-rat Alexa Fluor-488 (Molecular Probes A21208, 1/2,000) and streptavidin Alexa Fluor-546 (Molecular Probes S11225, 1/2,000), washed again and finally mounted in Fluoromount medium (SouthernBiotech). Images were acquired with an Eclipse 90i microscope (Nikon).

Mass spectrometry analysis

After immunoprecipitation, proteins were separated on SDS-PAGE gels (Invitrogen) and stained with colloidal blue staining (LabSafe GEL Blue™ GBiosciences). Gel slices were excised and proteins were reduced with 10 mM DTT prior to alkylation with 55 mM

iodoacetamide. After washing and shrinking the gel pieces with 100% MeCN, in-gel digestion was performed using trypsin (Promega) overnight in 25 mM NH₄HCO₃ at 30°C.

Peptides were extracted and analyzed by nano-LC-MS/MS using an Ultimate 3000 system (Dionex S.A.) coupled to an Orbitrap Fusion mass spectrometer (Q-OT-qIT, Thermo Fisher Scientific). Samples were loaded on a C18 precolumn (300 µm inner diameter × 5 mm; Dionex) at 20 µl/min in 5% MeCN, 0.1% TFA. After a desalting for 3 min, the precolumn was switched on the C18 column (75 µm i.d. × 50 cm, packed with C18 PepMap™, 3 µm, 100 Å; LC Packings) equilibrated in solvent A (2% MeCN, 0.1% HCO₂H). Bound peptides were eluted using a 150 min linear gradient (from 5 to 30% (v/v)) of solvent B (80% MeCN, 0.085% HCO₂H) at a 150 nl/min flow rate and an oven temperature of 40°C. We acquired Survey MS scans in the Orbitrap on the 400–1500 m/z range with the resolution set to a value of 240,000 and a 4 × 10⁵ ion count target. Each scan was recalibrated in real time by co-injecting an internal standard from ambient air into the C-trap. Tandem MS was performed by isolation at 1.6 Th with the quadrupole, HCD fragmentation with normalized collision energy of 35, and rapid scan MS analysis in the ion trap. The MS2 ion count target was set to 104 and the max injection time was 200 ms. Only those precursors with charge state 2–7 were sampled for MS2. The dynamic exclusion duration was set to 60 s with a 10 ppm tolerance around the selected precursor and its isotopes. The instrument was run in top speed mode with 3 s cycles.

Data were acquired using the Xcalibur software (v 3.0.63) and the resulting spectra were interrogated by the Mascot™ Software through Proteome Discoverer (v 1.4.0.288, Thermo Scientific) with the SwissProt Mus musculus database (20140402, 16,671 sequences). We set carbamidomethyl cysteine, oxidation of methionine and N-terminal acetylation as variable modifications. We set specificity of trypsin digestion and allowed 2 missed cleavage sites and we set the mass tolerances in MS and MS/MS to 2 ppm and 0.5 Da, respectively. The resulting Mascot files were further processed by using myProMS (v 3.0)¹⁵ and the estimated false discovery rate (FDR) by automatically filtering the Mascot score of all peptide identifications was less than 0.5%.

Results

Dataset 1. Quantitative PCR and mass spectrometry data of Otx2 homeo protein in the mouse visual cortex

<http://dx.doi.org/10.5256/f1000research.4869.d33384>

qPCR data: This excel file ("brain qpcr otx data") contains two worksheets. One worksheet has the Ct values for GAPDH, Otx2 and Otx1 obtained from extracts of different brain structures (LGN, Visual Cortex, Superior Colliculus and Cerebellum). Also included are the delta-Ct calculations. The second sheet contains the normalisation calculations and corresponding graph of expression values.

Mass spectroscopy data: This Mascot .dat file ("F022130_part.dat") contains the experiment description and the data pertaining to the 2 peptides that match Otx2 protein.

Otx1 locus but not Otx2 locus is active in the mouse visual cortex

We compared the expression of the *Otx1* and *Otx2* loci of various brain regions by performing quantitative PCR (Figure 1A). While both mRNA were detected in thalamic and cerebellar structures, only *Otx1* mRNA was found in the visual cortex. This result confirms the previously reported absence of GFP expression in the visual cortex of *Otx2*^{+/GFP} knockout mice and lack of signal in the visual cortex of wild type mice after *Otx2 in situ* hybridization, even though *Otx2* antibodies label FSPV cells⁴. These cells are enveloped by a dense extracellular matrix called perineuronal nets (PNNs) when localized to layer IV of the cortex (Figure 1B). *Otx1* locus is active in layer V⁸. In immunohistochemical analysis of *Otx1* knockout mouse, almost all *Otx* signal is lost in layer V while signal from *Otx2* protein continues in PNN-labeled cells (Figure 1B). However, we cannot preclude that some *Otx1* is also present in layer IV cells in wild type mice.

Otx2 protein but not Otx1 protein is found in granular and supragranular layers of visual cortex

In order to analyze *Otx* homeoprotein distribution in the visual cortex, we turned to immunoprecipitation (IP) experiments. We first performed IP on whole visual cortex extracts, which showed both *Otx1* and *Otx2* protein (Figure 2A). This result is confirmed by IP of choroid plexus, which strongly expresses *Otx2* but only very weakly expresses *Otx1*. To analyze granular and supragranular content, we dissected and extracted the superior layers of posterior adult mouse cortex (Figure 2B) and performed IP by using cross-linked magnetic beads. Immunoblot analysis detected only *Otx2* but not *Otx1* protein (Figure 2C). This result was confirmed by mass spectrometry analysis on these extracts, which identified 2 *Otx2* peptides (6.6% coverage with 100% specificity, Figure 2D). While this number of peptides is low, it was expected given that *Otx2* is predicted to be poorly ionizable. Furthermore, identification has an error of 1 protein in 20,000 (19,999 are true) with a FDR of less than 0.5% for 2 peptides in our database of 16,671 sequences, thus the chance of misinterpretation is very low. These peptides are 100% specific for *Otx2* and they have the same ions as the reference spectrum from a sample of purified *Otx2* protein. While the second peptide is also specific for *Otx1* and *Crx* homeoproteins, the first peptide is unique for *Otx2* (Figure 2D). These results confirm that the homeoprotein localized in granular and supragranular FSPV cells is indeed *Otx2* and not *Otx1*.

Discussion

The non-cell autonomous activity of homeoprotein transcription factors is now well established. There are clear phenotypes with recently developed *in vivo* single-chain secreted antibodies that neutralize extracellular homeoproteins yet leave intact cell autonomous activities^{9–11}. Non-autonomy can also be demonstrated by comparing mRNA and protein expression. Indeed, the absence of mRNA in presence of the protein argues in favor of non-cell autonomy. However, when the receiving territory is a short distance from the producing territory, one could invoke the possibility of cell migration or mRNA instability to bring into question the reality of homeoprotein transfer.

Figure 1. Expression of Otx1 and Otx2 in adult mouse brain. (A) Analysis of Otx1 and Otx2 expression by quantitative RT-PCR on extracts from lateral geniculate nucleus (LGN), visual cortex (V Cx), superior colliculus (S Col) and cerebellum (Cb). The fold-difference in expression is calculated relative to Otx1 in LGN. The *Otx2* locus is silent in visual cortex. **(B)** Non-cell autonomous Otx2 is found in visual cortex. Immunostaining in wild type mice reveals Otx1/2 cells in layers IV and V of visual cortex, including cells with perineuronal nets (stained by WFA lectin) enriched in layer IV. Staining for Otx2 persists in *Otx1* null mice (*Otx1* KO). Scale bar, 50 μ m.

In the visual system, Otx2 protein is found in the visual cortex far from two potential sources of Otx2 (where the *Otx2* locus is active), namely the eye and the choroid plexus^{5,12}. Indeed, the *Otx2* locus is not active in the adult cerebral cortex as verified by using the *Otx2*^{+/GFP} mouse, quantitative RT-PCR and *in situ* hybridization⁴. In addition, conditional *Otx2* ablation in the choroid plexus reduces its content in FSPV cells, further supporting non-cell autonomy⁵.

However, since most Otx1 and Otx2 antibodies are pan-Otx antibodies, it was still conceivable that some of the protein seen in FSPV cells by immunohistochemistry could correspond to Otx1 expressed in layer V of the cerebral cortex and transferred into PV cells. The present study shows that the staining in layer IV is maintained in the *Otx1* knockout mouse and that IP experiments of layers I-IV give immunoblot bands with expected Otx2 size and can be used to identify Otx2 by mass spectrometry. These results confirm that FSPV cells in granular and supragranular layers of the cerebral cortex only contain non-cell autonomous Otx2 and do not contain Otx1.

It may seem surprising that Otx1 expressed in layer V is not secreted and internalized by FSPV cells. Indeed the protein presents a homeodomain nearly identical to that of Otx2 and thus contains the two sequences necessary for internalization and secretion (for review see⁷). However, previous studies have demonstrated that homeoproteins are transported from the basolateral to the apical side of polarized cells and thus into the axon^{7,13}. Given their polarity and orientation, the pyramidal cells of layer V that express Otx1 are thus very unlikely to release it at the level of FSPV cells. In contrast, the choroid plexus epithelial cells present their apical surface toward the ventricles allowing Otx2 secretion into the cerebral spinal fluid. In conclusion, this study demonstrates that Otx2 is the only non-cell autonomous Otx family protein in the granular and supragranular FSPV cells.

Data availability

F1000Research: Dataset 1. Quantitative PCR and mass spectrometry data of Otx2 homeoprotein in the mouse visual cortex, [10.5256/f1000research.4869.d33384](https://doi.org/10.5256/f1000research.4869.d33384)¹⁶

Figure 2. Otx2 protein in the granular and supragranular layers of adult mouse visual cortex. (A) Immunoblots for Otx1/2 of immunoprecipitation (IP) experiments on extracts from visual cortex and choroid plexus. (B) Diagram of finely dissected region for extracts containing granular (IV) and supragranular (I-III) layers of visual cortex. (C) Immunoblot for Otx1/2 of samples from IP using cross-linked magnetic beads and finely dissected extracts. Choroid plexus (Ch PI) extract was used to control for Otx2 migration velocity. (D) The peptides (red, bold) matching Otx2 protein identified by high-resolution mass spectrometry. Only the homeodomain sequence of Otx2 (amino acids 38–97) is shown. The amino acid differing in Otx1 is highlighted in green, while amino acids that differ in Crx are highlighted in green and in yellow.

Author contributions

NK and AAD carried out experimental work

DA and AS made Otx1 knockout mice

DF carried out the mass spectroscopy experimental work

LD supervised mass spectroscopy and proteomic data analysis

AP and AAD conceived the ideas of the study, designed protocols, and drafted the manuscript

All authors read, critically revised, and approved the final manuscript

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by the Région Ile-de-France, the FRM, GRL Program n°2009-00424, ANR grant BRAINEVER n° 11-BLAN-069467, and ERC Advanced Grant HOMEOSIGN n°339379.

References

1. Hensch TK: **Critical period plasticity in local cortical circuits.** *Nat Rev Neurosci.* 2005; **6**(11): 877–88.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Morishita H, Hensch TK: **Critical period revisited: impact on vision.** *Curr Opin Neurobiol.* 2008; **18**(1): 101–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Hensch TK, Fagiolini M: **Excitatory-inhibitory balance and critical period plasticity in developing visual cortex.** *Prog Brain Res.* 2005; **147**: 115–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Sugiyama S, Di Nardo AA, Aizawa S, *et al.*: **Experience-dependent transfer of Otx2 homeoprotein into the visual cortex activates postnatal plasticity.** *Cell.* 2008; **134**(3): 508–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Spatazza J, Lee HHC, Di Nardo AA, *et al.*: **Choroid-plexus-derived Otx2 homeoprotein constrains adult cortical plasticity.** *Cell Rep.* 2013; **3**(6): 1815–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Beurdeley M, Spatazza J, Lee HHC, *et al.*: **Otx2 binding to perineuronal nets persistently regulates plasticity in the mature visual cortex.** *J Neurosci.* 2012; **32**(27): 9429–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Spatazza J, Di Lullo E, Joliot A, *et al.*: **Homeoprotein signaling in development, health, and disease: a shaking of dogmas offers challenges and promises from bench to bed.** *Pharmacol Rev.* 2013; **65**(1): 90–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Frantz GD, Weimann JM, Levin ME, *et al.*: **Otx1 and Otx2 define layers and regions in developing cerebral cortex and cerebellum.** *J Neurosci.* 1994; **14**(10): 5725–40.
[PubMed Abstract](#)
9. Lesaffre B, Joliot A, Prochiantz A, *et al.*: **Direct non-cell autonomous Pax6 activity regulates eye development in the zebrafish.** *Neural Dev.* 2007; **2**: 2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Wizenmann A, Brunet I, Lam JSY, *et al.*: **Extracellular Engrailed participates in the topographic guidance of retinal axons *in vivo*.** *Neuron.* 2009; **64**(3): 355–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Layalle S, Volovitch M, Mugat B, *et al.*: **Engrailed homeoprotein acts as a signaling molecule in the developing fly.** *Development.* 2011; **138**(11): 2315–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Sugiyama S, Prochiantz A, Hensch TK: **From brain formation to plasticity: insights on Otx2 homeoprotein.** *Dev Growth Differ.* 2009; **51**(3): 369–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Dupont E, Prochiantz A, Joliot A: **Identification of a signal peptide for unconventional secretion.** *J Biol Chem.* 2007; **282**(12): 8994–9000.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Acampora D, Mazan S, Avantaggiato V, *et al.*: **Epilepsy and brain abnormalities in mice lacking the Otx1 gene.** *Nat Genet.* 1996; **14**(2): 218–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Pouillet P, Carpentier S, Barillot E: **myProMS, a web server for management and validation of mass spectrometry-based proteomic data.** *Proteomics.* 2007; **7**(15): 2553–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Kim N, Acampora D, Dingli F, *et al.*: **Quantitative PCR and mass spectrometry data of Otx2 homeoprotein in the mouse visual cortex.** *F1000Research.* 2014.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 08 August 2014

<https://doi.org/10.5256/f1000research.5197.r5636>

© 2014 Bovolenta P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Paola Bovolenta

Centro de Biología Molecular Severo Ochoa, CSIC-UAM, Universidad Autonoma de Madrid, Madrid, Spain

Studies, mostly from A. Prochiantz's laboratory, have demonstrated that the transcription factor (TF) Otx2, produced by external sources such as the choroid plexus, is internalized into parvalbumin neurons of layers II/III and IV of the visual cortex, thereby controlling its plasticity. So far these studies had not ruled out the possibility that part of this effect could be mediated by Otx1, a closely related TF, which may act redundantly with Otx2. In this study, the authors combined the use of *Otx1*^{-/-} mice with immunoprecipitation from layers I-IV extracts and high-resolution mass spectrometry analysis to demonstrate the sole presence of Otx2 peptides in these layers. The study thus demonstrates that Otx2 is the only non-cell autonomous Otx family member functional in fast-spiking parvalbumin cells of layers I-IV of the visual cortex.

This is a clear and well-executed study that solves an existing concern.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 04 August 2014

<https://doi.org/10.5256/f1000research.5197.r5640>

© 2014 Gage F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fred Gage

Laboratory of Genetics, Salk Institute for Biological Studies, La Jolla, CA, USA

The authors address an issue that is unresolved in the literature. Is the Otx found in the cortex, and responsible in part of cortical plasticity, cortical derived Otx1 or choroid plexus derived Otx2? The problem exists because the antibodies used previously can not distinguish between the two. In the present study using *Otx1* KO mice and a variety of measures, the authors prove Otx2 is the protein in the cortical cells and that this therefore is a cell non- autonomous phenomenon. The one question I have relates to figure 1B, which suggests there is greater Otx2 staining in the *Otx1* KO mouse cortex. If this is a reliable observation the authors should comment. The low peptide coverage in the MASS SPEC could be a concern but the authors address that and in addition the other supporting data decrease the concern.

This is a clear, straightforward manuscript that adds important and definitive information to the existing literature

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Comments on this article

Version 1

Author Response () 20 Apr 2015

Ariel Di Nardo, College de France, 11 place Marcelin Berthelot, 75231 Paris Cedex 05, France

We thank Pierre Godement for his insightful reading of our article.

We are well aware that the rat antibody recognizes both Otx1 and Otx2 (as it was produced in our laboratory and kindly provided to Pierre Godement through Marion Wassef). Indeed, the *Otx1* knockout mouse contains a truncated form of Otx1. In a previous study to which Pierre Godement makes reference, protein staining with an Otx1-specific antibody is extremely faint and non-nuclear throughout all cortical layers of the homozygous *Otx1* knockout mouse (Weimann *et al.*, 1999). While it is possible that our antibody also recognizes a truncated form of Otx1, Figure 1B clearly shows that staining in layer V (a bona fide site of *Otx1* expression) is decreased in the knockout as compared to wild type. Furthermore, we continue to find nuclear staining in other cortical layers, contrary to what was found with the Otx1 antibody. This [link](#) contains the raw microscope images and includes an image of the heterozygous *Otx1* mouse visual cortex, which shows similar staining to wild type. (Ratio of Otx cells in infragranular versus supragranular layers in these images is 1.43, 1.46, and 1.11 for WT, het, and homozygote mice, respectively.)

We have not stained for PV (since it was not necessary in the context of this paper) but have used the WFA lectin (that recognizes the PNNs surrounding PV-cells). Contrary to what Pierre Godement

writes, we did not say in Sugiyama *et al.* (2008) that the Otx2 antibody stains "all parvalbumin positive" cells. Figure 1J and page 3 of the article clearly state that 71.4 +/- 0.5% of Otx2-positive cells are PV-positive and that 78.9 +/- 1.7% of the PV-positive cells are Otx2-positive.

One must indeed be cautious with pan-Otx antibodies; genetic evidence for non-cell autonomous cortical Otx2 is paramount. In Sugiyama *et al.* (2008), in addition to using 6 different Otx2 antibodies, we recombined *Otx2* by crossing *Otx2* floxed mouse with a *CamKII-Cre* mouse and lost Otx2 staining in PV-cells. Because CamKII is not expressed in PV-cells, this demonstrated the non-cell autonomous accumulation of Otx2. In addition, we have identified the choroid plexus as a source of Otx2. Recombining the *Otx2* locus specifically in this structure reduced the amount of Otx2 in cortical PV-cells and reopened plasticity in the adult (Spatazza *et al.*, 2013). It is correct that immunoprecipitation (IP) gels show Otx2 with an apparent MW just under 40 kD. This is where Otx2 runs under our conditions (NuPAGE 4%-12%, MES buffer). This [link](#) contains the full western blots corresponding to Figures 2A and 2C. Along with the IPs from visual cortex and choroid plexus, the full blot corresponding to Figure 2A includes an extract from retina (a primary structure strongly expressing Otx2). The velocity for Otx2 is identical for all 3 structures. MW markers and samples from an IP with another pan-Otx antibody (Millipore AB9566) are also included.

No *in situ* studies have shown Otx1 mRNA expression above layer V in the cortex, but the presence of Otx1 protein due to transfer between the layers is a possibility. Our mass spectroscopy analysis of supragranular layer IPs found two peptides corresponding to Otx1 and Otx2, but also one peptide unique for Otx2 (Dataset 1 and Figure 2D). In Figure 2C (supragranular layers), we only find one band migrating with the Otx2 velocity and recognized by Otx antibodies (IP and western blot). It is based on this result that we conclude there is no Otx1 in these layers. The questions of Otx2 amount, location and provenance are not within the scope of this article.

Finally, Pierre Godement raises two additional points. First is that he has made plasticity-related observations concerning axon guidance based on the regulation of extracellular matrix protein expression by Otx2 (Nguyen Ba-Charvet *et al.*, 1998). We agree with this potential impact on cortical plasticity and indeed cited this work in Sugiyama *et al.* (2008).

Second, Pierre Godement proposes that it is Otx1 that regulates the termination of the critical period and not Otx2. We do not exclude that Otx1 and many other proteins participate in critical period regulation but we maintain that Otx2 is a primary factor in the regulation of postnatal and adult cortical plasticity. Whether Otx1 is involved still awaits demonstration.

In conclusion, we sincerely thank Pierre Godement for giving us the opportunity to discuss some points that may not have been explained with enough clarity.

Competing Interests: No competing interests were disclosed.

Reader Comment 15 Apr 2015

Pierre Godement, IGFL, Ecole Normale Supérieure de Lyon, UMR5242 CNRS, France

The data reported here could lead to alternative conclusions in accordance to previous studies.

In the Otx1 knockout mice used here (Acampora *et al.*, 1996) it has been shown that a truncated OTX1 RNA and protein are expressed throughout all layers in the cerebral cortex (Weimann *et al.*, 1999). The positive nuclei in the KO in Figure 1B could correspond to this residual OTX1 protein rather than to OTX2, which cannot be distinguished one from another using the antibody. I also find – but maybe I am in error – that the magnifications (or is it the contrast?) of the left (WT) and the right (KO) panels in Figure 1B appear different (higher mag on the right side) therefore I think the authors could make available raw pictures, which would also provide more detail. One would also like to know if in the KO cells labeled with the OTX2 antibody are all parvalbumin positive as claimed previously (Sugiyama *et al.*, 2008) - from the picture this seems unlikely and this does not support the conclusions.

As for the specificity problem the authors know from already published studies (Fossat *et al.*, 2007) as well as from my own unpublished findings that the rat anti-OTX2 antibody used here labels OTX1 as well as OTX2 and it is too bad that this was not addressed in the original publication from this group (Sugiyama *et al.*, 2008) and specificity checks did not include western blots against the 3 proteins of the OTX family. This would have stimulated the critical sense of the general reader. But neither can they be found in the present one. In the original publication which stimulated these studies (Nothias *et al.*, 1998) it was already evident in my view that the “OTX2” labeling described in the cortex of juvenile mice was in fact OTX1 protein, visible in some large layer 5 pyramidal cells.

Another specific comment I have is from Figure 2A immunoprecipitated OTX2 appears to run at least at 39 kd (this is a 4-10% acrylamide gel) which is very odd – OTX2 runs as a doublet at 32-35 kd and OTX1 runs at 42-44 kd (see for instance Acampora *et al.*, 1998; Fossat *et al.*, 2007). In both this figure and Figure 2C the protein ladder is missing. There also seem to exist some uncertainties as concerns specificity for the Abcam anti-OTX2 antibody (ab21990) used in the IPs as judged from the many users' questions on the manufacturer's website.

The fact remains that so far in situ studies have not detected mRNA for OTX1 above layer 5 in cortex, where OTX immunopositivity is detected – though they did not exclude it (Frantz *et al.*, 1994). But I think it is dangerous to make strong conclusions from negative results especially as concerns in situ studies in adult brain – such labeling might be hard to conclusively demonstrate and OTX mRNAs and proteins are known to be expressed at very variable levels, sometimes quite low.

Nevertheless, the spectrometry data appear to show that some OTX2 protein can be found in cortex. But this would not give a clue as to the amount, location of the protein nor its provenance.

I would like to suggest that it is OTX1 protein, endogenously expressed in several neuronal populations throughout cortex, and which has extensive functional homology to OTX2 (Acampora *et al.*, 1999) which is involved in the phenomena – plasticity, regulation in particular termination of the critical period – which were attributed to transported OTX2. Several years ago we suggested that OTX2 might be involved in restricting axon growth in the embryonic forebrain inside narrow conduits which are bordered by areas with high proteoglycan expression (as is also a particularity of perineuronal nets around parvalbumin-positive, fast-spiking cortical neurons) (Nguyen Ba-Charvet *et al.*, 1998). Conceptually but also mechanistically one might find parallels in both

phenomena (a restriction in growth, and plasticity) – finding transcriptional targets of the OTX proteins shared by these two phenomena is now a subject for future studies.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research