Check for updates

RESEARCH ARTICLE

The proteasome activity reporter GFP-CL1 is degraded by autophagy in the aging model *Podospira anserina* [version 1; peer review: 2 approved, 2 approved with reservations]

Matthias Wiemer, Heinz D. Osiewacz

Institute of Molecular Biosciences and Cluster of Excellence Frankfurt Macromolecular Complexes; Department of Biosciences, J W Goethe University, Frankfurt, 60438, Germany

V1 First published: 30 Sep 2014, 3:230
<https://doi.org/10.12688/f1000research.5337.1>

Latest published: 30 Sep 2014, 3:230
<https://doi.org/10.12688/f1000research.5337.1>

Abstract

The degradation of damaged proteins is an important vital function especially during aging and stress. The ubiquitin proteasome system is one of the major cellular machineries for protein degradation. Health and longevity are associated with high proteasome activity. To demonstrate such a role in aging of *Podospira anserina*, we first analyzed the transcript and protein abundance of selected proteasome components in wild-type cultures of different age. No significant differences were observed. Next, in order to increase the overall proteasome abundance we generated strains overexpressing the catalytic proteasome subunits PaPRE2 and PaPRE3. Although transcript levels were strongly increased, no substantial effect on the abundance of the corresponding proteins was observed. Finally, the analysis of the *P. anserina* strains expressing the sequence coding for the CL1 degron fused to the *Gfp* gene revealed no evidence for degradation of the GFP-CL1 fusion protein by the proteasome. Instead, our results demonstrate the degradation of the CL1-degron sequence via autophagy, indicating that basal autophagy appears to be a very effective protein quality control pathway in *P. anserina*.

Open Peer Review

Approval Status

	1	2	3	4
version 1
30 Sep 2014	view	view	view	view
1. Nektarios Tavernarakis , Foundation for Research and Technology - Hellas, Heraklion, Greece				
2. Suresh Rattan , Aarhus University, Aarhus, Denmark				
3. Thorsten Hoppe , University of Cologne, Cologne, Germany Martina Liebl , University of Cologne, Cologne, Germany				
4. Tilman Grune , Friedrich-Schiller-University Jena, Jena, Germany				

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Heinz D. Osiewacz (osiewacz@bio.uni-frankfurt.de)

Competing interests: No competing interests were disclosed.

Grant information: In part, this work was supported by a grant of the European Commission (Acronym: Proteomage, FP6-51830) to HDO.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2014 Wiemer M and Osiewacz HD. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Wiemer M and Osiewacz HD. **The proteasome activity reporter GFP-CI1 is degraded by autophagy in the aging model *Podospora anserina* [version 1; peer review: 2 approved, 2 approved with reservations]** F1000Research 2014, **3**:230 <https://doi.org/10.12688/f1000research.5337.1>

First published: 30 Sep 2014, **3**:230 <https://doi.org/10.12688/f1000research.5337.1>

Introduction

The degradation of proteins, in particular of those that are damaged or are present in excess, is an important vital function of biological systems and is implicated in several cellular processes such as cell cycle control, proliferation, differentiation, apoptosis and protein quality control¹. Impairments in protein degradation lead to the formation of protein aggregates^{2,3}, promote the aging process^{4,5} and convey the development of neurodegenerative diseases like Alzheimer's or Parkinson's disease⁶. There are two major pathways involved in protein degradation: autophagy and degradation by the ubiquitin proteasome system (UPS)⁷. Autophagy is effective in nutrient recycling and protein degradation. During autophagy proteins or whole organelles are engulfed by a double membrane forming autophagosomes that deliver their cargo to the lysosome in animals and the vacuole in plants and fungi⁸. The UPS consists of a large number of different ubiquitin ligases that act jointly with the proteasome, a multi-protein complex with proteolytic activities. The ubiquitin ligases identify and mark proteins that need to be removed, by formation of a chain of ubiquitin on the target protein⁹. A ubiquitin chain linked at K-48 is recognized by the 26S proteasome¹⁰. The 26S proteasome consists of two subcomplexes, the catalytic 20S core particle and the 19S regulatory particle. The 19S regulatory particle conveys the identification, deubiquitination, unfolding and transport of the substrate into the proteolytic chamber. The core particle is responsible for the degradation of the target proteins. It is composed of four stacked rings, which enclose the proteolytic chamber. The inner rings consist of 7 β -subunits, including the proteolytic active PRE3 (β 1), PUP1 (β 2) and PRE2 (β 5). The three catalytic subunits are the first substrates of the proteasome. Each contains a prosequence that is removed during assembly of the proteasome by an autocatalytic mechanism^{11,12}. The assembled β -subunits are framed by rings of seven α -subunits, blocking the entrance to the proteolytic chamber, if no regulatory particle is bound (reviewed in:¹³).

Previous studies revealed that aging reduces the expression of genes coding for proteasome subunits and the activity of the proteasome in several model systems^{14–16}. Also, several studies indicate a health and lifespan prolonging effects of high proteasome activity. For example, the proteasome activity is elevated in human fibroblast cell cultures derived from centenarians¹⁴ and in the liver of the naked mole rat^{17,18}, a long-living rodent. Moreover, the overexpression of genes coding for proteasome subunit β 1 or β 5 in human fibroblasts was reported to lead to an increase in overall proteasome abundance and activity, resulting in an increased capacity to cope with stress¹⁹. Another component influencing proteasome activity is the proteasome assembly protein UMP1. *Saccharomyces cerevisiae* overexpressing *ScUmp1* shows increased lifespan and viability in response to oxidative stress²⁰. In *S. cerevisiae*, high levels of proteasome subunit ScRPN4 were reported to increase UPS capacity, enhance resistance to proteotoxic stress and increase replicative lifespan²¹. Overall, it appears that the proteasome is a relevant target for aging research. The data suggest that keeping protease activity high during aging can lead to an increase in the healthy lifespan of biological systems.

We use the filamentous ascomycete *Podospora anserina* as a model organism to investigate the mechanisms of aging including the role

of different quality control pathways (for recent reviews see:^{22–24}). In this study we investigated the impact of protein degradation by the UPS and autophagy. Although we could not demonstrate a role of the UPS, we established that the degradation of GFP-CL1 protein, that was expected to be a target of the proteasome, occurred via autophagy.

Materials and methods

P. anserina strains and cultivation

P. anserina was grown on plates with M2 medium (0.25 g/l KH_2PO_4 (Merck Cat# 5099.1000), 0.3 g/l K_2HPO_4 (Roth Cat# P749.1), 0.25 g/l $\text{MgSO}_4 \times 7 \text{H}_2\text{O}$ (Merck Cat# 1.05886.0500), 0.5 g/l urea (Merck Cat# 1.08487.0500) and 10 g/l yellow dextrin (Roth Cat# 6777.1), supplemented with 2.5 mg/l biotin (Serva Cat# 15060), 50 mg/l thiamine (Serva Cat# 36020), 5 mg/l citric acid $\times 1 \text{H}_2\text{O}$ (Sigma-Aldrich Cat# C-0759), 5 mg/l $\text{ZnSO}_4 \times 7 \text{H}_2\text{O}$ (Merck Cat# Z-0625), 1 mg/l $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \times 6 \text{H}_2\text{O}$ (Merck Cat# 1.03861.0250), 2.5 mg/l $\text{CuSO}_4 \times 5 \text{H}_2\text{O}$ (Merck Cat# 2790.1000), 25 mg/l $\text{MnSO}_4 \times 1 \text{H}_2\text{O}$ (Serva Cat# 28405), 50 mg/l $\text{Na}_2\text{MoO}_4 \times 2 \text{H}_2\text{O}$ (Serva Cat# 30207) and 50 mg/l H_3BO_3 (Merck Cat# 100165.5000) after sterilization of the basal medium) or in shaking Erlenmeyer flasks with CM-Medium (70 mM NH_4Cl (Merck Cat# 1.01145.5000), 7.3 mM KH_2PO_4 (Merck Cat# 1.04873.100), 6.7 mM KCl (Merck Cat# 1.04936.1000), 2 mM MgSO_4 (Merck Cat# 1.05886.0500), 1% glucose (Sigma Cat# G-5400), 0.2% tryptone (Otto Nordwald Cat# 211701), 0.2% yeast extract (DIFCO Cat# 0127-07), 5 mM $\text{FeCl}_2 \times 7 \text{H}_2\text{O}$ (Merck Cat# 13478-10-9), 3.5 mM ZnSO_4 (Merck Cat# 108883), 6.2 mM MnCl_2 (Merck Cat# 5934.0100), pH 6.5) under constant light at 27°C. For germination, spores were incubated for two days in the dark on standard cornmeal agar supplemented with 60 mM ammonium acetate (Merck Cat# 1116.1000)²⁵. Pieces of the mycelium derived from germinated spores were transferred on M2 medium to obtain cultures of specific age.

Quantitative Real-time PCR (qRT-PCR)

After germination of spores, pieces of the mycelium were directly used or grown at 27°C and constant light on M2 medium for 13 – 16 days (middle-aged) or 21 – 24 days (senescent), depending on the lifespan of the specific individual, to obtain cultures of specific age. A piece of the growth front was subsequently spread on a fresh M2 plate covered with cellophane (BioRad Cat# 1650963) and grown for 3 days. RNA was extracted with RNA-Plant kit (Machery-Nagel Cat# 740.949.250) and cDNA synthesis was performed using iScript kit (BioRad Cat# 170-8891). After dilution of cDNA to a concentration of 10 ng/ μ l, 20 ng was used per qRT-PCR reaction (IQ SybrGreen SuperMix, BioRad cat# 170-8882). The primers summarized in Table 1 were used to perform the qRT-PCR with three technical replicates per sample. A specific culture was compared to the mean CP of the juvenile cultures. Relative expression was normalized to *PaPorin* with the following formula²⁶.

$$\text{ratio} = \frac{(E_{\text{target}})^{\Delta\text{CP}_{\text{target}}(\text{control-sample})}}{(E_{\text{Porin}})^{\Delta\text{CP}_{\text{Porin}}(\text{control-sample})}}$$

E = PCR-Efficiency; CP = crossing point

Table 1. Primers used for qRT-PCR.

PaNo.	Gene	Primer	Sequence
Pa_5_4560	<i>PaPre3</i>	Pre3 for	ATGGAATTCGGTACATCGGG
		Pre3 rev	GAGGATAACGCCGTCTTTGA
Pa_1_12250	<i>PaPre2</i>	rTPre2for	TTGTTACAGAGCAGGAGCAG
		rTPre2rev	CGATCTTGATGGGGCAGT
Pa_1_24350	<i>PaUmp1</i>	ump1for	TCAACCGAACTCCGATACTC
		ump1rev2	TGGCCTCCCACTGTTTAAAG

Western blot analysis

To obtain total protein extracts, fungi of specific age were spread on a cellophane foil covered M2 surface for 3 days. Proteins were extracted as described in²⁴. Briefly, the mycelia were harvested, ground under liquid nitrogen, mixed with extraction buffer and centrifuged at 14,000 g at 4°C for 10 min. The supernatant was recovered and used for the experiments. The protein extracts were fractionated by 2-phase SDS-PAGE (14% separating gels) according to standard protocol²⁷. Proteins were subsequently transferred to PVDF membranes (Millipore Cat# IPFL00010). Blocking, antibody incubation and washing was performed according to western blot analysis handbook (LIC-OR Bioscience, Bad Homburg, Germany). The following primary antibodies were used: anti-PaPRE2 (rabbit, 1:500 dilution, raised against the specific peptide [H]-WKTKLE-KGEFSNVT-[OH]; Sigma), anti-PaPRE3 (1:2500 dilution, raised against the specific synthetic peptide [H]-LYLPDPTYKVRHEN-[OH]; Sigma), anti-PaPUP1 (rabbit, 1:5000 dilution, raised against the specific peptide Ac-CLKRNYIKPNERT-amid, NEP), anti-HSP60 (mouse, 1:4000 dilution, Biomol, Cat# SPA-807), Anti-GFP (mouse, 1:10000 dilution, Sigma-Aldrich Cat# G6795 RRID:AB_563117). Secondary antibodies conjugated with IRDye 680 (1:15000 dilution, goat anti-mouse 680RD: LI-COR Biosciences Cat# 926-68070 RRID:AB_10956588) or IRDye CW 800 (1:15000 dilution, goat anti-rabbit 800: LIC-OR Biosciences, Cat# 926-3221) were used. After western transfer, the polyacrylamid gels were stained 1 h with coomassie blue as additional loading control.

Cloning procedures and generation of *P. anserina* mutants

The vector pExMthph²⁸ was used as backbone for the generation of *PaPre2*, *PaPre3* and *PaUmp1* overexpression plasmids. For the assembly of pPaPre2Ex1, pExMthph was cut with BamHI and XbaI. The *PaPre2* gene and terminator were amplified with the primers Pre2ExpFor (AAGGATCCATGGACACCCTCGTTGCG; restriction sites are underlined) and Pre2ExpXbaRev (AAAGATCTTG-GCCCTCCTTACTAGAC), cut with BamHI and XbaI and ligated with the backbone. For the generation of pPaPre3Ex1, the *PaPre3* gene and terminator were amplified with the primers PaPre3FwdBam (TTGGATCCATGGAATTCGGTACATCGGG) and PaPre3RevPst (TTCTGCAGCCCAACAACCAGAACCTTTTAC) cut with BamHI and PstI and ligated with the similarly restricted vector pExMthph. pPaUmp1Ex1 was generated by amplification of *PaUmp1*, including the terminator, with the primers PaUmp1FwdBamHI (TTG-GATCCATGGTAAGTTGCAGCCAACC) and PaUmp1RevPstI (TTCTGCAGGCTCCCGTGAGGGCAGGAC), restriction of the product with BamHI and PstI, and ligation into the similarly cut

vector pExMthph. The generation of a *Gfp-Cli* overexpression plasmid was performed by 3 fragment ligation. The *Gpd* promoter from *Aspergillus nidulans*, the *eGfp* gene and the first part of the *Cli*-sequence were amplified by PCR with the plasmid pSM5 (based on pSM2²⁹) as template and with primers eGfp-Pgpd-cl1for (CCTCGAGGTCGACGGTATCGATAAGCTTGATATCGAATT) containing a HindIII restriction site and eGfp-Pgpd-cl1rev (GTG-GCTAGC *GCTGCTGAACCAGTTCCTGCAGGC* CTTGTACA-GCTCGTCCAT), containing half of the *Cli* sequence (italic) optimized for codon usage of *P. anserina* and a Eco47III restriction site. The second half of the *Cli*-sequence was amplified with the *TrpC* terminator in a similar manner with the primers Ttrpc-cl1for (CTTCAGCGAG *CTCAGCCACTTCGTCATCCACCTCTA* ATC-CACTTAACGTTACTGA) containing an Eco53kI restriction sites (underlined) and Ttrpc-cl1rev (CCACCGCGGTGGCGGC-CGCTCTAGAAAGAAGGATTACCTC) containing a XbaI restriction site. The PCR products were ligated into the purified backbone of pSM5, previously cut with HindIII and XbaI. The plasmids were used to transform *P. anserina* wild-type spheroplasts according to²⁴. pSM5 was used to generate a strain expressing *Gfp* without degren sequence. Briefly, mycelium of wild type "s" was blended and the cell wall digested with an enzyme solution. After filtration and concentration of sphaeroblasts by centrifugation, the sphaeroblasts were mixed with 10 µg plasmid DNA. Subsequently, polyethylene glycole (Serva Cat# 33136) was added to the sphaeroblasts. Transformants were selected for hygromycin B (Calbiochem Cat# 400051) resistance and the number of integrations was verified by Southern blot analysis.

Fluorescence microscopy

A piece of 2 day old mycelium was grown on a glass slide with a piece of PASM-medium²⁷ covered with a coverslip and incubated at 27°C for 2 days. Heat stressed samples were incubated at 27°C for 24 h followed by an incubation at 37°C for 24 h. The cover slip with hyphae on it was visualized using a fluorescence microscope (DM LB, Leica, Wetzlar, Germany) with the appropriate excitation and emission filter to detect the GFP signal and a digital camera system (DC500, Leica, Wetzlar, Germany).

Results

Regulation of proteasome components during aging and oxidative stress

In order to address the role of the UPS on aging of *P. anserina*, we investigated the expression of the genes coding for proteolytic subunits *PaPre2* (β5) and *PaPre3* (β1) and of the proteasome assembly

factor *PaUmp1*. First, we determined the abundance of transcript by using total RNA of juvenile, middle-aged and senescent cultures (Figure 1A). No significant changes in mRNA levels were observed in cultures of different age although mRNA abundance of all three genes was slightly reduced in senescent cultures. Next we analyzed protein levels of proteasome subunits in cultures of different age grown in standard growth medium and in medium to which paraquat was added as an inducer of oxidative stress. An increased abundance of mitochondrial HSP60 verified an increase in oxidative stress in senescent and in paraquat treated cultures (Figure 1B). However, no changes in the abundance of subunits PaPRE2, PaPRE3 and

PaPUP1 ($\beta 3$) of the proteasome were observed in the corresponding *P. anserina* cultures. We thus were unable to demonstrate a role of the ubiquitin proteasome system in counteracting adverse effects on cellular proteins in aged cultures and in cultures challenged with exogenous oxidative stress.

Overexpression of *PaPre2* or *PaPre3* does not increase the total amount of proteasome

High activity of the proteasome has been linked to increased health and lifespan^{30–33}. In human cell cultures, the overexpression of the subunits $\beta 5$ and $\beta 1$ was found to increase the overall abundance of the proteasome, as well as its activity and resistance to oxidative stress³⁴. To investigate whether or not such an effect is also observed in *P. anserina*, strains overexpressing the homolog subunits *PaPre2* ($\beta 5$) and *PaPre3* ($\beta 1$) were generated. First, plasmids conveying *PaPre2* and *PaPre3* overexpression were constructed and transformed into *P. anserina* spheroplasts (Table 2). Subsequently, overexpression of the genes was verified by qRT-PCR. *PaPre3* expression was increased by factor 140 to 380 in the respective overexpression strain compared to wild type (Figure 2A). The *PaPre2* overexpression strain exhibited a 94 times higher *PaPre2* expression than the wild type (Figure 2B). In the next step, we evaluated protein levels in the overexpression strains by western blot immunodetection. The analysis of three independent *PaPre3* overexpressors revealed two strains with unchanged protein abundance and one strain (*PaPre3_OEx2*) in which increased PaPRE3 signals occurred (Figure 2C). However, the detected signals are larger or smaller than expected for processed PaPRE3 and probably represent unprocessed PaPRE3 and a degradation product. A strain overexpressing *PaPre2* showed no increase in PaPRE2 abundance compared to the wild type (Figure 2D). Thus, despite the strong increase in mRNA abundance, no substantial change in protein levels of the two investigated proteasome subunits was observed in the generated strains.

The proteasome activity reporter GFP-CL1 is degraded by autophagy

The activity of the UPS is not exclusively defined by the abundance of proteasome subunits but influenced by various factors like ubiquitin ligases, deubiquitin ligases, ATP-level, the regulatory particle, oxidative stress and post-translational modifications. In order to evaluate the efficiency of the UPS during aging of *P. anserina*, we generated two *Gfp-cl1* strains with similar properties and a

Figure 1. Gene expression and protein abundance of proteasome subunits during aging and PQ-stress. (A) The expression of *PaPre2*, *PaPre3* and *PaUmp1* transcripts in juvenile, middle-aged and senescent samples is depicted relative to the juvenile wild type as mean $\pm$ SEM (5 – 7 biological replicates). **(B)** Western blot analysis of 50 μ g total protein extracts of 5 d, 13 d and 21 d old wild type cultures grown on medium with and without the addition of 5 μ M paraquat. Used antibodies are indicated on the right. The polyacrylamid gel stained with coomassie after blotting is used as loading control.

Table 2. Number of ectopic integrations.

Strain	Number of integrations
<i>Gfp</i>	2
<i>Gfp-cl1-1</i>	1
<i>Gfp-cl1-2</i>	1
<i>PaPre2_OEx</i>	1
<i>PaPre3_OEx1</i>	1
<i>PaPre3_OEx2</i>	3
<i>PaPre3_OEx3</i>	2

Figure 2. Overexpression of catalytic 20S subunits *PaPre2* or *PaPre3* does not alter the level of processed protein. The expression of *PaPre3* (A) and *PaPre2* (B) in the respective overexpression strain was examined by qRT-PCR. (C, D) Total protein extracts of *PaPre3* (50 µg) and *PaPre2* (60 µg) overexpression strains were probed with α-*PaPre3* and α-*PaPre2* for the amount of processed proteasome subunits. The polyacrylamide gel stained with coomassie after blotting is displayed as loading control.

Gfp strain. Successful transformation of wild type was verified by Southern blot analysis (Table 2). The introduced genes are under the control of the constitutive *Gpd* promoter of *Aspergillus nidulans*. *Gfp-cl1* codes for a protein containing the CL1 degron sequence fused to GFP. The CL1 sequence is a part of the *Saccharomyces cerevisiae* genome. It was first described in a *ScURA3-CL1* fusion protein, which is unstable in wild type, but stable in strains lacking the ubiquitin ligases *ScUbc6* and *ScUbc7*. Due to these characteristics the CL1 degron has been used to monitor proteasome activity in various species including fly³⁵, mouse³⁶, rat^{37,38} and human cell cultures^{39–44}.

In our work, we investigated the degradation of the CL1 degron fused to GFP. Fluorescence microscopy revealed diffuse fluorescence in whole cells of strains expressing *Gfp-cl1* and *Gfp*, respectively, indicating a cytoplasmic localization (Figure 3A). Significantly, after applying heat stress, the *Gfp-cl1* strain revealed a vacuolar localization of the GFP signal in some parts of the mycelium (Figure 3A). Western-blot analysis revealed two distinct GFP signals in *Gfp-cl1* strains (Figure 3B). One signal corresponds to a protein with a size of 28.8 kDa expected for GFP-CL1 fusion protein while the other has the size of free GFP (26.9 kDa). This result was surprising, because proteasomal degradation should result in total decomposition of GFP-CL1 and provided a first clue for the degradation of the CL1 degron sequence by autophagy since the GFP part remains and

is not, or only slowly, degraded by vacuolar proteases^{17,18}. To verify the degradation of the CL1 degron by autophagy, we generated a *P. anserina* strain lacking *PaAtg1*⁴⁵ and expressing *Gfp-cl1* by crossing of single mutants and selection of the double mutant. *PaATG1* is necessary for autophagy and $\Delta PaAtg1$ -strains are not able to transport proteins to the vacuole for degradation⁴⁵. Western blot analysis revealed that the double mutant contains only GFP-CL1 and no free GFP (Figure 3C), demonstrating that GFP-CL1 is at least partially degraded via autophagy in the wild type of *P. anserina*. This conclusion is supported by the accumulation of green fluorescence in the vacuoles of *Gfp-cl1* overexpressing strains after the induction of autophagy by heat stress (Figure 3A).

Raw data of qRT-PCR and western blot analyses of proteasome subunits and GFP-CL1 degradation in *Podospora anserina*

14 Data Files

<http://dx.doi.org/10.6084/m9.figshare.1177910>

Discussion

In the current study, we investigated the role of the proteasome in aging of *P. anserina*. Contrary to the mammalian model systems, we did not detect significant reduction of transcript or protein

Figure 3. GFP-CL1 is degraded by autophagy. (A) Fluorescence microscopy analysis of *Gfp* and *Gfp-cl1-1* mutant. Fungi were grown for 2d at 27°. Heat stressed samples were incubated at 37°C for the last 24 h. (B) Western-blot analysis of 60 μg total protein extracts from *Gfp-CL1*-1 and *Gfp* strains. Proteins were detected by western-blot-analysis with α-GFP antibody. The corresponding polyacrylamid gel was stained with coomassie after blotting as loading control. (C) Western-blot analysis of 45 μg total protein extracts from *Gfp-cl1-2* strains and from a *Gfp-cl1-2/ΔPaAtg1* double mutant. Proteins were detected by western-blot-analysis with α-GFP antibody. The polyacrylamid gel was stained with coomassie after blotting.

levels of proteasome subunits during aging⁴⁶. Moreover, attempts to modulate the abundance of selected proteasomal subunits failed, although transcript abundance was strongly increased in the generated overexpression strains. It appears that in *P. anserina* the biosynthesis of the investigated proteasome subunits is under a strong post-transcriptional control.

One aim of our study was the development of an assay to study proteasomal activity. In other systems such assays are based on the microscopic monitoring of fluorescence changes resulting from the degradation of a reporter protein, termed degron, which is fused to GFP. The degron becomes rapidly ubiquitinated and subsequently the whole fusion protein is delivered to the proteasome where it is degraded. A widely used degron is CL1 derived from *S. cerevisiae* and consisting of 15 hydrophobic amino acids. Although this sequence was successfully used to detect proteasome activity in a wide range of organisms including yeast⁴⁷, fly³⁵, mouse³⁶, rat^{37,38} and human cell cultures^{39–44}, our experiments did not reveal a clear degradation of the whole GFP-CL1 fusion protein as it would be expected for degradation by the proteasome. Beside other reasons, it may be that CL1 is not recognized by the *P. anserina* ubiquitination system and thus does not constitute a functional degron. On the other hand, the sequence may be recognized by both the UPS

and the autophagy machinery. Under the investigated conditions autophagy may be by far more efficient than ubiquitination and proteasomal degradation. An overlap of UPS and autophagy substrates has been shown previously^{48–51}. The degradation of this reporter by autophagy may indeed be a severe problem for the establishment of a reporter gene based proteasome activity assay in filamentous fungi because they seem to be characterized by high level of basal autophagy. In *Aspergillus oryzae*, mitochondria, peroxisomes and nuclei of basal hyphae are degraded during normal growth in an autophagy dependent manner to use the nutrients to support growth⁵². Previous work in *P. anserina* also detected a high basal autophagy level under non-starved standard growth conditions⁴⁵. On the other hand, basal autophagy in yeast and mammalian cell cultures appears to be low^{53,54}. Another complication of the system may be that, as previously shown in *Caenorhabditis elegans* and neuronal rat cells, CL1 fused to GFP can form toxic aggregates if the expression level exceeds the capacity of the degradation system⁵⁵. In our experiments this latter problem appears not to be valid since the fluorescence signal is distributed throughout the cell although we detected small condensed signals in some cells, which could indicate the formation of protein aggregates. Since such aggregates were only very small spots compared to those demonstrated in the mentioned studies with *C. elegans* and rat cells, the formation of

toxic GFP-CL1 aggregates appears to be negligible under the chosen expression conditions.

Data availability

figshare: Raw data of qRT-PCR and western blot analyses of proteasome subunits and GFP-CL1 degradation in *Podospora anserina*. DOI: [10.6084/m9.figshare.1177910](https://doi.org/10.6084/m9.figshare.1177910)⁵⁶

Author contributions

MW performed experiments. HDO initiated and supervised this study. HDO and MW wrote the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

In part, this work was supported by a grant of the European Commission (Acronym: Proteomage, FP6-51830) to HDO.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank Dr. J. Servos (Frankfurt) for providing plasmids used in this study and Dr. A. Hamann for discussion of data.

References

- Naujokat C, Hoffmann S: **Role and function of the 26S proteasome in proliferation and apoptosis.** *Lab Invest.* 2002; **82**(8): 965–980.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cummings CJ, Reinstein E, Sun Y, *et al.*: **Mutation of the E6-AP ubiquitin ligase reduces nuclear inclusion frequency while accelerating polyglutamine-induced pathology in SCA1 mice.** *Neuron.* 1999; **24**(4): 879–892.
[PubMed Abstract](#) | [Publisher Full Text](#)
- He L, Lu XY, Jolly AF, *et al.*: **Spongiform degeneration in *mahoganoid* mutant mice.** *Science.* 2003; **299**(5607): 710–712.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morimoto RI, Cuervo AM: **Proteostasis and the aging proteome in health and disease.** *J Gerontol A Biol Sci Med Sci.* 2014; **69**(Suppl 1): S33–S38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lindner AB, Demarez A: **Protein aggregation as a paradigm of aging.** *Biochim Biophys Acta.* 2009; **1790**(10): 980–996.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Powers ET, Morimoto RI, Dillin A, *et al.*: **Biological and chemical approaches to diseases of proteostasis deficiency.** *Annu Rev Biochem.* 2009; **78**: 959–991.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lilienbaum A: **Relationship between the proteasomal system and autophagy.** *Int J Biochem Mol Biol.* 2013; **4**(1): 1–26.
[PubMed Abstract](#) | [Free Full Text](#)
- Kroemer G, Mariño G, Levine B: **Autophagy and the integrated stress response.** *Mol Cell.* 2010; **40**(2): 280–293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hershko A, Ciechanover A: **The ubiquitin system.** *Annu Rev Biochem.* 1998; **67**: 425–479.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pickart CM, Fushman D: **Polyubiquitin chains: polymeric protein signals.** *Curr Opin Chem Biol.* 2004; **8**(6): 610–616.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Seemuller E, Lupas A, Baumeister W: **Autocatalytic processing of the 20S proteasome.** *Nature.* 1996; **382**(6590): 468–471.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen P, Hochstrasser M: **Autocatalytic subunit processing couples active site formation in the 20S proteasome to completion of assembly.** *Cell.* 1996; **86**(6): 961–972.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jung T, Catalgol B, Grune T: **The proteasomal system.** *Mol Aspects Med.* 2009; **30**(4): 191–296.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chondrogianni N, Petropoulos I, Franceschi C, *et al.*: **Fibroblast cultures from healthy centenarians have an active proteasome.** *Exp Gerontol.* 2000; **35**(6–7): 721–728.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Carrard G, Bulteau AL, Petropoulos I, *et al.*: **Impairment of proteasome structure and function in aging.** *Int J Biochem Cell Biol.* 2002; **34**(11): 1461–1474.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chondrogianni N, Gonos ES: **Proteasome dysfunction in mammalian aging: steps and factors involved.** *Exp Gerontol.* 2005; **40**(12): 931–938.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pérez VI, Buffenstein R, Masamsetti V, *et al.*: **Protein stability and resistance to oxidative stress are determinants of longevity in the longest-living rodent, the naked mole-rat.** *Proc Natl Acad Sci U S A.* 2009; **106**(9): 3059–3064.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rodriguez KA, Osmulski PA, Pierce A, *et al.*: **A cytosolic protein factor from the naked mole-rat activates proteasomes of other species and protects these from inhibition.** *Biochim Biophys Acta.* 2014; **1842**(11): 2060–2072.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chondrogianni N, Stratford FL, Trougakos IP, *et al.*: **Central role of the proteasome in senescence and survival of human fibroblasts: induction of a senescence-like phenotype upon its inhibition and resistance to stress upon its activation.** *J Biol Chem.* 2003; **278**(30): 28026–28037.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen Q, Thorpe J, Dohmen JR, *et al.*: **Ump1 extends yeast lifespan and enhances viability during oxidative stress: central role for the proteasome?** *Free Radic Biol Med.* 2006; **40**(1): 120–126.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kruegel U, Robison B, Dange T, *et al.*: **Elevated proteasome capacity extends replicative lifespan in *Saccharomyces cerevisiae*.** *PLoS Genet.* 2011; **7**(9): e1002253.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fischer F, Hamann A, Osiewacz HD: **Mitochondrial quality control: an integrated network of pathways.** *Trends Biochem Sci.* 2012; **37**(7): 284–292.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Osiewacz HD, Brust D, Hamann A, *et al.*: **Mitochondrial pathways governing stress resistance, life, and death in the fungal aging model *Podospora anserina*.** *Ann NY Acad Sci.* 2010; **1197**: 54–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Osiewacz HD, Hamann A, Zintel S: **Assessing organismal aging in the filamentous fungus *Podospora anserina*.** *Methods Mol Biol.* 2013; **965**: 439–462.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Esser K: ***Podospora anserina*.** In: *Handbook of Genetics*, King, R.C., ed, Plenum Press, New York. 1974; pp. 531–551.
- Pfaffl MW: **A new mathematical model for relative quantification in real-time RT-PCR.** *Nucleic Acids Res.* 2001; **29**(9): e45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brust D, Hamann A, Osiewacz HD: **Deletion of *PaAif2* and *PaAmd2*, two genes encoding mitochondrial AIF-like oxidoreductases of *Podospora anserina*, leads to increased stress tolerance and lifespan extension.** *Curr Genet.* 2010; **56**(3): 225–235.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Averbeck NB, Borghouts C, Hamann A, *et al.*: **Molecular control of copper homeostasis in filamentous fungi: increased expression of a metallothionein gene during aging of *Podospora anserina*.** *Mol Gen Genet.* 2001; **264**(5): 604–612.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pöggeler S, Masloff S, Hoff B, *et al.*: **Versatile EGFP reporter plasmids for cellular localization of recombinant gene products in filamentous fungi.** *Curr Genet.* 2003; **43**(1): 54–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tonoki A, Kuranaga E, Tomioka T, *et al.*: **Genetic evidence linking age-dependent attenuation of the 26S proteasome with the aging process.** *Mol Cell Biol.* 2009; **29**(4): 1095–1106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vilchez D, Boyer L, Morante I, *et al.*: **Increased proteasome activity in human embryonic stem cells is regulated by PSMD11.** *Nature.* 2012; **489**(7415): 304–308.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Katsiki M, Chondrogianni N, Chinou I, *et al.*: **The olive constituent oleuropein exhibits proteasome stimulatory properties *in vitro* and confers life span extension of human embryonic fibroblasts.** *Rejuvenation Res.* 2007; **10**(2): 157–172.
[PubMed Abstract](#) | [Publisher Full Text](#)

33. Vilchez D, Morante I, Liu Z, *et al.*: **RPN-6 determines *C. elegans* longevity under proteotoxic stress conditions.** *Nature*. 2012; **489**(7415): 263–268.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Chondrogianni N, Tzavelas C, Pemberton AJ, *et al.*: **Overexpression of proteasome beta5 assembled subunit increases the amount of proteasome and confers ameliorated response to oxidative stress and higher survival rates.** *J Biol Chem*. 2005; **280**(12): 11840–11850.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Pandey UB, Nie Z, Batlevi Y, *et al.*: **HDAC6 rescues neurodegeneration and provides an essential link between autophagy and the UPS.** *Nature*. 2007; **447**(7146): 859–863.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Liu Y, Hettinger CL, Zhang D, *et al.*: **The proteasome function reporter GFPu accumulates in young brains of the APPsw/PS1dE9 Alzheimer's disease mouse model.** *Cell Mol Neurobiol*. 2014; **34**(3): 315–322.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Tian Z, Wang C, Hu C, *et al.*: **Autophagic-lysosomal inhibition compromises ubiquitin-proteasome system performance in a p62 dependent manner in cardiomyocytes.** *PLoS One*. 2014; **9**(6): e100715.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Ranek MJ, Terpstra EJ, Li J, *et al.*: **Protein kinase g positively regulates proteasome-mediated degradation of misfolded proteins.** *Circulation*. 2013; **128**(4): 365–376.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Bence NF, Sampat RM, Kopito RR: **Impairment of the ubiquitin-proteasome system by protein aggregation.** *Science*. 2001; **292**(5521): 1552–1555.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Menendez-Benito V, Verhoef LG, Masucci MG, *et al.*: **Endoplasmic reticulum stress compromises the ubiquitin-proteasome system.** *Hum Mol Genet*. 2005; **14**(19): 2787–2799.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Bett JS, Cook C, Petrucelli L, *et al.*: **The ubiquitin-proteasome reporter GFPu does not accumulate in neurons of the R6/2 transgenic mouse model of Huntington's disease.** *PLoS One*. 2009; **4**(4): e5128.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Nonaka T, Hasegawa M: **A cellular model to monitor proteasome dysfunction by alpha-synuclein.** *Biochemistry*. 2009; **48**(33): 8014–8022.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Tydlacka S, Wang CE, Wang X, *et al.*: **Differential activities of the ubiquitin-proteasome system in neurons versus glia may account for the preferential accumulation of misfolded proteins in neurons.** *J Neurosci*. 2008; **28**(49): 13285–13295.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Hope AD, de Silva R, Fischer DF, *et al.*: **Alzheimer's associated variant ubiquitin causes inhibition of the 26S proteasome and chaperone expression.** *J Neurochem*. 2003; **86**(2): 394–404.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Knuppertz L, Hamann A, Pampaloni F, *et al.*: **Identification of autophagy as a longevity-assurance mechanism in the aging model *Podospora anserina*.** *Autophagy*. 2014; **10**(5): 822–834.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Chondrogianni N, Sakellari M, Lefaki M, *et al.*: **Proteasome activation delays aging *in vitro* and *in vivo*.** *Free Radic Biol Med*. 2014; **71**: 303–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Gilon T, Chomsky O, Kulka RG: **Degradation signals for ubiquitin system proteolysis in *Saccharomyces cerevisiae*.** *EMBO J*. 1998; **17**(10): 2759–2766.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. De Domenico I, Vaughn MB, Li L, *et al.*: **Ferroportin-mediated mobilization of ferritin iron precedes ferritin degradation by the proteasome.** *EMBO J*. 2006; **25**(22): 5396–5404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Webb JL, Ravikumar B, Atkins J, *et al.*: **Alpha-Synuclein is degraded by both autophagy and the proteasome.** *J Biol Chem*. 2003; **278**(27): 25009–25013.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Kabuta T, Suzuki Y, Wada K: **Degradation of amyotrophic lateral sclerosis-linked mutant Cu,Zn-superoxide dismutase proteins by macroautophagy and the proteasome.** *J Biol Chem*. 2006; **281**(41): 30524–30533.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Ravikumar B, Duden R, Rubinstein DC: **Aggregate-prone proteins with polyglutamine and polyalanine expansions are degraded by autophagy.** *Hum Mol Genet*. 2002; **11**(9): 1107–1117.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Shoji JY, Kikuma T, Arioka M, *et al.*: **Macroautophagy-mediated degradation of whole nuclei in the filamentous fungus *Aspergillus oryzae*.** *PLoS One*. 2010; **5**(12): e15650.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Abeliovich H, Klionsky DJ: **Autophagy in yeast: mechanistic insights and physiological function.** *Microbiol Mol Biol Rev*. 2001; **65**(3): 463–479.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Mizushima N, Yoshimori T, Levine B: **Methods in mammalian autophagy research.** *Cell*. 2010; **140**(3): 313–326.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Link CD, Fonte V, Hiestler B, *et al.*: **Conversion of green fluorescent protein into a toxic, aggregation-prone protein by C-terminal addition of a short peptide.** *J Biol Chem*. 2006; **281**(3): 1808–1816.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Wiemer M, Osiewicz HD: **Raw data of qRT-PCR and western blot analyses of proteasome subunits and GFP-CL1 degradation in *Podospora anserina*.** *figshare*. 2014.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 29 December 2014

<https://doi.org/10.5256/f1000research.5697.r6929>

© 2014 Grune T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tilman Grune

Department of Nutritional Toxicology, Friedrich-Schiller-University Jena, Jena, Germany

This is an interesting investigation studying proteostasis in *Podospora* during aging. I am not a specialist in *Podospora*, but can judge the work on aging and the proteolytic systems.

I believe that this paper is lacking one substantial part: this is the measurement of activities of the proteolytic systems. Without this it is hard to draw any conclusion.

Major points:

- In the introduction the authors refer to references reporting "reduced expression of genes for proteasome subunits and the activity of the proteasome" in other aging models. This is true, but the authors should note that the only consistent finding in the literature is a reduced proteasomal activity. Other authors also reported minimal or no changes in proteasomal gene expression, but a reduced activity of the proteasome. Therefore, the measurement of the proteasomal activity is of utmost importance.
- The conclusion (page 4) that the proteasome has no "role...counteracting adverse effects on cellular proteins" is not supported by the data at all, without activity measurement, inhibition of the proteasome and so on.
- In Fig.2 the authors refer to "...processed protein" What is this? The number of translation products produced? How can this be concluded, since the authors just study the steady state level?
- Another statement on page 5: "...proteasomal degradation should result in a total decomposition of GFP-CL1". In the light of the work of DeMartino/Thomas ([Liu et al. 2003](#)), I am not sure, whether this is true under all conditions.
- The knockout of ATG1 will reduce the macroautophagy process. (Are there alternative pathways in *Podospora*?) Is in the ATG1 k.o. the proteasome still active? Activities should be measured here, again.

- The formation of GFP-based aggregates is discussed by the authors. However, also here it is required to show that the proteasomal system is active, since low amounts of protein aggregates might be able to block/inhibit substantial parts of the proteasome.

Minor points:

- In the introduction the authors refer to autophagy as parts of the cell “engulfed by a double membrane”. This is only true for macroautophagy, not for chaperone-mediated autophagy and microautophagy. I do not know, whether these other forms exist in *Podospora*, but since the authors in their argumentation are jumping in the whole paper from model to model, this should be corrected.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 16 Jan 2015

Heinz Osiewacz, J W Goethe University, Frankfurt, Germany

We thank you very much for careful reading of the manuscript and your comments and suggestions. We agree that it would be very helpful to include the biochemical *in vitro* assay to measure proteasome activity in protein extracts. Ideally, both *in vitro* and *in vivo* studies about the activity of the proteasome should be performed. In fact, the isolation of functional proteasomes in filamentous fungi with a rigid cell wall appears to be a problem. We therefore tried to use the *in vivo* degradation of the CL1-degron in our study. We would like to stress here that we do not state that protein degradation via the proteasome does NOT play a role in *P. anserina*. We just cannot demonstrate this with the available methods, and this is what we wanted to report. Apart from this, we identify that the GFP-CL1-degron is degraded to GFP in an ATG1-dependent manner. The resulting free GFP is resistant to degradation. The knock-out of an essential autophagy component, like ATG1, is the accepted method to demonstrate a role of autophagy. We show that GFP-fluorescence ends up in the vacuole. These are new findings that, together with our recent work ¹, stress the importance of autophagy in the *Podospora anserina* system.

References

1. Knuppertz L, Hamann A, Pampaloni F, Stelzer E, et al.: Identification of autophagy as a longevity-assurance mechanism in the aging model *Podospora anserina*. *Autophagy*. 2014; **10** (5): 822-834 [PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)

Competing Interests: No competing interests were disclosed.

Reviewer Report 09 December 2014

<https://doi.org/10.5256/f1000research.5697.r6891>

© 2014 Hoppe T et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Thorsten Hoppe

Department of C elegans Genetics and Development, University of Cologne, Cologne, Germany

Martina Liebl

Department of C elegans Genetics and Development, University of Cologne, Cologne, Germany

It is well established that proteasomal degradation declines during aging in different cell culture systems and model organisms. Therefore, it is of general interest whether such activity changes are conserved throughout evolution. Here, the authors analyzed mRNA levels of the 20S proteasomal core subunits PaPre2, PaPre3, and the proteasomal assembly factor PaUmp1 as well as protein levels of PaPRE2, PaPRE3, and PaPUP1 in the fungus *P. anserina*. The experiments are convincing and well described. Surprisingly, the authors did not find any age-dependent differences with respect to the investigated proteasomal factors. However, as measuring proteasomal activity directly did not work, it remained unclear whether a decline in protein turnover is directly linked to aging in this model organism. As highlighted by the authors, other factors including ubiquitin ligases, deubiquitylation enzymes, the 19S regulatory particle, or post-translational modifications of these factors might be even more limiting for ubiquitin-dependent degradation and regulated with age in the fungus. In this context the authors could have discussed the finding that in *C. elegans* and human stem cells overexpression of the regulatory subunit RPN-6 is sufficient to reduce the amount of toxic protein aggregates and increase life span (Vilchez *et al.* 2012).

The authors state that protein levels of PaPUP1 did not change during aging and paraquat treatment. However, western blot analysis of figure 1B shows increased PaPUB1 levels in 21 day old cultures that were treated with paraquat compared to either paraquat treated young cultures or untreated old cultures. This does not necessarily reflect an age- or stress-dependent increase in PaPUB1 levels in *P. anserina*, but could be explained by a selective survival of fungi with higher PaPUB1 levels under toxic stress conditions.

This study nicely confirms the impression that *P. anserine* exhibits high basal rates of autophagy in contrast to other organisms and thus might serve as an excellent model organism for studying the role of autophagy under different physiological conditions. Given the elevated level of autophagy, this organism however might not be optimal for the investigation of age-dependent changes in the ubiquitin/proteasome system that is relevant for higher organisms.

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 03 December 2014

<https://doi.org/10.5256/f1000research.5697.r6542>

© 2014 Rattan S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Suresh Rattan

Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark

This is a very well written paper dealing with the determination of the proteasome activity in the ageing model *Podospora*. The experimental work is technically sound and is performed diligently. Although the results obtained, especially with respect to the lack of change in the proteasomal activity during ageing of *Podospora*, appear to be contrary to the ones reported for mammalian systems and for *Drosophila*, it does not decrease the significance of this study which once again underlies the fact that age-related changes can be both universal (public) or specific (private). Actually, this is the important point that needs to be discussed in this paper as to why such differences are observed (see some papers by George Martin on public and private mechanisms of ageing). Authors should therefore comment on the usefulness and limitations of using model systems such as *Podospora* in the context of modulating human ageing and longevity, and for testing/screening potential ageing-modulatory strategies. Furthermore, it will be useful if the implications of enhanced autophagy-related pathways in the context of ageing are discussed a bit more.

On a minor note, just to be up-to-date with the literature on proteasome activity during ageing of various systems (refs 14-16), authors may include and cite a paper on *Drosophila* ([Hansen et al., 2012](#)) which deals with the sex-specific differences in proteasome activity changes during ageing etc.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 16 Jan 2015

Heinz Osiewacz, J W Goethe University, Frankfurt, Germany

We appreciate very much your careful reading of the manuscript and the submitted comments. We totally agree that work on model systems has to be taken with caution. But this is inherent to ANY model including whole organisms (both lower as well as higher systems like mice) as well as *in vitro* systems like mammalian cell cultures. Nevertheless, models can provide important general information about the aging process in humans. This has been demonstrated in *P. anserina* which in the past has proven to generate pioneer knowledge on age-related processes like mitochondrial DNA instabilities ^{1,2}, the role of

mitochondrial dynamics³, or mitochondrial ultrastructure⁴.

However, we would like to emphasize that the aim of our work reported in this *F1000Research* paper was to evaluate the role of the proteasome in *P. anserina*. Following different approaches, we could not demonstrate such a role. We are aware that we do not exclude a role of the proteasome (in fact, we believe that it also plays a role). The key information we wanted to pass on to the scientific community is that autophagy acts on a substrate, the CL1 degren, that in other systems has been shown to be degraded by the proteasome and that autophagy plays is highly active in the system. We strongly believe that the work we presented here exactly meets the aims and scope of *F1000Research*.

References

1. Esser K, Tudzynski P, Stahl U, Kuck U: A Model to Explain Senescence in the Filamentous Fungus *Podospira anserina* by the Action of Plasmid Like DNA. *Molec gen Genet*. 1980; **178** (1): 213-216 [Reference Source](#)
2. Stahl U, Lemke PA, Tudzynski P, Kuck U, et al.: Evidence for plasmid like DNA in a filamentous fungus, the ascomycete *Podospira anserina*. *Molec Gen Genet*. 1978; **162** (3): 341-343 [PubMed Abstract](#)
3. Scheckhuber CQ, Erjavec N, Tinazli A, Hamann A, et al.: Reducing mitochondrial fission results in increased life span and fitness of two fungal ageing models. *Nat Cell Biol*. 2007; **9** (1): 99-105 [PubMed Abstract](#) | [Publisher Full Text](#) | [Reference Source](#)
4. Daum B, Walter A, Horst A, Osiewacz HD, et al.: Age-dependent dissociation of ATP synthase dimers and loss of inner-membrane cristae in mitochondria. *Proc Natl Acad Sci USA*. 2013; **110** (38): 15301-15306 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#) | [Reference Source](#)

Competing Interests: No competing interests were disclosed.

Reviewer Report 14 October 2014

<https://doi.org/10.5256/f1000research.5697.r6268>

© 2014 Tavernarakis N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nektarios Tavernarakis

Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology - Hellas, Heraklion, Crete, Greece

The manuscript aims to address the role of the proteasomal activity in aging and health span, using *Podospira anserina* as a model organism. The authors show that gene expression and protein abundance of proteasome subunits and proteasome assembly factor remain stable during

aging and oxidative stress. The authors used strains overexpressing the catalytic proteasome subunits PaPRE2 and PaPRE3 to increase proteasomal activity, but found no difference in lifespan, suggesting that proteasome activity does not affect lifespan. The study would benefit by including proteasomal activity controls to confirm activity increase. This is particularly relevant since the positive effect of the proteasome in lifespan, senescence and healthspan has been convincingly demonstrated in other systems. The authors suggest that the reporter they use, GFP-Cl1, is processed by the autophagic machinery. It would be interesting to examine whether overexpression of this construct causes aggregation of GFP which in turn triggers autophagy that subsequently clears the aggregates which cannot be cleared by the proteasome. As the authors state in their discussion toxic aggregates could be formed and this possibility should be experimentally evaluated. For instance the authors should evaluate whether ERAD (endoplasmic reticulum associated protein degradation) takes place, since accumulation of protein aggregates could initiate ER stress and ERAD. It is known that under various stress conditions proteasomal activity is enhanced. Perhaps the authors should try other conditions in addition to oxidative stress (ER stress, heat stress etc.).

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research