EDITORIAL

It's never too late to publish an abandoned trial [version 1; peer review: not peer reviewed]

Celia Shiles¹, Julia Sinclair²

¹Southampton General Hospital, Southampton, Hampshire, SO16 6YD, UK

²Faculty of Medicine, University of Southampton, Southampton, Hampshire, SO17 1BJ, UK

V1 First published: 15 May 2015, 4:120
<https://doi.org/10.12688/f1000research.6519.1>
Latest published: 15 May 2015, 4:120
<https://doi.org/10.12688/f1000research.6519.1>

Abstract

It is estimated that half of all trials have never been published which can lead to patients being denied the most effective treatment and being exposed to unnecessary side effects. Furthermore the trial participants have been misinformed since the trial results have not contributed to the care of future patients.

However the non-publication of trials is often not due to a deliberate decision to cover up results. Commonly in academia it is due to more understandable reasons such as researchers having busy clinical posts, moving onto other more demanding projects, changing research areas or starting a family. This is called the “file drawer” problem.

The examples in this editorial demonstrate that it is possible to go back, even decades later, and make the results available to inform future evidence based medicine. We call on others to look into their “file drawer” for unpublished trials.

Keywords

Abandoned trials, Science publishing, Meta-analyses

Not Peer Reviewed

This article is an Editorial and has not been subject to external peer review.

Any comments on the article can be found at the end of the article.

This article is included in the **All trials matter** collection.

Corresponding author: Celia Shiles (c.j.shiles@gmail.com)

Competing interests: No competing interests were disclosed.

Grant information: Celia Shiles is supported by the Wessex Deanery and Julia Sinclair by the Higher Education Funding Council.

Copyright: © 2015 Shiles C and Sinclair J. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Shiles C and Sinclair J. **It's never too late to publish an abandoned trial [version 1; peer review: not peer reviewed]** F1000Research 2015, 4:120 <https://doi.org/10.12688/f1000research.6519.1>

First published: 15 May 2015, 4:120 <https://doi.org/10.12688/f1000research.6519.1>

Article

The process by which a medical intervention is thought to be effective is now well described. Data from trials are collected together after a systematic search of the literature. These are then pooled in a meta-analysis to give an overall indication of efficacy. These results are then collated to inform policy documents, such as guidelines produced by the National Institute for Health and Care Excellence (NICE) in England and Wales. However any meta-analysis is only as good as the trials which it includes. Missing data such as that which comes from unpublished trials is a serious threat to the integrity of any systematic review.

It is estimated that half of all trials have never been published and the incidence of non-publication is roughly similar whether a trial is funded by industry, governments or academia¹. Therefore unpublished trials can lead to patients being denied the most effective treatment, or being exposed to ineffective treatments, or to unnecessary side effects. Furthermore the trial participants, who invested their time and placed themselves at potential risk of side effects, being on placebo or an ineffective treatment, have been misinformed since the trial results have not contributed to the care of future patients.

However the non-publication of trials is often not due to a deliberate decision to cover up results. Commonly in academia it is due to more understandable reasons such as researchers having busy clinical posts, moving onto other more demanding projects, changing research areas or starting a family. This is called the “file drawer” problem.

Many steps are currently being taken to address the non-publication of trials, such as enforcing trial registration. However implementing this type of action prospectively will not affect the current evidence base for some time. Therefore reducing publication bias partly depends on researchers pursuing the publication of trials that are sitting in the “file drawer”.

Doshi *et al.* have made a call for action². They have asked investigators of unpublished trials to either publish their trial within one year or allow their trial data to be made public so that it can be prepared for publication by others. This concept is called ‘Restoring Invisible and Abandoned Trials’ (RIAT). Furthermore the paper sets out a minimum set of criteria for responsible publication of abandoned trials.

The first paper to be published under the RIAT initiative was a trial that was conducted 20 years ago and involved 1447 bowel cancer

patients³. Treasure *et al.* contacted the trial investigators to ask for their permission to re-analyse the results and publish the trial. The trial investigators supported this work and now finally the efforts of the trial participants have contributed to the evidence base.

A further example of a recovered trial is a clinical trial on polycythaemia that was not published due to several setbacks including researchers retiring, data being lost during a relocation of its administrative centre, and journals refusing to publish a partially completed trial. Despite all these issues, the results of the trial have now been reported as an attachment to a commentary in the journal *Trials* which is specifically interested in unpublished trials⁴.

A final example is a randomised controlled trial of Brief Alcohol Interventions⁵. This trial was not published for 17 years due to researchers moving abroad and taking on other demanding jobs. Therefore, I was asked by the researchers to prepare the trial for publication by updating the introduction and discussion. As a medical student with plenty of time, I was very keen to assist. Now finally the results are available to contribute to future meta-analyses.

These examples demonstrate that it is possible to go back, even decades later, and make the results available to inform future evidence based medicine. We call on others to look into their “file drawer” for unpublished trials. There is plenty of assistance available to make publication as easy as possible, for example by asking others to update the manuscript or publishing through dedicated journals such as *Trials* or *F1000 Research*. It is never too late to publish an abandoned trial.

Author contributions

Both authors were involved in writing the manuscript and have agreed to its final contents.

Competing interests

No competing interests were disclosed.

Grant information

Celia Shiles is supported by the Wessex Deanery and Julia Sinclair by the Higher Education Funding Council.

Acknowledgements

I would like to thank Ian Bushfield for his advice on writing this editorial.

References

1. Chan AW, Song F, Vickers A, *et al.*: **Increasing value and reducing waste: addressing inaccessible research.** *Lancet.* 2014; **383**(9913): 257–266.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Doshi P, Dickersin K, Healy D, *et al.*: **Restoring invisible and abandoned trials: a call for people to publish the findings.** *BMJ.* 2013; **346**: f2865.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Treasure T, Monson K, Fiorentino F, *et al.*: **Operating to remove recurrent colorectal cancer: have we got it right?** *BMJ.* 2014; **348**: g2085.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Richards SM, Burrett JA: **A proposal for reducing the effect of one of many causes of publication bias.** *Trials.* 2013; **14**: 41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Shiles CJ, Canning UP, Kennell-Webb SA, *et al.*: **Randomised controlled trial of a brief alcohol intervention in a general hospital setting.** *Trials.* 2013; **14**: 345.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research