RESEARCH NOTE

Relationships between activation level, knowledge, self-efficacy, and self-management behavior in heart failure patients discharged from rural hospitals [version 1; peer review: 2 approved]

Van Do¹, Lufei Young², Sue Barnason², Hoang Tran³

¹Department of Health Services Research and Administration, College of Public Health, University of Nebraska Medical Center, Omaha, NE, 68198, USA

²College of Nursing-Lincoln Division, University of Nebraska Medical Center, Omaha, NE, 68198, USA

³Department of Epidemiology, College of Public Health, University of Nebraska Medical Center, Omaha, NE, 68198, USA

V1 First published: 11 Jun 2015, 4:150
<https://doi.org/10.12688/f1000research.6557.1>
 Latest published: 11 Jun 2015, 4:150
<https://doi.org/10.12688/f1000research.6557.1>

Abstract

Non-adherence to self-management guidelines accounted for 50% of hospital readmissions in heart failure patients. Evidence showed that patient activation affects self-management behaviors in populations living with chronic conditions. The purpose of this study was to describe patient activation level and its relationship with knowledge, self-efficacy and self-management behaviors in heart failure patients discharged from rural hospitals. Our study populations were recruited from two hospitals in rural areas of Nebraska. We found that two-thirds of the participants reported low activation levels (e.g., taking no action to manage their heart failure condition). In addition, low patient activation levels were associated with inadequate heart failure knowledge ($p=.005$), low self-efficacy ($p<.001$) and low engagement in heart failure self-management behaviors ($p<.001$) after discharge from hospital.

Keywords

Rural Populations, Cardiac Failure, Patient Activation, Patient Engagement, Self-Efficacy, Self-Management, Knowledge

Open Peer Review

Approval Status

	1	2
version 1
11 Jun 2015	view	view
1. Hoa L. Nguyen , Baylor Scott & White Health, Dallas, USA		
2. George Sokos , Allegheny General Hospital, Pittsburgh, USA		
Megan Clarke , Allegheny General Hospital, Pittsburgh, USA		

Any reports and responses or comments on the article can be found at the end of the article.

Associated Study Protocol

Young L, Barnason S and Do V » Promoting self-management through adherence among heart failure patients discharged from rural hospitals: a study protocol, F1000Research 2015, 3:317 (<https://doi.org/10.12688/f1000research.5998.2>)

Corresponding authors: Van Do (van.do@unmc.edu), Lufei Young (lyoun1@unmc.edu)

Competing interests: No competing interests were disclosed.

Grant information: Research reported in this publication was supported by the National Institutes Nursing Research of the National Institutes of Health under award number 1R15NR 13769-01A1. The sponsor had no role in conducting the study, preparing and disseminating the study results. The authors are the recipients of the funding provided by the National Institutes Nursing Research of the National Institutes of Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2015 Do V *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Do V, Young L, Barnason S and Tran H. **Relationships between activation level, knowledge, self-efficacy, and self-management behavior in heart failure patients discharged from rural hospitals** [version 1; peer review: 2 approved] F1000Research 2015, 4:150 <https://doi.org/10.12688/f1000research.6557.1>

First published: 11 Jun 2015, 4:150 <https://doi.org/10.12688/f1000research.6557.1>

Heart failure is a major public health problem in the United States and worldwide. In the United States, heart failure affected 5.1 million patients with 12–15 million office visits, 6.5 million hospital days and cost approximately 32 billion U.S. dollars in 2010¹. Despite the recent declining trend of cardiovascular disease-related mortality in the US, the rehospitalization rate for heart failure patients remains 30% within 60–90 days after discharge². Rural populations exhibit higher prevalence of heart failure³ and rural patients are more likely to be readmitted due to heart failure^{4,5} compared to those in urban areas.

Among all causes for heart failure-related readmission, non-adherence to self-management is the most common reason, accounting for 50% of readmissions in heart failure patients^{6,7}. Patient self-management is one of the key concepts in the Chronic Care Model developed by Edward H. Wagner⁸. Self-management behaviors refer to the practice of activities that individuals initiate and perform on their own behalf in the interest of maintaining life, health, continuing personal development, and well-being⁹. Self-management behaviors in heart failure patients primarily involve monitoring daily weight, following a restricted sodium diet, fluid restriction, taking prescribed medications, exercising regularly, and keeping scheduled follow-up appointments⁴.

Prior nationwide studies among adults with chronic diseases showed that a high level of patient knowledge, efficacy and activation level were associated with good self-management^{10–12} and ultimately led to fewer hospitalizations and emergency department visits^{10,13}. Few studies have reported the relationship between disease specific knowledge, self-efficacy, patient activation and self-management behaviors in rural heart failure patients. Self-efficacy for heart failure self-management is defined as how confidently individuals can achieve specific functions or control various aspects of their heart failure¹⁴.

Patient engagement with self-management is critical for heart failure patients. Patient activation is the concept that can be applied as a

means to determine patient engagement with self-management. The patient's activation level, which is measured by Hibbard's Patient Activation Measure (PAM), reflects the degree to which the person is ready, willing and able to engage in managing her or his health conditions^{12,15}. Based on the PAM score, a person's activation level is graded into 4 levels, from low to high: (1) Patients in level one believe they are responsible for managing their health; (2) patients in level two feel confident and knowledgeable regarding managing their health; (3) patients in level three actively engage in managing their health; and (4) patients in level four consistently engage in activities to manage their health and maintain those actions even under stress. The advancement in patient activation levels reflects the progress of the patient from being a passive care receiver to a more confident care manager^{12,13,15}.

Considering the low engagement in self-management of patients living with chronic diseases in the rural areas^{16,17}, it is critical to examine whether there are positive relationships between knowledge, self-efficacy, patient activation and self-management behaviors. To our knowledge, these relationships have not been well studied in the rural heart failure population. Our study will contribute knowledge in understanding the impact of patient activation on self-management, which could inform the development of effective interventions to promote self-management in rural heart failure population. For this purpose, our study populations were recruited from hospitals in rural areas of Nebraska.

Conceptual framework

As a cultural aspect, many rural patients endorse the importance of personal responsibility, productivity, and self-reliance in terms of health practice¹⁸. Based on the rural culture belief and health practice, Bandura's social cognitive theory¹⁹, Hibbard's Patient Activation Theory^{12,20}, and the Chronic Care Model^{8,21}, we proposed a conceptual framework (Figure 1). Our framework assumes that by attaining self-management knowledge and efficacy, patients will advance their activation to higher levels, leading to long-term engagement in self-management behaviors.

Figure 1. Patient activation concept framework.

Methods

Research design

We conducted a secondary analysis to evaluate the relationships between levels of patient activation and heart failure self-management knowledge, self-efficacy, and self-management behaviors in heart failure patients discharged from rural critical access hospitals to home. The data for this paper formed the baseline data from a NIH-funded randomized controlled trial titled “Patient AcTivated Care at Home (PATCH)” which aims to examine the feasibility of a 12-week home-based intervention to improve heart failure self-management adherence. This trial can be found on <https://www.clinicaltrials.gov/ct2/show/NCT01964053>.

Sample and setting

Participants were recruited from October 2013 to December 2014 from two rural critical access hospitals in southeast Nebraska. Patients were eligible for the study if they: 1) were age 21 or older; 2) had heart failure as one of their discharge diagnoses; 3) had New York Heart Association (NYHA) class II to IV heart failure or had NYHA class I heart failure and had at least one heart failure-related hospitalization or emergency department visit in the previous year; 4) were discharged to home; 5) passed the Mini-Cog screen test screening for dementia²²; 6) understood English; and 7) had access to a phone.

We excluded patients who: 1) had depressive symptoms (received a score of 3 or above on the Patient Health Questionnaire-2 (PHQ-2)²³; 2) were diagnosed with liver cirrhosis; 3) were diagnosed with chronic renal failure; and 4) were diagnosed with other end stage and/or terminal illness (e.g. cancer) which limited the patient’s ability to perform self-management behaviors. The study setting is described in more detail in the study protocol²⁴.

Measurements

We collected socio-demographic characteristics including age, gender, educational attainment, race/ethnicity, annual household income, marital status, and smoking status with a structured questionnaire. Clinical characteristics included comorbidities, echocardiographic ejection fraction (EF), and New York Heart Association (NYHA) Functional Classification.

We measured patient activation using the Short Form of the Patient Activation Measure (13-item version, available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1361231/table/tb11/>) which has similar reliability and validity to the long form (22 item version) across different ages, genders and health condition status¹⁵. Each item of the form was scored on the 5-point Likert response scale. For the ease of interpretation, the raw scores were transformed from the original metric to a 0–100 metric with higher scores indicating higher activation levels. Based on the patient activation score, patients were categorized into four levels: level 1 (score <47.0), level 2 (score 47.1–55.1), level 3 (score 55.2–67.0), and level 4 (score >67.0)^{12,15}.

We measured self-efficacy using the Self-Care of Heart Failure Index (SCHFI) which has six questions on a 4 point Likert scale¹⁴. The SCHFI was used to assess the degree of achievement in self-care maintenance and management²⁵. The raw scores were standardized to a 100 point scale, with higher scores indicating higher

efficacy. This measurement tool is reliable to assess self-efficacy with Cronbach’s alpha coefficient of .83¹⁴.

We measured patients’ heart failure knowledge using the 27 multiple-choice questions of the Atlanta Heart Failure Knowledge Test (AHFKT)²⁶. Each correct answer scored 1 point and the total score ranged from 0 to 27. This questionnaire established high reliability to evaluate heart failure knowledge with a Cronbach alpha coefficient of .84²⁶.

We used the 29-item Revised Heart Failure Self-Care Behavior Scale (RHFSCS) to assess patients’ behavior in six heart failure care domains: (1) seeking medical help, (2) being aware of the effects of heart failure, (3) prevention of complications, (4) awareness of deleterious effects of medical care, (5) accepting heart failure, and (6) learning to live with heart failure²⁷. Each response is granted a score from 0 (none of the time) to 5 (all of the time)²⁷. The internal reliability of this questionnaire is consistent with a Cronbach alpha coefficient of .84²⁷.

Statistical analysis

Statistical analyses were conducted using SPSS version 20, with a p value less than .05 considered significant. We reported mean and standard deviation for continuous variables. Frequency and percentage were used to report categorical variables. Since patient activation level was an ordered variable, the Chi-square test for trend was used to assess correlations between four activation levels and the categorical variables (e.g. patient’s demographic and clinical characteristics). To compare continuous variables (e.g. behaviors, heart failure knowledge, Self-efficacy) across four activation levels, we used the ANOVA test for normally-distributed variables and the Kruskal-Wallis test for the non-normally distributed variables.

Ethics

The study protocol was approved by the University of Nebraska Medical Center Institutional Review Board (IRB) and hospital ethical committees (IRB PROTOCOL # 228-13-EP).

Results

Demographic and clinical characteristics

There were 101 patients enrolled in this study. Table 1 describes socio-demographic and clinical characteristics. The mean age was 70±12.1 years. The majority of participants were women (63%). Two-thirds of participants had an annual household income of less than \$50,000 or an educational attainment of high school graduate or lower. All the patients had multiple comorbidities including more than 2 of these chronic conditions: Hypertension (98.0%), Coronary artery disease (94.1%), Dyslipidemia (83.2%), Diabetes mellitus (41.6%), COPD/asthma (39.4%), CVA/stroke (17.8%), and cancer (6.9%). The majority of participants are at NYHA class II and III with the average EF of 55.8±11.1.

Patient activation level

The mean patient activation score was 57 and the median was 52 (95% CI 53–61). The patient activation score did not distribute normally with a positively skewed pattern. About 40% of participants believed that they were responsible for caring for their health and illness but failed to take action to manage their heart failure

Table 1. Patient characteristics (n=101).

	N (%)	PAM score (SD)	p-value for the correlations with patient activation score
Gender			.806
Male	37 (36.6%)	58 (18.5)	
Female	64 (63.4%)	56 (18.9)	
Age[§]			.224
65 years or younger	39 (38.6%)	60 (18.3)	
Older than 65 years	62 (61.4%)	55 (18.9)	
Race/ Ethnicity			.121
Caucasian	96 (95.0%)	58 (18.9)	
Non-Caucasian	5 (5.0%)	44 (6.5)	
Educational attainment			.030
Lower or High school graduate	64 (63.4%)	54 (17.6)	
Beyond high school	37 (36.6%)	62 (19.7)	
Marital status			.637
Married/Lived with partner	51 (50.5%)	58 (17.5)	
Not married/lived with partner	50 (49.5%)	56 (20.0)	
Annual household income[†]			.400
Under \$50,000	67 (71.3%)	57 (19.8)	
\$50,000 – \$75,000	9 (8.9%)	63 (17.9)	
Greater than \$75,000	8 (7.9%)	61 (14.9)	
Refuse to response	10 (9.9%)	49 (14.9)	
Smoking status			.772
Not smoking	88 (87.1%)	57 (18.7)	
Current smoking	13 (12.9%)	56 (19.4)	
New York Heart Classification			.083
Class I	9 (8.9%)	58 (16.9)	
Class II	49 (48.5%)	61 (19.9)	
Class III	43 (42.6%)	52 (16.8)	
Ejection fraction (SD)[‡]			.977
< 40%	9 (10.1%)	64 (24.7)	
≥ 40%	80 (89.9%)	57 (18.2)	
Patient activation measure			
Level 1	39 (39%)	--	--
Level 2	23 (23%)	--	--
Level 3	19 (19%)	--	--
Level 4	19 (19%)	--	--

Significant results in bold face[§]missing data from 1 participant, n=100[†]missing data from 7 participants, n=94[‡]missing data from 12 participants, n=89

--: not applicable

condition (level 1). A quarter of participants felt confident and knowledgeable enough to manage their health but failed to take action (level 2). Less than 40% of participants actually took some actions to manage their condition (level 3 and 4).

Patient activation levels were not significantly different across age group, gender, marital status, household income or smoking status. There was an association between activation level and educational attainment. Participants with educational attainment beyond high school had higher patient activation scores on the PAM compared to patients who only graduated high school or lower (62 vs. 54, $p=.03$).

Patients' self-management knowledge, efficacy and behavior

Using the Chi-square test for trend, we found an increasing trend between knowledge and self-efficacy across the four levels of patient activation. Patients with higher heart failure knowledge or greater self-efficacy were at a higher level of patient activation ($p=.005$ and $p<.001$, respectively). Patients who were at a higher level of patient activation also had higher scores of self-management behavior ($p<.001$). Table 2 describes the average score of self-management knowledge, efficacy, and behavior among 4 levels of patient activation level.

Dataset 1. Heart failure (HF) patients discharged from rural hospitals: demographic, clinical, patient activation level, HF knowledge and self-management behavior characteristics

<http://dx.doi.org/10.5256/f1000research.6557.d49205>

Data was collected by questionnaires through patient interview and medical record review.

Details of labelling and abbreviation of the variables were included in the dataset⁴⁰.

Discussion

This is one of the first studies to assess self-management knowledge, self-efficacy, activation level, and self-management behavior in American rural heart failure patients. The mean patient activation score of our study was 57, lower than that in Shively's study (61.7) of heart failure patients or Green's in the general chronic patient population (66.4)^{10,28}. In addition, fewer of our rural heart failure

participants were actively managing their condition compared to participants in other studies of both rural and urban areas^{10,28}. According to our study, a lower percentage of heart failure patients (38%) had high activation levels and consistently took action to manage their health compared to diabetic patients in Begum's (69.9%) and Rask's studies (82.9%)^{29,30}. However, patient activation scores in our study were higher than that in Evangelista's study (37.3–39.3) in which the population consisted of heart failure patients in a palliative care setting³¹. Similarly to Marshall's study³², the distribution of patient activation scores in our study was not different across various socio-demographic groups except for educational attainment. A higher educational attainment was associated with a higher level of activation in our patients. This finding was consistent with other studies which also found educational attainment to be the most powerful predictor for patient activation in the chronic disease population^{33–36}. It is plausible that patients with higher educational levels are more likely to achieve better health literacy, giving them more awareness, skills and confidence to take self-management actions. In contrast to several studies, we did not find a variance of patient activation score among different age groups^{31,37}. A possible explanation is that our participants seemed to be older and have narrower age variation (mean age 70, rank from 40 to 93) compared other studies (mean 53.7³¹ and age range from 18 to older than 75³⁷). Evidence shows that an increase in patient activation level results in improved self-management behavior, leading to better health outcomes^{13,38}. The overall low patient activation score in our heart failure participants indicates the need of developing interventions to enhance activation and self-management behavior in rural heart failure patients.

The associations between patient activation and heart failure knowledge, self-efficacy, and self-management behavior support our conceptual framework, as well as previous studies^{34,37}. Patients with higher activation levels tended to have more knowledge to manage their heart failure, more confidence in self-management of heart failure, and were more likely to engage in self-management behaviors such as regular exercise, watching their diet and fluid intake, and adherence to medication. From a qualitative study, Dixon *et al.* reported that patients at lower activation levels indicated a lack of knowledge and lack of confidence as barriers for them to self-manage their health conditions³⁹. Our findings indicate that we can potentially boost patients' activation by increasing their self-management knowledge and efficacy to take care of their own health. These findings confirmed our original hypothesis that strategies

Table 2. Self-management knowledge, efficacy, and behavior among 4 levels of Patient activation level (n=100).

	Overall	Level 1	Level 2	Level 3	Level 4	p-value
	Mean (SD)					
Knowledge	20 (3.0)	20 (3.1)	19 (2.8)	21 (2.2)	22 (2.9)	.005*
Efficacy	47 (23.8)	31 (18.5)	47 (16.4)	52 (17.8)	75 (18.4)	<.001*
Behavior	89 (19.7)	82 (17.2)	93 (20.9)	91 (16.5)	98 (22.4)	<.001*

Significant results in bold face

* df=1, Chi-square for trend

to enhance activation levels should be included in the intervention to promote heart failure self-management behaviors.

Limitations

Our study had some limitations. We used only baseline data; therefore, we are unable to demonstrate the changes of patient activation scores, knowledge scores, self-efficacy and self-management behaviors over time. As a result, we could not establish the temporal relationship between knowledge, efficacy, patient activation level and their impact on behavior in this article, which will be reported in future manuscripts. Secondly, the patients who agreed to participate in the original randomized controlled trial might be more motivated than patients who refused to participate, therefore creating a pool of heart failure patients with higher activation levels compared to the general rural heart failure population. If this assumption is valid, the overall activation level in general rural heart failure population might be even lower than what is reported in this article, which indicates a great need to conduct interventions to enhance rural heart failure activation to engage in self-management behaviors.

Clinical implication

Using the short version of PAM (13 items) is a feasible way to assess a patient's activation level and identify those with low levels in both inpatient and outpatient settings. The assessment results allow clinicians to develop tailored interventions to support those high risk patients. Additionally, the patients with high activation levels could be encouraged and recruited as behavior coaches and/or peer supporters for those with low activation scores.

Conclusions

Self-management plays a vital role in improving health outcomes and reducing healthcare costs in the heart failure population. Patient activation level is significantly associated with self-management behavior. However, the activation level in our rural heart failure patients was relatively low compared to patients in other studies, which suggest the need for interventions to improve activation levels in the rural heart failure population.

Data availability

F1000Research: Dataset 1. Heart failure (HF) patients discharged from rural hospitals: demographic, clinical, patient activation level, HF knowledge and self-management behavior characteristics, [10.5256/f1000research.6557.d49205](https://doi.org/10.5256/f1000research.6557.d49205)⁴⁰

Consent

Written informed consent for publication of clinical details was obtained from all the participants.

Author contributions

Dr. Do and Dr. Young developed the research question and the conceptual framework. Dr. Do and Dr. Tran analyzed data and prepared for tables and figures. Dr. Do, Dr. Tran, Dr. Young and Dr. Barnason prepared the manuscript. The co-author, Dr. Young, has full access to the study data and takes responsibility for the integrity and the accuracy of the data analysis. All authors were involved in the review of the final draft and agreed to the content of the final submission.

Competing interests

No competing interests were disclosed.

Grant information

Research reported in this publication was supported by the National Institutes Nursing Research of the National Institutes of Health under award number 1R15NR 13769-01A1. The sponsor had no role in conducting the study, preparing and disseminating the study results. The authors are the recipients of the funding provided by the National Institutes Nursing Research of the National Institutes of Health.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Go AS, Mozaffarian D, Roger VL, *et al.*: **Heart disease and stroke statistics--2013 update: a report from the American Heart Association.** *Circulation*. 2013; **127**(1): e6–e245.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gheorghiade M, Vaduganathan M, Fonarow GC, *et al.*: **Rehospitalization for heart failure: problems and perspectives.** *J Am Coll Cardiol*. 2013; **61**(4): 391–403.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pearson TA, Lewis C: **Rural epidemiology: insights from a rural population laboratory.** *Am J Epidemiol*. 1998; **148**(10): 949–957.
[PubMed Abstract](#)
- Jessup M, Abraham WT, Casey DE, *et al.*: **2009 focused update: ACCF/AHA Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation.** *Circulation*. 2009; **119**(14): 1977–2016.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jin Y, Quan H, Cujec B, *et al.*: **Rural and urban outcomes after hospitalization for congestive heart failure in Alberta, Canada.** *J Card Fail*. 2003; **9**(4): 278–285.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Evangelista LS, Shinnick MA: **What do we know about adherence and self-care?** *J Cardiovasc Nurs*. 2008; **23**(3): 250–257.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- van der Wal MH, Jaarsma T, Moser DK, *et al.*: **Qualitative examination of compliance in heart failure patients in The Netherlands.** *Heart Lung*. 2010; **39**(2): 121–130.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bodenheimer T, Wagner EH, Grumbach K: **Improving primary care for patients with chronic illness.** *JAMA*. 2002; **288**(14): 1775–1779.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Orem DE, Taylor SG, Renpenning KM: **Nursing: Concepts of Practice.** 6th ed., Mosby St. Louis, 2001.
[Reference Source](#)
- Greene J, Hibbard JH: **Why does patient activation matter? An examination of**

- the relationships between patient activation and health-related outcomes. *J Gen Intern Med*. 2012; **27**(5): 520–526.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Mosen DM, Schmittiel J, Hibbard J, *et al.*: Is patient activation associated with outcomes of care for adults with chronic conditions? *J Ambul Care Manage*. 2007; **30**(1): 21–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
 12. Hibbard JH, Stockard J, Mahoney ER, *et al.*: Development of the patient activation measure (PAM): Conceptualizing and measuring activation in patients and consumers. *Health Serv Res*. 2004; **39**(4 Pt 1): 1005–1026.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Hibbard JH, Greene J, Tusler M: Improving the outcomes of disease management by tailoring care to the patient's level of activation. *Am J Manag Care*. 2009; **15**(6): 353–360.
[PubMed Abstract](#)
 14. Riegel B, Lee CS, Dickson VV, *et al.*: An update on the self-care of heart failure index. *J Cardiovasc Nurs*. 2009; **24**(6): 485–497.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. Hibbard JH, Mahoney ER, Stockard J, *et al.*: Development and testing of a short form of the patient activation measure. *Health Serv Res*. 2005; **40**(6 Pt 1): 1918–1930.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Bell RA, Arcury TA, Snively BM, *et al.*: Diabetes foot self-care practices in a rural triethnic population. *Diabetes Educ*. 2005; **31**(1): 75–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Coronado GD, Thompson B, Tejeda S, *et al.*: Sociodemographic factors and self-management practices related to type 2 diabetes among Hispanics and non-Hispanic whites in a rural setting. *J Rural Health*. 2007; **23**(1): 49–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
 18. Bardach SH, Tarasenko YN, Schoenberg NE: The role of social support in multiple morbidity: self-management among rural residents. *J Health Care Poor Underserved*. 2011; **22**(3): 756–771.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Bandura A: Perceived self-efficacy in cognitive development and functioning. *Educational Psychologist*. 1993; **28**(2): 117–148.
[Publisher Full Text](#)
 20. Hibbard JH, Mahoney ER, Stock R, *et al.*: Do increases in patient activation result in improved self-management behaviors? *Health Serv Res*. 2007; **42**(4): 1443–1463.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Wagner EH, Austin BT, Davis C, *et al.*: Improving chronic illness care: translating evidence into action. *Health Aff (Millwood)*. 2001; **20**(6): 64–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Borson S, Scanlan J, Brush M, *et al.*: The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry*. 2000; **15**(11): 1021–1027.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. Li C, Friedman B, Conwell Y, *et al.*: Validity of the Patient Health Questionnaire 2 (PHQ-2) in identifying major depression in older people. *J Am Geriatr Soc*. 2007; **55**(4): 596–602.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. Young L, Barnason S, Do V: Promoting self-management through adherence among heart failure patients discharged from rural hospitals: a study protocol [v2; ref status: indexed, <http://f1000r.es/5c7>]. *F1000Res*. 2015; **3**.
[Publisher Full Text](#)
 25. Riegel B, Carlson B, Moser DK, *et al.*: Psychometric testing of the self-care of heart failure index. *J Card Fail*. 2004; **10**(4): 350–360.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Reilly CM, Higgins M, Smith A, *et al.*: Development, psychometric testing, and revision of the Atlanta Heart Failure Knowledge Test. *J Cardiovasc Nurs*. 2009; **24**(6): 500–509.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Artinian NT, Magnan M, Sloan M, *et al.*: Self-care behaviors among patients with heart failure. *Heart Lung*. 2002; **31**(3): 161–172.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Shively MJ, Gardetto NJ, Kodiath MF, *et al.*: Effect of patient activation on self-management in patients with heart failure. *J Cardiovasc Nurs*. 2013; **28**(1): 20–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Begum N, Donald M, Ozolins IZ, *et al.*: Hospital admissions, emergency department utilisation and patient activation for self-management among people with diabetes. *Diabetes Res Clin Pract*. 2011; **93**(2): 260–267.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Rask KJ, Ziemer DC, Kohler SA, *et al.*: Patient activation is associated with healthy behaviors and ease in managing diabetes in an indigent population. *Diabetes Educ*. 2009; **35**(4): 622–630.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Evangelista LS, Liao S, Motie M, *et al.*: On-going palliative care enhances perceived control and patient activation and reduces symptom distress in patients with symptomatic heart failure: a pilot study. *Eur J Cardiovasc Nurs*. 2014; **13**(2): 116–123.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Marshall R, Beach MC, Saha S, *et al.*: Patient activation and improved outcomes in HIV-infected patients. *J Gen Intern Med*. 2013; **28**(5): 668–674.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Lubetkin EI, Lu WH, Gold MR: Levels and correlates of patient activation in health center settings: Building strategies for improving health outcomes. *J Health Care Poor Underserved*. 2010; **21**(3): 796–808.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Nijman J, Hendriks M, Brabers A, *et al.*: Patient activation and health literacy as predictors of health information use in a general sample of dutch health care consumers. *J Health Commun*. 2014; **19**(8): 955–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Rademakers J, Nijman J, van der Hoek L, *et al.*: Measuring patient activation in The Netherlands: translation and validation of the American short form Patient Activation Measure (PAM13). *BMC Public Health*. 2012; **12**: 577.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Saha S, Koley M, Mahoney ER, *et al.*: Patient activation measures in a government homeopathic hospital in India. *J Evid Based Complementary Altern Med*. 2014; **19**(4): 253–259.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Hendriks M, Rademakers J: Relationships between patient activation, disease-specific knowledge and health outcomes among people with diabetes; a survey study. *BMC Health Serv Res*. 2014; **14**(1): 393.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Hibbard JH, Greene J: What the evidence shows about patient activation: better health outcomes and care experiences; fewer data on costs. *Health Aff (Millwood)*. 2013; **32**(2): 207–214.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Dixon A, Hibbard J, Tusler M: How do People with Different Levels of Activation Self-Manage their Chronic Conditions? *Patient*. 2009; **2**(4): 257–268.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Do V, Young L, Barnason S: Dataset 1 in: Relationships between activation level, knowledge, self-efficacy, and self-management behavior in heart failure patients discharged from rural hospitals. *F1000Research*. 2015.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 09 July 2015

<https://doi.org/10.5256/f1000research.7041.r9428>

© 2015 Sokos G et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

George Sokos

Cardiovascular Institute, Allegheny General Hospital, Pittsburgh, USA

Megan Clarke

Section of Heart Failure/Transplant and Pulmonary Hypertension, Allegheny General Hospital, Pittsburgh, PA, USA

1. Page 2, paragraph 1: Consider adding information about CMS limited reimbursement for heart failure readmissions as this is a significant current concern in US hospitals.
2. Figure 1 and throughout entire paper: Consider simplifying terms "self-management knowledge" to "Knowledge", "Self management efficacy" to "efficacy". Many different variations of these terms are used throughout which reduces the readability of the paper. Table 2 on page 5 is organized well and has these terms outlined more simply.
3. Page 2, paragraph 3, last sentence: Consider changing "self-efficacy for heart failure self-management" to "Efficacy for heart failure self-management" as the terms can be confusing if the reader is trying to understand the concept using Figure 1. This may be considered throughout the paper.
4. Page 2, paragraph 4: It might be more helpful for the reader to see the PAM scores 1-4 in a table to allow for easy referencing when reading the paper.
5. Page 2, paragraph 5, first sentence: Consider changing "self-efficacy" to "efficacy" (see comment number 2 above).
6. Page 2, paragraph 6, last sentence: Consider removing "long-term" as this specific study make correlations but does not address the length of time the patient's behavior will be impacted.
7. Page 3, paragraph 1, sentence 1: Consider changing term "self efficacy" to efficacy (see comment number 2 above).

8. Page 3, paragraph 6, sentence 1: Consider changing term “self efficacy” to efficacy (see comment number 2 above).
9. Page 3, Measurement section: Consider arranging paragraphs to match flow in Figure 1 (Knowledge, Efficacy, Patient Activation, and Self-management behavior).
10. Page 3, paragraph 8: Change “patients behavior” to “patients self-management behavior”.
11. Table 1: Consider additional category of stage of heart failure (A,B,C,D) if available.
12. Page 5, paragraph 3 (first paragraph of discussion section): The statement regarding this study’s similarity to other findings which show educational attainment as the “most powerful predictor for patient activation” may led readers to believe that knowledge, efficacy, and behavior were compared and that education outweighed the others. I would consider re-wording and removing the term “most powerful.”.
13. Page 5, paragraph 4, sentence 2 (second paragraph of discussion section): Can “confidence” be changed to “efficacy” here to keep language consistent?
14. Page 5, paragraph 4, sentence 4: This sentence infers that the intervention in this study was to increase knowledge and efficacy to improve patient activation, but from my understanding this was a correlation and not an intervention.

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 22 June 2015

<https://doi.org/10.5256/f1000research.7041.r8999>

© 2015 **Nguyen H.** This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hoa L. Nguyen

Department of Epidemiology, Baylor Scott & White Health, Dallas, TX, USA

The paper potentially provides important information about the relationship between activation level, knowledge, self-efficacy and self-management behavior in the rural population with heart failure to inform future interventions.

I have several minor comments that the authors may consider in the revised version:

1. Abstract: The authors may consider adding sample size, and several basic characteristics of the study population (mean age, sex distribution, etc.) in the abstract.

2. Since the sample size of the study is relatively small ($n=101$), and the authors stated that the distribution of the patient activation score was skewed (page 3, last paragraph), they may consider reporting medians (inter quartile ranges) in the tables and text in addition to means (SDs).
3. Statistical analysis: The authors mentioned Chi-square, Anova, and Kruskal-Wallis tests were used to compare patients' characteristics across 4 activation levels, but it is unclear where the results were reported in the results section.

In table 1, patients' characteristics for all patients were presented (not according to activation levels) and means (SDs) PAM score by patients characteristics were reported.

The authors may consider stating clearly in the statistical analysis how they treated the activation score (the main outcome of interest) as continuous or as categorical variable or both approaches and report the results accordingly.

4. Discussion: The author wrote "the distribution of patient activation scores in our study was not different across various socio-demographic groups except for educational" (page 5). This may be due to small sample size issue; the study may be underpowered to detect potential differences (e.g. the mean activation score difference by race/ethnicity was 14 (58 vs. 44), however the difference was not statistically significant). This is one of the study limitations.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research