Check for updates

CASE REPORT

REVISED Case Report: A case report of unstable Hangman fracture in a eighty year old male [version 2; peer review: 2 approved, 1 not approved]

Sunil Munakomi , Binod Bhattarai

Department of Neurosurgery, College of Medical Sciences, Bharatpur, Nepal

V2 First published: 23 Jul 2015, 4:337
<https://doi.org/10.12688/f1000research.6799.1>

Latest published: 17 Dec 2015, 4:337
<https://doi.org/10.12688/f1000research.6799.2>

Abstract

Herein we discuss a rare variant of hangman's fracture in an eighty year old male with good Karnofsky performance score. We performed X-ray and magnetic resonance imaging (MRI) of the cervical spine to confirm the diagnosis. The patient was placed on a gentle cervical traction which showed good reduction. Despite being on a resource limited setup, we performed posterior occipitocervical fusion with bone graft fusion followed by early mobilization. A postoperative scan showed good reduction and purchase of the screws. This case highlights the importance of choosing the correct therapeutic attitude for the management of the geriatric population especially in those who do not have any significant co-morbid conditions.

Keywords

hangman's fracture , unstable , management

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 17 Dec 2015		view	view
version 1 23 Jul 2015	view	view	view

1. **Rajiv R. Ratan**, Weil Medical College of Cornell, White Plains, USA
2. **Virendra Deo Sinha**, S.M.S. Medical College, Jaipur, India
Amit Chakrabarty, Eternal Hospital, Jaipur, India
3. **George Wittenberg**, University of Maryland School of Medicine, Baltimore, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Sunil Munakomi (sunilmunakomi@gmail.com)

Competing interests: No competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2015 Munakomi S and Bhattarai B. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Munakomi S and Bhattarai B. **Case Report: A case report of unstable Hangman fracture in a eighty year old male [version 2; peer review: 2 approved, 1 not approved]** F1000Research 2015, 4:337 <https://doi.org/10.12688/f1000research.6799.2>

First published: 23 Jul 2015, 4:337 <https://doi.org/10.12688/f1000research.6799.1>

REVISED Amendments from Version 1

We herein address the issues as per the advice of our referees. We try to focus on a tailored made approach to manage spine injuries in elderly patients especially in those having a good Karnofsky performance score. Despite being on a resource restrained setup, we can achieve good outcome with valid alternative limited approaches.

See referee reports

Introduction

Rigid immobilization alone is sufficient for most cases of hangman's fracture (defined as traumatic spondylolisthesis of C2) classified as Effendi type I and some of type II. Effendi type III fractures are very rare and invariably have neurological deficits because of impingement due to the facet dislocation on the spinal cord posteriorly¹. Fracture instability is the presence of complete disruption of the annular and/or posterior ligament with forward and/or rotatory vertebral body slip of axis². Surgical stabilization and rigid immobilization together is recommended in such cases, such as Levine-Edwards type IIa and III fractures. Here we discuss the management of an unstable type III hangman's fracture in an aged patient without any neurological deficits. Most doctors choose traction and prolonged immobilization in a halo

vest due to associated medical comorbidities and the anesthetic risks involved in this group^{3,4}. However there is a high risk of nonunion, instability, persistent pain and a need for a prolonged period of halo immobilization⁵. Since our patient had a good Karnofsky performance score⁶, we opted for only posterior fusion so as to minimize the anesthetic risk involved with both anterior and posterior approaches. However, we chose a long segment occipitocervical screw and graft fusion so as to aid the healing process in the aged bone.

Case report

An 80 year old man from the Tarai region of Nepal was brought to emergency with the chief complaint of falling from a swing after being pushed by his grandson 2 days prior. He complained of pain at the nape of his neck. The patient was neurologically intact. He was placed in a cervical collar and an urgent X-ray of the cervical spine revealed presence of spondylolisthesis of the axis with significant translation and angulation (Figure 1). Magnetic resonance imaging (MRI) of the cervical spine revealed a type III hangman's fracture with presence of pinching effect on the cord without any significant signal changes (Figure 2).

The patient was an ex-army serviceman and was in good health with good Karnofsky performance score⁶. There was no significant past

Figure 1. X-ray of the cervical spine showing Hangman's fracture with significant translation and angulation.

Figure 2. MRI (t2 sequence) of the cervical spine revealing presence of pinching effect but no signal changes in the cord.

medical or surgical illnesses. He had a habit of smoking marijuana previously. However, routine screening echocardiography revealed a cardiac ejection fraction of only 33%. Because the fracture was an unstable type III variant, the decision of surgical fixation was taken. The best option in such a situation would have been anterior cervical disectomy and Casper plate fixation thereby avoiding prolonged prone positioning. However such armamentarium for the procedure was not available with us. After explaining the disease condition, treatment options available and the risks involved, the patient was placed on minimal cervical traction so as to avoid the risk of iatrogenic hanging. We looked for the level of realignment that was possible with the guarded traction. Stringent care was taken to observe for features of over distraction. Repeat imaging showed good realignment and normal canal diameter. Therefore we decided to go for occipito-cervical fusion so as to minimize the anesthetic risk imposed to the patient from both anterior and posterior approaches. A DEXA scan for assessing bone density would have been justified prior to occipito-cervical fusion since such procedure would further lead to severe motion restriction in such an elderly spine. However such facility was not available to us. Intra-operatively there was fracture of the pars and the lamina of C2. Since there was no atlantoaxial dislocation, we opted for occipital and C1 and C3 lateral mass fixation. There is evidence of good results with short fixation of C1 and C3 only, but keeping in mind the risk of osteoporosis in this case, we wanted further anchorage from occipital fusion as well. Since there was good posterior realignment of the spinal lines after traction (Figure 3) and intra-operatively, we choose the posterior approach only to minimize the added risk of the anterior approach. Lateral mass screws were placed in C1 and C3 (Figure 4 and Figure 5). Bone graft harvested from iliac bone was placed in the C1 and

Figure 3. X-ray spine after traction showing realignment of the posterior and the spino-laminar lines.

C2 inter-space to further enhance the fusion process. The patient was started on dexamethasone (8 mg intravenously and then rapidly tapered off in the following 2 days). The patient was safely extubated. Neurological examination was normal. The patient was in complete bed rest for a week and then mobilized with support. A CT spine check after one week revealed good screw purchase (Figure 6) and good reduction of fracture segment (Figure 7). The patient was restricted to light weight bearing and was advised to keep the cervical collar for at least 6 weeks. The patient was started on calcium supplementation (tablet calcium 500 mg orally every 12 hours). The patient followed up in the outpatient department after 1.5 months walking on his own without any deficits.

Discussion

“Hangman’s fracture”, a traumatic spondylolisthesis of C2, first coined by Schneider *et al.* in 1965⁷ results from hyperextension of

Figure 4. CT spine showing projection of screws through lateral mass of C1.

Figure 5. CT spine showing projection of screw through lateral mass of C3.

Figure 6. CT spine reconstruction showing projection and final alignment of the construct.

Figure 7. CT spine showing good reduction of the posterior and the spino-laminar lines and normal canal.

the upper cervical spine. There is fracture of the lateral mass and the pedicle of the axis with simultaneous disruption of the anterior longitudinal ligament allowing C2-C3 listhesis. Traumatic hangman's fracture, in contrast to the judicial hangman's fracture, is caused from extension and compression of the upper cervical spine with rare cord injury⁸.

The most widely used classification for hangman's fractures was firstly described by Effendi *et al.*⁹ and later modified by Levine *et al.*^{10,11}. Anterior approaches include anterior cervical disectomy and graft fusion¹²; posterior approaches include lateral mass, pedicle or transarticular screw placement¹³.

Anterior disectomy and screw plate fixation is an effective, but not very popular technique due to difficulty in exposing the C2-C3 region¹⁴ and the elimination of C2-C3 rotation¹⁵. Direct screw fixation of C2 pars adds to the risk of injury to the vertebral artery¹⁵ and also there is the need for complete manual reduction of the fracture intra-operatively^{15,16}.

Fusion of lateral masses of C1 and C3 for hangman's fractures minimizes risk of vertebral artery injury and displacement of fractured segments into the canal. The efficacy of this approach has been validated in a biomechanical study by Chittiboina *et al.*¹⁷

This study hereby highlights the importance of the treatment algorithm chosen for the management of unstable hangman's fracture in geriatric patients. Patients with good Karnofsky performance score would benefit from long segment posterior fusion, rather than both anterior and posterior approaches which might increase the intra-operative risk. Managing such patients with a prolonged period of immobilization in a halo imposes a higher risk of nonunion.

Conclusion

Age alone should not determine a doctor's approach to the treatment of geriatric patients. By taking only age into account when deciding on treatment, we risk compromising effective management in elderly patients. Karnofsky performance scale⁶ is one reliable marker that helps in making such treatment decisions. So despite being on a resource limited setup, we can tailor ourselves into adopting other viable options.

Consent

Both written and verbal informed consent for publication of images and clinical data related to this case was sought and obtained from the son of the patient.

Author contributions

SM reviewed the literature, designed the study and formatted the paper. BB revised and edited the final format.

Competing interests

No competing interests.

Grant information

No funding was involved in supporting this work.

References

1. Choi WG, Vishteh AG, Baskin JJ, *et al.*: **Completely dislocated hangman's fracture with a locked C2-3 facet. Case report.** *J Neurosurg.* 1997; **87**(5): 757–760.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Moon MS, Moon JL, Moon YW, *et al.*: **Traumatic spondylolisthesis of the axis: 42 cases.** *Bull Hosp Jt Dis.* 2001–2002; **60**(2): 61–66.
[PubMed Abstract](#)
3. Hadley MN, Browner C, Sonntag VK, *et al.*: **Axis fractures: a comprehensive review of management and treatment in 107 cases.** *Neurosurgery.* 1985; **17**(2): 281–290.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Schneider RC, Livingston KE, Cave AJ, *et al.*: **"Hangman's fracture" of the cervical spine.** *J Neurosurg.* 1965; **22**: 141–154.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Vaccaro AR, Madigan L, Bauerle WB, *et al.*: **Early halo immobilization of displaced traumatic spondylolisthesis of the axis.** *Spine (Phila Pa 1976).* 2002; **27**(20): 2229–2233.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Péus D, Newcomb N, Hofer S: **Appraisal of the Karnofsky Performance Status and proposal of a simple algorithmic system for its evaluation.** *BMC Med Inform Decis Mak.* 2013; **13**: 72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Boullosa JL, Colli BO, Carlotti CG Jr, *et al.*: **Surgical management of axis' traumatic spondylolisthesis (Hangman's fracture).** *Arq Neuropsiquiatr.* 2004; **62**(3B): 821–826.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Williams TG: **Hangman's fracture.** *J Bone Joint Surg Br.* 1975; **57**(1): 82–88.
[PubMed Abstract](#)
9. Effendi B, Roy D, Cornish B, *et al.*: **Fractures of the ring of the axis. A classification based on the analysis of 131 cases.** *J Bone Joint Surg Br.* 1981; **63-B**(3): 319–327.
[PubMed Abstract](#)
10. Levine AM, Edwards CC: **The management of traumatic spondylolisthesis of the axis.** *J Bone Joint Surg Am.* 1985; **67**(2): 217–226.
[PubMed Abstract](#)
11. Levine AM, Rhyne AL: **Traumatic spondylolisthesis of the axis.** *Semin Spine Surg.* 1991; **3**: 47–60.
12. Duggal N, Chamberlain RH, Perez-Garza LE, *et al.*: **Hangman's fracture: a biomechanical comparison of stabilization techniques.** *Spine (Phila Pa 1976).* 2007; **32**(2): 182–187.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Papagelopoulos PJ, Currier BL, Hokari Y, *et al.*: **Biomechanical comparison of C1-C2 posterior arthrodesis techniques.** *Spine (Phila Pa 1976).* 2007; **32**(13): E363–370.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Traynelis VC, Fontes RB: **Anterior fixation of the axis.** *Neurosurgery.* 2010; **67**(3 Suppl Operative): ons229–36; discussion ons236.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Bristol R, Henn JS, Dickman CA: **Pars screw fixation of a Hangman's fracture: Technical case report.** *Neurosurgery.* 2005; **56**(1 Suppl): E204; discussion E204.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. ElMiligui Y, Koptan W, Emran I: **Transpedicular screw fixation for type II Hangman's fracture: A motion preserving procedure.** *Eur Spine J.* 2010; **19**(8): 1299–305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Chittiboina P, Wylen E, Ogden A, *et al.*: **Traumatic spondylolisthesis of the axis: A biomechanical comparison of clinically relevant anterior and posterior fusion techniques.** *J Neurosurg Spine.* 2009; **11**(4): 379–87.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 20 January 2016

<https://doi.org/10.5256/f1000research.7957.r11993>

© 2016 Sinha V. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Virendra Deo Sinha

Department of Neurosurgery, S.M.S. Medical College, Jaipur, Rajasthan, India

The authors have successfully amended the article to the queries and reservations which we had made earlier

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 30 December 2015

<https://doi.org/10.5256/f1000research.7957.r11617>

© 2015 Wittenberg G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

George Wittenberg

Department of Neurology, University of Maryland School of Medicine, Baltimore, MD, USA

The authors have been very responsive to the comments. My statements before stand about the overall value of the paper. There is some idiosyncratic use of English, including the concluding statement about being able to "tailor ourselves into adopting other viable options." One could rewrite it as:

"So despite having limited resources, we can explore viable options."

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 03 December 2015

<https://doi.org/10.5256/f1000research.7308.r9611>

© 2015 Wittenberg G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

George Wittenberg

Department of Neurology, University of Maryland School of Medicine, Baltimore, MD, USA

I think this is a well-written and interesting case report. As a single case report, it cannot be used to guide treatment. The outcome could have been different even though the authors suggest a very reasonable approach to the situation. I would agree with Dr. Ratan's comments, although I may be less of an expert on the surgical management of hangman's fracture. I think there would be a small benefit to medical knowledge to indexing this paper, as the management of this condition would likely be guided by both exam, radiological findings, and patient medical condition.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 06 November 2015

<https://doi.org/10.5256/f1000research.7308.r11119>

© 2015 Sinha V et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Virendra Deo Sinha

Department of Neurosurgery, S.M.S. Medical College, Jaipur, Rajasthan, India

Amit Chakrabarty

Neurosurgery Department, Eternal Hospital, Jaipur, Rajasthan, India

We have read with interest the case report "A case report of unstable Hangman fracture in a

eighty year old male" by Munakomi *et al* .The authors need to be congratulated for highlighting their view in managing unstable Hangman fracture in the geriatric age group.

The patient in your report was put on traction with six kilogram loading. It is an accepted fact that patients with type 3 Hangman fracture should not be subjected to traction due to risk of "iatrogenic hanging".

Secondly, the radiology images published do not point towards severe osteoporosis. A DEXA scan could have been more informative as the decision to do occipito-cervical fusion in Hangman fracture exposes the patient to severe motion restriction in an already spondylotic spine.

Finally, I presume C2 -C3 anterior cervical discectomy and fixation with casper plates would have been preferable than to subjecting the patient with an ejection fraction of 33% to surgery in prone position.

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Reviewer Report 11 August 2015

<https://doi.org/10.5256/f1000research.7308.r9613>

© 2015 Ratan R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rajiv R. Ratan

Burke Medical Research Institute, Weil Medical College of Cornell, White Plains, NY, USA

This report describes a case of a Hangman's fracture without associated neurological symptoms. There are several problems with this report that limit its usefulness and message,

First, a Hangman's fracture, which is defined as fractures of both pedicles or pars interarticularis of the C2 vertebra (axis), is not defined in the abstract. There is a classification protocol that has been developed by Levine and Edwards that is based on etiology and guides treatment, this is also not mentioned.

Second, neurological impairment is only seen in 25% of cases, so the absence of neurological symptoms is the rule not the exception as implied by this abstract.

I am not convinced that this case report adds significantly to our understanding of how to diagnose and treat Hangman's fracture.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 11 Aug 2015

Sunil Munakomi, College of Medical Sciences, Bharatpur, Nepal

Thank you for the report. We totally agree with your comments, but the purpose of our paper was to highlight the clinical importance of choosing the correct surgical management, barring the age factor, which do significantly play a major role in decision making in developing countries like ours. So, we wanted to highlight the importance of neurological status and the Karnofsky score in making such decisions in geriatric patients.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research