Check for updates

OBSERVATION ARTICLE

Fused embryos and pre-metamorphic conjoined larvae in a broadcast spawning reef coral [version 1; peer review: 1 approved, 1 approved with reservations]

Lei Jiang¹⁻³, Xin-Ming Lei¹, Sheng Liu¹, Hui Huang^{1,2}

¹Key Laboratory of Tropical Marine Bio-resources and Ecology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, 510301, China

²Tropical Marine Biological Research Station in Hainan, Chinese Academy of Sciences, Sanya, 572000, China

³University of Chinese Academy of Sciences, Beijing, 100049, China

V1 First published: 13 Feb 2015, 4:44
<https://doi.org/10.12688/f1000research.6136.1>

Latest published: 05 Mar 2015, 4:44
<https://doi.org/10.12688/f1000research.6136.2>

Abstract

Fusion of embryos or larvae prior to metamorphosis is rarely known to date in colonial marine organisms. Here, we observed for the first time the embryos of the broadcast spawning coral *Platygyra daedalea* could fuse during blastulation and further develop into conjoined larvae, and the settlement of conjoined larvae immediately resulted in inborn juvenile colonies. Considering the frequent record of synchronous spawning events and spawn slicks in the field, fusion of embryos might be a naturally occurring phenomenon, and act as an adaptive strategy to form pre-metamorphic united larvae and larger recruits, thereby promoting early survival. However, whether fusion of embryos is common in spawning corals and its implications deserve further investigation.

Keywords

fusion , conjoined larvae , spawn slicks , inborn colonies , *Platygyra daedalea*

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
05 Mar 2015		
version 1		
13 Feb 2015	view	view

1. **Andrew H. Baird** , James Cook University, Townsville, Australia

2. **Baruch Rinkevich**, National Institute of Oceanography, Haifa, Israel

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Hui Huang (huanghui@scsio.ac.cn)

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Public Science and Technology Research Funds Projects of Ocean (201305030-3) and the National Natural Science Foundation of China (41306144 and U1301232). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2015 Jiang L *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Jiang L, Lei XM, Liu S and Huang H. **Fused embryos and pre-metamorphic conjoined larvae in a broadcast spawning reef coral [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2015, 4:44 <https://doi.org/10.12688/f1000research.6136.1>

First published: 13 Feb 2015, 4:44 <https://doi.org/10.12688/f1000research.6136.1>

Introduction

In sessile colonial marine invertebrates (e.g., sponges, cnidarians, bryozoans and ascidians), fusion among conspecifics during early ontogeny could immediately lead to a marked increase in juvenile size, thereby enhancing the performance in growth, survival and competition^{1,2}. In addition, the allogenic fusion is expected to form chimeras which possess greater genetic variability and wider ranges of physiological resistance¹. Larvae of colonial marine organisms tend to settle in a gregarious manner^{3–7} and their juveniles often come into physical contact through growth and then fuse^{8–10}. These life history traits increase the opportunities for fusion, and important rates of chimerism due to allogenic fusion have been detected in wild natural populations of corals and ascidians^{11,12}. Nevertheless, fusion of embryos or larvae during planktonic and dispersive phase (i.e. prior to settlement and metamorphosis) is rarely known to date.

Modular marine invertebrates like sponges and cnidarians usually spawn in a high synchrony and the embryos also tend to aggregate after release^{13–15}, thus also providing the chance of contact and fusion among embryos or larvae. For instance, sticky eggs released by the oviparous sponge *Cliona celata* were found to adhere to each other and form flattened egg mass, within which larvae fused in twos or threes. The compound larvae metamorphosed into sponges with single oscula, indicating the cytotoxic fusion among embryos or larvae¹³. More recently, larvae of viviparous sponge *Haliclona* sp. have been demonstrated to fuse and generate swimming chimeras¹⁶. Furthermore, sexually produced embryos of a non-colonial sea anemone *Urticina felina* were observed to fuse naturally during internal brooding, generating pre-metamorphic cytotoxic and sectorial chimeras^{17,18}. These findings suggested that the direct contact between embryos and larvae would facilitate fusion either during internal brooding or pelagic phase.

For broadcast spawning corals, synchronous spawning events usually result in billions of naked embryos floating at the sea surface in the form of spawn slicks^{19,20}. The direct contact between naked embryos highlights the possibility of fusion of coral embryos while sticking together in slicks. Moreover, previous studies have demonstrated there is a window in ontogeny, before allorecognition system matures, when newly settled polyps can fuse²¹. Time for allorecognition maturation in reef corals varied from 4 months following settlement in brooding species²², to 1–3 years in spawning species^{9,10}. This further supports the possibility of fusion at embryonic stage when allorecognition may be weak in corals. As yet, the possible occurrence of fused embryos and conjoined larvae in broadcast spawning corals has not been investigated.

Here, we happened to test this unexplored probability of fusion of embryos in broadcast spawning reef corals. We experimentally mimicked spawn slicks using gametes collected from 4 mature colonies of *Platygyra daedalea*, and followed the fate and development of embryos within lab-generated slicks.

Materials and methods

Ten gravid colonies of *P. daedalea* (20–30 cm in diameter) were collected at depth between 2–4 m from Luhuitou fringing reef in Sanya, China (18°12'N, 109°28'E). Corals were maintained in an

outdoor tank with flowing sand-filtered seawater in Tropical Marine Biological Research Station in Hainan, Sanya. Four colonies spawned around 22:00 on May 18, 2014 (5 nights after full moon). Egg-sperm bundles were collected using pipettes, then mixed and gently agitated to facilitate bundle disintegration and cross-fertilization. Fertilization was allowed to take place for about 2 hours, after which eggs (ca. 300, 000) were washed two times with fresh seawater and suspended in a 15 cm-diameter jar. Because of the logistical constraints, eggs were left undisturbed and they formed dense slicks on the seawater surface. The next morning around 08:30, embryos were inspected under a dissecting microscope and we accidentally discovered that some embryos fused. Embryos were washed and seawater was changed twice daily thereafter. Two days after fertilization, 500 larvae were randomly sampled to count the proportion of each type of fusion. Seven days after fertilization, chips of crustose coralline algae *Hydrolithon onkodes* were used to induce the settlement of larvae and the recruits were reared in the lab at 28°C until June 26.

Results

Embryos became bowl shaped (cushion stage) 8 h after fertilization. Notably, some embryos fused (Figure 1A) and a substantial proportion even stuck together into dense aggregates (Figure 1B). It could be deduced that fusion of embryos took place some time during blastulation. Mortality of embryos within the first 2 days was extremely high (>50%) and the dense aggregates all died and decomposed. Unitary larvae became pear-shaped and began to rotate actively 20 h after fertilization, while conjoined larvae were highly variable in shape. Bi-fused larvae were dominantly peanut-shaped, and multi-fused larvae were arranged in chains or triangles, or in the form of the letter “T” or “L” (Figure 1C, D).

Of the 500 randomly sampled larvae, 174 (34.8%) were conjoined with 2–4 partners. Conjoined larvae clearly showed their spatial arrangement after elongation and fusion was apparently without polarity. Larvae either joined at the aboral end (Figure 1E, F), or united side by side (Figure 1G), or even fused perpendicularly (Figure 1K). Furthermore, 56 out of the 174 conjoined larvae (32.2%) united at the aboral extremity and only these larvae were potentially competent to metamorphose normally into inborn colonies (Figure 1H, I), which were prominently larger in size than the single settlers (Figure 1J). In contrast, perpendicularly bi-fused larvae settled incompletely, with one partner metamorphosing and firmly attaching while the other still being parallel to the substrate and not able to settle (Figure 1K), ultimately leading to the death of the whole entity 3 days later. Since the coralline algae provided here was not suitable for the settlement of *P. daedalea* larvae, only 12 inborn colonies were obtained in total and they persisted for 26 days post-settlement when the study ended (Figure 1L).

Discussion

The present study documented for the first time the fusion of embryos, the subsequent formation of conjoined larvae and inborn colonies in a broadcast spawning coral. Fusion of *P. daedalea* embryos was spontaneous, resulting simply from the aggregation and contact of embryos in mimicked slicks, which was analogous with that in sponge *C. celata*¹³. While unlike the cytotoxic compound larvae in sponge *C. celata*, partners within the conjoined

Figure 1. *Platygyra daedalea*. (A) Fused embryos (arrows point to the fusing areas). (B) A dense aggregate comprising more than 20 embryos. (C–G) Unitary (asterisks) and conjoined larvae. (H, I) Inborn colonies. (J) Single settlers. (K) Incomplete settlement of perpendicularly bi-fused larvae, with the left partner being parallel to the substrate. (L) An inborn colony 26 days post-settlement. Roman numerals indicate visible individuals within inborn colonies. Scale bars 250 µm.

P. daedalea larvae still retained a degree of individuality, suggesting sectorial fusion of coral embryos and supporting the assumption that corals typically exhibit sectorial fusion¹.

Corals often spawn their gametes during seasonally calm periods and low-amplitude tides^{19,20,23} and spawn slicks extending up to few km in length were often observed in the field^{20,24}. Given that slicks remained aggregated 1–2 d after spawning^{19,20} and embryos can fuse during embryogenesis within 8 h post-fertilization, fusion of coral embryos is highly favored *in situ*. On the other hand, although mass coral spawning events usually involved several species, significant temporal differences in spawning to ensure fertilization and reproductive isolation have been demonstrated for many sympatric species^{25–27}, which considerably increase the encounters between embryos of the same species in slicks. Taken together, fusion of coral embryos might be a natural phenomenon. However, the density of embryos here was 1700 cm⁻² and likely to be much higher than that in the field. Therefore, the extent to which fusion of embryos occurs in natural slicks and whether there is a density effect on the formation of conjoined larvae remain to be fully evaluated.

At last, approximately 12 percent of all the larvae had the potential to form inborn colonies, which persisted for one month and exhibited no sign of rejection. The inborn colonies here originated from fusion of embryos and settlement of conjoined larvae, contrasting the traditional concept that the asexual budding of the primary polyp leads to the formation of a young coral colony²⁸, and thus fusion of embryos could be an unexpected shortcut to colony formation in reef corals. These facts raise questions as to the ecological implications of inborn colonies formed as a consequence of

fusion of embryos in corals. Firstly, larger coral colonies composed of multiple fused partners are known to yield remarkable gained benefits, such as enhanced survival and growth^{5,8}. Hence, the larger initial size and the status of multi-polyp at settlement may confer these inborn colonies better capacities to compete for space and survive partial mortality.

Fusion of coral embryos also shed new light on the chimerism in scleractinian corals, which was often attributed to fusion of gregariously settling larvae^{5,7}, or of juveniles that come into contact through growth^{7,9,10}. However, our study documented fusion between individuals in *P. daedalea* occurred at the embryonic stage, earlier than any other corals studied to date. It should be pointed out that though embryos here were produced sexually from 4 parent colonies, it did not denote they were genetically distinct. Therefore, the possibility of isogenic fusion can not be totally ruled out⁷. However, it is reasonable to assume the presence of allogenic fusion, suggesting fusion of embryos might add a novel mechanism to form chimeric larvae and colonies in scleractinian corals. In that case, the increased genetic diversity within these inborn colonies may translate into versatile physiological qualities, thus enabling them to better cope with environmental changes unless negative interactions occur^{1,29,30}.

Overall, this is the first report of fused embryos and pre-metamorphic conjoined larvae in reef corals. Fusion of coral embryos could be an adaptive strategy to form larger and chimeric recruits, thereby promoting growth and survival during the vulnerable early stages^{5,8}. Clearly, future studies are required to explore whether fusion of embryos is common in spawning corals and fully evaluate its biological and ecological implications.

Ethics statement

Coral sampling was permitted by the Administration of Sanya Coral Reef National Nature Reserve, the Department of Ocean and Fisheries of Hainan Province.

Author contributions

LJ conceived and performed the study. All authors wrote the manuscript and gave final consent for publication.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by Public Science and Technology Research Funds Projects of Ocean (201305030-3) and the National Natural Science Foundation of China (41306144 and U1301232).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Rinkevich B, Weissman IL: **Chimeras in colonial invertebrates: a synergistic symbiosis or somatic and germ-cell parasitism?** *Symbiosis*. 1987; **4**(1–3): 117–134.
[Reference Source](#)
- Buss LW: **Competition within and between encrusting clonal invertebrates.** *Trends Ecol Evol*. 1990; **5**(11): 352–356.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Keough MJ: **Kin-recognition and the spatial distribution of larvae of the bryozoan *Bugula neritina* (L.).** *Evolution*. 1984; **142**–147.
[Publisher Full Text](#)
- Barki Y, Gateño D, Graur D, *et al.*: **Soft-coral natural chimerism: a window in ontogeny allows the creation of entities comprised of incongruous parts.** *Mar Ecol Progr Ser*. 2002; **231**: 91–99.
[Publisher Full Text](#)
- Amar KO, Chadwick NE, Rinkevich B: **Coral kin aggregations exhibit mixed allogeneic reactions and enhanced fitness during early ontogeny.** *BMC Evol Biol*. 2008; **8**(1): 126–136.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Westerman E, Dijkstra J, Harris L: **High natural fusion rates in a botryllid ascidian.** *Mar Biol*. 2009; **156**(12): 2613–2619.
[Publisher Full Text](#)
- Puill-Stephan E, van Oppen MJ, Pichavant-Rafini K, *et al.*: **High potential for formation and persistence of chimeras following aggregated larval settlement in the broadcast spawning coral, *Acropora millepora*.** *Proc Biol Sci*. 2012; **279**(1729): 699–708.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Raymundo LJ, Maypa AP: **Getting bigger faster: Mediation of size-specific mortality via fusion in juvenile coral transplants.** *Ecol Appl*. 2004; **14**(1): 281–295.
[Publisher Full Text](#)
- Nozawa Y, Hirose M: **When does the window close?: The onset of allogeneic fusion 2–3 years post-settlement in the scleractinian coral, *Echinophyllia aspera*.** *Zool Stud*. 2011; **50**: 396.
[Reference Source](#)
- Puill-Stephan E, Willis BL, Abrego D, *et al.*: **Allorecognition maturation in the broadcast-spawning coral *Acropora millepora*.** *Coral Reefs*. 2012; **31**(4): 1019–1028.
[Publisher Full Text](#)
- Dorothea S, Bishop JDD: **Random amplified polymorphic DNA (RAPD) analysis reveals extensive natural chimerism in a marine protochordate.** *Mol Ecol*. 1999; **8**(5): 885–890.
[Publisher Full Text](#)
- Puill-Stephan E, Willis BL, van Herwerden L, *et al.*: **Chimerism in wild adult populations of the broadcast spawning coral *Acropora millepora* on the Great Barrier Reef.** *PLoS One*. 2009; **4**(11): e7751.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Warburton FE: **Reproduction of fused larvae in the boring sponge, *Cliona celata* Grant.** *Nature*. 1958; **181**(4607): 493–494.
[Publisher Full Text](#)
- Harrison PL, Babcock RC, Bull GD, *et al.*: **Mass spawning in tropical reef corals.** *Science*. 1984; **223**(4641): 1186–1189.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mariani S, Piscitelli M, Uriz MJ: **Temporal and spatial co-occurrence in spawning and larval release of *Cliona viridis* (Porifera: Hadromerida).** *J Mar Biol Assoc Uk*. 2001; **81**(04): 565–567.
[Reference Source](#)
- McGhee KE: **The importance of life-history stage and individual variation in the allorecognition system of a marine sponge.** *J Exp Mar Biol Ecol*. 2006; **333**(2): 241–250.
[Publisher Full Text](#)
- Mercier A, Sun Z, Hamel JF: **Internal brooding favours pre-metamorphic chimerism in a non-colonial cnidarian, the sea anemone *Urticina felina*.** *Proc Biol Sci*. 2011; **278**(1724): 3517–3522.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sun Z, Hamel JF, Mercier A: **Marked shifts in offspring size elicited by frequent fusion among siblings in an internally brooding marine invertebrate.** *Am Nat*. 2012; **180**(5): E151–E160.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gilmour JP, Smith LD, Brinkman RM: **Biannual spawning, rapid larval development and evidence of self-seeding for scleractinian corals at an isolated system of reefs.** *Mar Biol*. 2009; **156**(6): 1297–1309.
[Publisher Full Text](#)
- Oliver JK, Willis BL: **Coral-spawn slicks in the Great Barrier Reef: preliminary observations.** *Mar Biol*. 1987; **94**(4): 521–529.
[Publisher Full Text](#)
- Rinkevich B: **Allorecognition and xenorecognition in reef corals: a decade of interactions.** In *Coelenterate Biology 2003*, D. Fautin, *et al.*, Editors. Springer Netherlands, 2004; p. 443–450.
[Publisher Full Text](#)
- Frank U, Oren U, Loya Y, *et al.*: **Alloimmune maturation in the coral *Stylophora pistillata* is achieved through three distinctive stages, 4 months post-metamorphosis.** *Proc Biol Sci*. 1997; **264**(1378): 99–104.
[Publisher Full Text](#) | [Free Full Text](#)
- van Woessik R: **Calm before the spawn: global coral spawning patterns are explained by regional wind fields.** *Proc Biol Sci*. 2010; **277**(1682): 715–722.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harrison PL: **Coral spawn slicks at Lord Howe Island, Tasman Sea, Australia; the world's most southerly coral reef.** *Coral Reefs*. 2008; **27**(1): 35.
[Publisher Full Text](#)
- Van Oppen MJH, Willis BL, Van Rheede T, *et al.*: **Spawning times, reproductive compatibilities and genetic structuring in the *Acropora aspera* group: evidence for natural hybridization and semi-permeable species boundaries in corals.** *Mol Ecol*. 2002; **11**(8): 1363–1376.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fukami H, Omori M, Shimoi K, *et al.*: **Ecological and genetic aspects of reproductive isolation by different spawning times in *Acropora* corals.** *Mar Biol*. 2003; **142**(4): 679–684.
[Publisher Full Text](#)
- Leviton DR, Fukami H, Jara J, *et al.*: **Mechanisms of reproductive isolation among sympatric broadcast-spawning corals of the *Montastrea annularis* species complex.** *Evolution*. 2004; **58**(2): 308–323.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Harrison PL, Wallace CC: **Reproduction, dispersal and recruitment of scleractinian corals.** In *Ecosystems of the world: coral reefs*, Z. Dubinsky, Editor. Elsevier: Amsterdam, 1990; 133–207.
[Reference Source](#)
- Pineda-Krch M, Lehtilä K: **Costs and benefits of genetic heterogeneity within organisms.** *J Evol Biol*. 2004; **17**(6): 1167–1177.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rinkevich B, Yankelovich I: **Environmental split between germ cell parasitism and somatic cell synergism in chimeras of a colonial urochordate.** *J Exp Biol*. 2004; **207**(Pt 20): 3531–3536.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 17 February 2015

<https://doi.org/10.5256/f1000research.6575.r7679>

© 2015 Rinkevich B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Baruch Rinkevich

Israel Oceanography & Limnological Research, National Institute of Oceanography, Haifa, Israel

This short research article by Jiang *et al.* is timely, adds to our understanding on the magnitude of natural chimerism, and meets research standards with **methodological details that made available** to allow others to replicate this study. I recommend indexation pending responding on the below minor editorial/clarification points:

- Abstract (line 2)- An unclear sentence. Suggest revising from: 'Here, we observed for the first time the embryos....' to 'Here, we document for the first time that embryos...'. See also below the note appended to the citation of Mizrahi et al., 2014 and define your statement, 'for the first time'.
- 'Conjoined larvae' (as from Abstract line 4 and along the manuscript). This term and the following terms 'chimera' and 'fusion' are mutually used, irrespective to the somehow different biological statuses they present (for example, 'conjoined' is a very sterile term indicating being, coming, or brought together so as to meet, touch, overlap, or unit). I would suggest to use the term 'Conjoined larvae' in the Results section when first describing the process of larval joining and then use the terms 'fusion' and 'chimerism' in all other parts of the manuscript to state the biological outcome.
- Related to the aforementioned- it will be most valuable to show evidence for the 'fusion' outcomes, such as histological sections, or that no disjoining between the conjoined larvae occurred following the employment of physical force.
- Introduction and Discussion- There are four papers that I would suggest the authors to add and to cite. The first and the most important is the study by Mizrahi *et al.* (2014). In this paper the authors showed that *Tubastraea coccinea* offspring can metamorphose and aggregate/fuse in groups of up to eight polyps in the water column, before settlement. This may also change the way Jiang *et al.* refer to their finding (statements such as 'the first time' should be revised). The second paper is the study on sponge chimeras by Gauthier and Degnan (2008). This paper reveals the potential intermixing of cells between fused partners. The third paper is the old report by Duerden, the first observation documenting 'aggregated colonies' (Duerden, 1902). It will also be valuable to cite Rinkevich (2011).

- Introduction 2nd paragraph, 1st sentence: 'Modular marine invertebrates .... tend to aggregate after release' is redundant to the former introduction text. Delete or rephrase.
- M & M: '... then mixed and gently agitated...' Specify if the mixed eggs/sperm were from the same genotype or from all/several genotypes.
- M & M: '...to count the proportion of each type of fusion'. Which 'types' of fusion the authors imply to? Detail/explain.
- Discussion, below Fig 1: '...*P. daedalea* larvae still retained a degree of individuality...'. What is the meaning of 'individuality' here? What is the meaning of 'a degree'? Morphologically? Physiologically? Define and rephrase.
- Discussion, the sentence: 'It should be pointed out that though embryos here were produced sexually from 4 parent colonies, it did not denote they were genetically distinct'. Something is wrong with the sentence. If larvae are sexually produced they should be genetically distinct from each other. Rewrite.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 26 Feb 2015

Lei Jiang, Chinese Academy of Sciences, Guangzhou, China

- We revised the sentence in abstract as suggested. Our observation presented the embryonic chimeras as a result of fusion of embryos in a broadcast spawning coral, whereas the results of Mizrahi *et al.*, (2014) revealed that the larvae of brooding coral *Tubastraea coccinea* could metamorphose and aggregate in swimming groups. Thus we did document for the first time the fusion between individuals at the embryonic stage in reef corals. Moreover, we followed the settlement and growth of these chimeric larvae. Overall, it is proper and discreet to state "for the first time".
- We have followed the reviewer's advice to use "conjoined larvae" in the Results and Abstract sections where described the larval joining.
- This was only an accidental observation and it was a pity that we did not preserve samples for histological sections, nor did we employ physical force to see whether the conjoined larvae would disjoint.
- As aforementioned, we did observe the embryonic chimeras and follow the settlement and post-settlement growth of chimeric larvae in a broadcast spawning coral for the first time. Mizrahi *et al.* (2014) revealed that the swimming polyp clusters survived longer and did not confirm whether these clusters can settle. Therefore, the

findings of Mizrahi *et al.* (2014) were not contradictory to our statement of “the first time”. We have added the suggested references. But as for Rinkevich (2011), we think it is more suitable to cite the classical reference of Rinkevich *et al.* (1987) which mainly focused on the chimerism in colonial marine invertebrates.

- We deleted the latter part of this sentence and made it more specific as to the synchrony in gametes release.
- We used the word “combined” to show that the bundles from 4 colonies were brought together for cross fertilization.
- We replaced “each type of fusion” with “chimeric larvae”.
- We made it more specifically that these conjoined larvae were multi-headed to demonstrate the sectorial fusion.
- We were sorry for this mistake because we were a little confused at first when referring to the results of Puill-Stephan *et al.* (2012). They found that fusion between sexually produced larvae resulted in non-chimeric colonies of one genotype. Thus we got a misunderstanding that the sexually produced larvae were not genetically distinct absolutely. However, they only use nine microsatellite loci to genotype. We have corrected and rewritten this part.

Competing Interests: No competing interests were disclosed.

Reviewer Report 16 February 2015

<https://doi.org/10.5256/f1000research.6575.r7666>

© 2015 Baird A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Andrew H. Baird

School of Marine Biology and Aquaculture, James Cook University, Townsville, QLD, Australia

This is well written article and an interesting observation although some of the literature cited is a little dated (e.g. Harrison & Wallace 1990). To my knowledge this is first time that anyone has followed these embryonic chimeras through to settlement and that fact that they can settle and grow is an important observation. However, embryonic chimera formation is an artifact of the experimental conditions i.e. the very high densities of embryos in still water in the bowls. Indeed, embryonic chimera formation is quite common, particularly when working with larvae of merulinds at higher temperatures. It is very rare to see chimeras in spawn slicks or on settlement tiles, at least on soaks of short duration, although this has yet to be quantified. Therefore,

embryonic chimera formation is highly unlikely to be of much ecological or evolutionary significance. Indeed, a phenomenon of much more ecological significance is the fact that many embryos break up during development under conditions likely to prevail in the wild (see Heyward and Negri, 2012).

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 26 Feb 2015

Lei Jiang, Chinese Academy of Sciences, Guangzhou, China

Spawn slicks are commonly observed in the field and more importantly, they remained aggregated 1-2 days after spawning (Oliver *et al.*, 1987; Gilmour *et al.*, 2009). It is true that these conditions are favorable for the formation of embryonic chimeras. Moreover, the high density and absence of water turbulence in this study might trigger the fusion of embryos. Under similar experimental conditions, larvae of *A. millepora* tended to settle in aggregation (Puill-Stephan *et al.*, 2012), and larvae of *T. coccinea* formed swimming polyp clusters (Mizrahi *et al.* (2014)). Therefore, though fusion of coral embryos has never been reported in the wild, we speculate it might occur in the field and remains to be determined especially corals often broadcast spawn their gametes during calm periods and low-amplitude tides.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research