OPINION ARTICLE

Hypothesis: Is frequent, commercial jet travel by the general public a risk factor for developing cutaneous melanoma?

[version 1; peer review: 2 approved, 1 approved with reservations]

Harvey Arbesman¹⁻³

¹Department of Epidemiology and Environmental Health, University at Buffalo School of Public Health and Health Professions, Buffalo, NY, 14214-8001, USA

²Division of Cancer Prevention and Population Sciences, Roswell Park Cancer Institute, Buffalo, NY, 14263, USA

³Department of Dermatology, Clinical and Translational Research Center, University at Buffalo School of Medicine and Biomedical Sciences, Buffalo, NY, 14203, USA

V1 First published: 24 Aug 2015, 4:620
<https://doi.org/10.12688/f1000research.6962.1>
 Latest published: 24 Aug 2015, 4:620
<https://doi.org/10.12688/f1000research.6962.1>

Abstract

Melanoma incidence has been increasing worldwide over the past 50 years and various risk factors have been identified. Interestingly, multiple studies have shown a multifold increased risk of developing melanoma in jet pilots and airline crew. There has also been a dramatic increase in the availability and frequency of jet travel by the general population during this time period.. Therefore, it is hypothesized that frequent commercial jet travel may represent an additional risk factor for the development of cutaneous melanoma in susceptible individuals of the general public.

Keywords

melanoma , risk factor , epidemiology , jet travel , cosmic radiation

Open Peer Review

Approval Status

	1	2	3
version 1 24 Aug 2015	view	view	view

1. **Desiree Ratner**, Icahn School of Medicine at Mount Sinai, New York, USA
2. **Robyn Lucas**, Australian National University, Canberra, Australia
3. **Gabriella Fabbrocini**, University of Naples Federico II, Naples, Italy

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Harvey Arbesman (arbesman@buffalo.edu)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2015 Arbesman H. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Arbesman H. **Hypothesis: Is frequent, commercial jet travel by the general public a risk factor for developing cutaneous melanoma?** [version 1; peer review: 2 approved, 1 approved with reservations] F1000Research 2015, 4:620 <https://doi.org/10.12688/f1000research.6962.1>

First published: 24 Aug 2015, 4:620 <https://doi.org/10.12688/f1000research.6962.1>

Background

The incidence of cutaneous melanoma in light pigmented individuals worldwide has been steadily increasing over the past 4–5 decades^{1–4}. Sun exposure, including intermittent exposure, is an important environmental risk factor for melanoma, along with history of sunburn, residence in equatorial latitudes and tanning bed usage^{5–11}. Genetic risk factors include red hair, family history of melanoma, dysplastic nevi, lightly pigmented skin, tendency to burn, inability to tan, and DNA repair defects¹². Phenotypic expressions of gene/environmental interactions are risk factors and include melanocytic nevi (increased total number, multiple atypical [dysplastic], and congenital [particularly large axial lesions with multiple satellites]), ephelides, and personal history of melanoma^{12,13}. Non-solar occupational risk factors have also been shown to be associated with the development of melanoma¹⁴. Since no specific currently recognized risk factor adequately explains the rapidly increasing incidence among the general population, the identification of new risk factors that could play a role in melanoma prevention is needed.

Studies from many countries have shown a significant increase in the risk of melanoma in commercial and military pilots; these range from approximately 2 to 10 fold^{15–20}. An increased risk has also been shown in some studies of cabin crew^{21–23}. Recently a meta-analysis was published that also showed an increased risk in pilots and cabin crew²⁴.

Though it is unclear why aircrew have an increased risk of melanoma, one proposed explanation regarding their increased risk has focused on the exposure to cosmic ionizing radiation present at cruising altitudes of 30,000 feet or higher^{25–34}. Currently, commercial jets frequently cruise at an altitude that results in cosmic radiation exposure^{26,28,30–33}. In addition, the cosmic radiation exposure is increased with higher altitude flights and long-haul routes²⁵. Cosmic ionizing radiation contains multiple particles that can damage DNA^{28,32}. Epidemiologic studies have reported a possible relationship between melanoma and exposure to ionizing radiation in other occupational settings^{35–37}. A recent systematic review regarding cosmic radiation and cancer assessed the role cosmic radiation plays in the development of cancer as compared with other lifestyle factors²⁵.

Another proposed explanation for the documented increased risk among pilots is their increased exposure to UV light during flight^{24,38}. Recently an analysis was done and found greater amounts of UVA exposure through the windows of jets³⁹. At this time, it is unclear whether increased UVA exposure, cosmic ionizing radiation, circadian rhythm disturbance and/or other undetermined factors are contributing to this increased risk of melanoma in pilots and cabin crew²⁶.

Presentation of the hypothesis

It is hypothesized that individuals of the general population who frequently travel by jet plane may also have an increased risk of developing cutaneous melanoma in a similar fashion as pilots and cabin crew. This hypothesized risk factor may play a role in the development of melanoma due to a variety of potentially harmful exposures associated with frequent jet travel that could interact synergistically with other known genetic and environmental risk factors in susceptible individuals. This hypothesis is based on the findings that both pilots and cabin crew, generally healthier than the general public¹⁸, and with a prevalence of skin cancer risk factors similar to that of the general public^{21,40}, have a dramatically higher risk of melanoma.

Since the late 1950s, commercial jets have begun to cruise at an altitude of 30,000 feet or higher. This major change in the flying altitude of air travel is consistent with the temporal nature of the rapid increase in the incidence of melanoma. During the past 50 years, the availability and frequency of jet travel among the general public have increased dramatically. In addition, with the deregulation of the airline industry in the 1970s, and the resulting decrease in fares, a higher percentage of the overall population began to experience jet travel. An ecologic study found an association between accessibility to air travel and the incidence rate of melanoma⁴¹.

A widely accepted epidemiologic finding consistent with the proposed hypothesis is that intermittent sun exposure is an independent risk factor for melanoma^{5,11}. The correlation between melanoma incidence and ‘sun holidays’^{42–46} has been primarily interpreted as secondary to intermittent sun exposure. The proposed hypothesis suggests that the increased risk associated with ‘sun holidays,’ may also be related to jet travel and the resultant exposures associated with jet travel to those vacation destinations. Higher socioeconomic status (SES) is also a risk factor for melanoma development^{42,47}; and higher SES is associated with increased jet travel and ‘sun holidays’^{44,45,48} all findings consistent with the hypothesis.

Testing the hypothesis

A case-control methodology could be utilized to test this hypothesis. One would assess jet travel histories in melanoma patients and comparable controls, controlling for known risk factors such as age, skin type, genetic host factors and sun exposure history. One would obtain assessments of subjects’ jet travel history in terms of frequency, duration of flights and altitude^{49,50}. Assessment of travel routes, season of travel and increased sunspot activity during flights should also be undertaken. It would be necessary to disentangle intermittent sun exposure history from jet travel history.

Implications of the hypothesis

It is hypothesized that frequent commercial jet travel by the general public may increase the risk of developing melanoma due to various harmful exposures associated with frequent jet travel. This melanoma and jet travel hypothesis has potential for reducing melanoma-associated morbidity and mortality and warrants properly designed analytic epidemiologic evaluation to assess the validity of this hypothesis. In addition, if demonstrated to be a risk factor, evaluation of the underlying mechanisms behind this increased risk may lead to the expansion of basic science research of etiologic factors of melanoma and of cancer in general.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgements

I thank Joshua Arbesman, MD, James Marshall, PhD and Thomas Helm, MD for fruitful discussions about the hypothesis and reading drafts of the manuscript.

References

1. Erdmann F, Lortet-Tieulent J, Schüz J, *et al.*: **International trends in the incidence of malignant melanoma 1953-2008—are recent generations at higher or lower risk?** *Int J Cancer*. 2013; **132**(2): 385–400.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Holterhues C, Hollestein LM, Nijsten T, *et al.*: **Burden of disease due to cutaneous melanoma has increased in the Netherlands since 1991.** *Br J Dermatol*. 2013; **169**(2): 389–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Erdei E, Torres SM: **A new understanding in the epidemiology of melanoma.** *Expert Rev Anticancer Ther*. 2010; **10**(11): 1811–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. de Vries E, Bray FI, Coebergh JW, *et al.*: **Changing epidemiology of malignant cutaneous melanoma in Europe 1953-1997: rising trends in incidence and mortality but recent stabilizations in western Europe and decreases in Scandinavia.** *Int J Cancer*. 2003; **107**(1): 119–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Chang YM, Barrett JH, Bishop DT, *et al.*: **Sun exposure and melanoma risk at different latitudes: a pooled analysis of 5700 cases and 7216 controls.** *Int J Epidemiol*. 2009; **38**(3): 814–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Gandini S, Sera F, Cattaruzza MS, *et al.*: **Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure.** *Eur J Cancer*. 2005; **41**(1): 45–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Solomon CC, White E, Kristal AR, *et al.*: **Melanoma and lifetime UV radiation.** *Cancer Causes Control*. 2004; **15**(9): 893–902.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Lea CS, Scotto JA, Buffler PA, *et al.*: **Ambient UVB and melanoma risk in the United States: a case-control analysis.** *Ann Epidemiol*. 2007; **17**(6): 447–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Gandini S, Stanganelli I, Magi S, *et al.*: **Melanoma attributable to sunbed use and tan seeking behaviours: an Italian survey.** *Eur J Dermatol*. 2014; **24**(1): 35–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Boniol M, Autier P, Boyle P, *et al.*: **Cutaneous melanoma attributable to sunbed use: systematic review and meta-analysis.** *BMJ*. 2012; **345**: e4757.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Elwood JM, Jopson J: **Melanoma and sun exposure: an overview of published studies.** *Int J Cancer*. 1997; **73**(2): 198–203.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Gandini S, Sera F, Cattaruzza MS, *et al.*: **Meta-analysis of risk factors for cutaneous melanoma: III. Family history, actinic damage and phenotypic factors.** *Eur J Cancer*. 2005; **41**(14): 2040–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Gandini S, Sera F, Cattaruzza MS, *et al.*: **Meta-analysis of risk factors for cutaneous melanoma: I. Common and atypical naevi.** *Eur J Cancer*. 2005; **41**(1): 28–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Fortes C, de Vries E: **Nonsolar occupational risk factors for cutaneous melanoma.** *Int J Dermatol*. 2008; **47**(4): 319–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Yong LC, Pinkerton LE, Yin JH, *et al.*: **Mortality among a cohort of U.S. commercial airline cockpit crew.** *Am J Ind Med*. 2014; **57**(8): 906–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Hammer GP, Auvinen A, De Stavola BL, *et al.*: **Mortality from cancer and other causes in commercial airline crews: a joint analysis of cohorts from 10 countries.** *Occup Environ Med*. 2014; **71**(5): 313–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. dos Santos Silva I, De Stavola B, Pizzi C, *et al.*: **Cancer incidence in professional flight crew and air traffic control officers: disentangling the effect of occupational versus lifestyle exposures.** *Int J Cancer*. 2013; **132**(2): 374–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Sykes AJ, Larsen PD, Griffiths RF, *et al.*: **A study of airline pilot morbidity.** *Aviat Space Environ Med*. 2012; **83**(10): 1001–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Buja A, Lange JH, Perissinotto E, *et al.*: **Cancer incidence among male military and civil pilots and flight attendants: an analysis on published data.** *Toxicol Ind Health*. 2005; **21**(10): 273–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Rafnsson V, Hrafnkelsson J, Tulinius H: **Incidence of cancer among commercial airline pilots.** *Occup Environ Med*. 2000; **57**(3): 175–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Kojo K, Helminen M, Pukkala E, *et al.*: **Risk factors for skin cancer among Finnish airline cabin crew.** *Ann Occup Hyg*. 2013; **57**(6): 695–704.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Pukkala E, Helminen M, Haldorsen T, *et al.*: **Cancer incidence among Nordic airline cabin crew.** *Int J Cancer*. 2012; **131**(12): 2886–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Buja A, Mastrangelo G, Perissinotto E, *et al.*: **Cancer incidence among female flight attendants: a meta-analysis of published data.** *J Womens Health (Larchmt)*. 2006; **15**(1): 98–105.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Sanlorenzo M, Wehner MR, Linos E, *et al.*: **The risk of melanoma in airline pilots and cabin crew: a meta-analysis.** *JAMA Dermatol*. 2015; **151**(1): 51–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Di Trollo R, Di Lorenzo G, Fumo B, *et al.*: **Cosmic radiation and cancer: is there a link?** *Future Oncol*. 2015; **11**(7): 1123–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Grajewski B, Pinkerton LE: **Exposure assessment at 30 000 feet: challenges and future directions.** *Ann Occup Hyg*. 2013; **57**(6): 692–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Schüz J: **Airline crew cohorts: is there more to learn regarding their cancer risk?** *Occup Environ Med*. 2014; **71**(5): 307.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Sigurdson AJ, Ron E: **Cosmic radiation exposure and cancer risk among flight crew.** *Cancer Invest*. 2004; **22**(5): 743–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Lim MK: **Cosmic rays: are air crew at risk?** *Occup Environ Med*. 2002; **59**(7): 428–32; discussion 432–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Friedberg W, Copeland K, Duke FE, *et al.*: **Radiation exposure during air travel: guidance provided by the Federal Aviation Administration for air carrier crews.** *Health Phys*. 2000; **79**(5): 591–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Kendall GM: **Factors affecting cosmic-ray doses at aircraft altitudes.** *Health Phys*. 2000; **79**(5): 560–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Goldhagen P: **Overview of aircraft radiation exposure and recent ER-2 measurements.** *Health Phys*. 2000; **79**(5): 526–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Townsend LW: **Invited editorial: Radiation exposures of aircrew in high altitude flight.** *J Radiol Prot*. 2001; **21**(1): 5–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Menzel HG, O'Sullivan D, Beck P, *et al.*: **European measurements of aircraft crew exposure to cosmic radiation.** *Health Phys*. 2000; **79**(5): 563–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Fink CA, Bates MN: **Melanoma and ionizing radiation: is there a causal relationship?** *Radiat Res*. 2005; **164**(5): 701–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Freedman DM, Sigurdson A, Rao RS, *et al.*: **Risk of melanoma among radiologic technologists in the United States.** *Int J Cancer*. 2003; **103**(4): 556–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Caldwell GG, Kelley D, Zack M, *et al.*: **Mortality and cancer frequency among military nuclear test (Smoky) participants, 1957 through 1979.** *JAMA*. 1983; **250**(5): 620–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Chorley AC, Evans BJ, Benwell MJ: **Civilian pilot exposure to ultraviolet and blue light and pilot use of sunglasses.** *Aviat Space Environ Med*. 2011; **82**(9): 895–900.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Sanlorenzo M, Vujic I, Posch C, *et al.*: **The risk of melanoma in pilots and cabin crew: UV measurements in flying airplanes.** *JAMA Dermatol*. 2015; **151**(4): 450–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Rafnsson V, Hrafnkelsson J, Tulinius H, *et al.*: **Risk factors for cutaneous malignant melanoma among aircrews and a random sample of the population.** *Occup Environ Med*. 2003; **60**(11): 815–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Agredano YZ, Chan JL, Kimball RC, *et al.*: **Accessibility to air travel correlates strongly with increasing melanoma incidence.** *Melanoma Res*. 2006; **16**(1): 77–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Idorn LW, Wulf HC: **Socioeconomic status and cutaneous malignant melanoma in Northern Europe.** *Br J Dermatol*. 2014; **170**(4): 787–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Veierød MB, Adami HO, Lund E, *et al.*: **Sun and solarium exposure and melanoma risk: effects of age, pigimentary characteristics, and nevi.** *Cancer Epidemiol Biomarkers Prev*. 2010; **19**(1): 111–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Benthem G, Aase A: **Incidence of malignant melanoma of the skin in Norway, 1955-1989: associations with solar ultraviolet radiation, income and holidays abroad.** *Int J Epidemiol*. 1996; **25**(6): 1132–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Westerdaal J, Olsson H, Ingvar C, *et al.*: **Southern travelling habits with special reference to tumour site in Swedish melanoma patients.** *Anticancer Res*. 1992; **12**(5): 1539–42.
[PubMed Abstract](#)

46. Osterlind A, Tucker MA, Stone BJ, *et al.*: **The Danish case-control study of cutaneous malignant melanoma. II. Importance of UV-light exposure.** *Int J Cancer*. 1988; **42**(3): 319–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Clegg LX, Reichman ME, Miller BA, *et al.*: **Impact of socioeconomic status on cancer incidence and stage at diagnosis: selected findings from the surveillance, epidemiology, and end results: National Longitudinal Mortality Study.** *Cancer Causes Control*. 2009; **20**(4): 417–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Veierød MB, Weiderpass E, Thörn M, *et al.*: **A prospective study of pigmentation, sun exposure, and risk of cutaneous malignant melanoma in women.** *J Natl Cancer Inst*. 2003; **95**(20): 1530–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Hammer GP, Zeeb H, Tveten U, *et al.*: **Comparing different methods of estimating cosmic radiation exposure of airline personnel.** *Radiat Environ Biophys*. 2000; **39**(4): 227–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Kojo K, Aspholm R, Auvinen A: **Occupational radiation dose estimation for Finnish aircraft cabin attendants.** *Scand J Work Environ Health*. 2004; **30**(2): 157–63.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 30 November 2015

<https://doi.org/10.5256/f1000research.7498.r11326>

© 2015 Fabbrocini G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gabriella Fabbrocini

Department of Dermatology, University of Naples Federico II, Naples, Italy

The hypothesis put forward by the author is interesting, yet hard to prove. It is extremely difficult to separate the weight of individual exposures: the cosmic radiation at high altitudes and that related to sun exposure of the trip. The bibliography is extensive and still valid. Considering the bases from which originates the article, or the increased incidence of melanoma in the last decades and the multifactoriality, it might be interesting to start the case control study that the author sets out to accomplish. It would be necessary to consider some essential factors, including the frequency of air travel, the duration of the flight, altitude, intermittent sun exposure of patients during their trip.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 29 September 2015

<https://doi.org/10.5256/f1000research.7498.r10379>

© 2015 Lucas R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Robyn Lucas

National Centre for Epidemiology and Population Health, Australian National University, Canberra, Australia

The authors raise the hypothesis that commercial jet travel by the general public may increase the

risk of developing cutaneous melanoma through increased exposure to cosmic radiation or UVA radiation. Previous studies have shown an increased risk of melanoma in flight crews and pilots. The hypothesis in relation to UVA seems unlikely, given that most air travellers are well-clothed. The hypothesis in relation to cosmic radiation needs stronger evidence of the measured levels within aircraft. The author proposes a case-control study, but this seems unlikely to provide valid results - the most likely explanation of an increased risk of melanoma in flight crews and pilots is higher sun exposure at the locations that they fly to, especially of the intermittent high dose pattern that confers particular risk of melanoma. However sun exposure is poorly reported (as would be the measure of exposure in a case-control study) while air travel likely to be well-reported (and able to be validated). It will be impossible to clearly separate these "exposures". There is likely to be a very high correlation between a history of air travel and of sunny holidays in people with melanoma, such that a history of air travel is a proxy for high dose intermittent sun exposure. Findings resulting from such a study are highly likely to be spurious associations. It is an interesting hypothesis - a start would be to measure levels of cosmic radiation inside passenger aircraft, more clearly make the link between cosmic radiation and melanoma, and assess the body sites affected in cabin crew and pilots who develop melanoma.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 02 September 2015

<https://doi.org/10.5256/f1000research.7498.r10189>

© 2015 Ratner D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Desiree Ratner

Department of Dermatology, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

The hypothesis, "Is frequent, commercial jet travel by the general public a risk factor for developing cutaneous melanoma?", provides an interesting perspective on possible reasons for the increasing incidence of melanoma worldwide. The author cites multiple studies documenting a significant increase in melanoma risk among commercial and military pilots as well as cabin crew. Possible reasons for this increase include exposure to cosmic ionizing radiation at cruising altitudes of 30,000 feet or higher, particularly on long haul routes, as well as increased exposure to UV light during flight. Given that a significant portion of the world's population now takes advantage of the increased availability and frequency of jet travel, it is possible that the increased incidence of melanoma may be, in part, due to increased exposure to cosmic rays and/or UV exposure during high altitude jet travel.

The author suggests that a case-control methodology be used to test his hypothesis, by assessing jet travel histories in melanoma patients and comparable controls, controlling for known risk

factors such as age, skin type, and exposure history. It would be necessary to look at frequency of air travel, duration of flights, and altitude, and to disentangle patients' intermittent sun exposure history from their jet travel history. If the hypothesis that frequent commercial jet travel represents an additional risk factor for developing cutaneous melanoma proves to be true, not only will it be necessary to look at the mechanisms behind this increased risk, but it will also be necessary to develop appropriate screening and preventive measures to decrease the morbidity and mortality to the greatest degree possible from air travel associated melanoma.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Comments on this article

Version 1

Reader Comment 01 Sep 2015

Ian Burgess, Medical Entomology Centre, UK

With regard to aircrew engaged in commercial intercontinental services it should be noted that the general rostering arrangements are that they have a number of rest days between flights. In the case of crews travelling to warmer parts of the world it has certainly been the case historically that a good proportion of those "off" days were spent lounging around the hotel swimming pool, with associated exposure to higher levels of UV. Surely this must be a factor if there is detected higher incidence of radiation associated carcinomas in all grades of aircrew rather than just flight crew, who would be the only ones exposed directly to UV through cabin windows.

Competing Interests: No competing interests

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research