RESEARCH ARTICLE

REVISED Pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction: A short report [version 2; peer review: 1 approved, 1 approved with reservations, 1 not approved]

Hoa L. Nguyen^{1,2}, Dat T. Phan³, Duc A. Ha⁴, Quang N. Nguyen³, Robert J. Goldberg⁵

¹Department of Quantitative Sciences , Baylor Scott & White Health, Dallas, Texas, USA

²Institute of Population, Health and Development, Ha Noi, Vietnam

³Viet Nam National Heart Institute, Ha Noi, Vietnam

⁴Ministry of Health, Ha Noi, Vietnam

⁵Department of Quantitative Health Sciences, University of Massachusetts Medical School, Worcester, Massachusetts, USA

V2 First published: 27 Aug 2015, 4:633
<https://doi.org/10.12688/f1000research.6943.1>

Second version: 01 Aug 2017, 4:633
<https://doi.org/10.12688/f1000research.6943.2>

Latest published: 15 Jan 2018, 4:633
<https://doi.org/10.12688/f1000research.6943.3>

Abstract

Background: Administration of coronary reperfusion therapy to patients with an acute myocardial infarction (AMI) within the proper timeframe is essential in avoiding clinical complications and death. However, the extent of pre-hospital delay is unexplored in Vietnam. This report aims to describe the duration of pre-hospital delay of Hanoi residents hospitalized with a first AMI at the Vietnam National Heart Institute .

Methods: A total of 103 Hanoi residents hospitalized at the largest tertiary care medical center in the city for first AMI, who have information on prehospital delay was included in this report.

Results: One third of the study sample was women and mean age was 66 years. The mean and median pre-hospital delay duration were 14.9 hours and 4.8 hours, respectively. The proportion of patients who delayed <6 , 6-<12, and ≥ 12 hours were 45%, 13%, and 42%, respectively.

Conclusions: Our data shows that a prolonged pre-hospital delay is often observed in patients with a first AMI in Vietnam. In order to confirm these preliminary descriptive findings, a full-scale investigation of all Hanoi residents hospitalized with first AMI is needed. Increasing public awareness about AMI treatment is vital in encouraging patients to seek medical care timely after experiencing AMI symptoms such that received treatment is most effective.

Open Peer Review

Approval Status

	1	2	3
version 3 (revision) 15 Jan 2018			 view
version 2 (revision) 01 Aug 2017		 view	 view
version 1 27 Aug 2015	 view	 view	

1. **Felix Alvaro Medina Palomino**, Cayetano Heredia Peruvian University, Lima, Peru
2. **Dao Thi Mihn An**, Hanoi Medical University, Hano, Vietnam
3. **Polly W.C. Li**, The Chinese University of Hong Kong, Shatin, Hong Kong

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

Acute myocardial infarction, pre-hospital delay, outcomes, epidemiology, Vietnam

Corresponding author: Hoa L. Nguyen (hoanguyen@phad.org)

Author roles: **Nguyen HL:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Phan DT:** Data Curation, Investigation, Project Administration, Writing – Review & Editing; **Ha DA:** Data Curation, Investigation, Project Administration, Writing – Review & Editing; **Nguyen QN:** Data Curation, Investigation, Project Administration, Writing – Review & Editing; **Goldberg RJ:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: The authors have declared that no competing interests exist.

Grant information: This study was partially funded by the Global Health Office, University of Massachusetts Medical School, Worcester, MA, USA (Drs. Goldberg, Nguyen, and Ha). Additional support was provided by internal funding.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Nguyen HL *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Nguyen HL, Phan DT, Ha DA *et al.* **Pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction: A short report [version 2; peer review: 1 approved, 1 approved with reservations, 1 not approved]** F1000Research 2017, 4:633 <https://doi.org/10.12688/f1000research.6943.2>

First published: 27 Aug 2015, 4:633 <https://doi.org/10.12688/f1000research.6943.1>

REVISED Amendments from Version 1

In the version 2 of this paper, we have attempted to address each of the concerns raised about our submission as follow:

We have added information on our sample restriction in the method section. We restricted our patient population to those living in the city of Hanoi and with information on prehospital delay recorded in hospital medical records. We have also removed data on prehospital delay in the Methods sections and added them into the Results section.

We have modified the Results section to reflect our approaches to the analysis of these data. The present report is based on 103 patients with independently confirmed AMI. Data on duration of prehospital delay (continuous outcome) were available in only 65 patients, whereas the other 38 patients had data available only for a time interval (categorical outcome). Inasmuch, we have analyzed duration of prehospital delay as a continuous outcome in 65 patients and as a categorical outcome in 103 patients.

We have included one more limitation in the Discussion section. Due to the design of this retrospective study, we were unable to have systematically collected and/or recorded information about the time of onset of symptoms suggestive of AMI in hospital medical records. Data on duration of prehospital delay were missing in approximately two thirds of Hanoi residents hospitalized with AMI in 2010, which was mainly due to missing data on the time of onset of acute symptoms suggestive of AMI, and not ER arrival time. We have emphasized that since the sample size is small, we were unable to examine factors associated with prolonged delay, information which is important for designing and implementing various educational intervention approaches to improve patient's care seeking behavior.

See referee reports

Introduction

Vietnam is a low-middle income country, and has been in the midst of an important epidemiological transition. Over the past two decades, the overall morbidity and mortality rates resulting from non-communicable diseases (NCDs) has been rising rapidly and presents itself as a major problem in Vietnam. One quarter of all deaths annually in Vietnam are now the result of cardiovascular diseases, making it the leading cause of death.

Timely administration of reperfusion therapy to patients with an evolving acute myocardial infarction (AMI) is of utmost importance in reducing clinical complications and death. Previous research studies have convincingly proved that reperfusion treatment is most effective if patients with ST-segment elevation myocardial infarction (STEMI) are treated in a timely fashion, particularly within one hour of acute symptom onset^{1,2}; the relation between extent of pre-hospital delay and outcomes after non-STEMI has not been firmly established.

Despite the importance of prolonged pre-hospital delay on the timely receipt of effective treatments and short-term outcomes, so far we have found no evidence of studies conducted to examine the extent of pre-hospital delay among adult patients hospitalized with AMI in Vietnam. This short report hence aims to describe the extent of pre-hospital delay in Hanoi residents who were non-transferred, hospitalized with a first AMI at the Vietnam National Heart Institute in 2010.

Methods

Ethical statement

This study was approved by the Institutional Review Board at the Institute of Population, Health and Development. A waiver of patient consent was approved by the Institutional Review Board. All patients' personal information was de-identified before analysis.

Study setting

This study was conducted at the Vietnam National Heart Institute (VNHI)^{3,4}. This hospital is a 250 bed tertiary care medical center in Hanoi (2009 census = 6.5 million) which manages the majority of Hanoi residents hospitalized with acute coronary disease and other NCDs.

Case ascertainment

Computerized printouts of discharged patients from the VNHI in 2010 with possible AMI were collected and International Classification of Disease codes for possible AMI (I20–I25) were reviewed. Two cardiologists validated these cases utilizing predefined criteria developed by the World Health Organization. They include having a suggestive medical history, serum enzyme increasing above the hospital's reference range, and serial electrocardiographic (ECG) findings during hospitalization consistent with the presence of AMI; at least two of these three criteria needed to be present for an AMI to have occurred⁵. There were 315 Hanoi residents, who satisfied the criteria for AMI were identified. However, 13 (4%) patients with history of AMI and 199 (66%) patients transferred from other facilities were excluded from the present analysis. Patients were then further divided into groups of STEMI and non-STEMI, utilizing standard classification techniques⁶. All ECGs of potential AMI patients were reviewed by a physician under the supervision of a senior cardiologist. Due to reviewed ECGs' inadequate quality, nine patients with AMI could not be classified as STEMI or non-STEMI.

Data collection

For all AMI-validated cases through the aforementioned process of independent adjudication utilizing eligibility criteria, information on social demographics, clinical, medical management and hospital discharge status were collected from medical record reviews and recorded onto a standardized case report form by trained study physicians.

Pre-hospital delay is the primary outcome investigated by this study. The delay time was defined as the time interval between the onset of symptoms suggestive of AMI and the patient's arrival at medical centers^{7,8}. Our trained physicians reviewed any information they could seek in medical records, which described the duration of pre-hospital delay from emergency personnel, nurses, and physicians notes. Information on pre-hospital delay was collected in minutes (as a continuous variable) then was further categorized according to cut-points of 6 and 12 hours, which were commonly applied in prior studies, and based on the data distribution. Patients, in whom no exact time of symptom onset was reported, and only had a delay time interval recorded in their medical record (e.g. <6, 6–<12, and ≥12 hours), were incorporated into our analysis of delay time when

it was constructed as a categorical outcome. We restricted our patient population to those with information available on pre-hospital delay (either exact time noted or time interval) in their hospital medical records.

Data analysis

The overall mean (standard deviation-SD) and median (inter quartile range-IQR) duration of pre-hospital delay was calculated according to standard methods. Data were shown as percentages for categorical variables and compared between patients who delayed ≥ 6 hours, and patients who sought medical care earlier, using chi-square or Fisher exact tests; medians (inter quartile range-IQR) for continuous variables were calculated and compared using the Wilcoxon sum rank tests. Since the sample size in this study was small, no regression analysis was performed. All analyses were performed using STATA 11.0 (StataCorp. TX).

Results

A total of 103 non-transferred Hanoi residents hospitalized with a first AMI at VNHI in 2010, and had information available on pre-hospital delay recorded in their hospital charts were included in the report. The study sample had an average age of 66 years with a standard deviation of 13 years. One third of the study sample was women, and 69% were classified as STEMI.

Extent of pre-hospital delay

Information on pre-hospital delay was collected in minutes (as a continuous variable) for 63% (n=65) of patients. In 37% of patients (n=38), an exact time of symptom onset was not reported, and only a delay time interval was recorded in patient's medical record (e.g. <6, 6-<12, and ≥ 12 hours); these patients were included in the analysis of delay time when it was constructed as a categorical outcome.

Analysis of prehospital delay as a continuous outcome. Within the group of 65 patients reporting an exact time of symptom onset, the overall mean and median durations of pre-hospital delay were 14.9 hours (SD: 23 hours) and 4.8 hours (IQR: 3–10 hours), respectively (Figure 1). Among patients with STEMI (n=42), mean and median delay times were 16.4 hours (SD: 26 hours) and 4.8 hours (IQR: 3.0–6.2 hours), respectively. Among patients with non-STEMI (n=19), mean and median delay times were 14.0 hours (SD: 19.2 hours) and 8.3 hours (IQR: 2.5–12.3 hours), respectively.

Analysis of prehospital delay as a categorical outcome. When combining data from these patients (n=65), with more non-specific data from patients who reported duration of delay in a time interval (n=38), the proportion of patients who delayed <6 hours,

6-<12 hours, and ≥ 12 hours were 45%, 13%, and 42%, respectively (Figure 2). Among patients with STEMI (n=65), these proportions were 51%, 12% and 37%, respectively. Among patients with non-STEMI (n=29), these proportions were 31%, 17% and 52%, respectively.

Patient characteristics associated with prolonged pre-Hospital delay

Patients who had a delay time over 6 hours were more likely to be female compared to their counterparts, who had a delay time of less than 6 hours (Table 1).

Figure 1. Mean and Median of Pre-hospital Delay in Patients Hospitalized with an initial Acute Myocardial Infarction.

Figure 2. Distribution of Duration of Pre-hospital Delay in Patients Hospitalized with an initial Acute Myocardial Infarction.

Table 1. Characteristics of Patients with an Initial Acute Myocardial Infarction (AMI) according to Extent of Delay in Seeking Medical Care.

Characteristic	< 6 hours (n=46)	≥ 6 hours (n=57)	P-value
Age (mean, SD), years	65 (13)	67 (12)	0.51
Age (n,%), years			
<60	16(34.8)	17(29.8)	0.86
60-69	11(23.9)	14(24.6)	
≥70	19(41.3)	26(45.6)	
Female (%)	11(23.9)	25(43.9)	0.04
Ethnicity (n,%)			
Kinh*	44(95.7)	56(98.2)	0.44
Minority	2(4.3)	1(1.8)	
Have medical insurance (n,%)	22(47.8)	26(45.6)	0.82
Medical history (n,%)			
Atrial fibrillation	1(2.2)	0(0)	0.45
Coronary heart disease	5(10.9)	0(0)	NA
Diabetes mellitus	8(17.4)	11(19.3)	0.80
Heart failure	0(0)	0(0)	NA
Hypertension	30(65.2)	34(59.6)	0.56
Hyperlipidemia	0(0)	3(5.3)	NA
Stroke	5(10.9)	5(8.8)	0.72
AMI characteristics (n,%)†			
STEMI	33(78.6)	32(61.5)	0.08
Non-STEMI	9(21.4)	20(38.5)	
Acute symptoms (n,%)			
Chest pain	43(93.5)	49(86.0)	0.22
Shortness of breath	21(45.7)	21(36.8)	0.37
Nausea	5(10.9)	1(1.8)	0.06
Diaphoresis	9(19.6)	12(21.1)	0.85
Fatigue	5(10.9)	7(12.3)	0.82
Clinical presentation (median, IQR)			
Heart rate, beats/min	83(67-96)	82(67-100)	0.84
Systolic blood pressure, mmHg	120(110-140)	120(110-140)	0.46
Diastolic blood pressure, mmHg	80(70-80)	80(70-90)	0.64
Laboratory findings (median, IQR)			
Total cholesterol, mmol/L	4.5(3.9-5.0)	4.6(4.0-5.2)	0.39
LDL cholesterol, mmol/L	2.5(2.3-3.1)	2.7(2.3-3.0)	0.49
eGFR (ml/min/1.73m ²)	71(57-79)	68(56-89)	0.56

Pre-hospital delay was defined as the duration from onset of acute symptoms suggestive of AMI to hospital arrival.

P- values from chi square or Fisher exact tests for categorical variables, and t-tests or Wilcoxon-sum rank tests for continuous variables.

SD: Standard deviation; IQR: Inter quartile range; STEMI: ST-segment elevation MI; LDL: Low-density lipoprotein; eGFR: estimated glomerular filtration rate

*The Kinh people are the majority ethnic group in Vietnam, comprising 87% of the population (census 2009).

†Missing data in 9 patients

Dataset 1. Raw dataset for Nguyen *et al.*, 2015 'Pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction: A short report'

<http://dx.doi.org/10.5256/f1000research.6943.d100892>

For all AMI-validated cases, information on social demographics, clinical, medical management and hospital discharge status were collected from medical record reviews and recorded onto a standardized case report form by trained study physicians. GFR: Glomerular filtration rate; STEMI: ST elevated myocardial infarction; LDL: Low density lipoprotein.

Discussion

The results of this study indicated that Vietnamese patients experienced a considerable pre-hospital delay after the onset of AMI symptoms (median: 4.8 hours), with more than half experiencing a delay time of 6 hours or longer. Women delayed seeking care to a greater extent than men.

This report is, to our knowledge, the first in Vietnam to describe the duration of pre-hospital delays in patients hospitalized with a first AMI. In this study, patients experienced considerably longer pre-hospital delay compared with patients from both low-middle income countries and high-income countries^{7–16}. For example, a survey of 250 patients hospitalized with AMI in Shanghai, China in 2010 reported that the median time for patients to seek treatment was around 2 hours alongside another one half hour (median time) for transportation⁹. Another recent survey from 270 patients hospitalized with AMI in Bandar Abbas, Iran found that approximately 20% of patients experienced a delay time of 6 hours or more¹³. A study of more than 200 patients hospitalized with acute coronary syndrome in London, England between 2001 and 2004 reported that the median pre-hospital delay time was 2 hours¹⁵. Prior studies in the U.S. have shown that patients hospitalized with AMI delayed, on average, approximately 2 hours following the onset of symptoms suggestive of AMI^{11,16}. However, the duration of pre-hospital delay in our study was comparable to that reported from more than 60 African American patients hospitalized with AMI in the US in 2003–2004 (median: 4.3 hours)¹⁷ and from more than 400 patients hospitalized with AMI in Croatia in 2005 (median: 4.3 hours)¹⁸. We found that comparatively, female patients tended to seek treatment much later than male patients after the onset of suggestive AMI symptoms. This result is consistent with studies conducted previously^{9,10,19–22}, but contradictory to others, which reported that there was no association between sex and pre-hospital delay^{12–14}.

Our findings emphasize the need to implement intervention programs to increase awareness of the general population with regards to the importance of seeking treatment timely after experiencing AMI suggestive symptoms. Prolonged delay may be associated with an individual's risk of cardiac death, serious clinical complications, and delays in the receipt of effective treatments, primarily coronary reperfusion therapy. Moreover, to understand why these patients fail

to react promptly to their AMI symptoms, focus group discussions and/or in-depth interviews should be carried out in patients hospitalized with AMI focusing on their levels of cognition, knowledge, and attitudes toward hospitals and health care. Additional studies are warranted in exploring the effects of educational attainment, insurance coverage, neighborhood characteristics, psychosocial aspects, and other factors that may serve as either facilitators or obstacles for patients to the timelier seeking of medical care, particularly of factors that may be amenable to modify.

Study limitations

There are several limitations in this study. First, our data was collected from only one hospital, the VNHI in Hanoi, which limits the generalizability of our results to patients hospitalized in other medical centers across Hanoi. Second, due to our small sample size, we failed to identify potential factors that might relate to the excessive delay time, information which is important for designing and implementing various educational intervention approaches to improve patient's care seeking behavior. Third, we were only able to calculate total duration of pre-hospital delay, were not able to calculate patient delay time and transportation delay separately since the time when patients made a treatment-seeking decision was not recorded in hospital medical records. In Vietnam, patients with signs and symptoms suggestive of AMI normally call for an ambulance or reach the hospital by themselves, and extremely few patients have a primary care doctor to consult. In terms of transportation time, this study included only patients who reside in Hanoi. Forth, due to the retrospective design of the study, we were unable to have systematically collected and/or recorded information about the time of onset of symptoms suggestive of AMI in hospital medical records. Data on duration of prehospital delay were missing in approximately two thirds of Hanoi residents hospitalized with independently validated AMI, which was mainly due to missing data on the time of onset of acute symptoms suggestive of AMI, and not emergency department arrival time. Finally, we also were unable to obtain information on additional patient-associated characteristics (e.g., socioeconomic status [e.g., Education, occupation] and psychological factors [e.g., reluctance to call family members for help or having no one around when experiencing symptoms to ask for help]) which may have impacted on patient's medical care seeking behavior.

Our data show that patients admitted for initial AMI in Hanoi, Vietnam experienced a significant delay time in seeking medical attention after the onset of AMI suggestive symptoms. Additional full-scale data of hospitalized patients with AMI across Hanoi will be needed in order to confirm these preliminary descriptive findings. Public educational programs for the general population, specifically targeting towards women, is greatly needed to incentivize patients to seek medical assistance immediately to ensure maximum effectiveness of treatments. Further research is required to understand potential factors associated with the prolonged delay in Vietnamese patients hospitalized with AMI and the most effective ways to encourage patients to seek medical care in a timely manner.

Data availability

F1000Research: Dataset 1. Raw dataset for Nguyen *et al.*, 2015 ‘Pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction: A short report’, [10.5256/f1000research.6943.d100892](https://doi.org/10.5256/f1000research.6943.d100892)²³

Author contributions

HLN, RJG conceived and designed the study. HLN, DTP, DAH, QNN carried out the study. HLN, RJG analyzed the data and prepared the first draft of the manuscript. All authors were involved in the revision of the draft manuscript and have agreed to the final content.

Competing interests

The authors have declared that no competing interests exist.

Grant information

This study was partially funded by the Global Health Office, University of Massachusetts Medical School, Worcester, MA, USA (Drs. Goldberg, Nguyen, and Ha). Additional support was provided by internal funding.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank the doctors and nurses at the Vietnam National Heart Institute for their support and help.

References

1. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. *N Engl J Med*. 1993; 329(10): 673–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Newby LK, Rutsch WR, Califf RM, *et al.*: Time from symptom onset to treatment and outcomes after thrombolytic therapy. GUSTO-1 Investigators. *J Am Coll Cardiol*. 1996; 27(7): 1646–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Nguyen HL, Ha DA, Phan DT, *et al.*: Sex differences in clinical characteristics, hospital management practices, and in-hospital outcomes in patients hospitalized in a Vietnamese hospital with a first acute myocardial infarction. *PLoS One*. 2014; 9(4): e95631.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Nguyen HL, Nguyen QN, Ha DA, *et al.*: Prevalence of comorbidities and their impact on hospital management and short-term outcomes in Vietnamese patients hospitalized with a first acute myocardial infarction. *PLoS One*. 2014; 9(10): e108998.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Tunstall-Pedoe H, Kuulasmaa K, Mähönen M, *et al.*: Contribution of trends in survival and coronary-event rates to changes in coronary heart disease mortality: 10-year results from 37 WHO MONICA project populations. Monitoring trends and determinants in cardiovascular disease. *Lancet*. 1999; 353(9164): 1547–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. McManus DD, Gore J, Yarzebski J, *et al.*: Recent trends in the incidence, treatment, and outcomes of patients with STEMI and NSTEMI. *Am J Med*. 2011; 124(1): 40–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Goldberg RJ, Steg PG, Sadiq I, *et al.*: Extent of, and factors associated with, delay to hospital presentation in patients with acute coronary disease (the GRACE registry). *Am J Cardiol*. 2002; 89(7): 791–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Goldberg RJ, Yarzebski J, Lessard D, *et al.*: Decade-long trends and factors associated with time to hospital presentation in patients with acute myocardial infarction: the Worcester Heart Attack study. *Arch Intern Med*. 2000; 160(21): 3217–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Wang X, Hsu LL: Treatment-seeking delays in patients with acute myocardial infarction and use of the emergency medical service. *J Int Med Res*. 2013; 41(1): 231–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Sheifer SE, Rathore SS, Gersh BJ, *et al.*: Time to presentation with acute myocardial infarction in the elderly: associations with race, sex, and socioeconomic characteristics. *Circulation*. 2000; 102(14): 1651–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Nguyen HL, Gore JM, Saczynski JS, *et al.*: Age and sex differences and 20-year trends (1986 to 2005) in prehospital delay in patients hospitalized with acute myocardial infarction. *Circ Cardiovasc Qual Outcomes*. 2010; 3(6): 590–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Qian L, Ji KT, Nan JL, *et al.*: Factors associated with decision time for patients with ST-segment elevation acute myocardial infarction. *J Zhejiang Univ Sci B*. 2013; 14(8): 754–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Farshidi H, Rahimi S, Abdi A, *et al.*: Factors Associated With Pre-hospital Delay in Patients With Acute Myocardial Infarction. *Iran Red Crescent Med J*. 2013; 15(4): 312–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Khan MS, Jafari FH, Faruqi AM, *et al.*: High prevalence of lack of knowledge of symptoms of acute myocardial infarction in Pakistan and its contribution to delayed presentation to the hospital. *BMC Public Health*. 2007; 7: 284.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Perkins-Porras L, Whitehead DL, Strike PC, *et al.*: Pre-hospital delay in patients with acute coronary syndrome: factors associated with patient decision time and home-to-hospital delay. *Eur J Cardiovasc Nurs*. 2009; 8(1): 26–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Saczynski JS, Yarzebski J, Lessard D, *et al.*: Trends in prehospital delay in patients with acute myocardial infarction (from the Worcester Heart Attack Study). *Am J Cardiol*. 2008; 102(12): 1589–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Banks AD, Dracup K: Factors associated with prolonged prehospital delay of African Americans with acute myocardial infarction. *Am J Crit Care*. 2006; 15(2): 149–57.
[PubMed Abstract](#)
18. Novak K, Aljinovic J, Kostic S, *et al.*: Pain to hospital times after myocardial infarction in patients from Dalmatian mainland and islands, southern Croatia. *Croat Med J*. 2010; 51(5): 423–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. El-Menyar A, Zubaid M, Rashed W, *et al.*: Comparison of men and women with acute coronary syndrome in six Middle Eastern countries. *Am J Cardiol*. 2009; 104(8): 1018–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Goldberg RJ, Spencer FA, Fox KA, *et al.*: Prehospital Delay in Patients With Acute Coronary Syndromes (from the Global Registry of Acute Coronary Events [GRACE]). *Am J Cardiol*. 2009; 103(5): 598–603.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Nguyen HL, Saczynski JS, Gore JM, *et al.*: Age and sex differences in duration of prehospital delay in patients with acute myocardial infarction: a systematic review. *Circ Cardiovasc Qual Outcomes*. 2010; 3(1): 82–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. McGinn AP, Rosamond WD, Goff DC Jr, *et al.*: Trends in prehospital delay time and use of emergency medical services for acute myocardial infarction: experience in 4 US communities from 1987–2000. *Am Heart J*. 2005; 150(3): 392–400.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Nguyen HL, Phan D, Ha D, *et al.*: Dataset 1 in: Pre-hospital Delay in Vietnamese Patients Hospitalized with a First Acute Myocardial Infarction: A Short Report. *F1000Research*. 2015.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 11 December 2017

<https://doi.org/10.5256/f1000research.13251.r28540>

© 2017 Li P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Polly W.C. Li

The Nethersole School of Nursing, The Chinese University of Hong Kong, Shatin, Hong Kong

This report addressed an important gap in the literature about the care-seeking delay of Vietnamese in the context of acute myocardial infarction. The major limitations of the study were small sample size and retrospective nature of the study. Nevertheless, this is a good start to provide quick preliminary findings to support a more formal and large scale study to examine the phenomenon. I have several additional comments:

1. 69% of the sample suffered from STEMI, which is not consistent with the proportion of STEMI/NSTEMI in AMI population. This reflects that the sample is not representative and need to address in the discussion and limitation sections.
2. Retrospective medical record review is not a reliable data collection method for collecting data on pre-hospital delay. The inconsistency for healthcare professionals to determine the symptom onset time creates a significant bias to the results.
3. The very small number of NSTEMI patients made the subgroup analysis to be less appropriate.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 27 Dec 2017

Hoa L. Nguyen, Baylor Scott & White Health, Dallas, USA

We would like to thank the reviewer of our article for the helpful comments. We have attempted to address each of the concerns raised about our submission and have indicated in the revised text where changes have been made. We believe that the manuscript is considerably stronger than before.

1. 69% of the sample suffered from STEMI, which is not consistent with the proportion of STEMI/NSTEMI in AMI population. This reflects that the sample is not representative and need to address in the discussion and limitation sections.

Response: We have acknowledged this limitation in the revised discussion (Study limitations section), though we are unsure what the distribution of type of AMI may be in the Vietnamese population. In our conclusions section, we have emphasized the need for full-scale surveillance of Hanoi residents hospitalized with AMI at all Hanoi hospitals to confirm these preliminary descriptive findings.

2. Retrospective medical record review is not a reliable data collection method for collecting data on pre-hospital delay. The inconsistency for healthcare professionals to determine the symptom onset time creates a significant bias to the results.

Response: Due to the retrospective nature of this observational study, there was no systematic method established to record the time of onset of symptoms suggestive of AMI (e.g., chest pain) in a standardized manner by trained health professionals. While we would have preferred to have directly interviewed hospitalized patients with AMI in a standardized manner, as noted in response to a similar query by the first and second reviewers, this information was abstracted from notes written by different physicians and nurses and emergency personnel which may have been collected and recorded in a non-standardized manner. We have noted this limitation in the discussion section of the revised manuscript (Study limitations section).

3. The very small number of NSTEMI patients made the subgroup analysis to be less appropriate.

Response: We agree with the reviewer and have acknowledged this limitation in the revised discussion (Study limitations section). However, in this brief report, we would prefer to describe the extent of prehospital delay for patients with a STEMI and NSTEMI separately given the extremely limited data presently available on this topic in Vietnam and due to differences in the natural history of these conditions. In our conclusions, we have emphasized the need for full-scale surveillance of Hanoi residents hospitalized with AMI at all Hanoi hospitals to confirm these preliminary descriptive findings.

Competing Interests: None

Reviewer Report 13 September 2017

<https://doi.org/10.5256/f1000research.13251.r24711>

© 2017 Mihn An D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dao Thi Mihn An

Department of Epidemiology, Institute for Preventive Medicine and Public Health, Hanoi Medical University, Hanoi, Vietnam

I have read the Summary of revisions presented by the author of the paper "Pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction: A short report". Generally, the author has improved the weak points indicated by reviewers by adding information, discussing the limitation issues, or re-structuring within each section of the paper.

The main finding of this paper supplies evidence of pre-hospital delay among AMI patients in Vietnam that would probably result in further comprehensive study of this issue in the near future in Vietnam.

I agree with the revisions of the author, and approve this paper.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 03 August 2016

<https://doi.org/10.5256/f1000research.7478.r15408>

© 2016 Mihn An D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dao Thi Mihn An

Department of Epidemiology, Institute for Preventive Medicine and Public Health, Hanoi Medical University, Hanoi, Vietnam

There were not any official publications in Vietnam regarding issue of pre-hospital delay in Vietnamese patients hospitalized with a first acute myocardial infarction until now. Therefore, the short report on this issue written by Hoa *et al* is highly appreciated.

This study was conducted with a small and convenient sample of 103 AMI patients with their first AMI and the study approach was to review AMI patients' medical records after their discharge that must be embedded biases as it was a retrospective study. The inconsistency of asking and identifying time of delay by different health workers at the time of hospitalization and the different subjects who supplied information for health workers should be considered the main problems of the validity of data in this study beside 3 other limitations that the author has discussed on this paper. Moreover, it was likely that there were missing value regarding delay time occurred during process of extracting data from AMI patients' medical record.

However, this study initially indicated 2 very important findings that pre-hospital delay time for hospitalization among AMI patients was much longer than that of AMI patients in the local and worldwide and patients who had a delay time over 6 hours were more likely to be female compared to their counterparts. These findings make help in giving suggestion for further full-scale data of hospitalized patients with AMI across Hanoi in order to confirm these initial findings.

I would like very much to approve for this short report as it is the starting point for further large-

scale study in this issue in Vietnam where such information are still rare.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 24 Jul 2017

Hoa L. Nguyen, Institute of Population, Health and Development, Ha Noi, Vietnam

Response: Due to the retrospective nature of this observational study, there was no systematic method established to record the time of onset of symptoms suggestive of AMI (e.g., chest pain) in a standardized manner by trained health professionals. As noted in response to a similar query by the other reviewer, this information was abstracted from notes written by physicians and nurses and emergency personnel which may have been collected and recorded in a non-standardized manner. Duration of prehospital delay was missing in approximately two thirds of Hanoi residents hospitalized with AMI in 2010, which was mainly due to missing data on the time of onset of symptoms suggestive of AMI. We have noted this limitation in the discussion section of the revised manuscript.

Competing Interests: none

Reviewer Report 13 October 2015

<https://doi.org/10.5256/f1000research.7478.r10208>

© 2015 Palomino F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Felix Alvaro Medina Palomino

Alberto Hurtado School of Medicine, Cayetano Heredia Peruvian University, Lima, Peru

1. It is not clear how the author obtained their samples.
2. There is a very small population to study and to obtain valid conclusions from.
3. It is uncertain how the time of chest pain was recorded. It appears there is no system that allows for a correct determination of the precise moment the patient arrives at the ER, due to the fact that in many patients this particular data is missing.
4. The report is based on more than a hundred patients but in fact the author only presents data from 65 patients.
5. There is no explanation about the findings nor factors.
6. Because only a few patients were enrolled, it is not possible to determine factors related to early or delayed arrival at the hospital.
7. Finally some data, which are in fact results, are written in the methods.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 24 Jul 2017

Hoa L. Nguyen, Institute of Population, Health and Development, Ha Noi, Vietnam

1. It is not clear how the author obtained their samples.

Response: We have described the study population in greater detail in the Case Ascertainment section of the Methods (page 3, para 2). In brief, lists of patients discharged from the Vietnam National Heart Institute in 2010 with possible AMI were obtained and International Classification of Disease-10 codes for possible AMI (I20-I25) were reviewed. The diagnosis of AMI was made on the basis of criteria developed by the World Health Organization which includes a suggestive clinical history, serum enzyme elevations above the hospital's normal range, and serial electrocardiographic (ECG) findings during hospitalization consistent with the presence of AMI; at least 2 of these 3 criteria needed to be present for an AMI to have occurred.

We restricted our patient population to those living in the city of Hanoi and with information on prehospital delay recorded in hospital medical records. We have added this information in the revised manuscript.

2. There is a very small population to study and to obtain valid conclusions from.

Response: We have acknowledged this limitation in the revised discussion. In our conclusions, we have emphasized the need for full-scale surveillance of Hanoi residents hospitalized with AMI at all Hanoi hospitals to confirm these preliminary descriptive findings.

3. It is uncertain how the time of chest pain was recorded. It appears there is no system that allows for a correct determination of the precise moment the patient arrives at the ER, due to the fact that in many patients this particular data is missing.

Response: Due to the design of this retrospective study, whose information was based exclusively on information contained in hospital medical records, there was no systematic method to collect and record the time of onset of symptoms suggestive of AMI (e.g., chest pain) since these data were collected and recorded in hospital records by nurses and physicians with varying degrees of expertise and interest in collecting this information. Data on duration of prehospital delay were missing in approximately two thirds of Hanoi residents hospitalized with AMI in 2010, which was mainly due to missing data on the time of onset of acute symptoms suggestive of AMI, and not ER arrival time. We have included this limitation in the discussion section of the revised manuscript.

4. The report is based on more than a hundred patients but in fact the author only presents data from 65 patients.

Response: The present report is based on 103 patients with independently confirmed AMI. Data on duration of prehospital delay (continuous outcome) were available in only 65 patients, whereas the other 38 patients had data available only for a time interval (categorical outcome). Inasmuch, we have analyzed duration of prehospital delay as a continuous outcome in 65

patients and as a categorical outcome in 103 patients. We have modified the results section to reflect our approaches to the analysis of these data in the revised manuscript.

5. There is no explanation about the findings nor factors.

Response: In this brief report, our primary goal was to present descriptive data about the extent of prehospital delay in patients hospitalized with AMI at the major teaching hospital in Hanoi, information which is unknown and sorely lacking in Vietnam. Since the sample size is small, we were unable to examine factors associated with prolonged delay, information which is important for designing and implementing various educational intervention approaches to improve patient's care seeking behavior. We have emphasized this limitation and the need for future studies in the revised manuscript.

6. Because only a few patients were enrolled, it is not possible to determine factors related to early or delayed arrival at the hospital.

Response: Please see the above response (#5).

7. Finally some data, which are in fact results, are written in the methods.

Response: We have removed data on prehospital delay in the method sections and added them into the results section of the revised manuscript.

Competing Interests: none

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research