Check for updates

OPINION ARTICLE

Circulating nucleic acids: a new class of physiological mobile genetic elements [version 1; peer review: 2 approved]

Indraneel Mittra

Translational Research Laboratory, Advanced Centre for Treatment, Research and Education in Cancer, Tata Memorial Centre, Navi-Mumbai, India

V1 First published: 30 Sep 2015, 4:924
<https://doi.org/10.12688/f1000research.7095.1>
 Latest published: 30 Sep 2015, 4:924
<https://doi.org/10.12688/f1000research.7095.1>

Abstract

Mobile genetic elements play a major role in shaping biotic genomes and bringing about evolutionary transformations. Herein, a new class of mobile genetic elements is proposed in the form of circulating nucleic acids (CNAs) derived from the billions of cells that die in the body every day due to normal physiology and that act intra-corporeally. A recent study shows that CNAs can freely enter into healthy cells, integrate into their genomes by a unique mechanism and cause damage to their DNA. Being ubiquitous and continuously arising, CNA-induced DNA damage may be the underlying cause of ageing, ageing-related disabilities and the ultimate demise of the organism. Thus, DNA seems to act in the paradoxical roles of both preserver and destroyer of life. This new class of mobile genetic element may be relevant not only to multi-cellular organisms with established circulatory systems, but also to other multi-cellular organisms in which intra-corporeal mobility of nucleic acids may be mediated via the medium of extra-cellular fluid.

Keywords

Mobile genetic elements , horizontal gene transfer , circulating nucleic acids , circulating DNA , circulating chromatin , DNA damage , ageing

Open Peer Review

Approval Status

	1	2
version 1
30 Sep 2015	view	view

1. **Silvia Gravina**, Albert Einstein College of Medicine, Bronx, USA

2. **Manchanahalli Rangaswamy Satyanarayana Rao**, Jawaharlal Nehru Centre For Advanced Scientific Research, Bangalore, India

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Indraneel Mittra (indraneel.mittra@gmail.com)

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Department of Atomic Energy, Government of India, through its grant CTCTMC to Tata Memorial Centre awarded to IM.

Copyright: © 2015 Mittra I. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Mittra I. **Circulating nucleic acids: a new class of physiological mobile genetic elements [version 1; peer review: 2 approved]** F1000Research 2015, 4:924 <https://doi.org/10.12688/f1000research.7095.1>

First published: 30 Sep 2015, 4:924 <https://doi.org/10.12688/f1000research.7095.1>

Background

Barbara McClintock published her classic paper on mobile genetic elements (MGEs) in 1950¹, but it took the scientific community several decades to appreciate the enormity of her discovery. Today, it is recognized that MGEs occur widely in nature in prokaryotes, archaea and eukaryotes and play a major role in shaping their genomes and bringing about evolutionary transformations and adaptation^{2–5}. Their bizarre behavior of moving from one part of the genome to another distinguishes them from the functioning of conventional genetic elements.

MGEs belong to two classes *viz.*, intra-genomic and inter-genomic. Intra-genomic MGEs are transposable elements (TEs) or transposons which constitute nearly 50% of the human genome but are variable among species and comprise 1%–5% of prokaryotic genomes⁶. Inter-genomic transposable elements, on the other hand, underlie horizontal or lateral gene transfer (HGT) whereby segments of DNA are transferred from one organism to another^{7–9}. Although HGT is known to occur extensively in bacteria and are responsible for development of antibiotic resistance¹⁰, increasing evidence of HGT between other organisms is coming to light. For example, HGT between prokaryote and eukaryote, eukaryote and eukaryote, eukaryote and prokaryote has been reported^{7,11}. Although the initial claims of presence of bacterial DNA in human genome were dismissed as erroneous^{12,13}, recent evidence has confirmed the presence of bacteria DNA sequences in about one-third of healthy humans and in greater numbers in cancer cells¹⁴. A recent analysis of public databases of transcriptome sequences of multiple organisms discovered that human beings have picked up at least 145 genes from other species during the course of evolution¹¹. Thus, HGT results in what is called a ‘web of life’ rather than a steadily bifurcating evolutionary tree⁸.

Circulating nucleic acids as a new class of mobile genetic elements

Based on a recent finding¹⁵, a new class of mobile genetic elements is proposed *viz.*, circulating nucleic acids, which are produced as a result of normal physiology and operate intra-corporeally or within the body of an organism. Circulating nucleic acids (CNAs) in the form of fragmented DNA and chromatin (DNAfs and Cfs) are known to circulate in blood and are derived from the hundreds of billions of cells that die through apoptosis in the adult human body on a daily basis^{16,17}. These fragments have a size range of between 100bp–1000bp, have a half-life of 10–15 minutes and are ultimately removed by the liver^{18,19}. The presence of Cfs (nucleosomes) in blood can be detected by a sandwich ELISA assay¹⁵, but whether naked DNA circulates as such remains an open question since the possibility cannot be excluded that DNAfs isolated from plasma/serum are in fact products of the DNA purification process.

Results of a recent study summarized below have revealed that CNAs can act as mobile genetic elements¹⁵. DNAfs and Cfs isolated from blood of healthy volunteers and cancer patients are actively taken up by cells in culture whereupon they rapidly accumulate in their nuclei and associate themselves with their chromosomes. The intracellular DNAfs and Cfs trigger a DNA-damage-repair-response (DDR) with up-regulation of multiple pathways of DNA damage and repair that facilitate their integration into host cell genome.

Presence of human DNA in recipient mouse cell chromosomes could be detected by FISH while whole-genome sequencing uncovered tens of thousands of human reads in mouse cells. The integration of DNAfs and Cfs is stable and presence of extraneous DNA was demonstrable in single-cell clones developed from treated cells which had undergone numerous cell divisions. Genomic integration of DNAfs and Cfs results in phosphorylation of H2AX indicative of dsDNA breaks and up-regulation of apoptotic pathways in a proportion of cells. When injected intravenously into mice, DNAfs and Cfs integrate into cells of a variety of organs in the body, activate H2AX and the apoptotic marker active Caspase-3.

Whether genomic integration of CNAs occurs preferentially in a site-specific manner or is random is not known; but in either case, integration of CNAs would give rise to somatic mutations in the host genome. Since integration of CNAs occurred in all organs of the body examined¹⁵, it may not be far-fetched to imagine that CNA-integration also occurs in germ cells. Genomic integration of CNAs would lead to DNA rearrangements, translocations and deletions²⁰ – changes that are hallmarks of ageing, and large DNA rearrangements and cell to cell variations in gene expression are typical of ageing cells^{21,22}.

CNAs integrate into host-cell genomes by a unique mechanism

According to the model depicted in the [Figure](#), CNAs integrate into the genome by a unique mechanism in which activation of DDR plays a central role¹⁵. When DNAfs and Cfs enter into a cell, the latter mistakenly perceives the intracellular DNAfs and Cfs with dsDNA breaks in their two ends as damaged “self” DNA and activates DDR even before DNA damage has actually occurred. The activated DDR joins up multiple disparate DNAfs and Cfs into long concatemers by non-homologous-end-joining as a part of the repair process. It is the integration into the host cell genomes of the concatemers by homologous or non-homologous recombination that brings about damage to DNA. Thus, paradoxically, the activation of DDR brings about damage to DNA rather than preserving DNA integrity. This model of DNA damage and repair in which DDR precedes DNA damage is the reverse of the classical model based on damage induced by ionizing and UV-radiations and chemicals wherein DDR is activated after DNA damage. It is possible that this model of DNA damage and repair that facilitates CNAs integration may apply to horizontally transferred DNA in other organisms in nature.

Implications

Although xenobiotics and DNA damaging agents constantly damage human DNA, these are usually transient and do not inflict permanent damage. CNAs, on the other hand, are ubiquitous, physiological and continuously arising, inflicting repeated damage and mutations to the somatic DNA. This naturally suggests that the somatic genome is not stable but remains in a state of turbulence characterized by DNA damage, mutations and rearrangements leading to DNA mosaicism and cell-to-cell variation in genomic structure and function. Indeed, cell-to-cell variations are being increasingly uncovered in the human body and are related to ageing^{21,23–25}. The above events and the accompanying genomic instability may give rise to cancerous transformations which is compatible with the steep rise in the incidence of cancer with increasing age²⁶. CNAs may also play an etiological

Figure. A new mechanistic model of genomic integration of CNAs in which DDR precedes DNA damage. NHEJ = non-homologous end-joining; HR = homologous recombination; NHR = non-homologous recombination. Reproduced with permission from Mittra *et al.*, J Biosci. 2015. **40**: 91–111.

role in several other disease conditions which are characterized by elevated levels of DNAfs and Cfs. These include auto-immune disorders²⁷, and a host of acute and chronic human pathologies, namely, sepsis²⁸, trauma²⁹, burns³⁰, organ transplantation³¹, diabetes³², myocardial infarction³³, stroke³⁴ and renal failure³⁵.

Conclusions

CNAs are a new class of physiological, continuously arising intra-corporeal mutagenic agents that might be responsible for ageing, age-related disabilities and ultimately the demise of the organism. Thus, DNA seems to act in paradoxical roles of both preserver and destroyer of life. This new class of intra-corporeal mobile genetic elements may be relevant not only to multi-cellular organisms which have a developed circulatory system, but also to other

multi-cellular organisms in which intra-corporeal mobility of CNAs may be mediated via the medium of extra-cellular fluid.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by the Department of Atomic Energy, Government of India, through its grant CTCTMC to Tata Memorial Centre awarded to IM.

Acknowledgement

I thank Dr. Amit Dutt for helpful suggestions.

References

1. McClintock B: **The origin and behavior of mutable loci in maize.** *Proc Natl Acad Sci USA.* 1950; **36**(6): 344–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Frost LS, Leplae R, Summers AO, *et al.*: **Mobile genetic elements: the agents of open source evolution.** *Nat Rev Microbiol.* 2005; **3**(9): 722–732.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Deininger PL, Moran JV, Batzer MA, *et al.*: **Mobile elements and mammalian genome evolution.** *Curr Opin Genet Dev.* 2003; **13**(6): 651–658.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Kazazian HH Jr: **Mobile elements: drivers of genome evolution.** *Science.* 2004; **303**(5664): 1626–1632.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Brookfield JF: **Evolutionary genetics: Mobile DNAs as sources of adaptive change?** *Curr Biol.* 2004; **14**(9): R344–345.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Curcio MJ, Derbyshire KM: **The outs and ins of transposition: from mu to kangaroo.** *Nat Rev Mol Cell Biol.* 2003; **4**(11): 865–877.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Keeling PJ, Palmer JD: **Horizontal gene transfer in eukaryotic evolution.** *Nat Rev Genet.* 2008; **9**(8): 605–618.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Soucy SM, Huang J, Gogarten JP: **Horizontal gene transfer: building the web of life.** *Nat Rev Genet.* 2015; **16**(8): 472–482.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Levin HL, Moran JV: **Dynamic interactions between transposable elements and their hosts.** *Nat Rev Genet.* 2011; **12**(9): 615–627.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Polz MF, Alm EJ, Hanage W: **Horizontal gene transfer and the evolution of bacterial and archaeal population structure.** *Trends Genet.* 2013; **29**(3): 170–175.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Crisp A, Boschetti C, Perry M, *et al.*: **Expression of multiple horizontally acquired genes is a hallmark of both vertebrate and invertebrate genomes.** *Genome Biol.* 2015; **16**: 50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. International Human Genome Sequencing Consortium, Adekoya E, Ait-Zahra M, *et al.*: **Initial sequencing and analysis of the human genome.** *Nature.* 2001; **409**(6822): 860–921.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Stanhope MJ, Lupas A, Italia MJ, *et al.*: **Phylogenetic analyses do not support horizontal gene transfers from bacteria to vertebrates.** *Nature.* 2001; **411**(6840): 940–944.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Riley DR, Sieber KB, Robinson KM, *et al.*: **Bacteria-human somatic cell lateral gene transfer is enriched in cancer samples.** *PLoS Comput Biol.* 2013; **9**(6): e1003107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Mittra I, Khare NK, Raghuram GV, *et al.*: **Circulating nucleic acids damage DNA of healthy cells by integrating into their genomes.** *J Biosci.* 2015; **40**(1): 91–111.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Fliedner TM, Graessle D, Paulsen C, *et al.*: **Structure and function of bone marrow hemopoiesis: mechanisms of response to ionizing radiation exposure.** *Cancer Biother Radiopharm.* 2002; **17**(4): 405–426.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. van Nieuwenhuijze AE, van Lopik T, Smeenk RJ, *et al.*: **Time between onset of apoptosis and release of nucleosomes from apoptotic cells: putative implications for systemic lupus erythematosus.** *Ann Rheum Dis.* 2003; **62**(1): 10–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Elshimali YI, Khaddour H, Sarkissyan M, *et al.*: **The clinical utilization of circulating cell free DNA (CCFDNA) in blood of cancer patients.** *Int J Mol Sci.* 2013; **14**(9): 18925–18958.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Gauthier VJ, Tyler LN, Mannik M: **Blood clearance kinetics and liver uptake of mononucleosomes in mice.** *J Immunol.* 1996; **156**(3): 1151–1156.
[PubMed Abstract](#)
20. Nevers P, Saedler H: **Transposable genetic elements as agents of gene instability and chromosomal rearrangements.** *Nature.* 1977; **268**(5616): 109–115.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Vijg J, Dollé ME: **Large genome rearrangements as a primary cause of aging.** *Mech Ageing Dev.* 2002; **123**(8): 907–915.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Bahar R, Hartmann CH, Rodriguez KA, *et al.*: **Increased cell-to-cell variation in gene expression in ageing mouse heart.** *Nature.* 2006; **441**(7096): 1011–1014.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Jacobs KB, Yeager M, Zhou W, *et al.*: **Detectable clonal mosaicism and its relationship to aging and cancer.** *Nat Genet.* 2012; **44**(6): 651–658.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Laurie CC, Laurie CA, Rice K, *et al.*: **Detectable clonal mosaicism from birth to old age and its relationship to cancer.** *Nat Genet.* 2012; **44**(6): 642–650.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. McConnell MJ, Lindberg MR, Brennand KJ, *et al.*: **Mosaic copy number variation in human neurons.** *Science.* 2013; **342**(6158): 632–637.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Cancer Research UK. **All Cancers (C00-C97 Excl. C44) Average Number of New Cases per Year and Age-Specific Incidence Rates, UK, 2009–2011.**
[Reference Source](#)
27. Pisetsky DS, Ullal AJ: **The blood nucleome in the pathogenesis of SLE.** *Autoimmun Rev.* 2010; **10**(1): 35–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Rhodes A, Wort SJ, Thomas H, *et al.*: **Plasma DNA concentration as a predictor of mortality and sepsis in critically ill patients.** *Crit Care.* 2006; **10**(2): R60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Lam NY, Rainer TH, Chan LY, *et al.*: **Time course of early and late changes in plasma DNA in trauma patients.** *Clin Chem.* 2003; **49**(8): 1286–1291.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Chiu TW, Young R, Chan LY, *et al.*: **Plasma cell-free DNA as an indicator of severity of injury in burn patients.** *Clin Chem Lab Med.* 2006; **44**(1): 13–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Lui YY, Woo KS, Wang AY, *et al.*: **Origin of plasma cell-free DNA after solid organ transplantation.** *Clin Chem.* 2003; **49**(3): 495–496.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Butt AN, Shalchi Z, Hamaoui K, *et al.*: **Circulating nucleic acids and diabetic complications.** *Ann NY Acad Sci.* 2006; **1075**: 258–270.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Chang CP, Chia RH, Wu TL, *et al.*: **Elevated cell-free serum DNA detected in patients with myocardial infarction.** *Clin Chim Acta.* 2003; **327**(1–2): 95–101.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Tsai NW, Lin TK, Chen SD, *et al.*: **The value of serial plasma nuclear and mitochondrial DNA levels in patients with acute ischemic stroke.** *Clin Chim Acta.* 2011; **412**(5–6): 476–479.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Mittra I, Nair NK, Mishra PK: **Nucleic acids in circulation: Are they harmful to the host?** *J Biosci.* 2012; **37**(2): 301–312.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 11 November 2015

<https://doi.org/10.5256/f1000research.7639.r11047>

© 2015 Rao M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Manchanahalli Rangaswamy Satyanarayana Rao

Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre For Advanced Scientific Research, Bangalore, India

This article details a new paradigm in cancer genetics based on the recent original article published by the same group. The idea of the origin of the cancer phenotype based on the circulating nucleic acid/ chromatin fragments is really thought provoking and will generate a lot of interest in the cancer researchers. The manuscript is well written summarizing the salient features of their recently published article.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 13 October 2015

<https://doi.org/10.5256/f1000research.7639.r10595>

© 2015 Gravina S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Silvia Gravina

Department of Genetics, Albert Einstein College of Medicine, Bronx, NY, USA

In this Opinion Article entitled "[Circulating nucleic acids: a new class of physiological mobile genetic elements](#)", Indraneel Mittra investigated the topic of circulating nucleic acids as a new class of continuously arising DNA damaging agents.

The author discusses a recent work (Mittra *et al.*, 2015) which describes how free nucleic acids not only can enter cell nuclei freely, but they integrate in the host genome and activate proteins of DDR and apoptotic pathways. In particular, in the Opinion Article, the author hypothesizes that cell free DNA continuously arising induced damage can be the cause of aging and age related diseases.

The manuscript is certainly well written and attractive, and discusses a very novel yet important topic in the aging field. My only concern is that in two parts of the manuscript the author jumps to conclusions too quickly, and I would suggest rephrasing certain statements.

More specifically:

- pg2, mid right; "Whether genomic integration of CNAs occurs preferentially in a site-specific manner or is random is not known; but in either case, integration of CNAs would give rise to somatic mutations in the host genome."

This has to be demonstrated yet. We don't really know if CNA integration is a direct cause of somatic mutations in the host genome. I would suggest making this statement milder (i.e. "Whether genomic integration of CNAs occurs preferentially in a site-specific manner or is random is not known. It is reasonable to speculate that integration of CNAs may give rise to somatic mutations in the host genome." or "a fascinating hypothesis yet to be tested is that that integration of CNAs may give rise to somatic mutations in the host genome".)

- Same for pg2 bottom part; "CNAs, on the other hand, are ubiquitous, physiological and continuously arising, inflicting repeated damage and mutations to the somatic DNA". I would suggest here-as well- a milder statement (i.e., CNAs, on the other hand, are ubiquitous, physiological and continuously arising, inflicting repeated damage and potentially causing mutations to the somatic DNA").

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research