REVIEW

The critical importance of the fetal hypothalamus-pituitary-adrenal axis [version 1; peer review: 3 approved]

Charles E. Wood¹, Maureen Keller-Wood²

¹Department of Physiology and Functional Genomics, University of Florida College of Medicine, FL, USA

²Department of Pharmacodynamics, University of Florida College of Pharmacy, FL, USA

V1 First published: 28 Jan 2016, 5(F1000 Faculty Rev):115
<https://doi.org/10.12688/f1000research.7224.1>

Latest published: 28 Jan 2016, 5(F1000 Faculty Rev):115
<https://doi.org/10.12688/f1000research.7224.1>

Abstract

The fetal hypothalamus-pituitary-adrenal (HPA) axis is at the center of mechanisms controlling fetal readiness for birth, survival after birth and, in several species, determination of the timing of birth. Stereotypical increases in fetal HPA axis activity at the end of gestation are critical for preparing the fetus for successful transition to postnatal life. The fundamental importance in fetal development of the endogenous activation of this endocrine axis at the end of gestation has led to the use of glucocorticoids for reducing neonatal morbidity in premature infants. However, the choice of dose and repetition of treatments has been controversial, raising the possibility that excess glucocorticoid might program an increased incidence of adult disease (e.g., coronary artery disease and diabetes). We make the argument that because of the critical importance of the fetal HPA axis and its interaction with the maternal HPA axis, dysregulation of cortisol plasma concentrations or inappropriate manipulation pharmacologically can have negative consequences at the beginning of extrauterine life and for decades thereafter.

Keywords

fetal hypothalamus-pituitary-adrenal axis, HPA, cortisol

Open Peer Review

Approval Status

	1	2	3
version 1 28 Jan 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Peter W. Nathanielsz**, University of Wyoming, Laramie, USA
2. **Charles Ducsay**, Loma Linda University, Loma Linda, USA
3. **Dean A. Myers**, University of Oklahoma HSC, Oklahoma City, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Charles E. Wood (woodc@ufl.edu)

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Wood CE and Keller-Wood M. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Wood CE and Keller-Wood M. **The critical importance of the fetal hypothalamus-pituitary-adrenal axis [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):115 <https://doi.org/10.12688/f1000research.7224.1>

First published: 28 Jan 2016, 5(F1000 Faculty Rev):115 <https://doi.org/10.12688/f1000research.7224.1>

Cortisol is arguably the most important hormone for organizing maturation in late gestation. The discovery that, in sheep, interruption of the fetal hypothalamus-pituitary-adrenal (HPA) axis prevents spontaneous parturition^{1,2} initiated a series of important studies on the role that cortisol plays in the coordination of fetal readiness for extrauterine life. The mechanism of parturition and the role that the fetal adrenal plays in primate species are complicated compared with ruminants and other large animals. But, although cortisol does not directly initiate parturition in the human, it does have other critically important actions in the human: actions that were first discovered in the sheep model. Perhaps the most impactful discovery was that fetal sheep that were premature but otherwise treated with glucocorticoid were less likely to die in respiratory distress³. The mechanism of this effect involves a direct action of glucocorticoid on the fetal lung, accelerating terminal development and stimulating the production of pulmonary surfactant^{4–6}. This discovery, rapidly translated to the human^{7,8}, has resulted in the widespread antenatal administration of glucocorticoids to pregnant women threatening premature labor^{9,10}. This clinical practice has been an effective strategy for reduction or elimination of ventilator assist of the premature infant after birth. In addition to its effect on the lung, cortisol plays an important role in accelerating development of the fetal gastrointestinal tract¹¹ and liver¹². Recent evidence also supports a role for cortisol in the development of the fetal cardiovascular system¹³ and heart¹⁴. Similar to its effect in other systems, cortisol helps ready the fetal cardiovascular system for the transition to extrauterine life.

A central tenet of endocrinology is that circulating concentrations of hormones are controlled; that there is a set point around which negative feedback mechanisms maintain concentrations within a range^{15,16}. For a hormonal system as important as the fetal HPA axis, understanding those mechanisms that govern the increases in fetal adrenocorticotrophic hormone (ACTH) and cortisol becomes critically important for understanding the timing of birth, readiness for birth, and survival of the newborn. This most often involves the elucidation of influences, both positive and negative, that determine the trajectory of fetal ACTH secretion. For example, Thorburn and colleagues discovered a dynamic increase in the sensitivity of the fetal adrenal cortex to ACTH that can account for much of the increase in circulating cortisol concentrations prior to birth¹⁷. Several laboratories have investigated changes in circulating molecular forms of immunoreactive ACTH (resulting from partial or complete processing of the parent molecule pro-opiomelanocortin), altering the effective biological activity of this pituitary hormone^{18–21}. New evidence suggests that peripheral interconversion of cortisol and cortisone is also an important variable that can contribute to changes in circulating concentrations of cortisol²². Balancing the stimulation of the axis are mechanisms that prevent over-activity of the axis. The late-gestation fetal sheep has a cortisol negative feedback mechanism that is remarkably sensitive to small changes in plasma cortisol concentration^{23–27}. In the final days and hours of fetal life, this negative feedback sensitivity falls dramatically, allowing greater increases in fetal HPA axis activity²⁸.

Although nearly every variable within the fetal HPA axis has been studied to some extent, we are not close to fully understanding the interplay between the normal fetal developmental patterns, the influence of fetal and maternal stress, and the modulatory influence of

infection on the timing of parturition and the survival of the newborn. A danger in the translation of basic endocrinology to clinical practice has been an under-appreciation of untoward actions of hormones when they are present in unnaturally high concentrations. A clear example of this is the use, in practice, of antenatal steroid administration to women threatening preterm labor. First approved by a National Institutes of Health (NIH) consensus panel as a single treatment or a small number of repeated treatments⁷, the procedure was modified by practicing physicians so as to administer glucocorticoid weekly from as early as 22 to 24 weeks' gestation²⁹. Forgetting the basic rule of endocrinology—that control mechanisms prevent excessively high, as well as excessively low concentrations of hormones—can create unintended problems. Weekly or biweekly antenatal maternal betamethasone treatment resulted in the birth of babies that were more likely to be growth-restricted^{29–31}. Studies in animal models indicate that multiple doses of glucocorticoid can have negative neurodevelopmental outcomes. Using the sheep model, Newnham and colleagues clearly demonstrated that excessive treatment with glucocorticoid dramatically impacts fetal brain growth and development^{32,33}. This laboratory and others have also demonstrated negative effects of excessive glucocorticoid on fetal somatic growth^{34,35}. Recent evidence in the baboon model reveals sex-specific effects of antenatal betamethasone on learning and attention in the offspring³⁶. Recent work has suggested that the effects of excess glucocorticoid can be codified in the epigenome of the infant, theoretically with effects that could last a lifetime^{37,38}. On balance, however, the long-term biological cost of multiple treatments is not clear. Clinical trials have not demonstrated long-term growth or major neurosensory disabilities but have indicated an increased likelihood of attention deficit in children who were exposed to multiple doses of glucocorticoid before birth³⁹. Arterial stiffness, however, was increased in 14- to 26-year-old subjects exposed to two to nine weekly doses of betamethasone⁴⁰. In recognition of the potential risks of multiple antenatal glucocorticoid treatments, a second NIH consensus statement recommended a single treatment⁴¹. As a result of the revised recommendations, the incidence of multiple treatment has been reduced.

The fetus is an organism that is distinct from, yet dependent on, its mother. The circulation of the fetus does not admix with the circulation of the mother, and the fetus in late gestation has the capacity to synthesize its own hormones. However, by virtue of the fact that the fetal and maternal blood is separated by one or more layers of cells in the placenta (depending on the species and therefore the general structure of the placenta), there is a molecular communication between mother and fetus. This communication is most obvious with regard to blood gases. Fetal growth and development depend on the oxygen and nutrients supplied by the mother via trans-placental passage. However, in addition to hormones synthesized within the placenta (e.g., chorionic gonadotropin and chorionic somatomammotropin), there is endocrine communication. An important example of this communication is the influence of maternal adrenal cortical hormones on the fetus, perhaps a mechanism by which the fetus is informed about maternal stress.

The placentas of many species, including the human, rodent species, and the sheep, express the enzyme 11 β -hydroxysteroid dehydrogenase types 1 and 2 (encoded by the genes *HSD11B1* and *HSD11B2*),

which interconvert cortisol and cortisone (or, in rodent species, corticosterone and 11-dehydrocorticosterone)^{42–45}. The predominant reaction in placenta is oxidation (cortisol to cortisone and corticosterone to 11-dehydrocorticosterone, mediated by the type 2 enzyme isoform). Because cortisone and 11-dehydrocorticosterone have low affinity for the glucocorticoid and mineralocorticoid receptors (GR and MR, respectively), some of the active glucocorticoid in the maternal circulation is inactivated upon passage into the fetal circulation. Although this is a partial barrier for the natural glucocorticoids, it is not complete. Some of the cortisol that was converted to cortisone upon transplacental passage is converted back to cortisol by the type 1 11 β -HSD isoform in target tissues, such as lung^{46,47}, brain⁴⁸, and heart^{49–51}. As shown in the chronically catheterized fetal sheep model, elevation of maternal cortisol concentration to levels typical of stress increases fetal plasma cortisol concentrations and inhibits fetal ACTH secretion via a negative feedback mechanism⁵². Synthetic glucocorticoids, such as dexamethasone or betamethasone, are not substrates for 11 β -HSD and therefore are not inactivated when passing through the placenta.

The passage of natural glucocorticoid across placenta from mother to fetus raises the question of whether maternal stress is detrimental to the fetus. There is recent interest in the potential “programming” effects of maternal stress on the pattern of development in the fetus⁵³. There are known glucocorticoid effects on gene methylation and histone modification that can have long-lasting effects on the physiology of the offspring. Perinatal programming of the HPA axis has been reviewed elsewhere⁵⁴. If “programming” can result from alterations in the genome, however, can it also result from a more immediate effect of hypercortisolemia that bends the arc of development? In other words, what are the more immediate consequences of maternal stress?

The literature is rife with reports of poor pregnancy outcome when the pregnancy is complicated by maternal stress: low socioeconomic class, partner abuse, and violence^{55–57}. It is not known whether these negative effects on the pregnancy are caused entirely, or even partly, by the maternal cortisol stress response, but there is evidence that suggests that cortisol is at least partly to blame. For example, women with Cushing disease have a higher incidence of stillbirth that is associated with obstructive hypertrophic cardiomyopathy⁵⁸. The effect of the maternal hypercortisolemia can be reproduced in the sheep model (using cortisol administration rates that mimic the maternal stress response)⁵⁹. However, maternal cortisol also has an effect on fetal somatic growth, in part secondary to alterations in uterine blood flow⁶⁰. Increases in maternal cortisol also alter fetal pulmonary and renal fluid balance mechanisms⁶¹. How many of these modifications of the pattern of fetal growth and development result in increased incidence of morbidity and mortality in postnatal life? For example, does chronic maternal stress cause small-for-gestation babies⁶² that are more prone to infant mortality⁶³ or metabolic disease in adulthood? Does cortisol-induced alteration in fetal cardiac development underlie the increased incidence of coronary artery disease in adults who were small for gestational age at birth⁶⁴?

As it is detrimental to chronically elevate maternal plasma cortisol concentration, it is important to remember that there is a physiological set point for cortisol in maternal blood during pregnancy, and that reductions in maternal cortisol below that level could be detrimental. Circulating concentrations of cortisol in the human and in other species, such as the sheep, are naturally increased in the latter half of pregnancy^{65–67}. Stress, as previously discussed, further elevates the concentration of cortisol^{68,69}. But the fact that the normal set point for cortisol is increased in pregnancy raises the question of why this occurs and whether there are species differences with regard to mechanism⁶⁸.

The normal pregnancy-associated increase in maternal adrenal secretion of cortisol appears to play an important role on both sides of the placenta. Prior to the final stages of fetal life *in utero*, the majority of the cortisol circulating in the fetal blood derives from the maternal adrenal glands⁷⁰. Because cortisol plays an important role in developmental processes important for fetal readiness for birth, it is likely that the increase in maternal cortisol plasma concentrations provides an important source of cortisol to the fetus before the fetal HPA axis becomes fully competent. On the other hand, reduction in maternal cortisol concentration below the normal set point, down to the level normally observed in the nonpregnant state, disturbs blood pressure, fluid balance, and uterine blood flow, and slows fetal growth⁶⁶. Although the above-mentioned studies were performed in sheep, evidence from human pregnancies is consistent: untreated maternal adrenal insufficiency causes premature labor and neonatal morbidity^{71,72}.

It is important to recognize both the importance and the complexity of the HPA axis during pregnancy and during fetal development. The mechanisms controlling this endocrine axis during pregnancy (why is maternal cortisol normally increased during late gestation?) and fetal life (what causes the increase in fetal HPA axis activity that is important for neonatal survival?) have never been fully solved. For example, is the ontogenetic rise in fetal HPA axis activity caused secondary to the development of immune cells within the fetal brain⁷³? Are there placental signaling molecules (e.g., Corticotropin Releasing Hormone [CRH] in primate species) that stimulate fetal HPA axis activity⁷⁴? Are there programmed developmental events within the fetal brain that progress without modification by circulating endocrine signals? We argue that answering these basic questions will be important for reducing neonatal mortality and morbidity, for improvement of maternal health, and the design of better and smarter treatments for women threatening preterm labor, and for the babies that are admitted to neonatal intensive care units.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. Liggins GC, Kennedy PC, Holm LW: **Failure of initiation of parturition after electrocoagulation of the pituitary of the fetal lamb.** *Am J Obstet Gynecol.* 1967; **98**(8): 1080–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. McDonald TJ, Nathanielsz PW: **Bilateral destruction of the fetal paraventricular nuclei prolongs gestation in sheep.** *Am J Obstet Gynecol.* 1991; **165**(3): 764–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Liggins GC: **Adrenocortical-related maturational events in the fetus.** *Am J Obstet Gynecol.* 1976; **126**(7): 931–41.
[PubMed Abstract](#)
4. Ballard PL: **Hormonal regulation of pulmonary surfactant.** *Endocr Rev.* 1989; **10**(2): 165–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Ballard PL, Ballard RA: **Glucocorticoid receptors and the role of glucocorticoids in fetal lung development.** *Proc Natl Acad Sci U S A.* 1972; **69**(9): 2668–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Ballard PL, Hawgood S, Liley H, *et al.*: **Regulation of pulmonary surfactant apoprotein SP 28-36 gene in fetal human lung.** *Proc Natl Acad Sci U S A.* 1986; **83**(24): 9527–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. **Effect of corticosteroids for fetal maturation on perinatal outcomes. NIH Consensus Development Panel on the Effect of Corticosteroids for Fetal Maturation on Perinatal Outcomes.** *JAMA.* 1995; **273**(5): 413–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Ballard PL, Ballard RA: **Scientific basis and therapeutic regimens for use of antenatal glucocorticoids.** *Am J Obstet Gynecol.* 1995; **173**(1): 254–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Liggins GC, Howie RN: **A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory distress syndrome in premature infants.** *Pediatrics.* 1972; **50**(4): 515–25.
[PubMed Abstract](#)
10. Vidaeff AC, Mastrobattista JM: **Controversies in the use of antenatal steroids for fetal maturation.** *Semin Perinatol.* 2001; **25**(6): 385–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Lebenthal A, Lebenthal E: **The ontogeny of the small intestinal epithelium.** *JPN J Parenter Enteral Nutr.* 1999; **23**(5 Suppl): S3–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Fowden AL, Forhead AJ: **Adrenal glands are essential for activation of glucogenesis during undernutrition in fetal sheep near term.** *Am J Physiol Endocrinol Metab.* 2011; **300**(1): E94–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Unno N, Wong CH, Jenkins SL, *et al.*: **Blood pressure and heart rate in the ovine fetus: ontogenic changes and effects of fetal adrenalectomy.** *Am J Physiol.* 1999; **276**(1 Pt 2): H248–56.
[PubMed Abstract](#)
14. **Rog-Zielinska EA, Richardson RV, Denvir MA, *et al.*: Glucocorticoids and foetal heart maturation; implications for prematurity and foetal programming.** *J Mol Endocrinol.* 2014; **52**(2): R125–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
15. Akana SF, Shinsako J, Dallman MF: **Closed-loop feedback control of the nyctohemeral rise in adrenocortical system function.** *Fed Proc.* 1985; **44**(1 Pt 2): 177–81.
[PubMed Abstract](#)
16. Keller-Wood ME, Dallman MF: **Corticosteroid inhibition of ACTH secretion.** *Endocr Rev.* 1984; **5**(1): 1–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Jacobs RA, Young IR, Hollingworth SA, *et al.*: **Chronic administration of low doses of adrenocorticotropin to hypophysectomized fetal sheep leads to normal term labor.** *Endocrinology.* 1994; **134**(3): 1389–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Carr GA, Jacobs RA, Young IR, *et al.*: **Development of adrenocorticotropin-(1-39) and precursor peptide secretory responses in the fetal sheep during the last third of gestation.** *Endocrinology.* 1995; **136**(11): 5020–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Castro MI, Valego NK, Zehnder TJ, *et al.*: **The ratio of plasma bioactive to immunoreactive ACTH-like activity increases with gestational age in the fetal lamb.** *J Dev Physiol.* 1992; **18**(4): 193–201.
[PubMed Abstract](#)
20. Schwartz J, Kleftogiannis F, Jacobs R, *et al.*: **Biological activity of adrenocorticotropin hormone precursors on ovine adrenal cells.** *Am J Physiol.* 1995; **268**(4 Pt 1): E623–9.
[PubMed Abstract](#)
21. Valego N, Castro MI, Zehnder T, *et al.*: **The ratio of plasma bioactive to immunoreactive ACTH-like activity changes with stress in the late gestation ovine fetus.** *73rd Annual Meeting of the Endocrine Society.* 1991; 43.
22. **Guo C, Li C, Myatt L, *et al.*: Sexually dimorphic effects of maternal nutrient reduction on expression of genes regulating cortisol metabolism in fetal baboon adipose and liver tissues.** *Diabetes.* 2013; **62**(4): 1175–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. Saoud CJ, Wood CE: **Androstenedione alters cortisol-negative feedback sensitivity in late gestation fetal sheep.** *Proceedings of the 77th Annual Meeting of the Endocrine Society P3-526.* 1995.
24. Wood CE: **Does a decrease in cortisol negative feedback efficacy precede ovine parturition?** *Am J Physiol.* 1987; **252**(3 Pt 2): R624–7.
[PubMed Abstract](#)
25. Wood CE: **Negative-feedback inhibition of fetal ACTH secretion by maternal cortisol.** *Am J Physiol.* 1987; **252**(4 Pt 2): R743–8.
[PubMed Abstract](#)
26. Wood CE: **Sensitivity of cortisol-induced inhibition of ACTH and renin in fetal sheep.** *Am J Physiol.* 1986; **250**(5 Pt 2): R795–802.
[PubMed Abstract](#)
27. Wood CE, Rudolph AM: **Negative feedback regulation of adrenocorticotropin secretion by cortisol in ovine fetuses.** *Endocrinology.* 1983; **112**(6): 1930–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Wood CE: **Insensitivity of near-term fetal sheep to cortisol: possible relation to the control of parturition.** *Endocrinology.* 1988; **122**(4): 1565–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Newnham JP, Jobe AH: **Should we be prescribing repeated courses of antenatal corticosteroids?** *Semin Fetal Neonatal Med.* 2009; **14**(3): 157–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. **Peltoniemi OM, Kari MA, Hallman M: Repeated antenatal corticosteroid treatment: a systematic review and meta-analysis.** *Acta Obstet Gynecol Scand.* 2011; **90**(7): 719–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. **Wapner RJ, Sorokin Y, Thom EA, *et al.*: Single versus weekly courses of antenatal corticosteroids: evaluation of safety and efficacy.** *Am J Obstet Gynecol.* 2006; **195**(3): 633–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
32. Huang WL, Beazley LD, Quinlivan JA, *et al.*: **Effect of corticosteroids on brain growth in fetal sheep.** *Obstet Gynecol.* 1999; **94**(2): 213–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. **Moss TJ, Doherty DA, Nitsos I, *et al.*: Effects into adulthood of single or repeated antenatal corticosteroids in sheep.** *Am J Obstet Gynecol.* 2005; **192**(1): 146–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
34. Jensen EC, Gallaher BW, Breier BH, *et al.*: **The effect of a chronic maternal cortisol infusion on the late-gestation fetal sheep.** *J Endocrinol.* 2002; **174**(1): 27–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Quinlivan JA, Archer MA, Dunlop SA, *et al.*: **Fetal growth retardation, particularly within lymphoid organs, following repeated maternal injections of betamethasone in sheep.** *J Obstet Gynaecol Res.* 1998; **24**(3): 173–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Rodriguez JS, Zürcher NR, Keenan KE, *et al.*: **Prenatal betamethasone exposure has sex specific effects in reversal learning and attention in juvenile baboons.** *Am J Obstet Gynecol.* 2011; **204**(6): 545.e1–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. **Hompes T, Izzi B, Gellens E, *et al.*: Investigating the influence of maternal cortisol and emotional state during pregnancy on the DNA methylation status of the glucocorticoid receptor gene (NR3C1) promoter region in cord blood.** *J Psychiatr Res.* 2013; **47**(7): 880–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
38. **Oberlander TF, Weinberg J, Papsdorf M, *et al.*: Prenatal exposure to maternal depression, neonatal methylation of human glucocorticoid receptor gene (NR3C1) and infant cortisol stress responses.** *Epigenetics.* 2008; **3**(2): 97–106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
39. **Crowther CA, Doyle LW, Haslam RR, *et al.*: Outcomes at 2 years of age after repeat doses of antenatal corticosteroids.** *N Engl J Med.* 2007; **357**(12): 1179–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. **Norberg H, Stålnacke J, Nordenström A, *et al.*: Repeat antenatal steroid exposure and later blood pressure, arterial stiffness, and metabolic profile.** *J Pediatr.* 2013; **163**(3): 711–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
41. **Antenatal corticosteroids revisited: repeat courses.** *NIH Consensus Statement.* 2000; **17**(2): 1–18.
[PubMed Abstract](#) | [F1000 Recommendation](#)
42. Alifaidy N, Li W, MacIntosh T, *et al.*: **Late gestation increase in 11beta-hydroxysteroid dehydrogenase 1 expression in human fetal membranes: a novel intrauterine source of cortisol.** *J Clin Endocrinol Metab.* 2003; **88**(10): 5033–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Mark PJ, Augustus S, Lewis JL, *et al.*: **Changes in the placental glucocorticoid barrier during rat pregnancy: impact on placental corticosterone levels and regulation by progesterone.** *Biol Reprod.* 2009; **80**(6): 1209–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Pepe GJ, Burch MG, Albrecht ED: **Localization and developmental regulation of 11beta-hydroxysteroid dehydrogenase-1 and -2 in the baboon syncytiotrophoblast.** *Endocrinology.* 2001; **142**(1): 68–80.
[PubMed Abstract](#) | [Publisher Full Text](#)

45. Yang K, Smith CL, Dales D, *et al.*: **Cloning of an ovine 11 beta-hydroxysteroid dehydrogenase complementary deoxyribonucleic acid: tissue and temporal distribution of its messenger ribonucleic acid during fetal and neonatal development.** *Endocrinology*. 1992; **131**(5): 2120–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Murphy BE: **Cortisol production and inactivation by the human lung during gestation and infancy.** *J Clin Endocrinol Metab*. 1978; **47**(2): 243–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Wood CE, Srun R: **Ontogeny of 11 beta-hydroxysteroid dehydrogenase in ovine fetal kidney and lung.** *Reprod Fertil Dev*. 1995; **7**(5): 1329–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Sloboda DM, Moss TJ, Li S, *et al.*: **Expression of glucocorticoid receptor, mineralocorticoid receptor, and 11beta-hydroxysteroid dehydrogenase 1 and 2 in the fetal and postnatal ovine hippocampus: ontogeny and effects of prenatal glucocorticoid exposure.** *J Endocrinol*. 2008; **197**(2): 213–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Reini SA, Wood CE, Jensen E, *et al.*: **Increased maternal cortisol in late-gestation ewes decreases fetal cardiac expression of 11beta-HSD2 mRNA and the ratio of AT1 to AT2 receptor mRNA.** *Am J Physiol Regul Integr Comp Physiol*. 2006; **291**(6): R1708–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Reini SA, Wood CE, Keller-Wood M: **The ontogeny of genes related to ovine fetal cardiac growth.** *Gene Expr Patterns*. 2009; **9**(2): 122–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Thompson A, Han VK, Yang K: **Differential expression of 11beta-hydroxysteroid dehydrogenase types 1 and 2 mRNA and glucocorticoid receptor protein during mouse embryonic development.** *J Steroid Biochem Mol Biol*. 2004; **88**(4–5): 367–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Wood CE, Rudolph AM: **Can maternal stress alter fetal adrenocorticotropin secretion?** *Endocrinology*. 1984; **115**(1): 298–301.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Cottrell EC, Seckl JR, Holmes MC, *et al.*: **Foetal and placental 11 β -HSD2: a hub for developmental programming.** *Acta Physiol (Oxf)*. 2014; **210**(2): 288–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Wood CE: **Development and programming of the hypothalamus-pituitary-adrenal axis.** *Clin Obstet Gynecol*. 2013; **56**(3): 610–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. **F** Dominguez TP: **Race, racism, and racial disparities in adverse birth outcomes.** *Clin Obstet Gynecol*. 2008; **51**(2): 360–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. **F** Shah PS, Shah J; Knowledge Synthesis Group on Determinants of Preterm/LBW Births: **Maternal exposure to domestic violence and pregnancy and birth outcomes: a systematic review and meta-analyses.** *J Womens Health (Larchmt)*. 2010; **19**(11): 2017–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
57. **F** Weck RL, Paulose T, Flaws JA: **Impact of environmental factors and poverty on pregnancy outcomes.** *Clin Obstet Gynecol*. 2008; **51**(2): 349–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. **F** Fayol L, Masson P, Millet V, *et al.*: **Cushing's syndrome in pregnancy and neonatal hypertrophic obstructive cardiomyopathy.** *Acta Paediatr*. 2004; **93**(10): 1400–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Keller-Wood M, Feng X, Wood CE, *et al.*: **Elevated maternal cortisol leads to relative maternal hyperglycemia and increased stillbirth in ovine pregnancy.** *Am J Physiol Regul Integr Comp Physiol*. 2014; **307**(4): R405–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Jensen E, Wood CE, Keller-Wood M: **Chronic alterations in ovine maternal corticosteroid levels influence uterine blood flow and placental and fetal growth.** *Am J Physiol Regul Integr Comp Physiol*. 2005; **288**(1): R54–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Jensen E, Wood CE, Keller-Wood M: **Alterations in maternal corticosteroid levels influence fetal urine and lung liquid production.** *J Soc Gynecol Investig*. 2003; **10**(8): 480–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. **F** Brunton PJ: **Effects of maternal exposure to social stress during pregnancy: consequences for mother and offspring.** *Reproduction*. 2013; **146**(5): R175–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. McCormick MC: **The contribution of low birth weight to infant mortality and childhood morbidity.** *N Engl J Med*. 1985; **312**(2): 82–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Barker DJ, Winter PD, Osmond C, *et al.*: **Weight in infancy and death from ischaemic heart disease.** *Lancet*. 1989; **2**(8663): 577–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Carr BR, Parker CR Jr, Madden JD, *et al.*: **Maternal plasma adrenocorticotropin and cortisol relationships throughout human pregnancy.** *Am J Obstet Gynecol*. 1981; **139**(4): 416–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Jensen E, Wood C, Keller-Wood M: **The normal increase in adrenal secretion during pregnancy contributes to maternal volume expansion and fetal homeostasis.** *J Soc Gynecol Investig*. 2002; **9**(6): 362–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Jensen EC, Bennet L, Wood C, *et al.*: **Loss of the pregnancy-induced rise in cortisol concentrations in the ewe impairs the fetal insulin-like growth factor axis.** *Reprod Fertil Dev*. 2011; **23**(5): 665–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Keller-Wood M: **Reflex regulation of hormonal responses during pregnancy.** *Clin Exp Pharmacol Physiol*. 1995; **22**(2): 143–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Kutzler MA, Coksaygan T, Ferguson AD, *et al.*: **Maternally administered dexamethasone at 0.7 of gestation suppresses maternal and fetal pituitary and adrenal responses to hypoxemia in sheep.** *Pediatr Res*. 2004; **55**(5): 755–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Hennessy DP, Coghlan JP, Hardy KJ, *et al.*: **The origin of cortisol in the blood of fetal sheep.** *J Endocrinol*. 1982; **95**(1): 71–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. **F** Björnsdóttir S, Cnattingius S, Brandt L, *et al.*: **Addison's disease in women is a risk factor for an adverse pregnancy outcome.** *J Clin Endocrinol Metab*. 2010; **95**(12): 5249–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
72. **F** Fux Otta C, Szafryk de Mereshian P, Iraci GS, *et al.*: **Pregnancies associated with primary adrenal insufficiency.** *Fertil Steril*. 2008; **90**(4): 1199.e17–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
73. Rabaglino MB, Keller-Wood M, Wood CE: **Transcriptomics of the late gestation ovine fetal brain: modeling the co-expression of immune marker genes.** *BMC Genomics*. 2014; **15**(1): 1001.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
74. McLean M, Bisits A, Davies J, *et al.*: **A placental clock controlling the length of human pregnancy.** *Nat Med*. 1995; **1**(5): 460–3.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Dean A. Myers**

Department of Obstetrics and Gynecology, University of Oklahoma HSC, Oklahoma City, OK, USA

Competing Interests: No competing interests were disclosed.

2. **Charles Ducsay**

Center for Perinatal Biology, Loma Linda University, Loma Linda, CA, USA

Competing Interests: No competing interests were disclosed.

3. **Peter W. Nathanielsz**

Life-course Health Research Center, University of Wyoming, Laramie, WY, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research