Check for updates

CASE REPORT

REVISED Case Report: Frontalis sign for early bedside consideration of impending uncal herniation [version 2; peer review: 2 approved]

Previously titled 'Case Report: Frontalis sign for early bedside diagnosis of impending uncal herniation'

Sunil Munakomi , Bijoy Mohan Kumar

Department of Neurosurgery, College of Medical Sciences, Bharatpur, Nepal

V2 First published: 01 Feb 2016, 5:125
<https://doi.org/10.12688/f1000research.7871.1>
 Latest published: 04 Aug 2016, 5:125
<https://doi.org/10.12688/f1000research.7871.2>

Abstract

It is prudent to have early diagnosis and timely management of uncal herniation for better management of neurosurgical patients. There are several clinical and radiological armamentariums that aid in early recognition of the condition. Through this case report, we try to highlight a simple bedside clinical sign that can be a valuable adjunct in early recognition of the impending uncal herniation especially in scenarios wherein it is difficult to assess the pupillary size and reactivity correctly. The improvement in the sign also confirms the resolution of the mass effect in the postoperative period. This is especially helpful for doctors working in the periphery or in resource restrained areas, for a timely referral of the patient to tertiary centre.

Keywords

Trauma , herniation , sign

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
04 Aug 2016	 view	
version 1		
01 Feb 2016	view	view

1. **G. Bryan Young**, University of Western Ontario, London, Canada
2. **Osman Sinanović**, University Clinical Center of Tuzla, Tuzla, Bosnia and Herzegovina
Sanela Zukić, University Clinical Center Tuzla, Tuzla, Bosnia and Herzegovina

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Sunil Munakomi (sunilmunakomi@gmail.com)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Munakomi S and Mohan Kumar B. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Munakomi S and Mohan Kumar B. **Case Report: Frontalis sign for early bedside consideration of impending uncal herniation [version 2; peer review: 2 approved]** F1000Research 2016, 5:125 <https://doi.org/10.12688/f1000research.7871.2>

First published: 01 Feb 2016, 5:125 <https://doi.org/10.12688/f1000research.7871.1>

REVISED Amendments from Version 1

We have highlighted the role of 'Frontalis sign' only as an adjunct clinical marker in considering early impending herniation. Assessing pupillary size and reactivity still is the workhorse in diagnosing the condition. We have made appropriate changes as per the suggestions of our referees.

See referee reports

Introduction

Traumatic brain injury (TBI) is now a global epidemic¹. The prognosis of patients with head injury is dependent on many clinical parameters but one of the major determinants is the time lapsed for appropriate management². TBI has a significant impact not only on the patients and their relatives but also has a major influence on the health and socioeconomical status in the global arena. Thus, it is prudent to have clinical tools for early recognition of life threatening neurosurgical emergencies. Herein we discuss one such example: Frontalis sign for detecting early uncal herniation. This may be helpful for early referral of patients to tertiary care centres and for timely management of the same. It could therefore have a positive impact on these patients, resulting in a better outcome.

Case report

A 50-year-old male from Siraha, a distant village in Nepal, was referred to our neurosurgical centre following a road traffic accident after being hit by a speeding car. The patient had a brief loss of consciousness and a single episode of vomiting following the incident. There was no history of seizurogenic activity observed during the transfer. On arrival to the emergency department, his Glasgow coma scale (GCS) was E3M6V5 with no paucity in movement of any limbs. His vital parameters were within normal range with blood pressure of 130/90, pulse rate of 86/min and oxygen saturation of 99% in room air. It was difficult to assess differences in pupillary size as he had corneal opacity on the left eye, resulting from an injury sustained during his childhood. However, on close examination, we observed that there was prominence of the forehead wrinkles on the right half of his face especially when the patient was trying to open his eyes during conversation, which we termed as frontalis sign (Figure 1). The wrinkles on the contra lateral half were normal with no abnormal deviation of angle of the mouth dismissing the differential diagnosis of upper facial nerve palsy. Because of the finding, we suspected impending uncal herniation in the patient and thereby advised for an emergency computed tomography (CT) scan of the head. It revealed right sided huge temporo-parietal contusion with thin fronto-temporo-parietal subdural hematoma with features of uncal herniation (Figure 2). The condition was explained to his relatives and they were counseled for emergency evacuation of the hematoma. On their consent, we performed a craniotomy, evacuation of the subdural hematoma and removal of the contusion. Following the procedure, the brain was lax and pulsatile. The patient was extubated without any untoward events in the postoperative period. The frontalis sign diminished following the surgery (Figure 3). The post operative scan confirmed resolution of the mass effect and normalization of the cisternal anatomy (Figure 4).

Figure 1. Prominence of wrinkles on the right forehead during attempted eye opening, along with ptosis termed as frontalis sign.

Figure 2. CT head image showing evidence of right sided uncal herniation with obliteration of the ipsilateral crural and the ambient cisterns following temporo-parietal huge contusion.

The patient was started on Levitracetam 500 mg intravenously every 12 hours which was changed to oral medication after three days as seizure prophylaxis. The patient was discharged after suture removal on the 8th postoperative day. The patient followed up in the outpatient clinic 2 weeks later in sound health. Eye opening was near normal. The patient was advised for monthly follow up.

Figure 3. Resolution of the frontalis sign in the early postoperative period.

Figure 4. Postoperative CT scan confirmed resolution of the mass effect and normalization of the cisternal anatomy.

Discussion

Levator palpebrae superioris supplied by the third nerve helps in elevation of the lid during eye opening^{4,5}. However third nerve involvement due to uncal herniation weakens the muscle thereby restricting its action⁶. In order to compensate the deficits, the frontalis belly of the occipito-frontalis muscle supplied by the facial nerve, helps in elevation of the lid^{7,8}. This leads to prominence of forehead wrinkles in the same side on comparison to the other half. This is termed frontalis sign. This can act as a reliable bedside marker for diagnosing impending herniation. There are other routine signs for impending uncal herniation such as anisocoria. But there may be inter-observer bias in assessing the same⁹. There may

be other causes for anisocoria such as Horner's syndrome following carotid artery dissection, traumatic third nerve palsy, Marcus gun pupil (Relative afferent pupillary defect) and ocular perforations¹⁰. Sometimes drugs such as Ipratropium Bromide used for nebulisation in the intensive care unit can cause anisocoria. It is also difficult to observe the size and reaction of the pupils in patients with severe eye lid swellings¹¹. The frontalis sign can be used as an adjunct for consideration of uncal herniation and thereby initiating the correct management. This is even more valuable for proper patient referral from peripheral and resource limited setups, especially in developing countries like ours who are still far behind implementing the guidelines for managing patients with TBI¹².

Conclusion

The implications of the use of this simple bedside sign for early diagnosis of the uncal diagnosis can be influential in providing timely and correct therapeutic targets for patients with TBI. It can be a valuable adjunct to the present panoply of our armamentarium in diagnosis the traumatic cerebral herniation syndromes.

Consent

Written informed consent was obtained from the daughter of the patient for publication of this case report and any accompanying images and/or other details that could potentially reveal the patient's identity.

Author contributions

SM reviewed the literature and formatted the paper. BB suggested, revised and edited the final format.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Jennett B: **Epidemiology of head injury.** *J Neurol Neurosurg Psychiatry.* 1996; **60**(4): 362–369.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jennett B, Teasdale G, Braakman R, *et al.*: **Prognosis of patients with severe head injury.** *Neurosurgery.* 1979; **4**(4): 283–289.
[PubMed Abstract](#)
- Humphreys I, Wood RL, Phillips CJ, *et al.*: **The costs of traumatic brain injury: a literature review.** *Clinicoecon Outcomes Res.* 2013; **5**: 281–287.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ettl A, Priglinger S, Kramer J, *et al.*: **Functional anatomy of the levator palpebrae superioris muscle and its connective tissue system.** *Br J Ophthalmol.* 1996; **80**(8): 702–707.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ng SK, Chan W, Marcet MM, *et al.*: **Levator palpebrae superioris: an anatomical update.** *Orbit.* 2013; **32**(1): 76–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Maramattom BV, Wijdicks EF: **Uncal herniation.** *Arch Neurol.* 2005; **62**(12): 1932–1935.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bérzin F: **Occipitofrontalis muscle: functional analysis revealed by electromyography.** *Electromyogr Clin Neurophysiol.* 1989; **29**(6): 355–8.
[PubMed Abstract](#)
- Kushima H, Matsuo K, Yuzuriha S, *et al.*: **The occipitofrontalis muscle is composed of two physiologically and anatomically different muscles separately affecting the positions of the eyebrow and hairline.** *Br J Plast Surg.* 2005; **58**(5): 681–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Olson DM, Stutzman S, Saju C, *et al.*: **Interrater Reliability of Pupillary Assessments.** *Neurocrit Care.* 2016; **24**(2): 251–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Atkins EJ, Newman NJ, Biouesse V: **Post-traumatic visual loss.** *Rev Neurol Dis.* 2008; **5**(2): 73–81.
[PubMed Abstract](#) | [Free Full Text](#)
- Yalcin S, Pampal K, Erden A, *et al.*: **Do we really need to panic in all anisocoria cases in critical care?** *Indian J Anaesth.* 2010; **54**(4): 365–366.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- National Collaborating Centre for Acute Care (UK): **Head Injury: Triage, Assessment, Investigation and Early Management of Head Injury in Infants, Children and Adults.** London: National Collaborating Centre for Acute Care (UK), (NICE Clinical Guidelines, No. 56.), 2007.
[PubMed Abstract](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 05 September 2016

<https://doi.org/10.5256/f1000research.10092.r13404>

© 2016 Young G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

G. Bryan Young

Department of Clinical Neurological Sciences, University of Western Ontario, London, ON, Canada

The revision is well done and any queries have been addressed.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 29 July 2016

<https://doi.org/10.5256/f1000research.8472.r15318>

© 2016 Sinanović O et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Osman Sinanović

Department of Neurology, Acta Medica Saliniana, University Clinical Center of Tuzla, Tuzla, Bosnia and Herzegovina

Sanela Zukić

Department of Neurology, University Clinical Center Tuzla, Tuzla, Bosnia and Herzegovina

In this case report authors try to highlight a simple bedside clinical sign (frontalis sign) in early recognition of the impending uncal herniation.

Title "Frontalis sign for early bedside diagnosis of impending uncal herniation" is appropriate for content of the article but with a clinical sign we can not make a diagnosis. So, title "Frontalis sign for early bedside consideration of impending uncal herniation" may be more appropriate.

Article content including design, case report and discussion is correct.

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 May 2016

<https://doi.org/10.5256/f1000research.8472.r13921>

© 2016 Young G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

G. Bryan Young

Department of Clinical Neurological Sciences, University of Western Ontario, London, ON, Canada

The paper is of interest. The authors should give details of pupillary reactivity. With herniation there is usually a change in pupillary size and reactivity before the ptosis. Was this not the case here?

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 03 Jul 2016

Sunil Munakomi, College of Medical Sciences, Bharatpur, Nepal

We thank you for the report on our article. We acknowledge the role of assessing pupillary size in patients with trauma. Here, we are emphasizing on the adjunct role of frontalis sign in diagnosing early uncal herniation in cases wherein pupillary assessment is problematic as in severe eye lid swelling, corneal injuries (in our case), following use of sedatives or following post traumatic seizure.

Competing Interests: No competing interests.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research