REVIEW

Plant adaptation to drought stress [version 1; peer review: 3 approved]

Supratim Basu*, Venkategowda Ramegowda*, Anuj Kumar, Andy Pereira

Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, Arkansas, 72701, USA

* Equal contributors

V1 **First published:** 30 Jun 2016, 5(F1000 Faculty Rev):1554
<https://doi.org/10.12688/f1000research.7678.1>
Latest published: 30 Jun 2016, 5(F1000 Faculty Rev):1554
<https://doi.org/10.12688/f1000research.7678.1>

Abstract

Plants in their natural habitats adapt to drought stress in the environment through a variety of mechanisms, ranging from transient responses to low soil moisture to major survival mechanisms of escape by early flowering in absence of seasonal rainfall. However, crop plants selected by humans to yield products such as grain, vegetable, or fruit in favorable environments with high inputs of water and fertilizer are expected to yield an economic product in response to inputs. Crop plants selected for their economic yield need to survive drought stress through mechanisms that maintain crop yield. Studies on model plants for their survival under stress do not, therefore, always translate to yield of crop plants under stress, and different aspects of drought stress response need to be emphasized. The crop plant model rice (*Oryza sativa*) is used here as an example to highlight mechanisms and genes for adaptation of crop plants to drought stress.

Keywords

Adaptation , Drought tolerance , drought resistance , grain yield , rice, photosynthesis ,

Open Peer Review

Approval Status

	1	2	3
version 1
30 Jun 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Krishna Jagadish**, Kansas State University, Manhattan, USA
2. **Zhulong Chan**, Chinese Academy of Sciences, Wuhan, China
3. **Arvind Kumar**, International Rice Research Institute, Metro Manila, Philippines

Any comments on the article can be found at the end of the article.

Corresponding author: Andy Pereira (apereira@uark.edu)

Competing interests: The authors declare that they have no competing interests.

Grant information: This work was supported by the National Science Foundation award numbers DBI-0922747 and ABI1062472. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2016 Basu S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Basu S, Ramegowda V, Kumar A and Pereira A. **Plant adaptation to drought stress [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):1554 <https://doi.org/10.12688/f1000research.7678.1>

First published: 30 Jun 2016, 5(F1000 Faculty Rev):1554 <https://doi.org/10.12688/f1000research.7678.1>

Introduction

Drought stress is the most prevalent environmental factor limiting crop productivity¹, and global climate change is increasing the frequency of severe drought conditions². The sheer diversity of plant species grown across climatic regions that include extreme dry conditions suggests that, in nature, plants have evolved to endure drought stress with an array of morphological, physiological, and biochemical adaptations³. 'Drought resistance' (DR) is a broader term applied to plant species with adaptive features that enable them to escape, avoid, or tolerate drought stress⁴. 'Drought escape' is the ability of a plant species to complete its life cycle before the onset of drought. Thereby, plants do not experience drought stress, as they are able to modulate their vegetative and reproductive growth according to water availability, essentially through two different mechanisms: rapid phenological development and developmental plasticity⁵. Rapid phenological development involves rapid plant growth, producing a minimal number of seeds before the soil water depletes, and these plants are considered not to have any special morphological, physiological, or biochemical adaptations. Plants with mechanisms of developmental plasticity show little growth during the dry season, with very few flowers and seeds, but in wet seasons they grow indeterminately, producing a large amount of seed. 'Drought avoidance' is the ability of plants to maintain (relatively) higher tissue water content despite reduced water content in the soil⁴. This is achieved through a variety of adaptive traits involving the minimization of water loss (water savers) and optimization of water uptake (water spenders). Water spenders achieve higher tissue water status by maintaining the water uptake through increased rooting, hydraulic conductance, etc. under drought stress. In contrast, water savers use water effectively through reduced loss of water by reducing transpiration, transpiration area, radiation absorption, etc. under drought stress. 'Drought tolerance' (DT) is the ability of plants to endure low tissue water content through adaptive traits. These adaptive traits involve maintenance of cell turgor through osmotic adjustment and cellular elasticity, and increasing protoplasmic resistance⁶.

Improvement of yield and maintaining yield stability of crops, under normal as well as drought stress conditions, is essential for the food security of the growing global population. It is difficult to resolve the role of different components of DR in the stability of the crop yield as the major objective. However, there exist a variety of different mechanisms for drought escape, avoidance, or tolerance in natural populations that can improve DR and maintain grain yield in crop plants. In nature, extreme DR is found in resurrection plants^{7,8} which possess strong drought escape mechanisms. Resurrection plants can be exposed to severe drought for months, extending up to years, forcing them to optimize their growth for survival, but not for seed production, in the long term⁹. Therefore, the DR mechanisms that enable plants to merely survive longer lead to subsistence yield, which is much lower than that which is observed under normal conditions. Crop plants, on the other hand, are grown by humans in environments under conditions for high agricultural production and will be exposed to only a random short-term drought stress of days to weeks, from which they must quickly respond to limit the damage caused by

short-term drought stress while they continue to grow and yield in the stressful environments. Therefore, bringing in the drought adaptive mechanisms from plants adapted to grow in extreme dry conditions may not be a feasible option, as it may result in growth and/or yield penalty in crop plants under drought as well as normal conditions.

Although plant survival is very critical in the early growth stages, the mechanisms have little relevance to increasing grain yield directly. The emphasis to improve DR of crop plants should therefore be based on stability of yield components and not on plant survival alone. So far, most of the efforts to improve grain yield under drought stress were focused on secondary traits such as root architecture, leaf water potential, osmotic adjustment, and relative water content at the vegetative stage, which are often not highly correlated with grain yield^{10,11}. Looking forward in crops, the effective drought improvement approach should be selection for yield and its component traits under reproductive-stage drought stress¹². Additionally, little importance has also been given to simultaneous improvement of grain yield under normal and drought conditions. Selection for DT has been suggested to have a yield drag under normal conditions. It has been proposed that the yield potential of crop plants should be simultaneously selected for under favorable and environmental stress conditions, as there is a positive correlation between yield potential under normal and drought stress conditions¹³. Combining high yield potential under normal conditions with good yield under drought stress is the ideal trait. Identification of mechanisms, traits, and genes regulating yield under drought stress that are free from yield drag under normal conditions should be the focus. For example, regulation of yield under normal as well as drought stress conditions has been shown for three NAC family transcription factors (TFs). Transgenic plants expressing *OsNAC5*, *OsNAC9*, and *OsNAC10* TFs showed an increase in grain yield of 5-26% under normal conditions¹⁴⁻¹⁶. Nevertheless, in these studies, yield under normal conditions has been overlooked with more emphasis given to yield under drought stress. Two of our recent studies show the potential of simultaneously improving and stabilizing grain yield, both under normal as well as drought stress conditions, using two regulatory genes, namely *GUDK* and *HYR* in rice^{17,18}. These studies indicate that it might be advantageous to identify mechanisms and genes for increasing grain yield that are also stable or maintained under drought stress conditions.

Despite the complexity of DR, tremendous progress has been made in understanding the drought-adaptive mechanisms of plants^{1,19,20}. Adaptation through DR mainly involves morphophysiological alterations. These alterations in adaptive processes are controlled by molecular mechanisms that regulate the expression of genes²¹. There exists a large diversity in drought adaptation within a crop species, as some genotypes are able to cope with drought better than others. Genotypes that differ in drought adaptive mechanisms serve as an important resource to study the variation in drought adaption in crop plants. This natural variation needs to be exploited to simultaneously improve DR and yields of cultivated varieties through better understanding of the

underlying mechanisms and to aid in selection for these traits²². In the following sections, we describe the widely known morpho-physiological processes and recent molecular advances in regulating these drought-adaptive processes leading to increased yield in crop plants.

Photosynthesis

Drought stress is known to reduce photosynthesis by decreasing both leaf area and photosynthetic rate per unit leaf area. Reduced photosynthetic rate is mainly through stomatal closure or metabolic impairment²³. Continued photosynthetic light reactions during drought stress under limited intercellular CO₂ concentration results in the accumulation of reduced photosynthetic electron transport components, which can potentially reduce molecular oxygen, resulting in the production of reactive oxygen species (ROS). ROS can cause severe damage to the photosynthetic apparatus²⁴. The adaptive responses that plants have developed to reduce drought-induced damage to photosynthesis include thermal dissipation of light energy, the xanthophyll cycle, the water-water cycle, and dissociation of the light-harvesting complexes from photosynthetic reaction centers^{25–27}. The metabolic impairment during drought stress is mainly caused by changes in photosynthetic carbon metabolism²⁴. The biochemical efficiency of photosynthesis under drought stress mainly depends on ribulose-1,5-bisphosphate (RuBP) regeneration and the activity of ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)^{28,29}. Considerable progress has been made in improving the stomatal components for CO₂ diffusion, photosynthetic light reaction, and metabolic changes, including the expression of photosynthesis-related genes to regulate photosynthesis under drought towards the improvement of grain yield³⁰.

The C4 pathway of carbon assimilation has been suggested to be the major adaptation of the C3 pathway to limit water loss, reduce photorespiration, and improve photosynthetic efficiency under drought stress³¹. However, many important crops—including rice, wheat, soybean, and potato—use the C3 pathway of photosynthesis. Although the transfer of the C4 pathway into C3 crops is underway, so far its contribution to increased grain yield is very limited³². Photosynthetic adaptation of plants to drought stress involves a complex interaction of hormones, ROS, sugars, and other metabolic events³³. Combinations of computational models, which integrate the physiological and metabolic processes with gene expression data, along with modern breeding and transgenic technologies hold promise in improving photosynthesis and hence crop yield under normal as well as drought stress conditions.

In recent studies, we used a rice gene regulatory network to identify a TF termed HYR (HIGHER YIELD RICE), which was highly associated with primary carbon metabolism¹⁷, and on overexpression in rice enhanced photosynthesis under normal conditions as well as under drought and high temperature stress. HYR regulates several morpho-physiological processes leading to higher yield under normal and environmental stress conditions. Our study showed that HYR is a master regulator of photosynthesis, directly activating photosynthesis genes, cascades

of TFs, and other downstream genes involved in photosynthetic carbon metabolism, resulting in improved yield.

Hormonal regulation

Major phytohormones, such as abscisic acid (ABA), cytokinin (CK), gibberellic acid (GA), auxin, and ethylene, regulate diverse processes which enable plant adaptation to drought stress³⁴. Upon exposure of plants to drought stress, ABA is the major hormone synthesized in roots and translocated to leaves to initiate adaptation of plants to drought stress through stomatal closure and reduced plant growth³⁵. However, modulating the ABA-induced drought adaptation of plants for better yield remains a greater challenge because of the potential inadvertent reduction in carbon gain upon stomatal closure and ABA-induced senescence, especially if the drought occurs at the reproductive stage³⁶. There are ABA signaling genes, such as *OsNAP*, *OsNAC5*, and *DSM2*, which promote improved yield under reproductive drought^{37–40}. These ABA-induced non-stomatal adaptations of plants under drought stress can be exploited to improve grain yield under reproductive drought (Figure 1).

Under drought stress, CKs are known to delay premature leaf senescence and death, adaptive traits very useful for increasing grain yield. An increase in the endogenous levels of CK through expression of *isopentenyltransferase* (*IPT*), a CK biosynthetic pathway gene, leads to stress adaptation by delaying drought-induced senescence and an increase in yield^{41,42}. Generally, auxin has been shown to negatively regulate drought adaptation in plants. Decrease in indole-3-acetic acid (IAA) content was shown to be associated with up-regulation of genes encoding late embryogenesis abundant (LEA) proteins, leading to drought adaptation in plants^{43,44}. Recently, the *DEEPER ROOTING 1* (*DRO1*) gene determining a quantitative trait locus (QTL) controlling root growth angle was shown to be negatively regulated by auxin. Higher expression of *DRO1* in a shallow-rooting rice cultivar resulted in drought avoidance and high yield under drought⁴⁵. GA is suggested to positively regulate plant adaptation to drought stress. A rapid decline in levels of endogenous GA was observed in plants subjected to drought stress, resulting in growth inhibition⁴⁶. The role of GA in regulating grain yield of crop plants is thus an important area that can be further explored. Ethylene is a negative regulator of drought stress response by promoting leaf senescence and inhibiting root growth and development, shoot/leaf expansion, and photosynthesis^{47–51}. Ethylene can also directly affect yield by increasing embryo and grain abortion and reducing the grain-filling rate⁵². In addition to the major hormones, other hormones such as brassinosteroids, jasmonic acid (JA), salicylic acid (SA), and strigolactone also have an equally important role in plant growth and development. However, their function under drought stress is relatively less characterized. Tillering in rice has been suggested to be the outcome of an interaction among three hormones, CK, auxin, and strigolactone, with CK promoting branching and the other two inhibiting it^{53,54}, suggesting that all hormones do not act in isolation but instead interact and modulate each other's biosynthesis and responses. Therefore, the net outcome of the drought stress response is regulated by a balance between hormones that promote and those that inhibit the trait, rather than individual hormones.

Figure 1. The abscisic acid (ABA)-dependent gene regulatory pathway in rice. This pathway is required for drought stress tolerance and grain yield under drought. The drought stress ABA-dependent signal is shown perceived directly by the regulatory genes described in the text, followed by transcriptional regulation of downstream genes and underlying stress response mechanisms. Genes regulating drought tolerance (DT) at the vegetative stage are shaded green, and genes regulating DT and grain yield under drought are shaded orange. The resulting phenotypes are represented for DT at the vegetative level only (green diamonds), or DT and grain yield (orange diamonds). The genes described here are *OsGH3*⁹², *OsNAP*, *OsABI2*³⁹, *AP37*⁷³, *OsPP2C09*, *OsPP2C06*³⁴, *OsPYL/RCAR5*, *OsSIDP366*³⁵, *OsMYB48*³⁶, *OsRK1*³⁷, *OsHox22*³⁸, *SNAC2*^{39,100}, *OsOAT*¹⁰¹, *OsBZIP23*¹⁰², *SNAC1*³⁹, *OsEREBP1*¹⁰³, *OsBZIP71*¹⁰⁴, *OsBZIP46*¹⁰⁵, *OsABI5*¹⁰⁶, *DSM2*⁴⁰, *AREB2*¹⁰⁷, *OsSRO1c*¹⁰⁸, and *OsABA8OX3*¹⁰⁹.

Transpiration and stomatal conductance

The immediate response of plants on being exposed to drought stress is stomatal closure. However, stomatal closure not only diminishes water loss through transpiration but also reduces CO₂ and nutrient uptake, and hence alters metabolic pathways such as photosynthesis⁵⁵. Plants growing in dry areas have developed xeromorphic traits to reduce transpiration under drought stress. Reduction in transpiration under drought stress conditions can also be achieved through leaf shedding (i.e. deciduous species in drought) as well as decrease in leaf number, leaf size, and branching. Another adaptation to counter drought stress is sclerophylly, where plants form hard leaves that will not suffer permanent damage due to wilting and can be restored to full functionality when normal conditions resume⁵⁶. Recent research has shown that decreased stomatal conductance in response to drought stress is related not only to reduced expression of aquaporin genes but also to anatomical traits leading to reduction of chloroplast surface area exposed to intercellular space per unit leaf area^{57,58}. Several other factors, including leaf developmental stage and light availability, are also known to interact with drought in modulating mesophyll and chloroplast differentiation, ultimately affecting conductance and photosynthetic capacity⁵⁸. Reduction in stomatal size and

number on exposure to drought is another adaptation for survival under drought conditions. Previous studies have shown that while there is an increase in stomatal density under mild drought stress, there is a decrease during severe drought⁵⁹. Thus, all these adaptations in plants reduce the negative impacts of drought stress on photosynthesis and thereby have a positive effect on water use efficiency (WUE), which in turn will result in high yield potential and high yield⁶⁰. Such an adaptation was shown in rice by overexpression of the Arabidopsis AP2/ERF TF *HARDY* that improved WUE (the ratio of biomass produced to water used) by enhancing photosynthesis and reducing transpiration⁶¹. The above reported traits therefore exemplify adaptive mechanisms in plants to survive under drought stress without loss of productivity or yield.

Root morphology

In many agriculturally important crops, drought stress is perceived first by the root system, which continues to grow underneath the soil even though shoot growth is inhibited under these conditions⁶². Although the growth of the primary root is not affected by drought stress, the growth of lateral roots is significantly reduced, mainly by suppression of the activation of the lateral root meristems⁶³. The Arabidopsis R2R3-type MYB TF *MYB96* has been shown to

regulate activation of lateral root meristem through an ABA signaling cascade, with an activation-tagged mutant showing enhanced DR with reduced lateral root formation⁶⁴. The plant microRNA miR393 has also been shown to play a role in root-mediated adaptation to drought stress response through attenuation of auxin signaling⁶⁵. In addition to the lateral roots, the presence of small roots is also considered as an adaptive strategy to increase water uptake by providing more absorptive surface. Presence of specialized tissues like rhizodermis, with a thickened outer cell wall or suberized exodermis, or reduction in the number of cortical layers are considered an adaptive advantage for drought stress survival. Hydrotropism is another adaptive measure taken by plants to counter stress, where studies have shown that degradation of amyloplasts in the columella cells of plant roots on exposure to drought stress increases hydrotropism^{66,67}. Hormonal cross-talk mediated by auxin, CK, GA, and ABA has been implicated as a potential chemical signal in response to water stress to modulate root system architecture⁶⁸.

The expression of enzymes related to root morphology (e.g. xyloglucan endotransglucosylase) is induced upon mild drought stress, while other structural proteins are down-regulated, which is strongly correlated with root growth and hence an augmentation in the surface area for water uptake. The alterations in the expression of these proteins correlate positively with lateral development that in turn also affects photosynthesis⁶⁹. More lateral root and root hair formation was found in lines possessing a QTL, qDTY12.1, only when under drought⁷⁰. Such traits, which are expressed only under drought stress, have higher potential to increase grain yield under drought. Moreover, it has also been shown that drought stress triggers a wide variety of anatomical traits expressed to different levels and patterns in different species and even in different cultivars within species^{71–73}. For example, suberization and compaction of sclerenchyma layer cells were shown to decrease in rice under drought, which increases retention of water under drought stress⁷¹.

Osmotic adjustment

Osmotic adjustment (OA) is defined as a process of solute accumulation in dividing cells when the water potential is reduced, and thereby helps in maintaining the turgor⁷⁴. Cell enlargement and growth in plants is highly dependent on water availability and helps in maintaining the turgor. Turgor measurement in growing regions of plants, especially the leaves and stems, shows little or no reduction, though cell enlargement is inhibited during drought stress and is believed to be due to OA^{75,76}. Under conditions of drought stress, OA has been implicated in maintaining stomatal conductance, photosynthesis, leaf water volume, and growth^{74,77}. At times of drought stress, in addition to the reduction in water content, there are also other associated changes, such as increases in salt concentration and mechanical impedance⁷⁸. Inorganic cations, organic acids, carbohydrates, and free amino acids are the known predominant solutes that accumulate in response to water stress. Previous studies have shown that drought-resistant wheat varieties, with yield stability under drought stress, have a greater capacity for osmoregulation than less resistant varieties⁷⁶. The accumulation of compatible solutes such as proline and glycine betaine help in protecting the plants from detrimental effects of drought stress not

only by OA but also by detoxification of ROS, protection of membrane integrity, and stabilization of enzymes or proteins⁷⁹. Enzymes such as betaine aldehyde dehydrogenase (BADH), pyrroline-5-carboxylate reductase (P5CR), and ornithine δ -aminotransferase (OAT) have been shown to play major roles in OA. Overexpression of Arabidopsis *EDT1/HDG11* was shown to increase DT of poplar and cotton through increased accumulation of solutes such as proline and soluble sugars and also increase the yield of cotton in the field⁸⁰. However, there are some plants in which sugars are the main osmolytes that play a significant role in OA, including sucrose, trehalose, glucose, and fructose. Previous studies have shown that overexpression of the *sucrose:fructan-6-fructosyltransferase* (*6-SFT*) gene from *Psathyrostachys huashanica* in tobacco and the trehalose-6-phosphate phosphatase gene *OsTPP1* in rice confers abiotic stress tolerance^{81,82}. Researchers have also identified a QTL for OA on chromosome 8 in rice that is homeologous with a segment of wheat chromosome 7⁸³.

Source-sink relationships

Source-sink relationships largely determine the grain yield of cereal crops, with developing grains being primary sinks while the top two leaves, the flag leaf in particular, are the primary source^{84,85}. Drought stress affects the source-sink relationship by reducing the source strength, leading to yield reduction. Sufficient sugar supply through photosynthesis, transport, and conversion of sugars is regarded as the most critical component in determining the viability of reproductive organs in rice^{86–88}. Drought stress dramatically affects pollen viability due to abnormal starch accumulation⁸⁹. Insufficient starch synthesis and arrested pollen development have been linked to reduced invertase activity under drought stress⁸⁸. In addition to invertases, active grain filling involves other key enzymes, such as sucrose synthase, ADP glucose pyrophosphorylase (AGPase), and starch synthase as well as starch branching and debranching enzymes, which are also affected by drought stress⁸⁸. Therefore, enhancing rice yield through source-sink relationships involves not only sugar metabolism but also the regulated mobilization of metabolic resources from source to sink tissue.

Future research perspectives

Improving DR in crop plants is a challenge for plant breeders and crop physiologists, as it is a complex genetic trait with multiple pathways involved. Effective development of drought-resistant crop plants thus requires the pyramiding and interaction of many mechanisms, traits, and genes that are appropriate to individual crops and their growing environments. Success in this direction not only extends the growing area of crop plants but also achieves stable yield in drought-prone areas. Identifying genetic variation for DR is the first step towards development of drought-resistant crop plants. Such variation is often present in wild species and adapted genotypes that have evolved under natural selection and these are the best source of DR traits. Evaluation of these resources through an integrated phenotyping and genotyping approach under field conditions alongside identification of traits that are directly associated with yield is key to improving DR⁹⁰. Comparative omics analyses between the diverse germplasm could aid in bettering our understanding of the variety of crop adaptations to drought stress.

In addition, analysis of specific genes focussed on increasing DR while stabilizing the yield is crucial for understanding the broad basis of complex traits such as DR⁹¹. The development of high-yielding resilient crops⁷⁴ that maintain yield stability under drought and other environmental stresses due to climate change is also currently needed. Drought-resistant plants should combine a better root system, stomatal regulation, WUE, and hormonal balance while avoiding the negative effects on grain yield under both normal and drought stress conditions. Therefore, a holistic crop improvement strategy should involve the deployment of high crop yield potential and the utilization of a combination of morpho-physiological, biochemical, and anatomical adaptive responses to drought stress.

Author contributions

All authors contributed to writing the manuscript. Supratim Basu and Andy Pereira prepared [Figure 1](#).

Competing interests

The authors declare that they have no competing interests.

Grant information

This work was supported by the National Science Foundation award numbers DBI-0922747 and ABI1062472.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Bray EA: **Plant responses to water deficit**. *Trends Plant Sci.* 1997; 2(2): 48–54.
[Publisher Full Text](#)
- Dai A: **Increasing drought under global warming in observations and models**. *Nat Clim Chang.* 2012; 3: 52–8.
[Publisher Full Text](#)
- Bohnert HJ, Nelson DE, Jensen RG: **Adaptations to Environmental Stresses**. *Plant Cell.* 1995; 7(7): 1099–111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Levitt J: **Responses of plants to environmental stresses. Volume II. Water, radiation, salt, and other stresses**. (Academic Press), 1980.
[Reference Source](#)
- Jones MM, Turner NC, Osmond CB: **Mechanisms of drought resistance**. The physiology and biochemistry of drought resistance in plants. 1981; 1: 15–37.
- Morgan JM: **Osmoregulation and Water Stress in Higher Plants**. *Annu Rev Plant Physiol.* 1984; 35: 299–319.
[Publisher Full Text](#)
- Gaff DF: **Desiccation tolerant plants in South America**. *Oecologia.* 1987; 74(1): 133–6.
[Publisher Full Text](#)
- Gaff DF: **Desiccation-tolerant flowering plants in southern Africa**. *Science.* 1971; 174(4013): 1033–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Porembski S: **Granitic and gneissic outcrops (inselbergs) as centers of diversity for desiccation-tolerant vascular plants**. *Plant Ecol.* 2000; 151(1): 19–28.
[Publisher Full Text](#)
- Jongdee B, Fukai S, Cooper M: **Leaf water potential and osmotic adjustment as physiological traits to improve drought tolerance in rice**. *Field Crops Res.* 2002; 76(2–3): 153–63.
[Publisher Full Text](#)
- Pantawan G, Fukai S, Cooper M, et al.: **Yield response of rice (*Oryza sativa* L.) genotypes to drought under rainfed lowland: 3. Plant factors contributing to drought resistance**. *Field Crops Res.* 2002; 73(2–3): 181–200.
[Publisher Full Text](#)
- Venuprasad R, Lafitte HR, Atlin GN: **Response to Direct Selection for Grain Yield under Drought Stress in Rice**. *Crop Sci.* 2007; 47(1): 285–293.
[Publisher Full Text](#) | [F1000 Recommendation](#)
- Guan YS, Serraj R, Liu SH, et al.: **Simultaneously improving yield under drought stress and non-stress conditions: a case study of rice (*Oryza sativa* L.)**. *J Exp Bot.* 2010; 61(15): 4145–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeong JS, Kim YS, Redillas MC, et al.: ***OsNAC5* overexpression enlarges root diameter in rice plants leading to enhanced drought tolerance and increased grain yield in the field**. *Plant Biotechnol J.* 2013; 11(1): 101–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Redillas MC, Jeong JS, Kim YS, et al.: **The overexpression of *OsNAC9* alters the root architecture of rice plants enhancing drought resistance and grain yield under field conditions**. *Plant Biotechnol J.* 2012; 10(7): 792–805.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeong JS, Kim YS, Baek KH, et al.: **Root-specific expression of *OsNAC10* improves drought tolerance and grain yield in rice under field drought conditions**. *Plant Physiol.* 2010; 153(1): 185–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ambavaram MM, Basu S, Krishnan A, et al.: **Coordinated regulation of photosynthesis in rice increases yield and tolerance to environmental stress**. *Nat Commun.* 2014; 5: 5302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ramegowda V, Basu S, Krishnan A, et al.: **Rice *GROWTH UNDER DROUGHT KINASE* is required for drought tolerance and grain yield under normal and drought stress conditions**. *Plant Physiol.* 2014; 166(3): 1634–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yordanov I, Velikova V, Tsonev T: **Plant responses to drought, acclimation, and stress tolerance**. *Photosynthetica.* 2000; 38(2): 171–86.
[Publisher Full Text](#)
- Bray EA: **Molecular Responses to Water Deficit**. *Plant Physiol.* 1993; 103(4): 1035–40.
[PubMed Abstract](#) | [Free Full Text](#)
- Osakabe Y, Osakabe K, Shinozaki K, et al.: **Response of plants to water stress**. *Front Plant Sci.* 2014; 5: 86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Reynolds M, Tuberosa R: **Translational research impacting on crop productivity in drought-prone environments**. *Curr Opin Plant Biol.* 2008; 11(2): 171–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tezara W, Mitchell VJ, Driscoll SD, et al.: **Water stress inhibits plant photosynthesis by decreasing coupling factor and ATP**. *Nature.* 1999; 401: 914–7.
[Publisher Full Text](#)
- Lawlor DW, Cornic G: **Photosynthetic carbon assimilation and associated metabolism in relation to water deficits in higher plants**. *Plant Cell Environ.* 2002; 25(2): 275–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Niyogi KK: **PHOTOPROTECTION REVISITED: Genetic and Molecular Approaches**. *Annu Rev Plant Physiol Mol Biol.* 1999; 50: 333–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Demmig-Adams B, Adams WW III: **The role of xanthophyll cycle carotenoids in the protection of photosynthesis**. *Trends Plant Sci.* 1996; 1(1): 21–6.
[Publisher Full Text](#)
- Demmig-Adams B, Adams WW 3rd: **Photoprotection in an ecological context: the remarkable complexity of thermal energy dissipation**. *New Phytol.* 2006; 172(1): 11–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lawlor DW: **Limitation to photosynthesis in water-stressed leaves: stomata vs. metabolism and the role of ATP**. *Ann Bot.* 2002; 89(Spec No): 871–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Medrano H, Parry MAJ, Socias X, et al.: **Long term water stress inactivates Rubisco in subterranean clover**. *Ann Appl Biol.* 1997; 131(3): 491–501.
[Publisher Full Text](#)
- Chaves MM, Flexas J, Pinheiro C: **Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell**. *Ann Bot.* 2009; 103(4): 551–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Edwards G, Walker D: **C3, C4: mechanisms, and cellular and environmental regulation, of photosynthesis**. Univ of California Press, 1983.
[Reference Source](#)
- Gowik U, Westhoff P: **The path from C₃ to C₄ photosynthesis**. *Plant Physiol.* 2011; 155(1): 56–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pinheiro C, Chaves MM: **Photosynthesis and drought: can we make metabolic connections from available data?** *J Exp Bot.* 2011; 62(3): 869–82.
[PubMed Abstract](#) | [Publisher Full Text](#)

34. Wilkinson S, Kudoyarova GR, Veselov DS, *et al.*: **Plant hormone interactions: innovative targets for crop breeding and management.** *J Exp Bot.* 2012; **63**(9): 3499–509.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Wilkinson S, Davies WJ: **Drought, ozone, ABA and ethylene: new insights from cell to plant to community.** *Plant Cell Environ.* 2010; **33**(4): 510–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Ji X, Dong B, Shiran B, *et al.*: **Control of abscisic acid catabolism and abscisic acid homeostasis is important for reproductive stage stress tolerance in cereals.** *Plant Physiol.* 2011; **156**(2): 647–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Sperotto RA, Ricachenevsky FK, Duarte GL, *et al.*: **Identification of up-regulated genes in flag leaves during rice grain filling and characterization of OsNAC5, a new ABA-dependent transcription factor.** *Planta.* 2009; **230**(5): 985–1002.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. **F** Liang C, Wang Y, Zhu Y, *et al.*: **OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice.** *Proc Natl Acad Sci U S A.* 2014; **111**(27): 10013–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
39. **F** Chen X, Wang Y, Lv B, *et al.*: **The NAC family transcription factor OsNAP confers abiotic stress response through the ABA pathway.** *Plant Cell Physiol.* 2014; **55**(3): 604–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
40. Du H, Wang N, Cui F, *et al.*: **Characterization of the beta-carotene hydroxylase gene DSM2 conferring drought and oxidative stress resistance by increasing xanthophylls and abscisic acid synthesis in rice.** *Plant Physiol.* 2010; **154**(3): 1304–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Peleg Z, Reguera M, Tumimbang E, *et al.*: **Cytokinin-mediated source/sink modifications improve drought tolerance and increase grain yield in rice under water-stress.** *Plant Biotechnol J.* 2011; **9**(7): 747–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Peleg Z, Blumwald E: **Hormone balance and abiotic stress tolerance in crop plants.** *Curr Opin Plant Biol.* 2011; **14**(3): 290–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Xie Z, Jiang D, Cao W, *et al.*: **Relationships of endogenous plant hormones to accumulation of grain protein and starch in winter wheat under different post-anthesis soil water statuses.** *Plant Growth Regul.* 2003; **41**(2): 117–27.
[Publisher Full Text](#)
44. Zhang S, Li C, Cao J, *et al.*: **Altered architecture and enhanced drought tolerance in rice via the down-regulation of indole-3-acetic acid by TLD1/OsGH3.13 activation.** *Plant Physiol.* 2009; **151**(4): 1889–901.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. **F** Uga Y, Sugimoto K, Ogawa S, *et al.*: **Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions.** *Nat Genet.* 2013; **45**(9): 1097–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
46. Wang C, Yang A, Yin H, *et al.*: **Influence of water stress on endogenous hormone contents and cell damage of maize seedlings.** *J Integr Plant Biol.* 2008; **50**(4): 427–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Munné-Bosch S, Alegre L: **Die and let live: Leaf senescence contributes to plant survival under drought stress.** *Funct Plant Biol.* 2004; **31**(3): 203–216.
[Publisher Full Text](#)
48. **F** Fukao T, Xu K, Ronald PC, *et al.*: **A variable cluster of ethylene response factor-like genes regulates metabolic and developmental acclimation responses to submergence in rice.** *Plant Cell.* 2006; **18**(8): 2021–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
49. Perata P, Voesenek LA: **Submergence tolerance in rice requires Sub1A, an ethylene-response-factor-like gene.** *Trends Plant Sci.* 2007; **12**(2): 43–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Rajala A, Pelttonen-Sainio P: **Plant Growth Regulator Effects on Spring Cereal Root and Shoot Growth.** *Agron J.* 2001; **93**(4): 936–943.
[Publisher Full Text](#)
51. Sharp RE: **Interaction with ethylene: changing views on the role of abscisic acid in root and shoot growth responses to water stress.** *Plant Cell Environ.* 2002; **25**(2): 211–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Yang JC, Zhang JH, Ye YX, *et al.*: **Involvement of abscisic acid and ethylene in the responses of rice grains to water stress during filling.** *Plant Cell Environ.* 2004; **27**(8): 1055–64.
[Publisher Full Text](#)
53. **F** Gomez-Roldan V, Feras S, Brewer PB, *et al.*: **Strigolactone inhibition of shoot branching.** *Nature.* 2008; **455**(7210): 189–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
54. Xing Y, Zhang Q: **Genetic and molecular bases of rice yield.** *Annu Rev Plant Biol.* 2010; **61**: 421–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Xiong L, Zhu J: **Molecular and genetic aspects of plant responses to osmotic stress.** *Plant Cell Environ.* 2002; **25**(2): 131–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Micco VD, Aronne G: in *Plant Responses to Drought Stress.* (ed Aroca. Ricardo). Springer, 2002.
57. Miyazawa S, Yoshimura S, Shinzaki Y, *et al.*: **Deactivation of aquaporins decreases internal conductance to CO₂ diffusion in tobacco leaves grown under long-term drought.** *Funct Plant Biol.* 2008; **35**(7): 553–564.
[Publisher Full Text](#)
58. Tosens T, Niinemets U, Vislap V, *et al.*: **Developmental changes in mesophyll diffusion conductance and photosynthetic capacity under different light and water availabilities in *Populus tremula*: how structure constrains function.** *Plant Cell Environ.* 2012; **35**(5): 839–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Xu Z, Zhou G: **Responses of leaf stomatal density to water status and its relationship with photosynthesis in a grass.** *J Exp Bot.* 2008; **59**(12): 3317–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Blum A: **Drought resistance, water-use efficiency, and yield potential—are they compatible, dissonant, or mutually exclusive?** *Aust J Agric Res.* 2005; **56**(11): 1159–1168.
[Publisher Full Text](#)
61. Karaba A, Dixit S, Greco R, *et al.*: **Improvement of water use efficiency in rice by expression of *HARDY*, an *Arabidopsis* drought and salt tolerance gene.** *Proc Natl Acad Sci U S A.* 2007; **104**(39): 15270–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Spollen WG, Sharp RE: **Spatial distribution of turgor and root growth at low water potentials.** *Plant Physiol.* 1991; **96**(2): 438–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. Deak KI, Malamy J: **Osmotic regulation of root system architecture.** *Plant J.* 2005; **43**(1): 17–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Seo PJ, Park C: **Auxin homeostasis during lateral root development under drought condition.** *Plant Signal Behav.* 2009; **4**(10): 1002–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Chen H, Li Z, Xiong L: **A plant microRNA regulates the adaptation of roots to drought stress.** *FEBS Lett.* 2012; **586**(12): 1742–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Jaffe MJ, Takahashi H, Biro RL: **A pea mutant for the study of hydrotropism in roots.** *Science.* 1985; **230**(4724): 445–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Takahashi N, Yamazaki Y, Kobayashi A, *et al.*: **Hydrotropism interacts with gravitropism by degrading amyloplasts in seedling roots of *Arabidopsis* and radish.** *Plant Physiol.* 2003; **132**(2): 805–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. **F** Bilou I, Xu J, Wildwater M, *et al.*: **The PIN auxin efflux facilitator network controls growth and patterning in *Arabidopsis* roots.** *Nature.* 2005; **433**(7021): 39–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
69. **F** Sengupta D, Reddy AR: **Water deficit as a regulatory switch for legume root responses.** *Plant Signal Behav.* 2011; **6**(6): 914–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
70. Dixit S, Kumar Biswal A, Min A, *et al.*: **Action of multiple intra-QTL genes concerted around a co-localized transcription factor underpins a large effect QTL.** *Sci Rep.* 2015; **5**: 15183.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. Henry A, Cal AJ, Batoto TC, *et al.*: **Root attributes affecting water uptake of rice (*Oryza sativa*) under drought.** *J Exp Bot.* 2012; **63**(13): 4751–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Huang B, Fry JD: **Root Anatomical, Physiological, and Morphological Responses to Drought Stress for Tall Fescue Cultivars.** *Crop Sci.* 1998; **38**(4): 1017–1022.
[Publisher Full Text](#)
73. Steudle E: **Water uptake by roots: effects of water deficit.** *J Exp Bot.* 2000; **51**(350): 1531–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Chaves MM, Oliveira MM: **Mechanisms underlying plant resilience to water deficits: prospects for water-saving agriculture.** *J Exp Bot.* 2004; **55**(407): 2365–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. Meyer RF, Boyer JS: **Sensitivity of cell division and cell elongation to low water potentials in soybean hypocotyls.** *Planta.* 1972; **108**(1): 77–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Serraj R, Sinclair TR: **Osmolyte accumulation: can it really help increase crop yield under drought conditions?** *Plant Cell Environ.* 2002; **25**(2): 333–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Oosterhuis DM, Wulfschleger SD: **Osmotic Adjustment in Cotton (*Gossypium hirsutum* L.) Leaves and Roots in Response to Water Stress.** *Plant Physiol.* 1987; **84**(4): 1154–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Sauter A, Davies WJ, Hartung W: **The long-distance abscisic acid signal in the droughted plant: the fate of the hormone on its way from root to shoot.** *J Exp Bot.* 2001; **52**(363): 1991–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Ashraf M, Foolad MR: **Roles of glycine betaine and proline in improving plant**

- abiotic stress resistance. *Environ Exp Bot.* 2007; **59**(2): 206–16.
[Publisher Full Text](#)
80. **F** Yu LH, Wu SJ, Peng YS, *et al.*: **Arabidopsis ED1/HDG11 improves drought and salt tolerance in cotton and poplar and increases cotton yield in the field.** *Plant Biotechnol J.* 2016; **14**(1): 72–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 81. **F** He X, Chen Z, Wang J, *et al.*: **A sucrose: Fructan-6-fructosyltransferase (6-SFT) gene from *Psathyrostachys huashanica* confers abiotic stress tolerance in tobacco.** *Gene.* 2015; **570**(2): 239–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 82. Ge L, Chao D, Shi M, *et al.*: **Overexpression of the trehalose-6-phosphate phosphatase gene *OsTPP1* confers stress tolerance in rice and results in the activation of stress responsive genes.** *Planta.* 2008; **228**(1): 191–201.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Ahn S, Anderson JA, Sorrells ME, *et al.*: **Homoeologous relationships of rice, wheat and maize chromosomes.** *Mol Gen Genet.* 1993; **241**(5–6): 483–90.
[PubMed Abstract](#)
 84. Hirota O, Oka M, Takeda T: **Sink activity estimation by sink size and dry matter increase during the ripening stage of barley (*Hordeum vulgare*) and rice (*Oryza sativa*).** *Ann Bot.* 1990; **65**(4): 349–353.
[Reference Source](#)
 85. Gladun IV, Karpov EA: **Production and partitioning of assimilates between the panicle and vegetative organs of rice after flowering.** *Russ J Plant Physiol.* 1993; **40**(5): 629–633.
[Reference Source](#)
 86. **F** Li X, Lawas LMF, Malo R, *et al.*: **Metabolic and transcriptomic signatures of rice floral organs reveal sugar starvation as a factor in reproductive failure under heat and drought stress.** *Plant Cell Environ.* 2015; **38**(10): 2171–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 87. Nguyen GN, Hailstones DL, Wilkes M, *et al.*: **DROUGHT STRESS: Role of Carbohydrate Metabolism in Drought-Induced Male Sterility in Rice Anthers.** *J Agron Crop Sci.* 2010; **196**(5): 346–57.
[Publisher Full Text](#)
 88. Sheoran IS, Saini HS: **Drought-induced male sterility in rice: Changes in carbohydrate levels and enzyme activities associated with the inhibition of starch accumulation in pollen.** *Sexual Plant Reprod.* 1996; **9**(3): 161–9.
[Publisher Full Text](#)
 89. Jin Y, Yang H, Wei Z, *et al.*: **Rice male development under drought stress: phenotypic changes and stage-dependent transcriptomic reprogramming.** *Mol Plant.* 2013; **6**(5): 1630–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Tuberosa R: **Phenotyping for drought tolerance of crops in the genomics era.** *Front Physiol.* 2012; **3**: 347.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 91. Nguyen TT, Klueva N, Chamareck V, *et al.*: **Saturation mapping of QTL regions and identification of putative candidate genes for drought tolerance in rice.** *Mol Genet Genomics.* 2004; **272**(1): 35–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
 92. Du H, Wu N, Fu J, *et al.*: **A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice.** *J Exp Bot.* 2012; **63**(18): 6467–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 93. Oh SJ, Kim YS, Kwon CW, *et al.*: **Overexpression of the Transcription Factor AP37 in Rice Improves Grain Yield under Drought Conditions.** *Plant Physiol.* 2009; **150**(3): 1368–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 94. **F** He Y, Hao Q, Li W, *et al.*: **Identification and Characterization of ABA Receptors in *Oryza sativa*.** *PLoS One.* 2014; **9**(4): e95246.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. **F** Guo C, Luo C, Guo L, *et al.*: **OsSIDP366, a DUF1644 gene, positively regulates responses to drought and salt stresses in rice.** *J Integr Plant Biol.* 2016; **58**(5): 492–502.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 96. **F** Xiong H, Li J, Liu P, *et al.*: **Overexpression of OsMYB48-1, a Novel MYB-Related Transcription Factor, Enhances Drought and Salinity Tolerance in Rice.** *PLoS One.* 2014; **9**(3): e92913.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 97. Chae M, Lee JS, Nam MH, *et al.*: **A rice dehydration-inducible SNF1-related protein kinase 2 phosphorylates an abscisic acid responsive element-binding factor and associates with ABA signaling.** *Plant Mol Biol.* 2007; **63**(2): 151–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
 98. Zhang S, Haider I, Kohlen W, *et al.*: **Function of the HD-Zip I gene *Oshox22* in ABA-mediated drought and salt tolerances in rice.** *Plant Mol Biol.* 2012; **80**(6): 571–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 99. Hu H, Dai M, Yao J, *et al.*: **Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice.** *Proc Natl Acad Sci U S A.* 2006; **103**(35): 12987–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 100. Hu H, You J, Fang Y, *et al.*: **Characterization of transcription factor gene *SNAC2* conferring cold and salt tolerance in rice.** *Plant Mol Biol.* 2008; **67**(1–2): 169–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
 101. You J, Hu H, Xiong L: **An ornithine δ -aminotransferase gene *OsOAT* confers drought and oxidative stress tolerance in rice.** *Plant Sci.* 2012; **197**: 59–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
 102. Xiang Y, Tang N, Du H, *et al.*: **Characterization of OsbZIP23 as a key player of the basic leucine zipper transcription factor family for conferring abscisic acid sensitivity and salinity and drought tolerance in rice.** *Plant Physiol.* 2008; **148**(4): 1938–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 103. **F** Jisha V, Dampanaboina L, Vadassery J, *et al.*: **Overexpression of an AP2/ERF Type Transcription Factor *OsEREBP1* Confers Biotic and Abiotic Stress Tolerance in Rice.** *PLoS One.* 2015; **10**(6): e0127831.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 104. **F** Liu C, Mao B, Ou S, *et al.*: **OsbZIP71, a bZIP transcription factor, confers salinity and drought tolerance in rice.** *Plant Mol Biol.* 2014; **84**(1–2): 19–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 105. Tang N, Zhang H, Li X, *et al.*: **Constitutive activation of transcription factor *OsbZIP46* improves drought tolerance in rice.** *Plant Physiol.* 2012; **158**(4): 1755–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 106. Zou M, Guan Y, Ren H, *et al.*: **A bZIP transcription factor, OsABI5, is involved in rice fertility and stress tolerance.** *Plant Mol Biol.* 2008; **66**(6): 675–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
 107. Yoshida T, Fujita Y, Sayama H, *et al.*: **AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation.** *Plant J.* 2010; **61**(4): 672–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 108. You J, Zong W, Li X, *et al.*: **The *SNAC1*-targeted gene *OsSRO1c* modulates stomatal closure and oxidative stress tolerance by regulating hydrogen peroxide in rice.** *J Exp Bot.* 2013; **64**(2): 569–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 109. **F** Cai S, Jiang G, Ye N, *et al.*: **A key ABA catabolic gene, *OsABA8ox3*, is involved in drought stress resistance in rice.** *PLoS One.* 2015; **10**(2): e0116646.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. Arvind Kumar**
International Rice Research Institute, Metro Manila, Philippines
Competing Interests: No competing interests were disclosed.
- 2. Zhulong Chan**
Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan, China
Competing Interests: No competing interests were disclosed.
- 3. Krishna Jagadish**
Kansas State University, Manhattan, KS, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research