REVIEW

Recent advances in phototherapy for psoriasis [version 1; peer review: 2 approved]

Mio Nakamura¹, Benjamin Farahnik², Tina Bhutani¹

¹Department of Dermatology, University of California San Francisco Psoriasis and Skin Treatment Center, San Francisco, California, USA

²University of Vermont College of Medicine, Burlington, Vermont, USA

V1 First published: 13 Jul 2016, 5(F1000 Faculty Rev):1684
<https://doi.org/10.12688/f1000research.8846.1>
 Latest published: 13 Jul 2016, 5(F1000 Faculty Rev):1684
<https://doi.org/10.12688/f1000research.8846.1>

Abstract

Phototherapy involves repeated exposure of the skin to ultraviolet light to treat various inflammatory skin conditions such as psoriasis. Recent studies have identified specific immunologic effects of phototherapy that may underlie phototherapy efficacy. Furthermore, recent advancements have been made in developing safe and effective targeted phototherapy modalities for difficult-to-treat areas such as scalp psoriasis. Targeted phototherapy in the form of the excimer laser holds potential for more aggressive, effective treatment and long-lasting remission of psoriasis. Phototherapy is now also used successfully with biologic agents as combination therapy to treat recalcitrant psoriasis. Therefore, though one of the oldest therapeutic modalities for psoriasis, phototherapy remains a mainstay treatment with promise for further advancement.

Keywords

Psoriasis , Phototherapy , ultraviolet , UV

Open Peer Review

Approval Status

	1	2
version 1
13 Jul 2016		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Joel Gelfand**, University of Pennsylvania
Perelman School of Medicine, Philadelphia, USA
2. **Steven R Feldman**, Wake Forest School of Medicine, Winston-Salem, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Mio Nakamura (mio.nakamura@ucsf.edu)

Competing interests: Tina Bhutani is an advisor for Cutanea (Wayne, PA, USA) and conducts research for AbbVie (North Chicago, IL, USA), Janssen (Beerse, Belgium), PhotoMedex (Horsham, PA, USA), Merck/Sun Pharmaceutical Industries (Kenilworth, NJ, USA), and Novartis (Basel, Switzerland). Mio Nakamura and Benjamin Farahnik declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Nakamura M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Nakamura M, Farahnik B and Bhutani T. **Recent advances in phototherapy for psoriasis [version 1; peer review: 2 approved]** F1000Research 2016, 5(F1000 Faculty Rev):1684 <https://doi.org/10.12688/f1000research.8846.1>

First published: 13 Jul 2016, 5(F1000 Faculty Rev):1684 <https://doi.org/10.12688/f1000research.8846.1>

Introduction

Phototherapy involves repeated exposure of the skin to ultraviolet (UV) light to treat various inflammatory skin conditions such as psoriasis, eczema, and vitiligo. This therapy is one of the oldest treatment modalities in dermatology, dating back to the ancient Egyptians, who used natural light in combination with herbal extracts to treat skin disease¹. Phototherapy continues to be a highly preferred treatment by dermatologists²; however, its use has been decreasing in the past decades³. The cause of this decline is likely multifactorial, including lack of training⁴, low insurance reimbursement rates, and increase in cost of phototherapy treatments³. Developments have been made in the past several decades to make phototherapy an effective, convenient, and accessible treatment option for patients; however, despite research efforts, there is still much to be learned about phototherapy, including mechanism of action and how best to maximize effectiveness of treatments. This review will focus on recent advances in phototherapy for the treatment of psoriasis, including new information on the possible impact of UV light on the immunologic environment, new methods of delivering targeted light treatments by efficient yet safe methods, and using phototherapy in combination with biologic agents to treat recalcitrant psoriasis.

There are three main types of phototherapy used for the treatment of psoriasis: broadband ultraviolet B (BB-UVB), narrowband ultraviolet B (NB-UVB), and psoralen plus ultraviolet A (PUVA). BB-UVB was the first to be developed, in the 1940s, and emits wavelengths of light of between 290 and 313 nm. NB-UVB, which emits wavelengths of between 308 and 313 nm, was developed in the late 1980s when studies demonstrated that wavelengths of around 311 nm were more effective than broad-spectrum UVB in clearing psoriasis. UVA, with wavelengths of 320 to 400 nm, is used in combination with psoralen, a photosensitizing medication, which can be taken orally prior to UVA exposure in systemic PUVA or applied topically in soak PUVA⁵. NB-UVB is the most commonly used phototherapy modality today as it has a wider application across various dermatologic conditions, is easier to use, and has fewer side effects when compared with BB-UVB or PUVA⁶.

Phototherapy is most commonly administered in the office setting in stand-up light booths. The starting dose of UV light is determined by the individual patient's minimal erythema dose (MED) or by Fitzpatrick skin type, and subsequent dosing is increased as tolerated. Approximately 60% to 75% of patients with moderate to severe psoriasis achieve at least 75% improvement in the Psoriasis Area and Severity Index (PASI-75)⁷. The various modes of phototherapy, including BB-UVB, NB-UVB, systemic PUVA, and soak PUVA, appear to have comparable efficacy⁸. Phototherapy is relatively well tolerated, and the most common side effect is burning or itching. There is no overwhelming evidence for increased risk of skin cancers except in Caucasian patients who have undergone more than 250 treatments of systemic PUVA⁹.

Phototherapy treatments in a hospital or office setting are inconvenient and unfeasible for some patients. Many patients have difficulty committing to the phototherapy regimen in the office because of time restraints, transportation problems, mobility issues, and high insurance co-pays¹⁰. For these patients, home UV units have

been developed. The use of home phototherapy for patients with psoriasis was first reported in 1979 in a study from Sweden in patients who lived in remote areas far from their nearest hospital phototherapy unit¹¹. Home phototherapy is associated with lower burden of treatment and increased patient satisfaction¹².

Recent advances

The immunoregulatory effects of phototherapy: possible pathways

Psoriasis is caused by abnormal interactions among innate immune cells, T cells, and keratinocytes, leading to activation of the T helper cell type 1/T helper cell type 17 (Th1/Th17) immune axes and related cytokines. This contributes to the hyperproliferation and inflammation seen in psoriasis. There are various mechanisms by which phototherapy may be effective for psoriasis¹³. First, UV light induces apoptosis of keratinocytes and T cells in the epidermis and dermis¹⁴. Second, UV light promotes immunosuppression by promoting migration of Langerhans cells out of the epidermis¹⁵ as well as decreasing mast cell degranulation and histamine release¹⁶. Lastly, UV light induces alterations in the cytokine profile of psoriasis.

Research in the last 5 years has led to a better understanding of the specific pathways and cytokines that are altered by phototherapy. Phototherapy shifts the immune response away from the Th1/Th17 pathway, toward the counter-regulatory Th2 axis¹⁷. The Th1/Th17 pathway is suppressed by NB-UVB, leading to decreased levels of interleukin-12 (IL-12), IL-17, IL-20, IL-22, and IL-23¹⁸. These effects on cytokines appear to be systemic and not just localized to psoriatic lesions. PUVA and NB-UVB lower plasma levels of tumor necrosis factor- α , IL-17, IL-22, and IL-23 at the end of 6 weeks of treatment¹⁹. Furthermore, regulatory T (Treg) cells of patients with severe psoriasis display an enhanced propensity to convert into IL-17A-producing cells, which is linked to loss of forkhead box P3 (Foxp3)²⁰. UVB increases Foxp3-positive Treg cells in psoriatic skin lesions²¹. This increase in Foxp3 expression improves Treg cell stability and reduces pro-inflammatory Th1/Th17 cytokines in psoriatic skin lesions.

Handheld phototherapy: targeting difficult-to-treat psoriasis in the office and at home

Whole-body phototherapy is limited in that there can be needless exposure of uninvolved skin and no benefit to unexposed skin (such as the hair-covered scalp). In the last decade, various portable and lightweight handheld phototherapy units have become available for the treatment of localized psoriasis in the office and at home (Table 1)^{22,23}. These handheld devices, as compared with full-body irradiance in a booth or by a panel, have the added benefit of limiting skin exposure to UV light. The handheld devices are useful for the treatment of scalp psoriasis as well as recalcitrant localized psoriasis plaques. For example, the Dermalight 90 by National Biological Corporation has a comb attachment that permits direct application of light to scalp lesions. Although large-scale clinical trials are lacking, such light combs appear to be efficacious with longer remission compared with topical treatments^{24,25}. The handheld devices typically deliver NB-UVB, but some devices use BB-UVB or UVA. The Dualight by Theralight, Inc. and Psoria-Light by Psoria-Shield can deliver both UVB and UVA, but these devices are for use only in an office setting.

Table 1. Handheld phototherapy devices^{11,19}.

Series	Source	Treatment area, inches	Size, inches	Features
Dermalight 90 (National Biological Corporation)	NB-UVB (three lamps)	1.75 × 5	3.2 × 12.8 × 1.2	<ul style="list-style-type: none"> • Comb attachment for scalp treatment • Timer to monitor treatment length • Alert when energy from lamps gets low
Dermalume 2x (National Biological Corporation)	NB-UVB (two lamps)	3.3 × 5.4	4.1 × 7.5 × 2.8 (folded)	<ul style="list-style-type: none"> • Key lock • Timer that automatically turns lamps on/off • Emergency on/off switch
DermaPal (Daavlin)	NB-UVB UVA	8.25 × 1.75	2.75 × 9 × 5	<ul style="list-style-type: none"> • High-output lamps shorten treatment times • Highly accurate, integrated timer
DuaLight (Theralight) ^a	NB-UVB BB-UVB UVA	0.75 × 0.75	6.5 × 16.5 × 11.5	<ul style="list-style-type: none"> • Minimal erythema dose and minimal phototoxic dose phototest modes • Built-in timer and calibrating port • Square aperture handpiece provides unobstructed view • Flexible liquid light guide provided • Disposable tip for hygienic single-patient use prevents cross-contamination
Levia (Daavlin)	UVB	0.67 × 0.67	8 × 6 × 11.5	<ul style="list-style-type: none"> • Touch-screen control panel • Built-in calibration port • LiteSpot and LiteBrush attachments • Light administered in small squares that can be tiled to treat the entire surface
Psoria-Light (Psoria-Shield) ^a	NB-UVB, UVA	(0.45 square inches)	18 × 22 × 8.5	<ul style="list-style-type: none"> • LED (light emitting diode) source • Fast switching, on/off ability, not requiring a warm-up or cool-down period • High-definition digital camera to capture before and after photos of treatment sites • Touch technology to detect the patient's treatment site before enabling UV dosage administration
100 Series Handheld Phototherapy (SolarC Systems)	NB-UVB BB-UVB (two lamps)	2.5 × 5	3.5 × 7 × 2.25	<ul style="list-style-type: none"> • Includes a carrying case • Includes aperture plates for precise localized treatment • Positioning arm and UV brush • Acrylic panel covering lamps • Timer with maximum 20 minutes • Switch lock

^aOffice use only. BB-UVB, broadband ultraviolet B; NB-UVB, narrowband ultraviolet B; UV, ultraviolet; UVA, ultraviolet A; UVB, ultraviolet B.

Two concerns for home administration of phototherapy are patient adherence and the possibility of overuse, underuse, and inappropriate use²⁶. Fortunately, various features have been developed in recent years to increase safety of home phototherapy. Features such as an integrated timer and automated calibration can limit the length of time of treatment sessions and number of treatment sessions between follow-up visits with the physician. However, proper screening of candidates and patient education continue to be important to ensure safety and compliance²⁷. For the physician, an increasing number of resources are now available to increase knowledge and ease of prescribing home phototherapy units²³.

The excimer laser: a potential new indication and a novel dosimetry protocol

In addition to targeted therapy in the form of handheld devices, the 308 nm excimer laser was developed in 1997 as a targeted NB-UVB source for the treatment of psoriasis²⁸. The advantage to using the excimer laser is that because psoriasis plaques can take higher doses of light compared with normal skin, targeted treatment of psoriasis lesions using higher doses permits quicker time to clearance. In a multicenter open-label trial, 72% of patients with mild to moderate psoriasis achieved at least 75% improvement of the target plaque in an average of 6.2 treatments²⁹. Compared with traditional

whole-body phototherapy, the excimer laser required fewer visits²⁹. In another study, 13 out of 26 patients with plaque-type psoriasis had continued clearance or long-term improvement after 1 year³⁰.

These initial studies tested the excimer laser in patients with localized psoriasis and thus the excimer laser is currently US Food and Drug Administration indicated for the treatment of mild to moderate psoriasis. However, the excimer laser may also be safe and effective for use in generalized, moderate to severe psoriasis³¹. In a single-center pilot study, patients with greater than 10% but less than 30% body surface area involvement (moderate to severe psoriasis) were treated with excimer laser twice weekly for 12 weeks for a total of 24 treatment sessions. Fifty-four percent achieved PASI-75 and 83% achieved PASI-50, which was maintained without further treatment for 6 months³². However, the downside to treating moderate to severe psoriasis is the long duration required per session to treat a large body surface area, which may not be feasible in many dermatology office settings.

Another area of recent advancement for the excimer laser is in the dosimetry protocol. The current accepted starting dose is typically based on the patient's MED or Fitzpatrick skin type as well as location and thickness of the psoriatic lesion²⁸. The dose is increased as tolerated according to degree of erythema at each treatment session²⁸. A limitation to the current dosing method is that because the MED and Fitzpatrick skin type are based on normal skin, the starting dose is typically an underestimate. It typically takes 5 to 10 treatments of incremental dose increase just to reach the ideal maximum dose of the plaque. A recent study has attempted to develop a new dosing protocol called "plaque-based sub-blistering dosimetry", in which the patient's psoriatic plaque, rather than uninvolved skin, is tested with incrementally increasing doses to determine the dose at which blistering is observed³². The patients then are treated just below that dose. This allows plaques to be treated at the most aggressive possible dose (though still a tolerable one) to achieve faster clearing. The starting dose used in this protocol is 8 to 16 times the MED, compared with 2 to 4 times the MED of the currently accepted dosing protocol. The use of this "plaque-based sub-blistering dosimetry" can result in achievement of PASI-75 in a patient with moderate psoriasis after only two treatment sessions³².

Phototherapy and biologic agents: combination therapy for recalcitrant psoriasis

Although several biologic agents showing excellent efficacy in the treatment of moderate to severe psoriasis have been developed in the last decade, phototherapy appears to play an important role in a subset of patients with severe, recalcitrant psoriasis despite treatment with a biologic agent. Several studies have demonstrated the efficacy of using etanercept and NB-UVB in combination^{33–38}. These studies evaluated this combination therapy in patients who had not previously received treatment, patients who had an inadequate response with etanercept alone (50 mg once-weekly or 50 mg twice-weekly dosing), or patients who had an inadequate response to NB-UVB alone. Overall, combination therapy was superior and time to clearance was reduced. A study by Lynde *et al.* also demonstrated the importance of high adherence to the NB-UVB regimen in order to achieve significant clinical improvement³⁶. High adherence to the NB-UVB regimen was defined as missing not more than two treatments in any 4-week period.

One study to date has failed to establish efficacy of combination therapy of etanercept and NB-UVB. This head-to-head pilot study by Park *et al.*, who examined combination treatment with NB-UVB and etanercept 50 mg twice weekly compared with etanercept monotherapy, did not demonstrate significantly enhanced improvement with combination therapy³⁹. However, this study was limited by a small sample size of 13 patients. Furthermore, the patients all had a body mass index (BMI) of greater than 30, and studies have reported a suboptimal response to etanercept in psoriasis patients with a BMI of greater than 30^{40,41}.

Adalimumab or ustekinumab in combination with NB-UVB has also been investigated in a limited number of studies. A study by Bagel *et al.* evaluated the combination of adalimumab and NB-UVB⁴², and another study, by Wolf *et al.*, evaluated adalimumab and the excimer laser⁴³. Both studies demonstrated that phototherapy significantly accelerates therapeutic response and improves the clearance of psoriatic lesions in patients who received adalimumab treatment. One study to date has evaluated the combination of ustekinumab with the excimer laser⁴⁴. This was an intra-individual, half-body comparison study in which PASI-75 was achieved significantly more often on the UV-irradiated half than on the non-irradiated half at week 6 in patients on ustekinumab.

In general, combination therapy involving a biologic agent with NB-UVB phototherapy was very well tolerated. The most common side effect was erythema. Although long-term studies do not currently exist, no skin cancers were reported throughout the duration of the above-mentioned trials.

Conclusions

Although phototherapy is one of the oldest therapeutic modalities for psoriasis, it remains a mainstay treatment that holds promise for further advancement. Recent studies have delineated specific immunologic mediators that are affected by UV light. Such findings may lead to identification of targets for future psoriasis therapies.

Another area of recent advancement is in targeted phototherapy. Targeted phototherapy is advantageous in that it is a user-friendly tool that can irradiate difficult-to-treat areas as well as deliver higher doses of UV light compared with the traditional, whole-body phototherapy treatments. Currently, there are limited studies evaluating efficacy of handheld UVB use, and future studies should be conducted to evaluate long-term efficacy, cost effectiveness, and patient satisfaction. Interesting topics of future studies include the development of an effective dosimetry protocol for handheld UV devices that can be used safely both in the office setting and by patients at home.

Future areas of research for the excimer laser include large-scale, long-term studies evaluating its use for the treatment of moderate to severe psoriasis as a potential new indication. The evaluation of safety, efficacy, and practicality of the "plaque-based sub-blistering dosimetry" in a large-scale, long-term study is also of interest.

Lastly, NB-UVB appears to be a safe and effective treatment to be used concomitantly with biologic agents for recalcitrant psoriasis. Long-term, large-scale studies are needed to better understand safety of the use of this combination therapy, especially pertaining

to the risk of non-melanoma skin cancer. Similar studies using BB-UVB and PUVA are lacking and are of interest for future research. Cost effectiveness, patient satisfaction, and quality of life with this combination therapy should also be studied.

Abbreviations

BB-UVB, broadband ultraviolet B; BMI, body mass index; Foxp3, forkhead box P3; IL, interleukin; MED, minimal erythema dose; NB-UVB, narrowband ultraviolet B; PASI, Psoriasis Area and Severity Index; PUVA, psoralen plus ultraviolet A; Th1, T helper cell type 1; Th17, T helper cell type 17; Treg, regulatory T; UV, ultraviolet; UVA, ultraviolet A; UVB, ultraviolet B.

Competing interests

Tina Bhutani is an advisor for Cutanea (Wayne, PA, USA) and conducts research for AbbVie (North Chicago, IL, USA), Janssen (Beerse, Belgium), PhotoMedex (Horsham, PA, USA), Merck/Sun Pharmaceutical Industries (Kenilworth, NJ, USA), and Novartis (Basel, Switzerland). Mio Nakamura and Benjamin Farahnik declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Racz E, Prens EP: **Phototherapy and photochemotherapy for psoriasis.** *Dermatol Clin.* 2015; **33**(1): 79–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wan J, Abuabara K, Troxel AB, *et al.*: **Dermatologist preferences for first-line therapy of moderate to severe psoriasis in healthy adult patients.** *J Am Acad Dermatol.* 2012; **66**(3): 376–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Housman TS, Rohrbach JM, Fleischer AB Jr, *et al.*: **Phototherapy utilization for psoriasis is declining in the United States.** *J Am Acad Dermatol.* 2002; **46**(4): 557–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Danesh MJ, Butler DC, Beroukhi K, *et al.*: **A cross-sectional survey study to evaluate phototherapy training in dermatology residency.** *Photodermatol Photoimmunol Photomed.* 2015; **31**(5): 269–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Stern RS: **Psoralen and ultraviolet A light therapy for psoriasis.** *N Engl J Med.* 2007; **357**(7): 682–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hönigsmann H: **Phototherapy for psoriasis.** *Clin Exp Dermatol.* 2001; **26**(4): 343–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Almutawa F, Alnomair N, Wang Y, *et al.*: **Systematic review of UV-based therapy for psoriasis.** *Am J Clin Dermatol.* 2013; **14**(2): 87–109.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen X, Yang M, Cheng Y, *et al.*: **Narrow-band ultraviolet B phototherapy versus broad-band ultraviolet B or psoralen-ultraviolet A photochemotherapy for psoriasis.** *Cochrane Database Syst Rev.* 2013; (10): CD009481.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Wang E, Sasaki J, Nakamura M, *et al.*: **Cutaneous carcinogenic risk of phototherapy: an updated comprehensive review.** *J Psoriasis and Psoriatic Arth.* 2015 winter; **1**(1): 44–51.
[Reference Source](#)
- Rajpara AN, O'Neill JL, Nolan BV, *et al.*: **Review of home phototherapy.** *Dermatol Online J.* 2010; **16**(12): 2.
[PubMed Abstract](#)
- Larkö O, Swanbeck G: **Home solarium treatment of psoriasis.** *Br J Dermatol.* 1979; **101**(1): 13–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Koek MB, Buskens E, van Weelden H, *et al.*: **Home versus outpatient ultraviolet B phototherapy for mild to severe psoriasis: pragmatic multicentre randomised controlled non-inferiority trial (PLUTO study).** *BMJ.* 2009; **338**: b1542.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Wong T, Hsu L, Liao W: **Phototherapy in psoriasis: a review of mechanisms of action.** *J Cutan Med Surg.* 2013; **17**(1): 6–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Weatherhead SC, Farr PM, Jamieson D, *et al.*: **Keratinocyte apoptosis in epidermal remodeling and clearance of psoriasis induced by UV radiation.** *J Invest Dermatol.* 2011; **131**(9): 1916–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- DeSilva B, McKenzie RC, Hunter JA, *et al.*: **Local effects of TL01 phototherapy in psoriasis.** *Photodermatol Photoimmunol Photomed.* 2008; **24**(5): 268–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Danno K, Toda K, Horio T: **Ultraviolet-B radiation suppresses mast cell degranulation induced by compound 48/80.** *J Invest Dermatol.* 1986; **87**(6): 775–8.
[PubMed Abstract](#)
- Enk CD, Sredni D, Blauvelt A, *et al.*: **Induction of IL-10 gene expression in human keratinocytes by UVB exposure in vivo and in vitro.** *J Immunol.* 1995; **154**(9): 4851–6.
[PubMed Abstract](#)
- Johnson-Huang LM, Suárez-Fariñas M, Sullivan-Whalen M, *et al.*: **Effective narrow-band UVB radiation therapy suppresses the IL-23/IL-17 axis in normalized psoriasis plaques.** *J Invest Dermatol.* 2010; **130**(11): 2654–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Coimbra S, Oliveira H, Reis F, *et al.*: **Interleukin (IL)-22, IL-17, IL-23, IL-8, vascular endothelial growth factor and tumour necrosis factor-α levels in patients with psoriasis before, during and after psoralen-ultraviolet A and narrowband ultraviolet B therapy.** *Br J Dermatol.* 2010; **163**(6): 1282–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bovenschen HJ, van de Kerkhof PC, van Erp PE, *et al.*: **Foxp3+ regulatory T cells of psoriasis patients easily differentiate into IL-17A-producing cells and are found in lesional skin.** *J Invest Dermatol.* 2011; **131**(9): 1853–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Zhang D, Chen Y, Chen L, *et al.*: **Ultraviolet Irradiation Promotes FOXP3 Transcription via p53 in Psoriasis.** *Exp Dermatol.* 2016; **25**(7): 513–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Mudigonda T, Dabade TS, Feldman SR: **A review of targeted ultraviolet B phototherapy for psoriasis.** *J Am Acad Dermatol.* 2012; **66**(4): 664–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Anderson KL, Feldman SR: **A guide to prescribing home phototherapy for patients with psoriasis: the appropriate patient, the type of unit, the treatment regimen, and the potential obstacles.** *J Am Acad Dermatol.* 2015; **72**(5): 868–78.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Caccialanza M, Piccinno R, Cappio F, *et al.*: **[Phototherapy of psoriasis of the scalp. Results in 21 patients treated with a special portable ultraviolet rays lamp].** *G Ital Dermatol Venereol.* 1989; **124**(11–12): LXI–LXV.
[PubMed Abstract](#)
- Dotterud LK, Braun R: **[UV-B comb versus betamethasone solution in scalp psoriasis].** *Tidsskr Nor Lægeforen.* 2000; **120**(16): 1858–9.
[PubMed Abstract](#)
- Yarbrough C, Yentzer BA, Yelverton CB, *et al.*: **Continued use of home narrowband ultraviolet B light phototherapy for psoriasis after completion of a clinical trial.** *J Am Acad Dermatol.* 2009; **60**(5): 877–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Feldman SR, Clark A, Reboussin DM, *et al.*: **An assessment of potential problems of home phototherapy treatment of psoriasis.** *Cutis.* 1996; **58**(1): 71–3.
[PubMed Abstract](#)
- Mehta D, Lim HW: **Ultraviolet B Phototherapy for Psoriasis: Review of Practical Guidelines.** *Am J Clin Dermatol.* 2016; **17**(2): 125–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Feldman SR, Mellen BG, Housman TS, *et al.*: **Efficacy of the 308-nm excimer**

- laser for treatment of psoriasis: results of a multicenter study. *J Am Acad Dermatol.* 2002; **46**(6): 900–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Fikrie T, Pizinger K: [The use of the 308 nm excimer laser for the treatment of psoriasis]. *J Dtsch Dermatol Ges.* 2003; **1**(7): 559–63.
[PubMed Abstract](#)
 31. Gattu S, Pang ML, Pugashetti R, *et al.*: Pilot evaluation of supra-erythemogenic phototherapy with excimer laser in the treatment of patients with moderate to severe plaque psoriasis. *J Dermatolog Treat.* 2010; **21**(1): 54–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. **F** Debbaneh MG, Levin E, Sanchez Rodriguez R, *et al.*: Plaque-based sub-blistering dosimetry: Reaching PASI-75 after two treatments with 308-nm excimer laser in a generalized psoriasis patient. *J Dermatolog Treat.* 2015; **26**(1): 45–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 33. De Simone C, D'Agostino M, Capizzi R, *et al.*: Combined treatment with etanercept 50 mg once weekly and narrow-band ultraviolet B phototherapy in chronic plaque psoriasis. *Eur J Dermatol.* 2011; **21**(4): 568–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Kircik L, Bagel J, Korman N, *et al.*: Utilization of narrow-band ultraviolet light B therapy and etanercept for the treatment of psoriasis (UNITE): efficacy, safety, and patient-reported outcomes. *J Drugs Dermatol.* 2008; **7**(3): 245–53.
[PubMed Abstract](#)
 35. Wolf P, Hofer A, Legat FJ, *et al.*: Treatment with 311-nm ultraviolet B accelerates and improves the clearance of psoriatic lesions in patients treated with etanercept. *Br J Dermatol.* 2009; **160**(1): 186–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. **F** Lynde CW, Gupta AK, Guenther L, *et al.*: A randomized study comparing the combination of nbUVB and etanercept to etanercept monotherapy in patients with psoriasis who do not exhibit an excellent response after 12 weeks of etanercept. *J Dermatolog Treat.* 2012; **23**(4): 261–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 37. Gambichler T, Tigges C, Scola N, *et al.*: Etanercept plus narrowband ultraviolet B phototherapy of psoriasis is more effective than etanercept monotherapy at 6 weeks. *Br J Dermatol.* 2011; **164**(6): 1383–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. **F** Calzavara-Pinton PG, Sala R, Arisi M, *et al.*: Synergism between narrowband ultraviolet B phototherapy and etanercept for the treatment of plaque-type psoriasis. *Br J Dermatol.* 2013; **169**(1): 130–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 39. **F** Park KK, Wu JJ, Koo J: A randomized, 'head-to-head' pilot study comparing the effects of etanercept monotherapy vs. etanercept and narrowband ultraviolet B (NB-UVB) phototherapy in obese psoriasis patients. *J Eur Acad Dermatol Venereol.* 2013; **27**(7): 899–906.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 40. Gordon K, Korman N, Frankel E, *et al.*: Efficacy of etanercept in an integrated multistudy database of patients with psoriasis. *J Am Acad Dermatol.* 2006; **54**(3 Suppl 2): S101–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Strober B, Gottlieb A, Leonardi C, *et al.*: Levels of response of psoriasis patients with different baseline characteristics treated with etanercept. *J Am Acad Dermatol.* 2006; **54**(3 Suppl): AB220.
[Reference Source](#)
 42. **F** Bagel J: Adalimumab plus narrowband ultraviolet B light phototherapy for the treatment of moderate to severe psoriasis. *J Drugs Dermatol.* 2011; **10**(4): 366–71.
[PubMed Abstract](#) | [F1000 Recommendation](#)
 43. **F** Wolf P, Hofer A, Weger W, *et al.*: 311 nm ultraviolet B-accelerated response of psoriatic lesions in adalimumab-treated patients. *Photodermatol Photoimmunol Photomed.* 2011; **27**(4): 186–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 44. **F** Wolf P, Weger W, Legat FJ, *et al.*: Treatment with 311-nm ultraviolet B enhanced response of psoriatic lesions in ustekinumab-treated patients: a randomized intraindividual trial. *Br J Dermatol.* 2012; **166**(1): 147–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Steven R Feldman

Department of Dermatology, Wake Forest School of Medicine, Winston-Salem, NC, USA

Competing Interests: No competing interests were disclosed.

2. Joel Gelfand

Dermatology and Epidemiology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104, USA

Competing Interests: In the previous 12 months, Dr. Gelfand served as a consultant for Abbvie., Astrazeneca, Celgene Corp, Coherus, Eli Lilly, Janssen Biologics (formerly Centocor), Sanofi, Merck, Novartis Corp, Valeant, and Pfizer Inc., receiving honoraria; and receives research grants (to the Trustees of the University of Pennsylvania) from Abbvie, Amgen, Eli Lilly, Janssen, Novartis Corp, Regeneron, and Pfizer Inc.; and received payment for continuing medical education work related to psoriasis. Dr. Gelfand is a co-patent holder of resiquimod for treatment of cutaneous T cell lymphoma.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research