Check for updates

OPINION ARTICLE

A new hypothesis: some metastases are the result of inflammatory processes by adapted cells, especially adapted immune cells at sites of inflammation [version 1; peer review: 3 approved]

Leili Shahriyari

Mathematical Biosciences Institute, The Ohio State University, Columbus, OH, USA

V1 First published: 16 Feb 2016, 5:175
<https://doi.org/10.12688/f1000research.8055.1>

Latest published: 16 Feb 2016, 5:175
<https://doi.org/10.12688/f1000research.8055.1>

Abstract

There is an old hypothesis that metastasis is the result of migration of tumor cells from the tumor to a distant site. In this article, we propose another mechanism for metastasis, for cancers that are initiated at the site of chronic inflammation. We suggest that cells at the site of chronic inflammation might become adapted to the inflammatory process, and these adaptations may lead to the initiation of an inflammatory tumor. For example, in an inflammatory tumor immune cells might be adapted to send signals of proliferation or angiogenesis, and epithelial cells might be adapted to proliferation (like inactivation of tumor suppressor genes). Therefore, we hypothesize that metastasis could be the result of an inflammatory process by adapted cells, especially adapted immune cells at the site of inflammation, as well as the migration of tumor cells with the help of activated platelets, which travel between sites of inflammation. If this hypothesis is correct, then any treatment causing necrotic cell death may not be a good solution. Because necrotic cells in the tumor micro-environment or anywhere in the body activate the immune system to initiate the inflammatory process, and the involvement of adapted immune cells in the inflammatory processes leads to the formation and progression of tumors. Adapted activated immune cells send more signals of proliferation and/or angiogenesis than normal cells. Moreover, if there were adapted epithelial cells, they would divide at a much higher rate in response to the proliferation signals than normal cells. Thus, not only would the tumor come back after the treatment, but it would also grow more aggressively.

Keywords

Metastasis, Cancer, Chronic inflammation, Adapted immune cells, Inflammatory processes, Immune system, Platelets, Wound healing process.

Open Peer Review

Approval Status

	1	2	3
version 1			
16 Feb 2016	view	view	view

1. **Maryam Keshtkar Jahromi**, Johns Hopkins University School of Medicine, Baltimore, USA
2. **Patrice Forget**, Université Catholique de Louvain, Brussels, Belgium
3. **Alain Borgeat**, Orthopedic University Hospital Balgrist, Zurich, Switzerland

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Leili Shahriyari (shahriyari.1@mbi.osu.edu)

Competing interests: No competing interests were disclosed.

Grant information: This research has been supported in part by the Mathematical Biosciences Institute and the National Science Foundation under grant DMS 0931642.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Shahriyari L. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Shahriyari L. **A new hypothesis: some metastases are the result of inflammatory processes by adapted cells, especially adapted immune cells at sites of inflammation [version 1; peer review: 3 approved]** F1000Research 2016, 5:175 <https://doi.org/10.12688/f1000research.8055.1>

First published: 16 Feb 2016, 5:175 <https://doi.org/10.12688/f1000research.8055.1>

Many cancers arise from sites of chronic inflammation¹. Immune cells inside the chronic inflammation site initiate tumor progression by releasing reactive oxygen or nitrogen species, which lead to DNA damage in epithelial cells². Inflammation not only can cause mutation in epithelial cells³, but can also change their fitness⁴.

In chronic inflammation, T-cells might become adapted to send high levels of proliferation signals, and regulatory T-cells might have been changed to prevent their inhibition⁵. Effector T-cells also create an environment for tumor initiation and progression by releasing tumor-promoting cytokines IL-6².

These findings suggest that cells at the site of chronic inflammation are adapted to the wound healing process. Immune cells are adapted to send signals of proliferation or angiogenesis, and tissue cells are adapted to proliferation (like inactivation of tumor suppressor genes). These adaptations lead to the initiation of a tumor.

If there are adapted immune cells, then we can look at metastasis from a new perspective. Any site of inflammation might recruit adapted immune cells, and then they contribute to the inflammatory process there. If immune cells are adapted to send more signals of proliferation and angiogenesis, then new tumors would initiate at sites of inflammation. Below we gather some evidence that supports this hypothesis.

- There is a hypothesis that metastasis is the result of cancer cell migration from the tumor extracellular matrix to the bloodstream or lymphatic vessels as circulating tumor cells (CTCs), and then to the new site. CTCs only use lymphatic routes to migrate to the nearby lymph nodes, but not for traveling long distances⁶. Cell migration occurs when tumor epithelial (E) cells lose their cell-cell adhesion and become motile mesenchymal (M) cells, i.e. epithelial-mesenchymal transition (EMT). When these CTCs exit the bloodstream, they undergo a reverse process called mesenchymal-epithelial transition (MET) to continue their differentiation and develop a secondary tumor⁷⁻⁹.
- In inflammatory breast cancer, there is a correlation between immune activation and CTCs with EMT characteristics¹⁰. Cohen *et al.* hypothesized that EMT could be the result of inflammatory processes initiated by activated immune cells¹⁰. Epithelial cells from the colonic epithelium of patients with benign colon diseases also circulate in the blood¹¹. During wound healing, epithelial cells migrate and disperse as individual mesenchymal cells by down regulating cell-cell junctions¹². Thus, CTCs in blood could be the result of inflammation.
- By querying published available data sets¹³⁻¹⁷, we calculate the probability of not detecting any CTCs in blood from patients with metastatic breast cancer, and the result is 0.6. That means there might be other phenomena, beside CTCs, that cause metastasis. Since, no CTCs were detected in the blood of 29% of metastatic breast cancer patients starting a first or new line of therapy, it is unlikely that treatments are responsible for not observing CTCs in blood¹⁶.
- CTC-clusters are rare compared with single CTCs, however CTC-clusters, which usually include platelets and white blood cells¹⁸, significantly increase metastatic potential¹⁹. Moreover inhibition of platelet activation or platelet depletion decreases metastasis rates²⁰⁻²². There is a hypothesis that metastasis is reduced when the host is platelet depleted because platelets are protecting tumor cells from natural killing cells^{23,24}. Platelets in tumor cell clusters release transforming growth factor β (TGF- β)²⁵, and tumor cells that are bound to platelets highly express EMT-associated genes²⁶.
- Cancer of unknown primary origin (CUP) is defined by the presence of metastatic disease with no identified primary tumor. CUPs are 3.5% of all human malignancies, and one of the ten most frequent cancers worldwide²⁷. The major sites of CUP are bones, liver, lung, and lymph nodes²⁸, the sites where immune cells are most active.
- New studies show that recruited bone marrow progenitor cells change the host's environment to generate the "pre-metastatic niche" to which the tumor cells metastasize²⁹. Bone marrow-derived haematopoietic progenitor cells (HPCs) that express vascular endothelial growth factor receptor 1 form cellular clusters at pre-metastatic sites before appearance of a single tumor cell³⁰.
- There is evidence of metastasis to the site of injury. Two patients with squamous cell carcinoma of the lung developed distant localized metastatic disease at sites of physical injuries; one to the knee injured in an accidental fall six weeks earlier, and the other to portions of the liver injured in a mechanical fall two months earlier³¹. In a mice model of metastatic breast cancer, radiation-induced pulmonary injury lead to chronic inflammatory responses, and development of pre-metastatic niches³². In another mice model, hepatic ischemia-reperfusion injury increased the number of liver metastases of human pancreatic cancer (Capan-1) cells, which were injected into the mice spleen³³. Several studies show that lung injury induced by the chemotherapy drug, bleomycin, increases lung metastases; they also observed tumor cell adherence to extracellular matrix and fibrin at injured areas^{34,35}. Therefore we suggest that the sites of injuries are potential metastatic sites.

We hypothesize that chronic inflammation can cause adapted bone marrow derived cells (for example, adapted macrophages or T-cells) and/or adapted tissue cells (for instance, adapted epithelial cells or stromal fibroblasts) to lead tumor initiation and progression. If adapted immune cells are present, then the new site of inflammation might recruit these adapted immune cells and cause metastasis. Additionally, the new site of inflammation may recruit activated platelets. The activated platelets travel between sites of inflammation, including the site of inflammatory carcinoma. Tumor cells can link to adhesion receptors on platelets and travel to the new site of inflammation. The activated platelets start the wound healing process at the new site, which now includes some tumor cells. As the tumor cells respond to the wound healing signals more strongly than normal cells, new tumors would initiate in the site of inflammation (Figure 1).

Figure 1. Metastasis. Chronic inflammation leads to adapted tissue and/or adapted immune cells. These adaptations cause tumor initiation. A new inflammation site recruits these adapted immune cells. The adapted immune cells start the wound healing process at the inflammation site and send more inflammatory signals than normal immune cells. Thus, a new tumor initiates there. Also, the activated platelets travel between sites of inflammation, including the site of primary inflammatory tumor. Tumor cells can link to adhesion receptors on platelets and travel to the new site of inflammation with the help of platelets. The activated platelets start the inflammatory process at the new site, which now includes some tumor cells. Tumor cells respond to the inflammatory signals more strongly than normal cells. The transported tumor cells initiate a tumor at the inflammation site.

Competing interests

No competing interests were disclosed.

Grant information

This research has been supported in part by the Mathematical

Biosciences Institute and the National Science Foundation under grant DMS 0931642.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Balkwill F, Charles KA, Mantovani A: **Smoldering and polarized inflammation in the initiation and promotion of malignant disease.** *Cancer Cell.* 2005; **7**(3): 211–217.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Waldner MJ, Neurath MF: **Colitis-associated cancer: the role of T cells in tumor development.** *Semin Immunopathol.* 2009; **31**(2): 249–256.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hussain SP, Amstad P, Raja K, et al.: **Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease.** *Cancer Res.* 2000; **60**(13): 3333–3337.
[PubMed Abstract](#)
- Vermeulen L, Morrissey E, van der Heijden M, et al.: **Defining Stem Cell Dynamics in Models of Intestinal Tumor Initiation.** *Science.* 2013; **342**(6161): 995–998.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Erdman SE, Poutahidis T: **Roles for inflammation and regulatory T cells in colon cancer.** *Toxicol Pathol.* 2010; **38**(1): 76–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chaffer CL, Weinberg RA: **A perspective on cancer cell metastasis.** *Science.* 2011; **331**(6024): 1559–1564.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tsai JH, Yang J: **Epithelial-mesenchymal plasticity in carcinoma metastasis.** *Genes Dev.* 2013; **27**(20): 2192–2206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tam WL, Weinberg RA: **The epigenetics of epithelial-mesenchymal plasticity in cancer.** *Nat Med.* 2013; **19**(11): 1438–1449.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lu M, Jolly MK, Onuchic J, et al.: **Toward decoding the principles of cancer metastasis circuits.** *Cancer Res.* 2014; **74**(17): 4574–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cohen EN, Gao H, Anfossi S, et al.: **Inflammation Mediated Metastasis: Immune Induced Epithelial-To-Mesenchymal Transition in Inflammatory Breast Cancer Cells.** *PLoS One.* 2015; **10**(7): e0132710.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pantel K, Denève E, Nocca D, et al.: **Circulating epithelial cells in patients with benign colon diseases.** *Clin Chem.* 2012; **58**(5): 936–940.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zvaifler NJ: **Relevance of the stroma and epithelial-mesenchymal transition (EMT) for the rheumatic diseases.** *Arthritis Res Ther.* 2006; **8**(3): 210.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pierga JY, Bonneton C, Vincent-Salomon A, et al.: **Clinical significance of immunocytochemical detection of tumor cells using digital microscopy in peripheral blood and bone marrow of breast cancer patients.** *Clin Cancer Res.* 2004; **10**(4): 1392–1400.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Riethdorf S, Fritsche H, Müller V, et al.: **Detection of circulating tumor cells in peripheral blood of patients with metastatic breast cancer: a validation study of the CellSearch system.** *Clin Cancer Res.* 2007; **13**(3): 920–928.
[PubMed Abstract](#) | [Publisher Full Text](#)

15. Cristofanilli M, Budd GT, Ellis MJ, *et al.*: **Circulating tumor cells, disease progression, and survival in metastatic breast cancer.** *N Engl J Med.* 2004; **351**(8): 781–791.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Allard WJ, Matera J, Miller MC, *et al.*: **Tumor cells circulate in the peripheral blood of all major carcinomas but not in healthy subjects or patients with nonmalignant diseases.** *Clin Cancer Res.* 2004; **10**(20): 6897–6904.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Racila E, Euhus D, Weiss AJ, *et al.*: **Detection and characterization of carcinoma cells in the blood.** *Proc Natl Acad Sci U S A.* 1998; **95**(8): 4589–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Mitrugno A, Tormoen GW, Kuhn P, *et al.*: **The prothrombotic activity of cancer cells in the circulation.** *Blood Rev.* 2015; pii: S0268-960X(15)00056-9.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Aceto N, Bardia A, Miyamoto DT, *et al.*: **Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis.** *Cell.* 2014; **158**(5): 1110–1122.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Tzanakakis GN, Agarwal KC, Veronikis DK, *et al.*: **Effects of antiplatelet agents alone or in combinations on platelet aggregation and on liver metastases from a human pancreatic adenocarcinoma in the nude mouse.** *J Surg Oncol.* 1991; **48**(1): 45–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Amirkhosravi A, Mousa SA, Amaya M, *et al.*: **Inhibition of tumor cell-induced platelet aggregation and lung metastasis by the oral GpIb/IIa antagonist XV454.** *Thromb Haemost.* 2003; **90**(3): 549–554.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Palumbo JS, Talmage KE, Massari JV, *et al.*: **Platelets and fibrin(ogen) increase metastatic potential by impeding natural killer cell-mediated elimination of tumor cells.** *Blood.* 2005; **105**(1): 178–185.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Palumbo JS: **Mechanisms linking tumor cell-associated procoagulant function to tumor dissemination.** *Semin Thromb Hemost.* 2008; **34**(2): 154–160.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Nieswandt B, Hafner M, Echtenacher B, *et al.*: **Lysis of tumor cells by natural killer cells in mice is impeded by platelets.** *Cancer Res.* 1999; **59**(6): 1295–1300.
[PubMed Abstract](#)
25. Yu M, Bardia A, Wittner BS, *et al.*: **Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition.** *Science.* 2013; **339**(6119): 580–584.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Labelle M, Begum S, Hynes RO: **Direct signaling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis.** *Cancer Cell.* 2011; **20**(5): 576–590.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Pavlidis N, Fizazi K: **Carcinoma of unknown primary (CUP).** *Crit Rev Oncol Hematol.* 2009; **69**(3): 271–278.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Hainsworth JD, Greco FA: **Cancer of Unknown Primary Origin.** *Goldman's Cecil Medicine: Twenty Fourth Edition.* 2012; **1**: 1334–1337.
[Publisher Full Text](#)
29. Peinado H, Lavotshkin S, Lyden D: **The secreted factors responsible for pre-metastatic niche formation: old sayings and new thoughts.** *Semin Cancer Biol.* 2011; **21**(2): 139–146.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Kaplan RN, Riba RD, Zacharoulis S, *et al.*: **VEGFR1-positive haematopoietic bone marrow progenitors initiate the pre-metastatic niche.** *Nature.* 2005; **438**(7069): 820–827.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Walter ND, Rice PL, Redente EF, *et al.*: **Wound healing after trauma may predispose to lung cancer metastasis: review of potential mechanisms.** *Am J Respir Cell Mol Biol.* 2010; **44**(5): 591–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Gong HY, Hu WG, Hu QY, *et al.*: **Radiation-induced pulmonary injury accelerated pulmonary metastasis in a mouse model of breast cancer.** *Oncol Lett.* 2015; **10**(6): 3613–3618.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Yoshimoto K, Tajima H, Ohta T, *et al.*: **Increased E-selectin in hepatic ischemia-reperfusion injury mediates liver metastasis of pancreatic cancer.** *Oncol Rep.* 2012; **28**(3): 791–796.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Orr FW, Adamson IY, Young L: **Promotion of pulmonary metastasis in mice by bleomycin-induced endothelial injury.** *Cancer Res.* 1986; **46**(2): 891–7.
[PubMed Abstract](#)
35. Adamson IY, Orr FW, Young L: **Effects of injury and repair of the pulmonary endothelium on lung metastasis after bleomycin.** *J Pathol.* 1986; **150**(4): 279–87.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 27 April 2016

<https://doi.org/10.5256/f1000research.8666.r13490>

© 2016 Borgeat A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alain Borgeat

Orthopedic University Hospital Balgrist, Zurich, Switzerland

There is more and more evidence that cancer metastases and inflammation share a common pathway. Therefore this hypothesis has a relevant scientific background. Prospective clinical trials looking at this issue will be welcome.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 22 April 2016

<https://doi.org/10.5256/f1000research.8666.r13492>

© 2016 Forget P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Patrice Forget

Department of Anesthesiology, Université Catholique de Louvain, Brussels, Belgium

The inflammatory nature of cancer, and especially metastases development, becomes a paradigm shift. This text summarizes well some new hypotheses and future challenges. Next steps to translate this in clinical trials are then needed.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 18 April 2016

<https://doi.org/10.5256/f1000research.8666.r13279>

© 2016 Keshtkar Jahromi M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Maryam Keshtkar Jahromi

Division of Infectious Diseases, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA

This hypothesis has a potential to make a positive impact in cancer research and drug discovery. For cancer patients, it might help to accelerate both the development of promising therapies and the potential of personalized medicine — treatments based on an individual's unique immune deficiency. This is a very interesting and unique idea which needs more research.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research