REVIEW

Advances in gene therapy for muscular dystrophies [version 1; peer review: 2 approved]

Hayder Abdul-Razak, Alberto Malerba, George Dickson

School of Biological Sciences, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK

V1 First published: 18 Aug 2016, 5(F1000 Faculty Rev):2030
<https://doi.org/10.12688/f1000research.8735.1>

Latest published: 18 Aug 2016, 5(F1000 Faculty Rev):2030
<https://doi.org/10.12688/f1000research.8735.1>

Abstract

Duchenne muscular dystrophy (DMD) is a recessive lethal inherited muscular dystrophy caused by mutations in the gene encoding dystrophin, a protein required for muscle fibre integrity. So far, many approaches have been tested from the traditional gene addition to newer advanced approaches based on manipulation of the cellular machinery either at the gene transcription, mRNA processing or translation levels. Unfortunately, despite all these efforts, no efficient treatments for DMD are currently available. In this review, we highlight the most advanced therapeutic strategies under investigation as potential DMD treatments.

Open Peer Review

Approval Status

	1	2
version 1 18 Aug 2016

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Giulio Cossu**, University of Manchester, Manchester, UK
2. **Shin'ichi Takeda**, National Center of Neurology and Psychiatry, Tokyo, Japan

Any comments on the article can be found at the end of the article.

Corresponding author: George Dickson (g.dickson@rhul.ac.uk)

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Abdul-Razak H *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Abdul-Razak H, Malerba A and Dickson G. **Advances in gene therapy for muscular dystrophies [version 1; peer review: 2 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2030 <https://doi.org/10.12688/f1000research.8735.1>

First published: 18 Aug 2016, 5(F1000 Faculty Rev):2030 <https://doi.org/10.12688/f1000research.8735.1>

Introduction

Duchenne muscular dystrophy (DMD) is a rare, severe, degenerative X-linked myopathy caused by mutations in the gene (*DMD*) encoding the dystrophin protein. DMD affects about 1 in 3500 to 4000 boys globally^{1,2}, and one-third of cases is attributed to spontaneous new mutations³. Dystrophin plays a key role in joining the actin of the cytoplasmic cytoskeleton to the extracellular matrix surrounding the sarcolemma of muscle fibres directly binding a protein complex known as dystrophin-associated glycoprotein complex (DGC)⁴. Several proteins like alpha-dystrobrevin, syncoilin, synemin, sarcoglycan, dystroglycan, and sarcospan are located at the DGC, where they mediate the muscle-networking signals that are essential for both muscle cell function and the development and maintenance of membrane integrity^{2,5}. Mutations in the *DMD* gene usually cause partial or complete absence of dystrophin, leading to membrane instability and muscle cell death associated with progressive tissue degeneration, muscle weakness, joint contractures, and kyphoscoliosis^{6,7}. Both skeletal and cardiac muscles are affected in patients with DMD. In particular, the loss of dystrophin leads to respiratory failure⁸ because of the severely damaged diaphragm and to cardiomyopathy and heart failure⁹, which usually result in premature death by the age of 25 to 35^{10,11}, although some long-term survivors have been observed¹².

Despite more than 20 years of research and the clear understanding of the molecular basis of the disease, only limited advancement of therapeutic approaches has been obtained, and currently the only available treatment is a combination of physiotherapy and corticosteroids. Although these palliative treatments can provide a considerable improvement in affected boys, they can only slow the course of the disorder. In this article, we will discuss the relevance of three approaches for the treatment of DMD: the gene addition approach by triple trans-splicing (TTS), the use of gene editing to correct the *DMD* transcript, and the exon-skipping approach to re-frame the faulty *DMD* pre-mRNA. All of these methods are promising DMD treatments that tackle the cause of the disease and offer the potential to treat many *DMD* mutations.

Facing DMD challenges

The *DMD* gene is expressed mainly in skeletal and cardiac muscle. About 65% of the mutations affecting dystrophin are deletions of part of the gene with two predominant hotspots within the *DMD* sequence. Two types of mutations are associated with two allelic forms of the disease: the first leads to the expression of out-of-frame mRNA that abolishes dystrophin production and causes the most severe form-of-the-disease, the DMD. The second type of mutation induces the expression of an in-frame mRNA, resulting in a milder form of DMD known as Becker muscular dystrophy (BMD), in which a shorter but still partially functional dystrophin is produced¹³.

There are currently many pharmacological approaches that can hinder DMD symptoms by focusing on secondary effects, yet they would treat just one aspect of DMD pathogenesis and may be associated with possible side effects¹⁴. For a better understanding of the different potential therapies available for DMD, we firstly have to highlight which challenges they face. In this regard, one hurdle is that DMD is frequently caused by new mutations, meaning

that many patients have no family history¹⁵. Furthermore, more than 60% of *DMD* mutations are due to intragenic deletions of one or more exons, resulting in disruption of the correct *DMD* open reading frame (ORF)¹⁶. Thus, DMD mutations can vary in severity and the phenotype is often unpredictable.

Therefore, the most urgent interest is to develop a genetic strategy that can provide a treatment for all patients with DMD. The other crucial point is that the ideal DMD therapy should be sustainable and lifelong. Previous observations in patients with X-linked myopathy suggest that in order to prevent muscle weakness, at least 30% of dystrophin must be expressed in skeletal muscle¹⁷. Even if this may seem a low level compared with the one in normal tissues, a crucial issue is that muscles of boys with DMD are substantially wasted and present a significant amount of fat and connective tissue that makes the remaining muscle tissue hardly accessible by the therapeutic agents delivered to the bloodstream. Furthermore, although most therapeutic agents have proven to be successful in small animal models (for example, rodents), human muscles are significantly larger and this requires scaling up the therapeutic agents' manufacturing process, which usually is related to logistic and economical challenges. Nowadays, gene therapy has emerged as a promising applicable strategy, as DMD therapy can cure the genetic defect and not just its downstream effects.

DMD gene addition by trans-splicing

The vast majority of gene therapy-based clinical trials for other disorders are based on the gene addition "replacement" approach¹⁸. For DMD gene replacement, a point of major concern is the size of the very long *DMD* gene (exonic DNA >11 kb). Hence, a partially functional, intact, and usually shortened *DMD* copy is delivered into muscle cells in order to mimic the beneficial effect of a smaller but partially functional protein expressed in patients with BMD. To achieve that, many delivery systems have been studied for DMD gene replacement, yet so far the most efficient is the use of adeno-associated virus (AAV)^{19–21}. The main issue with AAV applications is that their limited packaging capacity of approximately 4.7 kb requires the delivery of substantially internally deleted (truncated) dystrophin expression cassettes. Additionally, when AAVs are used in large quantities, their capsid proteins could potentially activate an immunogenic response; consequently, strategies to attenuate the immune system are possibly needed for a clinically successful AAV approach³.

In preclinical tests, single AAV vectors were used to successfully deliver microdystrophins (<5 kb)—carrying about 30% of the coding sequence, both locally and systematically—and were associated with good improvement of muscle function^{22–25}. Nevertheless, some domains of dystrophin, like the critical parts of rod-and-hinge domains, should be included to increase dystrophin functionality and stabilise the membrane as they contribute to the recruitment of other components of DGC, such as neuronal nitric oxide synthase, syntrophin, and dystrobrevin^{26,27}, on the sarcolemma. In order to expand AAV packaging capacity and improve muscle functionality, trans-splicing dual AAV vectors were developed about 15 years ago to deliver less than 10 kb minidystrophin^{28–30}. In this approach, a large gene is divided, then packaged and delivered by two AAVs and ultimately expressed by dual trans-splicing. For this system to

work, the first vector (the 5' *DMD* part) is tail-tagged with splicing donor signal while the second vector (the 3' *DMD* part) provides a preceding splicing acceptor signal. By co-transduction of the two vectors, the AAV head-to-tail vector genome concatemerisation^{31,32} and the removal of viral sequences from the transcribed mRNA by the cellular splicing machinery allow the expression of the entire minidystrophin protein³³. In two different studies, the localised injections of trans-splicing dual AAVs into a dystrophic muscle of a mouse model of DMD have shown successful minidystrophin expression in myofibres of a single muscle^{31,34}. In another *in vivo* study, systemic injection led to high-level transduction within skeletal and cardiac muscles³². Another possible strategy is based on dual protein trans-splicing that was applied in a mouse model leading to therapeutic gene expression and improving dystrophic muscle morphology and histology³⁵. Although the proof of principle for this approach was successfully demonstrated and muscle functions were slightly improved, transduction efficiency was too low to achieve the therapeutic level for a functional protein as minidystrophins still lack some potentially essential domains in the final dystrophin conformation.

This strategy may be improved by developing a system to deliver the full-length exonic *DMD* sequence and express an entirely functional dystrophin. In this regard, recent research has examined the possibility of dividing the whole native exonic *DMD* (~11 kb) into three AAV vectors, creating the Triple transplating-AAV (TTS-AAV) system by using different inverted terminal repeats within the three vectors to favour the formation of the correct head-to-tail concatemers³⁶. The proof of concept was demonstrated in the DMD mouse model, showing the expression of a full-length dystrophin protein co-localising with the expression of a tagged protein (enhanced green fluorescent protein, or eGFP), whose coding sequence was at the 3' end of the gene³⁶. However, the efficiency of the TTS-AAV system was very low. Only one out of four treated mice showed evidence of splicing between the three vectors. Further optimisation of this approach is essential to overcome the low efficient expression by, for example, improving splicing elements, maximising gene expression, and directing the AAV vector concatemerisation more efficiently. The TTS-AAV system then could bypass the issue of the AAV vector-related size limitation and pave the way for similar gene therapy applications involving large defective genes.

DMD gene editing

Gene editing is the most exciting approach to treat DMD, as it harnesses the natural cellular repair mechanisms of non-homologous end joining (NHEJ) or homologous recombination (HR) to repair faulty genes at their endogenous loci. This approach would allow the gene to maintain its native regulation and to permanently correct the *DMD* defects. In somatic mammalian (including human) cells, the expected spontaneous NHEJ rate is approximately 1 in 10³ cells but for HR is approximately 1 in 10⁶ cells^{37,38}. Notably, these low spontaneous frequencies have limited both the experimental and the therapeutic gene-editing strategies. Nevertheless, these rates could be increased via induction of site-specific DNA double-strand breaks (DSBs) by introducing custom designer nucleases to specifically cleave the DNA and then leave the cellular repair machinery to be recruited and correct the cleavage.

In practice, engineered nucleases could be directed to target any defective gene within the genome by induction of site-specific DSBs that will be repaired by either NHEJ or HR. To date, nucleases such as meganucleases (MNs), zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and clustered regularly interspaced short palindromic repeats (CRISPRs) systems have been tested in several models of disease.

The feasibility of designer nuclease-mediated gene editing has been shown in muscle disorders^{39–43}. By applying MNs, the normal ORF of a dog microdystrophin containing a frame-shift mutation³⁹ was rescued while the expression of full-length dystrophin mRNA was restored in human patient myoblasts containing a deletion of exons 45–52⁴⁴. This was the first example of successful genome editing of the *DMD* locus. ZFNs were used to target the dystrophin gene producing INDELs (insertion-deletions) of different sizes and leading to the restoration of the normal ORF⁴⁰. Recently, ZFNs were applied to successfully remove exon 51 from the dystrophin transcript, restoring the dystrophin normal ORF in cells of patients with DMD. This approach may lead to the treatment of approximately 13% of mutations of patients with DMD⁴³. The proof of principle of using TALENs for gene editing in primary dermal fibroblasts that originated from a DMD patient carrying a deletion of exons 46–50 was also recently published⁴². Again, the *DMD* normal ORF was restored and functional dystrophin was ultimately expressed by targeting and removing exon 51.

It has lately been shown that AAVs can be used to successfully deliver CRISPR/Cas9 system into *mdx* mice to remove exon 23 from the dystrophin gene, leading to the expression of a partially functional dystrophin in skeletal myofibres and cardiac muscle, and also to improve muscle force⁴⁵. Moreover, it has recently been reported that TALENs and CRISPR/Cas9 systems can be applied side by side to correct the dystrophin gene in induced pluripotent stem cells derived from patients with DMD. In this case, the designer nucleases were used to disrupt the splicing acceptor to skip exon 45 of dystrophin, to induce small INDELs to correct the dystrophin ORF, and to knock in the exon 44 to restore full protein expression⁴⁶. In another study, TALENs and CRISPR nucleases were used to achieve a permanent restoration of *DMD* ORFs in patient-derived muscle cells. To achieve *DMD* restoration, short INDELs were incorporated at out-of-frame sequences to restore normal ORF, a splice acceptor was knocked out to skip exons permanently, and CRISPR-CRISPR or CRISPR-TALEN multiplexing was used to excise targeted exons⁴⁷.

Particularly important for a successful gene-editing approach in muscle disorders is to achieve and maintain a sustainable genetic correction by targeting both post-mitotic muscle tissue and muscle stem cells. Indeed, the correction of endogenous affected muscle stem cells would be advantageous, as their self-renewal feature would allow long-lasting regeneration of a patient's muscles with newly formed corrected muscle cells⁴⁸. Despite promising results of applying designer nuclease-mediated gene editing for DMD treatment, there are still some crucial challenges: an efficient nuclease-based system should ensure a robust cleavage of the targeted DNA but for a very short time in order to achieve the desired site-specific DSBs and to control post-delivery nuclease expression

at their target cells. Furthermore, possible related non-specificity and off-target cuttings (even at very low levels) are relevant concerns for the safety of this approach and preclude, at least with the currently available gene-editing tools, any *in vivo* application in humans.

DMD exon skipping

The exon-skipping approach for DMD is based mainly on delivering antisense oligonucleotides (AONs) targeting some sequences associated with the exons and normally recognized by the cellular splicing machinery. AONs mask these sequences so that specific exons are spliced out from the pre-mRNA and the dystrophin ORF are restored by the expression of shortened BMD-like dystrophin with partial function^{49,50}. In more than 80% of all DMD mutations¹³ (including the majority of out-of-frame deletions⁵¹), the skipping of one or two specific exons could lead to a correct *DMD* gene.

Since its first demonstration two decades ago⁵², exon skipping has witnessed many optimisations and modifications in murine^{53–55} and canine^{56,57} models of DMD. Two AON chemistries have been used in clinical trial so far: 2'-O-methyl-modified ribose molecules with a full-length phosphorothioate backbone (2'OMePS) and phosphorodiamidate morpholino oligomers (PMOs). The largest group (13%) of all DMD patients could be treated by skipping exon 51¹³. The first clinical trial, based on intramuscular delivery of PRO051, a 2'OMePS AON, to induce exon 51 skipping was a success; sarcolemmal dystrophin was restored in 64 to 97% of examined myofibres⁵⁸. In subsequent phase II/III clinical trials, GlaxoSmithKline (London, UK) and BioMarin (formerly Prosensa) (Novato, CA, USA) tested the systemic delivery of the same AON, called drisapersen (latterly Kyndrisa). However, the results failed to demonstrate a statistically significant improvement in some crucial tests (for example, 6-minute-walk test)⁵⁹, and early this year, the US Food and Drug Administration (FDA) did not approve drisapersen as a marketable drug and this resulted in discontinuation of the drug⁶⁰. In parallel, AVI BioPharma (now known as Sarepta Therapeutics, Cambridge, MA, USA) developed AVI-4658 based on PMO chemistry to skip the exon 51 of dystrophin. After intramuscular injection into boys with DMD, AVI-4658 showed a 44 to 79% increase of dystrophin expression⁶¹. The same AON tested systemically by Sarepta Therapeutics, renamed eteplirsen, increased dystrophin-positive fibres by 23% compared with placebo-injected controls⁶². Eteplirsen then was systemically injected in a phase II/III clinical trial but with results similar to those obtained by the 2'OMePS-based AON drisapersen⁶³, and its approval by the FDA is currently under debate⁶⁴.

Despite promising results of systemic delivery of AONs for exon skipping, both in animal models and in humans, the treatment suffers from relatively poor efficiency. Increasing drug effectiveness by elevating dose levels over chronic time periods may not be feasible owing to the risk of toxic side effects. One other option is to increase the relative dose effectiveness without unduly exacerbating the risk of side effects. Thus, more research needs to be done to find better AON chemistries and possibly more efficient strategies to deliver them. New chemistries have recently been tested as tricyclo-DNA, a DNA analog, which was systematically administered into two DMD mouse models, leading

to efficient dystrophin expression in both skeletal and cardiac muscles and, to a lesser extent, in the brain⁶⁵. Additionally, Gao *et al.* have reported that, in *mdx* mice, the repeated administration of peptide nucleic acid AONs, a synthetic chemistry, restores dystrophin in gastrocnemius, leading to amelioration of dystrophic pathology in DMD mice⁶⁶. Moreover, it has been demonstrated that systemic administration of PMO conjugated to cell-penetrating peptides resulted in high levels of dystrophin restoration in major respiratory muscles, including the diaphragm, and improved the cardiac function in *mdx* mice⁶⁷. Also, in an *mdx* model, it was recently demonstrated that dual exon skipping of dystrophin and myostatin pre-mRNAs using PMO conjugated with an arginine-rich peptide improved dystrophin expression and decreased muscle necrosis, particularly in the diaphragm⁶⁸. Likewise, the same principle was used to target dystrophin and Actvr2b to produce an internally deleted protein leading to comparable exon-skipping levels for both pre-mRNA targets when compared with individual PMO conjugates both *in vitro* and *in vivo* in *mdx* mice⁶⁹.

Alternatively, exon skipping of dystrophin exons could be achieved by using other approaches like CRISPR-Cas9 nuclease. In this regard, it has recently been shown that exon 23 could be skipped by using the CRISPR-Cas9 system when delivered to postnatal *mdx* mice intraperitoneally at postnatal day 1 (P1), intramuscularly at P12, and retro-orbitally at P18. Two CRISPR systems were used, preceding and following the mutated exon in the *DMD* gene, and delivered by AAV9. Following this genome editing treatment, restoration of dystrophin expression was detected at varied levels in both cardiac and skeletal muscles up to 12 weeks after injection⁵⁴. In a similar study, a dual CRISPR system was applied to induce a specific larger deletion across exons 50 and 54 of the *DMD* gene, resulting in fusion of the targeted exons. Whereas the *in vitro* experiments were performed in cultured 293T cells or DMD patient myoblasts (with deletion of exons 51–53), the *in vivo* experiments were done in the humanised hDMD/*mdx* mice. Study results revealed that significant levels of hybrid exon 50/54 were formed *in vitro* in DMD myoblasts and that in hDMD/*mdx* mice *in vivo* the corrected *DMD* ORF was partially restored⁵⁵.

The last two studies suggest that this gene-editing system may be advantageous over AON exon skipping, as a single injection leads to the permanent correction of the genome. However, no data on toxicity or off-target effects of using such gene-editing tools *in vivo* were reported. Those two studies highlight the importance of applying more than one approach to restore dystrophin. Indeed, a relevant amount of future research will focus on developing combined strategies to ameliorate the disease. As an example of this research direction, engineered CD133⁺ human DMD stem cells were transduced with lentivirus vectors that permanently delivered the cloned AONs rescuing murine dystrophin expression⁷⁰. Moreover, lately it was reported that co-administration of PMOs with glucose enhances exon-skipping activity in *mdx* mice⁵³.

Altogether, the *DMD* trans-splicing, gene-editing, and exon-skipping approaches, though still presenting some important issues, such as relatively low efficiency, possible associated toxic effects, or the need for chronic delivery, represent the most attractive and promising genetic treatments for DMD.

Abbreviations

2'OMePS, 2'-O-methyl-modified ribose molecules with a full-length phosphorothioate backbone; AAV, adeno-associated virus; AON, antisense oligonucleotide; BMD, Becker muscular dystrophy; CRISPR, clustered regularly interspaced short palindromic repeat; DGC, dystrophin-associated glycoprotein complex; DMD, Duchenne muscular dystrophy; DSB, double-strand break; FDA, US Food and Drug Administration; HR, homologous recombination; INDEL, insertion-deletion; MN, meganuclease; NHEJ, non-homologous end joining; ORF, open reading frame; P, postnatal day; PMO, phosphorodiamidate morpholino oligomer; TALEN,

transcription activator-like effector nuclease; TTS, triple trans-splicing; ZFN, zinc finger nuclease.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Mah JK, Korngut L, Dykeman J, *et al.*: **A systematic review and meta-analysis on the epidemiology of Duchenne and Becker muscular dystrophy.** *Neuromuscul Disord.* 2014; 24(6): 482–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Judge LM, Haraguchi M, Chamberlain JS: **Dissecting the signaling and mechanical functions of the dystrophin-glycoprotein complex.** *J Cell Sci.* 2006; 119(Pt 8): 1537–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ramos J, Chamberlain JS: **Gene Therapy for Duchenne muscular dystrophy.** *Expert Opin Orphan Drugs.* 2015; 3(11): 1255–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Konieczny P, Swiderski K, Chamberlain JS: **Gene and cell-mediated therapies for muscular dystrophy.** *Muscle Nerve.* 2013; 47(5): 649–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jarmin S, Kymalainen H, Popplewell L, *et al.*: **New developments in the use of gene therapy to treat Duchenne muscular dystrophy.** *Expert Opin Biol Ther.* 2014; 14(2): 209–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fairclough RJ, Wood MJ, Davies KE: **Therapy for Duchenne muscular dystrophy: renewed optimism from genetic approaches.** *Nat Rev Genet.* 2013; 14(6): 373–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Long C, McAnally JR, Shelton JM, *et al.*: **Prevention of muscular dystrophy in mice by CRISPR/Cas9-mediated editing of germline DNA.** *Science.* 2014; 345(6201): 1184–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Emery AEH, Muntoni F: **Duchenne Muscular Dystrophy.** OUP Oxford, 2003.
[Reference Source](#)
- Cox GF, Kunkel LM: **Dystrophies and heart disease.** *Curr Opin Cardiol.* 1997; 12(3): 329–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zhu X, Wheeler MT, Hadhazy M, *et al.*: **Cardiomyopathy is independent of skeletal muscle disease in muscular dystrophy.** *FASEB J.* 2002; 16(9): 1096–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Townsend D, Yasuda S, Li S, *et al.*: **Emergent dilated cardiomyopathy caused by targeted repair of dystrophic skeletal muscle.** *Mol Ther.* 2008; 16(5): 832–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Martone J, Briganti F, Legnini I, *et al.*: **The lack of the Celf2a splicing factor converts a Duchenne genotype into a Becker phenotype.** *Nat Commun.* 2016; 7: 10488.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Aartsma-Rus A, Fokkema I, Verschuuren J, *et al.*: **Theoretic applicability of antisense-mediated exon skipping for Duchenne muscular dystrophy mutations.** *Hum Mutat.* 2009; 30(3): 293–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fairclough RJ, Perkins KJ, Davies KE: **Pharmacologically targeting the primary defect and downstream pathology in Duchenne muscular dystrophy.** *Curr Gene Ther.* 2012; 12(3): 206–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Emery A: **Muscular dystrophy—the facts.** *Neuromuscul Disord.* 1995; 5(6): 521.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bladen CL, Salgado D, Monges S, *et al.*: **The TREAT-NMD DMD Global Database: analysis of more than 7,000 Duchenne muscular dystrophy mutations.** *Hum Mutat.* 2015; 36(4): 395–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Neri M, Torelli S, Brown S, *et al.*: **Dystrophin levels as low as 30% are sufficient to avoid muscular dystrophy in the human.** *Neuromuscul Disord.* 2007; 17(11–12): 913–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cavazzana-Calvo M, Fischer A: **Gene therapy for severe combined immunodeficiency: are we there yet?** *J Clin Invest.* 2007; 117(6): 1456–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Athanasopoulos T, Graham IR, Foster H, *et al.*: **Recombinant adeno-associated viral (rAAV) vectors as therapeutic tools for Duchenne muscular dystrophy (DMD).** *Gene Ther.* 2004; 11(Suppl 1): S109–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blankinship MJ, Gregorevic P, Chamberlain JS: **Gene therapy strategies for Duchenne muscular dystrophy utilizing recombinant adeno-associated virus vectors.** *Mol Ther.* 2006; 13(2): 241–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fang H, Lai NC, Gao MH, *et al.*: **Comparison of adeno-associated virus serotypes and delivery methods for cardiac gene transfer.** *Hum Gene Ther Methods.* 2012; 23(4): 234–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koo T, Malerba A, Athanasopoulos T, *et al.*: **Delivery of AAV2/9-microdystrophin genes incorporating helix 1 of the coiled-coil motif in the C-terminal domain of dystrophin improves muscle pathology and restores the level of α 1-syntrophin and α -dystrobrevin in skeletal muscles of mdx mice.** *Hum Gene Ther.* 2011; 22(11): 1379–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hayashita-Kinoh H, Yugeta N, Okada H, *et al.*: **Intra-amniotic rAAV-mediated microdystrophin gene transfer improves canine X-linked muscular dystrophy and may induce immune tolerance.** *Mol Ther.* 2015; 23(4): 627–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Wang Z, Storb R, Halbert CL, *et al.*: **Successful regional delivery and long-term expression of a dystrophin gene in canine muscular dystrophy: a preclinical model for human therapies.** *Mol Ther.* 2012; 20(8): 1501–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Bostick B, Shin JH, Yue Y, *et al.*: **AAV micro-dystrophin gene therapy alleviates stress-induced cardiac death but not myocardial fibrosis in >21-m-old mdx mice, an end-stage model of Duchenne muscular dystrophy cardiomyopathy.** *J Mol Cell Cardiol.* 2012; 53(2): 217–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mendell JR, Campbell K, Rodino-Klapac L, *et al.*: **Dystrophin immunity in Duchenne's muscular dystrophy.** *N Engl J Med.* 2010; 363(15): 1429–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mays LE, Wilson JM: **The complex and evolving story of T cell activation to AAV vector-encoded transgene products.** *Mol Ther.* 2011; 19(1): 16–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yan Z, Zhang Y, Duan D, *et al.*: **Trans-splicing vectors expand the utility of adeno-associated virus for gene therapy.** *Proc Natl Acad Sci U S A.* 2000; 97(12): 6716–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sun L, Li J, Xiao X: **Overcoming adeno-associated virus vector size limitation through viral DNA heterodimerization.** *Nat Med.* 2000; 6(5): 599–602.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nakai H, Storm TA, Kay MA: **Increasing the size of rAAV-mediated expression cassettes in vivo by intermolecular joining of two complementary vectors.** *Nat Biotechnol.* 2000; 18(5): 527–32.
[PubMed Abstract](#) | [Publisher Full Text](#)

31. Lai Y, Yue Y, Liu M, *et al.*: **Efficient *in vivo* gene expression by trans-splicing adeno-associated viral vectors.** *Nat Biotechnol.* 2005; 23(11): 1435–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Ghosh A, Yue Y, Long C, *et al.*: **Efficient whole-body transduction with trans-splicing adeno-associated viral vectors.** *Mol Ther.* 2007; 15(4): 750–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Duan D, Yue Y, Engelhardt JF: **Expanding AAV packaging capacity with trans-splicing or overlapping vectors: a quantitative comparison.** *Mol Ther.* 2001; 4(4): 383–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Yue Y, Liu M, Duan D: **C-terminal-truncated microdystrophin recruits dystrobrevin and syntrophin to the dystrophin-associated glycoprotein complex and reduces muscular dystrophy in symptomatic utrophin/dystrophin double-knockout mice.** *Mol Ther.* 2006; 14(1): 79–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Li J, Sun W, Wang B, *et al.*: **Protein trans-splicing as a means for viral vector-mediated *in vivo* gene therapy.** *Hum Gene Ther.* 2008; 19(9): 958–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Koo T, Popplewell L, Athanasopoulos T, *et al.*: **Triple trans-splicing adeno-associated virus vectors capable of transferring the coding sequence for full-length dystrophin protein into dystrophic mice.** *Hum Gene Ther.* 2014; 25(2): 98–108.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Gellhaus K, Cornu TI, Heilbronn R, *et al.*: **Fate of recombinant adeno-associated viral vector genomes during DNA double-strand break-induced gene targeting in human cells.** *Hum Gene Ther.* 2010; 21(5): 543–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Pruett-Miller SM, Connelly JP, Maeder ML, *et al.*: **Comparison of zinc finger nucleases for use in gene targeting in mammalian cells.** *Mol Ther.* 2008; 16(4): 707–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Chapdelaine P, Pichavant C, Rousseau J, *et al.*: **Meganucleases can restore the reading frame of a mutated dystrophin.** *Gene Ther.* 2010; 17(7): 846–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Rousseau J, Chapdelaine P, Boisvert S, *et al.*: **Endonucleases: tools to correct the dystrophin gene.** *J Gene Med.* 2011; 13(10): 522–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. **F** Benabdallah BF, Duval A, Rousseau J, *et al.*: **Targeted Gene Addition of Microdystrophin in Mice Skeletal Muscle via Human Myoblast Transplantation.** *Mol Ther Nucleic Acids.* 2013; 2(1): e68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
42. **F** Ousterout DG, Perez-Pinera P, Thakore PI, *et al.*: **Reading frame correction by targeted genome editing restores dystrophin expression in cells from Duchenne muscular dystrophy patients.** *Mol Ther.* 2013; 21(9): 1718–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. **F** Ousterout DG, Kabadi AM, Thakore PI, *et al.*: **Correction of dystrophin expression in cells from Duchenne muscular dystrophy patients through genomic excision of exon 51 by zinc finger nucleases.** *Mol Ther.* 2015; 23(3): 523–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. Popplewell L, Koo T, Leclerc X, *et al.*: **Gene correction of a duchenne muscular dystrophy mutation by meganuclease-enhanced exon knock-in.** *Hum Gene Ther.* 2013; 24(7): 692–701.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. **F** Nelson CE, Hakim CH, Ousterout DG, *et al.*: ***In vivo* genome editing improves muscle function in a mouse model of Duchenne muscular dystrophy.** *Science.* 2016; 351(6271): 403–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. **F** Li HL, Fujimoto N, Sasakawa N, *et al.*: **Precise correction of the dystrophin gene in duchenne muscular dystrophy patient induced pluripotent stem cells by TALEN and CRISPR-Cas9.** *Stem Cell Reports.* 2015; 4(1): 143–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
47. **F** Maggio I, Stefanucci L, Janssen JM, *et al.*: **Selection-free gene repair after adenoviral vector transduction of designer nucleases: rescue of dystrophin synthesis in DMD muscle cell populations.** *Nucleic Acids Res.* 2016; 44(3): 1449–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. Bertoni C: **Emerging gene editing strategies for Duchenne muscular dystrophy targeting stem cells.** *Front Physiol.* 2014; 5: 148.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Popplewell LJ, Malerba A, Dickson G: **Optimizing antisense oligonucleotides using phosphorodiamidate morpholino oligomers.** *Methods Mol Biol.* 2012; 867: 143–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Betts CA, Hammond SM, Yin HF, *et al.*: **Optimizing tissue-specific antisense oligonucleotide-peptide conjugates.** *Methods Mol Biol.* 2012; 867: 415–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Muntoni F, Wood MJ: **Targeting RNA to treat neuromuscular disease.** *Nat Rev Drug Discov.* 2011; 10(8): 621–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Dunkley MG, Eperon IC, Dickson G: **Modulation of pre-mRNA splicing in the Duchenne muscular dystrophy gene.** *Biochem Soc Trans.* 1996; 24(2): 276S.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. **F** Han G, Gu B, Cao L, *et al.*: **Hexose enhances oligonucleotide delivery and exon skipping in dystrophin-deficient *mdx* mice.** *Nat Commun.* 2016; 7: 10981.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
54. **F** Long C, Amoasii L, Mireault AA, *et al.*: **Postnatal genome editing partially restores dystrophin expression in a mouse model of muscular dystrophy.** *Science.* 2016; 351(6271): 400–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. **F** Iyombe-Engembe JP, Ouellet DL, Barbeau X, *et al.*: **Efficient Restoration of the Dystrophin Gene Reading Frame and Protein Structure in DMD Myoblasts Using the CinDel Method.** *Mol Ther Nucleic Acids.* 2016; 5: e283.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. Vulin A, Barthélemy I, Goyenvalle A, *et al.*: **Muscle function recovery in golden retriever muscular dystrophy after AAV1-U7 exon skipping.** *Mol Ther.* 2012; 20(11): 2120–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Yokota T, Nakamura A, Nagata T, *et al.*: **Extensive and prolonged restoration of dystrophin expression with vivo-morpholino-mediated multiple exon skipping in dystrophic dogs.** *Nucleic Acid Ther.* 2012; 22(5): 306–15.
[PubMed Abstract](#) | [Free Full Text](#)
58. **F** van Deutekom JC, Janson AA, Ginjaar IB, *et al.*: **Local dystrophin restoration with antisense oligonucleotide PRO051.** *N Engl J Med.* 2007; 357(26): 2677–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Lu Q, Cirak S, Partridge T: **What Can We Learn From Clinical Trials of Exon Skipping for DMD?** *Mol Ther Nucleic Acids.* 2014; 3(3): e152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. http://www.ukmhs.nhs.uk/applications/bdo/record_view_open.asp?newDrugID=4873. 2016.
61. Kinali M, Arechavala-Gomez V, Feng L, *et al.*: **Local restoration of dystrophin expression with the morpholino oligomer AVI-4658 in Duchenne muscular dystrophy: a single-blind, placebo-controlled, dose-escalation, proof-of-concept study.** *Lancet Neurol.* 2009; 8(10): 918–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. **F** Mendell JR, Rodino-Klapac LR, Sahenk Z, *et al.*: **Eteplirsen for the treatment of Duchenne muscular dystrophy.** *Ann Neurol.* 2013; 74(5): 637–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. **F** Mendell JR, Goemans N, Lowes LP, *et al.*: **Longitudinal effect of eteplirsen versus historical control on ambulation in Duchenne muscular dystrophy.** *Ann Neurol.* 2016; 79(2): 257–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
64. Hodgkinson L, Sorbera L, Graul AI: **Duchenne muscular dystrophy drugs face tough path to approval.** *Drugs Today (Barc).* 2016; 52(3): 199–202.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. **F** Goyenvalle A, Griffith G, Babbs A, *et al.*: **Functional correction in mouse models of muscular dystrophy using exon-skipping tricyclo-DNA oligomers.** *Nat Med.* 2015; 21(3): 270–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
66. **F** Gao X, Shen X, Dong X, *et al.*: **Peptide Nucleic Acid Promotes Systemic Dystrophin Expression and Functional Rescue in Dystrophin-deficient *mdx* Mice.** *Mol Ther Nucleic Acids.* 2015; 4(10): e255.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
67. **F** Betts CA, Saleh AF, Carr CA, *et al.*: **Implications for Cardiac Function Following Rescue of the Dystrophic Diaphragm in a Mouse Model of Duchenne Muscular Dystrophy.** *Sci Rep.* 2015; 5: 11632.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. Lu-Nguyen NB, Jarmin SA, Saleh AF, *et al.*: **Combination Antisense Treatment for Destructive Exon Skipping of Myostatin and Open Reading Frame Rescue of Dystrophin in Neonatal *mdx* Mice.** *Mol Ther.* 2015; 23(8): 1341–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. **F** Shabanpoor F, McClorey G, Saleh AF, *et al.*: **Bi-specific splice-switching PMO oligonucleotides conjugated via a single peptide active in a mouse model of Duchenne muscular dystrophy.** *Nucleic Acids Res.* 2015; 43(1): 29–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
70. **F** Meregalli M, Farini A, Sitzia C, *et al.*: **Stem Cell-Mediated Exon Skipping of the Dystrophin Gene by the Bystander Effect.** *Curr Gene Ther.* 2015; 15(6): 563–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Shin'ichi Takeda**

National Center of Neurology and Psychiatry, Tokyo, Japan

Competing Interests: No competing interests were disclosed.

2. **Giulio Cossu**

University of Manchester, Manchester, UK

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research