REVIEW

Pathophysiological links between traumatic brain injury and post-traumatic headaches [version 1; peer review: 3 approved]

Robert L. Ruff¹, Kayla Blake²

¹Case Western Reserve University, Cleveland, OH, USA

²Mayfield High School, Mayfield, OH, USA

V1 First published: 31 Aug 2016, 5(F1000 Faculty Rev):2116
<https://doi.org/10.12688/f1000research.9017.1>

Latest published: 31 Aug 2016, 5(F1000 Faculty Rev):2116
<https://doi.org/10.12688/f1000research.9017.1>

Abstract

This article reviews possible ways that traumatic brain injury (TBI) can induce migraine-type post-traumatic headaches (PTHs) in children, adults, civilians, and military personnel. Several cerebral alterations resulting from TBI can foster the development of PTH, including neuroinflammation that can activate neural systems associated with migraine. TBI can also compromise the intrinsic pain modulation system and this would increase the level of perceived pain associated with PTH. Depression and anxiety disorders, especially post-traumatic stress disorder (PTSD), are associated with TBI and these psychological conditions can directly intensify PTH. Additionally, depression and PTSD alter sleep and this will increase headache severity and foster the genesis of PTH. This article also reviews the anatomic loci of injury associated with TBI and notes the overlap between areas of injury associated with TBI and PTSD.

Keywords

Traumatic brain injury , post traumatic headaches , post traumatic stress disorder , neuroinflammation

Open Peer Review

Approval Status

	1	2	3
version 1 31 Aug 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Karen Barlow**, Cumming School of Medicine, University of Calgary, Calgary, Canada
2. **Mary Jo Pugh**, University of Texas Health Science Center, San Antonio, USA
Carlos Jaramillo, South Texas Veterans Health Care System, Texas, USA
3. **Ross Zafonte**, Harvard Medical School; Spaulding Rehabilitation Hospital; Brigham and Women's Hospital, Boston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Robert L. Ruff (robertlruff@aol.com)

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Ruff RL and Blake K. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ruff RL and Blake K. **Pathophysiological links between traumatic brain injury and post-traumatic headaches [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2116 <https://doi.org/10.12688/f1000research.9017.1>

First published: 31 Aug 2016, 5(F1000 Faculty Rev):2116 <https://doi.org/10.12688/f1000research.9017.1>

Introduction

The second International Classification of Headaches (ICHD-2) defines post-traumatic headaches (PTHs) as secondary headaches following traumatic brain injury (TBI)¹. No symptom-based criteria define PTH in the ICHD-2 criteria. PTH is more frequent after mild TBI (mTBI) compared with moderate or severe TBI² and can resemble one or more primary headache disorders^{3–9}. The most common PTH patterns resemble two primary headache disorders—migraine or probable migraine and tension-type headaches—and migraine-type headaches are more prevalent^{3–5,8,10,11}. In one prospective study of adult civilian TBI associated with motor vehicle accidents, 58% of subjects continued to have PTH one year after their TBI¹². Migraine-like PTHs usually resemble migraine without aura^{5,7,8,13,14}. Neck injury in conjunction with TBI can result in cervicogenic headaches or pain that may resemble occipital neuralgia^{11,15}. PTH can impair quality of life. Among veterans returning from Iraq or Afghanistan, the presence of PTH, especially if combined with one of the frequent TBI-associated psychological conditions of post-traumatic stress disorder (PTSD) or depression, diminishes the success of returning to work and other aspects of community reintegration¹⁶.

Cervicogenic pain and tension-type PTH following TBI are often attributed to musculoskeletal injury caused by the trauma that produced TBI^{11,15}. However, musculoskeletal trauma does not explain migraine-type PTH. This review describes possible mechanistic links between TBI and migraine to suggest how TBI can induce migraine-type PTH.

Traumatic brain injury-induced neuroinflammation

Many brain tissue responses following TBI enhance the likelihood of, and perhaps directly cause, migraine-type PTH. For example, cellular injury increases the concentration of extracellular potassium, which will trigger neuronal depolarization and release of neurotransmitters that promote headaches¹⁷. In addition, TBI triggers neuroinflammation, which enhances neuronal death and impairs recovery of function, but can also alter central nervous system (CNS) pain processing to induce migraine. Neuroinflammation is characterized by the activation of microglia; the release of pro-inflammatory chemicals, including chemokines, specific interleukins, and tumor necrosis factor- α (TNF- α); and possibly the invasion of the CNS by inflammatory lymphocytes and phagocytic white cells coming through cerebral blood vessels. Neuroinflammation is a central element in the development of chronic traumatic encephalopathy (CTE) associated with repeated sports-associated TBI events^{18–22} and includes headache as part of the symptom spectrum¹⁸. Non-hemorrhagic, closed-head TBI activates microglia, leading to the production of inflammatory and pro-inflammatory molecules^{20,23–28}, and even minor experimental TBI induces neuroinflammation by activating microglia²⁹. Moderate and severe TBI can disrupt the blood-brain barrier, allowing invasion of neutrophils from leaky blood vessels^{23,24,28}. Neuroinflammation develops quickly after TBI as shown by a post-mortem study of civilians in which the brains analyzed after death associated with acute TBI demonstrated elevated mRNA levels of interleukin-1 beta (IL-1 β), IL-6, and IL-8 and TNF- α within minutes of TBI³⁰.

Neuroinflammation can be modulated in several ways. Microglial activation is regulated by several factors, including poly (ADP-ribose) polymerase-1 (PARP-1) and the metabotropic glutamate receptor 5 (mGluR5). PARP-1 is present in neurons and glia. It is the most abundant member of a family of enzymes that attach ADP-ribose polymers onto proteins. PARP-1 resides in the nucleus and is activated by DNA strand breaks^{31,32}. At low levels of activation, PARP-1 facilitates DNA base excision repair by poly-ADP-ribosylation of proteins involved in DNA repair. PARP-1 recruits DNA repair proteins to DNA break sites to restore DNA integrity. However, PARP-1 activation can damage neurons and glia after TBI^{33,34}. PARP-1-mediated neuronal cell death may be partially due to PARP-1 activation depleting NAD⁺ and ATP³¹. Additionally, PARP-1 may initiate neuronal death through the release of apoptosis-inducing factor and impairing mitochondrial function and glycolytic metabolism^{32,35,36}. PARP-1-dependent nuclear factor-kappa-B activation induces the expression of pro-inflammatory genes and activates microglia^{37,38}. Activated microglia release multiple types of neurotoxic molecules. Therefore, strong activation of PARP-1 due to TBI may convert PARP-1 from being an agent of DNA repair to a trigger of neuronal injury and neuroinflammation. In support of PARP-1 contributing to CNS damage in TBI, PARP-1 inhibition reduces microglial activation and functional impairment and increases neuron survival in experimental models of TBI^{38–40}. The activation of mGluR5 decreases microglial activation and reduces microglial release of pro-inflammatory factors, which reduces neuronal loss and improves functional recovery after experimental TBI⁴¹.

Peroxisome proliferator-activated receptors (PPARs) are ligand-activated transcription factors that are members of the super family of nuclear hormone receptors^{42,43}. PPARs control many physiological functions, including glucose absorption, lipid balance, and cell growth and differentiation. Among the several PPAR isoforms, the activation of PPAR- α and PPAR- γ exerts neuroprotective actions in acute cerebral injury such as TBI^{44–47} and stroke⁴⁸ and reduces neuronal injury in animal models of cerebral degenerative disorders, including Parkinson's disease^{49–52}, amyotrophic lateral sclerosis⁵³, and Alzheimer's disease⁵⁴. Treatment with a PPAR agonist after experimental TBI prevents microglial activation, reduces cognitive impairment, and reduces neuron death⁴⁶.

The role of neuroinflammation in brain injury and recovery following TBI is complex. IL-6 is an inflammatory mediator. IL-6 is one of the first cytokines produced after TBI. Chemically suppressing IL-6, IL-1 β , and TNF- α in mice reduces brain edema and apoptosis⁵⁵. However, selective knockout of IL-6 in mice impaired functional recovery and enhanced the production of IL-1 β following TBI compared with controls⁵⁶. In children, IL-6 and nerve growth factor upregulation correlated with improved outcome after severe TBI⁵⁷. Activation of adenosine 2A receptors can have pro-inflammatory or anti-inflammatory actions in the brain depending on the cell types containing the receptor and the degree of receptor activation⁵⁸. Chemokines may have a complex role in brain response to TBI. Deletion of the chemokine ligand 2 receptor (CCR2) resulted in reduced monocyte infiltration of the cortex

along with diminished neuronal death and axon injury in an animal model of TBI²⁰. However, CCR2 deletion did not change microglia activation. Additionally, CCR2 deletion was associated with increased tau-protein mislocalization to the neuron cell bodies in the cortex and increased levels of phosphorylated tau in the hippocampus²⁰. The changes in tau associated with CCR2 deletion resemble findings described in CTE^{19,21}.

Physical exercise before experimental TBI reduces neuroinflammation and improves functional recovery following fluid-percussion TBI in rats⁵⁹. The protective effect of physical conditioning prior to TBI might influence recovery after TBI in athletes and active military personnel. The mechanisms of exercise reducing neuroinflammation are under investigation.

Possible links between neuroinflammation and migraine

Neuroinflammation alters the functions and sensitivities of meningeal and cerebrovascular structures⁶⁰ that are involved in the development and progression of head pain and other features of migraine⁶¹. Additionally, inflammation and other responses to injury can enhance neuronal excitability^{62–64}, which could potentiate the genesis of cortical spreading depression (CSD). CSD is an important element in migraine, particularly migraine aura⁶¹. Thus, altering the properties of cerebral, meningeal, and cerebrovascular structures involved in migraine could be a link between TBI and PTH.

Neuroinflammation may play a role in brain recovery after TBI; however, exaggerated inflammation may lead to a state of hyperexcitability involving structures that could potentiate the development of migraine and other headaches. The headache phase of migraine is associated with the release of vasoactive neuropeptides by the trigeminovascular system, vasodilation of the extracerebral intracranial arteries, and increased nociceptive transmission within the central trigeminocervical complex⁶⁵. Cady proposed that migraine symptoms evolve as the pathology of inflammation and neural sensitization progress⁶⁶. Tension headache symptoms appeared early in the course of inflammation. As inflammation and neural hyperexcitability advance, the pattern of headaches shifted from tension-like to migrainous headaches to classic migraine and other forms of migraine. This convergence model of headaches provides a conceptual framework to consider how PTH can result from post-TBI neuroinflammation. Several reviews addressed the connections between neuroinflammation and migraine^{67–69}.

Hyperexcitability of primary trigeminal afferent nociceptive neurons can initiate or aggravate neuroinflammation involving meningeal blood vessels. Hyperexcitability resulting from inflammation of trigeminal nerve branches mediates the throbbing head pain of migraine⁶¹. The throbbing character of migraine may reflect pain signals carried by afferent pain fibers that innervate pulsating meningeal blood vessels. CSD is involved in the genesis of migraine aura and activation of meningeal pain receptors⁶⁹. CSD may sensitize and activate meningeal nociceptors through the release of vasoactive pain-stimulating chemicals such as substance P, leading to the activation of mast cells and macrophages, which in turn release inflammatory mediators, including cytokines⁶⁹. Another

factor involved in migraine is the neuropeptide calcitonin gene-related peptide (CGRP)⁷⁰. CGRP modulates pain by peripheral and central mechanisms. Outside the brain, meningeal nociceptors are activated by CSD triggering the release of CGRP, which induces vasodilation and edema⁶⁸. CGRP can also sensitize meningeal nociceptors producing positive feedback that increases the activity of meningeal nociceptors. Additionally, glia in the trigeminal ganglion release a CGRP-like peptide called procalcitonin, which also induces neuroinflammation^{68,70}. The transient receptor potential V1 channel (TRPV1) potentiates the release of CGRP in nociceptive trigeminal ganglia neurons and TBI activates TRPV1⁷¹.

Traumatic brain injury can impair pain modulation

Pain intensity is based in part on the activity of intrinsic pain modulation systems that diminish pain transmission and alter the perceived intensity of painful stimuli⁷². Pain modulation is impaired in people with chronic PTH⁷³. Diminished ability to reduce the perceived intensity of pain could prolong the duration and increase the intensity of PTH⁷³. Impaired pain modulation in people with PTH results from disruption of the intrinsic pain modulation system and psychological changes that can alter pain perception. We will first consider TBI-induced alterations in the intrinsic pain modulation system.

Serotonergic (5-HT) neurons are important in pain modulation⁷², and migraine is associated with 5-HT pathways. The triptan class of migraine abortive medications are agonists of 5-HT(1B/1D) receptors⁷⁴. TBI could facilitate PTH by reducing the activity of 5-HT neurons that counter migraine. Activating 5-HT receptors inhibits the release of CGRP and other vasoactive neuropeptides from trigeminal nerves^{68,70}. 5-HT(1B/1D) receptors constrict painfully dilated cerebral blood vessels and inhibit nociceptive neurotransmission in trigeminal pathways⁶⁵. Trigeminal ganglion neurons project centrally to several locations, including pain pathways and the nucleus tractus solitarius (NTS), which is involved in the sensation of nausea⁷⁵.

TBI targets 5-HT pathways in the brain. Long serotonergic axons may be damaged by diffuse axonal injury (DAI) associated with TBI. TBI causes DAI because long axons are stretched by the physical distortion of the brain which occurs with all forms of diffuse TBI^{76,77}. An imaging study of people with chronic PTH demonstrated central white matter damage that correlated with the presence of chronic PTH⁷⁸. The imaging study could not identify whether the white matter damage involved components of the pain modulation system; however, the locations of white matter injury associated with chronic PTH included regions containing pain modulation fibers that interface with the amygdalae and other cerebral nuclei⁷².

Serotonergic neuron cell bodies are organized into two distinct groups: caudal (medulla and caudal pons) and rostral (midbrain and rostral pons)⁷⁹. Additionally, there are serotonergic neuron cell bodies in the medullary reticular formation (involved in alertness) and other non-raphe regions such as the hippocampus and the substantia nigra⁷⁹. There is a biphasic change in the levels of 5-HT in cerebral extracellular and perivascular spaces after TBI. For the first few days after TBI, 5-HT levels are elevated^{80–82}. Elevation of 5-HT

levels may result from stretch-induced firing of serotonergic nerve fibers and release of 5-HT from inflammatory cells. The acute elevation of 5-HT levels is associated with breakdown of the blood-brain barrier and cerebral edema⁸³. Levels of 5-HT and its metabolites decline several days after brain injury and can remain persistently reduced^{84,85}. Persistent damage to 5-HT pathways, particularly serotonergic innervation of cerebral blood vessels, may induce migraine following TBI by disrupting normal serotonergic suppression of patterns of neuronal and cerebrovascular activity associated with migraine⁸². Additionally, trigeminovascular afferent activation of NTS cells can induce symptoms such as nausea. Nausea can be suppressed by activating 5-HT(1B/1D) receptors that inhibit NTS cells associated with migraine symptoms such as nausea⁷⁵. Thus, somatic symptoms associated with migraine such as nausea may be potentiated by over-activity of injured trigeminal neurons and reduction in serotonergic tone following TBI.

Traumatic brain injury-associated psychological changes that may potentiate post-traumatic headache

Psychological states of depression and anxiety can increase the perceived intensity of pain such as PTH⁷². Among civilians with migraine, the frequency and severity of migraine attacks are associated with PTSD⁸⁶. The presence of mTBI may increase the likelihood that a psychologically traumatic event results in PTSD⁸⁷. In rats, explosions that produce mTBI induce PTSD-like behavior⁸⁸. Among combat veterans with mTBI-associated explosions, there is a dose-response relationship between the number of TBI events and the likelihood of an individual developing PTSD^{4,89,90}. A study of anxiety disorders in children following TBI found that 8.5% of children developed anxiety disorders, usually PTSD, within 6 months of the TBI and those with mTBI had the greatest likelihood of developing PTSD⁹¹. A national study of civilian trauma in Australia reported that the prevalence of new-onset depression was 9% after trauma. The presence of TBI with the trauma did not seem to influence the development of depression. PTSD prevalence following mTBI was 6%, which was about 1.9-fold higher than following trauma without TBI⁹². PTSD and mTBI are associated in combat personnel. About 40% of military personnel and veterans with combat-acquired mTBI had PTSD^{11,93}. The incidence rates of PTSD were 44% for soldiers recently deployed in Iraq who experienced an episode of loss of consciousness due to mTBI, 16% for soldiers with other injuries, and 9% for uninjured soldiers⁹³.

Depression and PTSD are frequently present in civilians with chronic PTH⁹⁴⁻⁹⁷. A study of psychological issues in civilians with chronic PTH found that 30% of subjects with PTH met criteria for PTSD and depression⁹⁸. Although experiencing chronic pain can alter an individual's psychological state, data suggest that PTSD is probably linked to the traumatic event rather than to chronicity of pain⁹⁹. A meta-analysis of studies of civilian and military TBI concluded that PTSD may modulate the intensity and severity of chronic PTH but that TBI independently correlated with the genesis of chronic PTH^{2,100}. A longitudinal study of Iraq and Afghanistan war veterans initially treated by the Department of Veterans Affairs in 2008 and followed through 2011 reported that PTH correlated with mTBI⁹⁵. Additionally, coexistence of PTSD, depression, or

both increased the likelihood of chronic PTH beyond that associated with mTBI alone⁹⁵.

PTSD and depression can intensify PTH by disrupting sleep and this reduces the threshold for pain. PTSD disrupts sleep, and the impaired sleep associated with PTSD intensifies pain symptoms¹⁰¹⁻¹⁰⁴. Improving the sleep of US combat veterans who sustained mTBI in Iraq or Afghanistan reduced PTSD severity and decreased the frequency and intensity of PTH episodes^{102,103}. Depression is associated with impaired sleep, and impaired sleep can exacerbate depression¹⁰⁵. Sleep deprivation and depression independently lower the pain threshold¹⁰⁶. Several studies found that sleep deprivation lowers pain thresholds¹⁰⁷⁻¹⁰⁹.

PTSD is associated with chronic pain disorders, and PTSD has complex interactions with pain thresholds¹¹⁰. One report indicated that PTSD increases the pain threshold but that supra-threshold pain was perceived as excessively intense¹¹¹. PTSD disrupts sleep by inducing nightmares^{4,101-103,112-116}. Prazosin, an alpha-1 adrenergic receptor antagonist that passes through the blood-brain barrier, reduces PTSD-associated nightmares^{102,103,112-117}, improves cognitive performance, and reduces PTH frequency and severity^{4,102,103}.

Possible relationships between mild traumatic brain injury and post-traumatic stress disorder

We suggest that cerebral injury associated with mTBI predisposes individuals to develop PTSD in response to psychologically traumatic events associated with the mTBI. A hypothesis on the neural basis of PTSD is that PTSD results from over-activation of the amygdalae due to loss of inhibitory regulation by the ventromedial prefrontal cortex and hippocampus¹¹⁸. Pitman *et al.*¹¹⁹ and Sherin and Nemeroff¹²⁰ discuss the roles of hippocampus and ventromedial prefrontal cortex regulation of the amygdalae in the genesis of PTSD. Brain areas with significantly altered levels of activity identified by functional imaging in PTSD include ventromedial prefrontal and medial temporal lobes¹²¹⁻¹²³ which are damaged in mTBI¹²⁴⁻¹³⁷. Single-photon emission tomography demonstrated reduced benzodiazepine receptor binding in the prefrontal cortex of people with PTSD¹³⁸. Prefrontal cortex activity in response to emotional stimuli is altered in PTSD¹³⁹. In combat-associated TBI with PTSD, PTSD severity was associated with the presence of magnetic resonance imaging lesions involving the prefrontal cortex¹⁴⁰.

In addition to imaging evidence that the ventromedial prefrontal cortex is altered in PTSD, the neurological examination finding of impaired olfaction supports altered function of the ventromedial prefrontal cortex in people with PTSD associated with mTBI^{3,4,89,92}. The initial site of cortical processing of olfactory information is in the ventromedial prefrontal cortex¹⁴¹. Impaired olfaction was the most frequently recognized chronic/persistent neurological deficit on clinical physical examination⁴. TBI compromises olfaction by shearing the olfactory nerves traversing the cribriform plate or ventromedial prefrontal cortex injury or both¹⁴¹. Injured olfactory nerve fibers can recover over time, leading to improvement of olfaction in the first year after TBI¹⁴². Veterans in the study of olfaction

after mTBI were evaluated more than one year after their last TBI; thus, the reported persistent reduction in olfaction likely resulted from damage to the ventromedial prefrontal cortex⁴.

A study of US military personnel who served in Iraq or Afghanistan and sustained combat TBI, predominantly mTBI, reported that 29% of subjects had white matter lesions in at least two areas of interest¹²⁷. The areas of injury were ventromedial prefrontal cortex, cingulum, and middle cerebellar peduncles. The finding of ventromedial prefrontal cortex injury associated with combat mTBI provides a structural explanation for impaired olfaction in combat mTBI^{4,92}. Further support for a structural basis for both impaired olfaction and PTSD with combat mTBI is that both impaired olfaction and the presence and severity of PTSD correlated with the number of mTBI events^{4,92}.

Animal studies suggest that the neural elements associated with PTSD, including the amygdalae, ventromedial prefrontal cortex, and hippocampus, have unique responses to psychological stress that may perpetuate and enhance the severity of PTSD. Placing rats in prolonged stressful environments enhanced dendritic arborization of neurons in the amygdalae^{143,144} and produced dendrite atrophy in the hippocampus¹⁴³. Consequently, continued stress, as might be produced by difficulty functioning within society because of chronic PTH, could lead to structural brain changes that perpetuate and intensify PTSD. For combat veterans, the frequency and severity of PTH are associated with PTSD severity⁴. Consequently, PTH and PTSD could augment each other via cerebral structural changes.

Studies of Vietnam-era veterans show that subtle neurological deficits can increase the risk of developing PTSD¹⁴⁵. In studies of monozygotic twins in which one twin was in combat and the other was not, among twin pairs where one had combat-associated PTSD, both twins had a higher prevalence of subtle neurological deficits¹⁴⁵ and gray matter abnormalities in the right hippocampus, pregenual anterior cingulate cortex, and left and right insular cortex¹⁴⁶. The twin studies showed that subtle, genetically based neurological deficits can potentiate the genesis of PTSD.

The twin studies contrast with the Vietnam Head Injury Program (VHIP), in which veterans with combat penetrating severe TBIs involving the right amygdala or bilateral ventromedial prefrontal cortex had reduced prevalence of PTSD¹¹⁹. One explanation for the differences between the VHIP and twin studies is that the subjects in the VHIP study had severe TBI compared with the subtle deficits of participants in the twin studies. Mild cerebral injury could foster PTSD genesis by impairing normal interactions among the amygdala, ventromedial prefrontal cortex, and hippocampus, whereas the more severe injuries in the VHIP study may prevent “the ‘super-normal’ levels of fear/anxiety that define PTSD”¹¹⁹.

Thus, mild injury to specific areas of the brain may potentiate the genesis of PTSD, whereas obliteration of these areas may prevent PTSD from developing. In childhood TBI, PTSD and other anxiety disorders correlated with mild injury to the superior frontal gyrus, anterior frontal white matter, and ventromedial prefrontal gray matter^{93,123}. Overall, the risk of developing PTSD following combat TBI is greater for mTBI compared with severe TBI¹⁴⁷. Children with mTBI are more likely to develop PTSD than children with severe TBI⁹³.

In summary, TBI can induce PTH by several mechanisms. TBI can induce neuroinflammation that activates structures associated with headache genesis, especially migraine. CNS damage resulting from TBI will impair pain modulation and enhance headache genesis. The CNS injury resulting from TBI can potentiate the development of depression and anxiety disorders, including PTSD that can intensify PTH. The areas of the brain injured in TBI overlap with the areas of impaired function associated with PTSD. The complex nature of PTH suggests that their treatment should include interventions that address psychological issues such as depression and PTSD and impaired sleep and that directly address pain symptoms associated with PTH.

Many questions remain regarding the genesis of PTH. We believe that an important issue to examine is whether some individuals are more susceptible to developing persisting PTH. Psychological conditions and secondary gain can influence the likelihood that an individual will develop PTH¹⁴⁸. Psychological resilience enhances the ability of individuals to recover from psychological trauma and appears to reduce the likelihood that an individual will recover from TBI with prolonged post-concussion symptoms, including PTH¹⁴⁹. The twin studies demonstrated that subtle, genetically based neurological deficits influence the genesis of PTSD^{145,146}. We suggest that genetically based or acquired alterations in brain structure may also influence the likelihood that an individual will develop PTH.

Author contributions

RR developed the initial draft of the manuscript, reviewed relevant literature, and contributed to manuscript revisions. KB contributed to the initial manuscript, literature review, and manuscript revisions.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. Headache Classification Subcommittee of the International Headache Society: **The International Classification of Headache Disorders: 2nd edition.** *Cephalgia*. 2004; 24(Suppl 1): 9–160.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. **F** Nampiaparampil DE: **Prevalence of chronic pain after traumatic brain injury: a systematic review.** *JAMA*. 2008; 300(6): 711–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
3. Ruff RL, Ruff SS, Wang XF: **Headaches among Operation Iraqi Freedom/Operation Enduring Freedom veterans with mild traumatic brain injury associated with exposures to explosions.** *J Rehabil Res Dev*. 2008; 45(7): 941–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Ruff RL, Riechers RG 2nd, Wang XF, *et al.*: **A case-control study examining whether neurological deficits and PTSD in combat veterans are related to episodes of mild TBI.** *BMJ Open*. 2012; 2(2): e000312.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Patil VK, St Andre JR, Crisan E, *et al.*: **Prevalence and treatment of headaches in veterans with mild traumatic brain injury.** *Headache*. 2011; 51(7): 1112–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Theeler BJ, Erickson JC: **Posttraumatic headache in military personnel and veterans of the Iraq and Afghanistan conflicts.** *Curr Treat Options Neurol*. 2012; 14(1): 36–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Lew HL, Lin PH, Fuh JL, *et al.*: **Characteristics and treatment of headache after traumatic brain injury: a focused review.** *Am J Phys Med Rehabil*. 2006; 85(7): 619–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Hoffman JM, Lucas S, Dikmen S, *et al.*: **Natural history of headache after traumatic brain injury.** *J Neurotrauma*. 2011; 28(9): 1719–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Lucas S, Hoffman JM, Bell KR, *et al.*: **Characterization of headache after traumatic brain injury.** *Cephalalgia*. 2012; 32(8): 600–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Theeler BJ, Flynn FG, Erickson JC: **Chronic daily headache in U.S. soldiers after concussion.** *Headache*. 2012; 52(5): 732–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Girona RJ, Clark ME, Ruff RL, *et al.*: **Traumatic brain injury, polytrauma, and pain: challenges and treatment strategies for the polytrauma rehabilitation.** *Rehabil Psychol*. 2009; 54(3): 247–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. **F** Lucas S, Hoffman JM, Bell KR, *et al.*: **A prospective study of prevalence and characterization of headache following mild traumatic brain injury.** *Cephalalgia*. 2014; 34(2): 93–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
13. Theeler BJ, Mercer R, Erickson JC: **Prevalence and impact of migraine among US Army soldiers deployed in support of Operation Iraqi Freedom.** *Headache*. 2008; 48(6): 876–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Theeler BJ, Erickson JC: **Mild head trauma and chronic headaches in returning US soldiers.** *Headache*. 2009; 49(4): 529–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Borkum JM: **Benign Secondary Headaches: Cervicogenic, Temporomandibular Dysfunction, Trigeminal Neuralgia and Posttraumatic Headache.** Chapter 8 in: *Chronic Headaches - Biology, Psychology, and Behavioral Treatment*. Mahwah, NJ, USA: Lawrence Erlbaum Associates, 2007; 101–207.
[Reference Source](#)
16. **F** Wu E, Graham DP: **Association of Chronic Pain and Community Integration of Returning Veterans With and Without Traumatic Brain Injury.** *J Head Trauma Rehabil*. 2016; 31(1): E1–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
17. Packard RC, Ham LP: **Pathogenesis of posttraumatic headache and migraine: a common headache pathway?** *Headache*. 1997; 37(3): 142–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. McKee AC, Cantu RC, Nowinski CJ, *et al.*: **Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury.** *J Neuropathol Exp Neurol*. 2009; 68(7): 709–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Ling H, Hardy J, Zetterberg H: **Neurological consequences of traumatic brain injuries in sports.** *Mol Cell Neurosci*. 2015; 66(Pt B): 114–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. **F** Gyoneva S, Kim D, Katsumoto A, *et al.*: **Ccr2 deletion dissociates cavity size and tau pathology after mild traumatic brain injury.** *J Neuroinflammation*. 2015; 12: 228.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
21. McKee AC, Daneshvar DH, Alvarez VE, *et al.*: **The neuropathology of sport.** *Acta Neuropathol*. 2014; 127(1): 29–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. McKee AC: **Association between Repetitive Head Impacts and Development of Chronic Traumatic Encephalopathy.** *The Bridge*. 2016; 46(1): 56–60.
[Reference Source](#)
23. Lu XM, Chen RW, Yao C, *et al.*: **NNZ-2566, a glypromate analog, improves functional recovery and attenuates apoptosis and inflammation in a rat model of penetrating ballistic-type brain injury.** *J Neurotrauma*. 2009; 26(1): 141–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Wei HH, Lu XM, Shear DA, *et al.*: **NNZ-2566 treatment inhibits neuroinflammation and pro-inflammatory cytokine expression induced by experimental penetrating ballistic-like brain injury in rats.** *J Neuroinflammation*. 2009; 6: 19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Shoji H, Kaneko Y, Mabuchi T, *et al.*: **Genetic and histologic evidence implicates role of inflammation in traumatic brain injury-induced apoptosis in the rat cerebral cortex following moderate fluid percussion injury.** *Neuroscience*. 2010; 171(4): 1273–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Kumar A, Loane DJ: **Neuroinflammation after traumatic brain injury: opportunities for therapeutic intervention.** *Brain Behav Immun*. 2012; 26(8): 1191–201.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. **F** Glushakova OY, Johnson D, Hayes RL: **Delayed increases in microvascular pathology after experimental traumatic brain injury are associated with prolonged inflammation, blood-brain barrier disruption, and progressive white matter damage.** *J Neurotrauma*. 2014; 31(13): 1180–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
28. **F** Shetty AK, Mishra V, Kodali M, *et al.*: **Blood brain barrier dysfunction and delayed neurological deficits in mild traumatic brain injury induced by blast shock waves.** *Front Cell Neurosci*. 2014; 8: 232.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
29. Shultz SR, MacFabe DF, Foley KA, *et al.*: **Sub-concussive brain injury in the Long-Evans rat induces acute neuroinflammation in the absence of behavioral impairments.** *Behav Brain Res*. 2012; 229(1): 145–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Frugier T, Morganti-Kossmann MC, O'Reilly D, *et al.*: **In situ detection of inflammatory mediators in post mortem human brain tissue after traumatic injury.** *J Neurotrauma*. 2010; 27(3): 497–507.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Ha HC, Snyder SH: **Poly(ADP-ribose) polymerase-1 in the nervous system.** *Neurobiol Dis*. 2000; 7(4): 225–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Kauppinen TM, Swanson RA: **The role of poly(ADP-ribose) polymerase-1 in CNS disease.** *Neuroscience*. 2007; 145(4): 1267–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Whalen MJ, Clark RS, Dixon CE, *et al.*: **Reduction of cognitive and motor deficits after traumatic brain injury in mice deficient in poly(ADP-ribose) polymerase.** *J Cereb Blood Flow Metab*. 1999; 19(8): 835–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Satchell MA, Zhang X, Kochanek PM, *et al.*: **A dual role for poly-ADP-ribosylation in spatial memory acquisition after traumatic brain injury in mice involving NAD⁺ depletion and ribosylation of 14-3-3gamma.** *J Neurochem*. 2003; 85(3): 697–708.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Koh DW, Dawson TM, Dawson VL: **Mediation of cell death by poly(ADP-ribose) polymerase-1.** *Pharmacol Res*. 2005; 52(1): 5–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Yu SW, Wang H, Dawson TM, *et al.*: **Poly(ADP-ribose) polymerase-1 and apoptosis inducing factor in neurotoxicity.** *Neurobiol Dis*. 2003; 14(3): 303–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Hassa PO, Hottiger MO: **The functional role of poly(ADP-ribose)polymerase 1 as novel coactivator of NF-kappaB in inflammatory disorders.** *Cell Mol Life Sci*. 2002; 59(9): 1534–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. **F** Tao X, Chen X, Mao X, *et al.*: **Protective effects of PARP inhibitor, PJ34, is related to down-regulation of calpain and NF-κB in a mouse model of TBI.** *Brain Inj*. 2016; 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
39. **F** d'Ávila JC, Lam TI, Bingham D, *et al.*: **Microglial activation induced by brain trauma is suppressed by post-injury treatment with a PARP inhibitor.** *J Neuroinflammation*. 2012; 9: 31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. **F** Stoica BA, Loane DJ, Zhao Z, *et al.*: **PARP-1 inhibition attenuates neuronal loss, microglia activation and neurological deficits after traumatic brain injury.** *J Neurotrauma*. 2014; 31(8): 758–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
41. Byrnes KR, Loane DJ, Stoica BA, *et al.*: **Delayed mGluR5 activation limits neuroinflammation and neurodegeneration after traumatic brain injury.** *J Neuroinflammation*. 2012; 9: 43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Kapadia R, Yi JH, Vemuganti R: **Mechanisms of anti-inflammatory and**

- neuroprotective actions of PPAR-gamma agonists. *Front Biosci.* 2008; 13: 1813–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Gillespie W, Tyagi N, Tyagi SC: **Role of PPARgamma, a nuclear hormone receptor in neuroprotection.** *Indian J Biochem Biophys.* 2011; 48(2): 73–81.
[PubMed Abstract](#)
 44. Hyong A, Jadhav V, Lee S, *et al.*: **Rosiglitazone, a PPAR gamma agonist, attenuates inflammation after surgical brain injury in rodents.** *Brain Res.* 2008; 1215: 218–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Yi JH, Park SW, Brooks N, *et al.*: **PPARgamma agonist rosiglitazone is neuroprotective after traumatic brain injury via anti-inflammatory and anti-oxidative mechanisms.** *Brain Res.* 2008; 1244: 164–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Sauerbeck A, Gao J, Readnower R, *et al.*: **Pioglitazone attenuates mitochondrial dysfunction, cognitive impairment, cortical tissue loss, and inflammation following traumatic brain injury.** *Exp Neurol.* 2011; 227(1): 128–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Semple BD, Noble-Haeusslein LJ: **Broad-spectrum neuroprotection against traumatic brain injury by agonism of peroxisome proliferator-activated receptors.** *Exp Neurol.* 2011; 229(2): 195–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Iwashita A, Muramatsu Y, Yamazaki T, *et al.*: **Neuroprotective efficacy of the peroxisome proliferator-activated receptor delta-selective agonists *in vitro* and *in vivo*.** *J Pharmacol Exp Ther.* 2007; 320(3): 1087–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Schintu N, Frau L, Ibba M, *et al.*: **PPAR-gamma-mediated neuroprotection in a chronic mouse model of Parkinson's disease.** *Eur J Neurosci.* 2009; 29(5): 954–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Garrido-Gil P, Joglar B, Rodriguez-Perez AI, *et al.*: **Involvement of PPAR- γ in the neuroprotective and anti-inflammatory effects of angiotensin type 1 receptor inhibition: effects of the receptor antagonist telmisartan and receptor deletion in a mouse MPTP model of Parkinson's disease.** *J Neuroinflammation.* 2012; 9: 38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. **F** Uppalapati D, Das NR, Gangwal RP, *et al.*: **Neuroprotective Potential of Peroxisome Proliferator Activated Receptor- α Agonist in Cognitive Impairment in Parkinson's Disease: Behavioral, Biochemical, and PBPK Profile.** *PPAR Res.* 2014; 2014: 753587.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 52. Swanson CR, Du E, Johnson DA, *et al.*: **Neuroprotective Properties of a Novel Non-Thiazolinedione Partial PPAR- γ Agonist against MPTP.** *PPAR Res.* 2013; 2013: 582809.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. **F** Joardar A, Menzl J, Podolsky TC, *et al.*: **PPAR gamma activation is neuroprotective in a Drosophila model of ALS based on TDP-43.** *Hum Mol Genet.* 2015; 24(6): 1741–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 54. Landreth GE, Heneka MT: **Anti-inflammatory actions of peroxisome proliferator-activated receptor gamma agonists in Alzheimer's disease.** *Neurobiol Aging.* 2001; 22(6): 937–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. Jin W, Kong J, Wang H, *et al.*: **Protective effect of tert-butylhydroquinone on cerebral inflammatory response following traumatic brain injury in mice.** *Injury.* 2011; 42(7): 714–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 56. Ley EJ, Clond MA, Singer MB, *et al.*: **IL6 deficiency affects function after traumatic brain injury.** *J Surg Res.* 2011; 170(2): 253–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 57. Chiaretti A, Antonelli A, Mastrangelo A, *et al.*: **Interleukin-6 and nerve growth factor upregulation correlates with improved outcome in children with severe traumatic brain injury.** *J Neurotrauma.* 2008; 25(3): 225–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 58. Dai SS, Zhou YG: **Adenosine 2A receptor: a crucial neuromodulator with bidirectional effect in neuroinflammation and brain injury.** *Rev Neurosci.* 2011; 22(2): 231–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 59. Mota BC, Pereira L, Souza MA, *et al.*: **Exercise pre-conditioning reduces brain inflammation and protects against toxicity induced by traumatic brain injury: behavioral and neurochemical approach.** *Neurotox Res.* 2012; 21(2): 175–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Miyauchi T, Wei EP, Povlishock JT: **Therapeutic targeting of the axonal and microvascular change associated with repetitive mild traumatic brain injury.** *J Neurotrauma.* 2013; 30(19): 1664–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Theeler B, Lucas S, Riechers RG 2nd, *et al.*: **Post-traumatic headaches in civilians and military personnel: a comparative, clinical review.** *Headache.* 2013; 53(6): 881–900.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Hains BC, Waxman SG: **Activated microglia contribute to the maintenance of chronic pain after spinal cord injury.** *J Neurosci.* 2006; 26(16): 4308–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Waxman SG, Hains BC: **Fire and phantoms after spinal cord injury: Na⁺ channels and central pain.** *Trends Neurosci.* 2006; 29(4): 207–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
 64. Zhao P, Waxman SG, Hains BC: **Extracellular signal-regulated kinase-regulated microglia-neuron signaling by prostaglandin E₂ contributes to pain after spinal cord injury.** *J Neurosci.* 2007; 27(9): 2357–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. Tepper SJ, Rapoport AM, Sheftell FD: **Mechanisms of action of the 5-HT_{1B/1D} receptor agonists.** *Arch Neurol.* 2002; 59(7): 1084–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Cady RK: **The convergence hypothesis.** *Headache.* 2007; 47(Suppl 1): S44–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Buzzi MG, Moskowitz MA: **The pathophysiology of migraine: year 2005.** *J Headache Pain.* 2005; 6(3): 105–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 68. Geppetti P, Rossi E, Chiarugi A, *et al.*: **Antidromic vasodilatation and the migraine mechanism.** *J Headache Pain.* 2012; 13(2): 103–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. Levy D: **Endogenous mechanisms underlying the activation and sensitization of meningeal nociceptors: the role of immuno-vascular interactions and cortical spreading depression.** *Curr Pain Headache Rep.* 2012; 16(3): 270–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. Raddant AC, Russo AF: **Calcitonin gene-related peptide in migraine: intersection of peripheral inflammation and central modulation.** *Expert Rev Mol Med.* 2011; 13: e36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. **F** Loyd DR, Weiss G, Henry MA, *et al.*: **Serotonin increases the functional activity of capsaicin-sensitive rat trigeminal nociceptors via peripheral serotonin receptors.** *Pain.* 2011; 152(10): 2267–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 72. Ossipov MH, Morimura K, Porreca F: **Descending pain modulation and chronification of pain.** *Curr Opin Support Palliat Care.* 2014; 8(2): 143–51.
[PubMed Abstract](#) | [Free Full Text](#)
 73. **F** Defrin R, Riabinin M, Feingold Y, *et al.*: **Deficient pain modulatory systems in patients with mild traumatic brain and chronic post-traumatic headache: implications for its mechanism.** *J Neurotrauma.* 2015; 32(1): 28–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 74. Loder E: **Triptan therapy in migraine.** *N Engl J Med.* 2010; 363(1): 63–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. Hoskin KL, Lambert GA, Donaldson C, *et al.*: **The 5-hydroxytryptamine_{1B/1D/1F} receptor agonists eletriptan and naratriptan inhibit trigeminovascular input to the nucleus tractus solitarius in the cat.** *Brain Res.* 2004; 998(1): 91–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 76. Andriessen TM, Jacobs B, Vos PE, *et al.*: **Clinical characteristics and pathophysiological mechanisms of focal and diffuse traumatic brain injury.** *J Cell Mol Med.* 2010; 14(10): 2381–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Shumskaya E, Andriessen TM, Norris DG, *et al.*: **Abnormal whole-brain functional networks in homogeneous acute mild traumatic brain injury.** *Neurology.* 2012; 79(2): 175–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. **F** Ghodadra A, Alhilali L, Fakhran S: **Principal Component Analysis of Diffusion Tensor Images to Determine White Matter Injury Patterns Underlying Postconcussive Headache.** *AJNR Am J Neuroradiol.* 2016; 37(2): 274–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 79. Azmitia EC: **Serotonin and brain: evolution, neuroplasticity, and homeostasis.** *Int Rev Neurobiol.* 2007; 77: 31–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
 80. Eghwudjakpor PO, Miyake H, Kurisaka M, *et al.*: **Central nervous system bioaminergic responses to mechanical trauma. An experimental study.** *Surg Neurol.* 1991; 35(4): 273–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. Busto R, Dietrich WD, Globus MY, *et al.*: **Extracellular release of serotonin following fluid-percussion brain injury in rats.** *J Neurotrauma.* 1997; 14(1): 35–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Ueda Y, Walker SA, Povlishock JT: **Perivascular nerve damage in the cerebral circulation following traumatic brain injury.** *Acta Neuropathol.* 2006; 112(1): 85–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Sharma HS, Winkler T, Stalberg E, *et al.*: **p-Chlorophenylalanine, an inhibitor of serotonin synthesis reduces blood-brain barrier permeability, cerebral blood flow, edema formation and cell injury following trauma to the rat brain.** *Acta Neurochir Suppl.* 2000; 76: 91–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Wester P, Bergstrom U, Eriksson A, *et al.*: **Ventricular cerebrospinal fluid monoamine transmitter and metabolite concentrations reflect human brain neurochemistry in autopsy cases.** *J Neurochem.* 1990; 54(4): 1148–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
 85. Meythaler JM, Depalma L, Devivo MJ, *et al.*: **Sertraline to improve arousal and alertness in severe traumatic brain injury secondary to motor vehicle crashes.** *Brain Inj.* 2001; 15(4): 321–31.
[PubMed Abstract](#) | [Publisher Full Text](#)

86. **F** Smitherman TA, Kolivas ED: **Trauma exposure versus posttraumatic stress disorder: relative associations with migraine.** *Headache*. 2013; 53(5): 775–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
87. Ruff RL, Riechers RG, Ruff SS: **Relationships between mild traumatic brain injury sustained in combat and post-traumatic stress disorder.** *F1000 Med Rep*. 2010; 2: 64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. Elder GA, Dorr NP, De Gasperi R, *et al.*: **Blast exposure induces post-traumatic stress disorder-related traits in a rat model of mild traumatic brain injury.** *J Neurotrauma*. 2012; 29(16): 2564–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
89. Kontos AP, Kotwal RS, Elbin RJ, *et al.*: **Residual effects of combat-related mild traumatic brain injury.** *J Neurotrauma*. 2013; 30(8): 680–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
90. Ruff RL: **Impaired Olfaction and Other Indicators of Neurological Damage Due to Mild TBI Associated with Combat.** *The Bridge*. 2016; 46(1): 76–78.
[Reference Source](#)
91. **F** Max JE, Keatley E, Wilde EA, *et al.*: **Anxiety disorders in children and adolescents in the first six months after traumatic brain injury.** *J Neuropsychiatry Clin Neurosci*. 2011; 23(1): 29–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
92. Bryant RA, O'Donnell ML, Creamer M, *et al.*: **The psychiatric sequelae of traumatic injury.** *Am J Psychiatry*. 2010; 167(3): 312–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
93. **F** Hoge CW, McGurk D, Thomas JL, *et al.*: **Mild traumatic brain injury in U.S. Soldiers returning from Iraq.** *N Engl J Med*. 2008; 358(5): 453–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
94. de Leeuw R, Schmidt JE, Carlson CR: **Traumatic stressors and post-traumatic stress disorder symptoms in headache patients.** *Headache*. 2005; 45(10): 1365–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
95. **F** Jaramillo CA, Eapen BC, McGeary CA, *et al.*: **A cohort study examining headaches among veterans of Iraq and Afghanistan wars: Associations with traumatic brain injury, PTSD, and depression.** *Headache*. 2016; 56(3): 528–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
96. Breslau N, Lipton RB, Stewart WF, *et al.*: **Comorbidity of migraine and depression: investigating potential etiology and prognosis.** *Neurology*. 2003; 60(8): 1308–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. Zwart J, Dyb G, Hagen K, *et al.*: **Depression and anxiety disorders associated with headache frequency. The Nord-Trøndelag Health Study.** *Eur J Neurol*. 2003; 10(2): 147–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
98. Chibnall JT, Duckro PN: **Post-traumatic stress disorder in chronic post-traumatic headache patients.** *Headache*. 1994; 34(6): 357–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
99. Ursano RJ, Fullerton CS, Epstein RS, *et al.*: **Acute and chronic posttraumatic stress disorder in motor vehicle accident victims.** *Am J Psychiatry*. 1999; 156(4): 589–95.
[PubMed Abstract](#)
100. **F** Peterlin BL, Nijjar SS, Tietjen GE: **Post-traumatic stress disorder and migraine: epidemiology, sex differences, and potential mechanisms.** *Headache*. 2011; 51(6): 860–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
101. Schoenfeld FB, Deviva JC, Manber R: **Treatment of sleep disturbances in posttraumatic stress disorder: a review.** *J Rehabil Res Dev*. 2012; 49(5): 729–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
102. Ruff RL, Ruff SS, Wang XF: **Improving sleep: initial headache treatment in OIF/OEF veterans with blast-induced mild traumatic brain injury.** *J Rehabil Res Dev*. 2009; 46(9): 1071–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
103. Ruff RL, Riechers RG 2nd, Wang XF, *et al.*: **For veterans with mild traumatic brain injury, improved posttraumatic stress disorder severity and sleep correlated with symptomatic improvement.** *J Rehabil Res Dev*. 2012; 49(9): 1305–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
104. **F** Lang KP, Veazey-Morris K, Andrasik F: **Exploring the role of insomnia in the relation between PTSD and pain in veterans with polytrauma injuries.** *J Head Trauma Rehabil*. 2014; 29(1): 44–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
105. Roberts RE, Shema SJ, Kaplan GA, *et al.*: **Sleep complaints and depression in an aging cohort: A prospective perspective.** *Am J Psychiatry*. 2000; 157(1): 81–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
106. Chiu YH, Silman AJ, Macfarlane GJ, *et al.*: **Poor sleep and depression are independently associated with a reduced pain threshold. Results of a population based study.** *Pain*. 2005; 115(3): 316–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
107. Onen SH, Alloui A, Gross A, *et al.*: **The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects.** *J Sleep Res*. 2001; 10(1): 35–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Lentz MJ, Landis CA, Rothermel J, *et al.*: **Effects of selective slow wave sleep disruption on musculoskeletal pain and fatigue in middle aged women.** *J Rheumatol*. 1999; 26(7): 1586–92.
[PubMed Abstract](#)
109. Kundermann B, Spornal J, Huber MT, *et al.*: **Sleep deprivation affects thermal pain thresholds but not somatosensory thresholds in healthy volunteers.** *Psychosom Med*. 2004; 66(6): 932–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
110. Moeller-Bertram T, Keltner J, Strigo IA: **Pain and post traumatic stress disorder - review of clinical and experimental evidence.** *Neuropharmacology*. 2012; 62(2): 586–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
111. Defrin R, Ginzburg K, Solomon Z, *et al.*: **Quantitative testing of pain perception in subjects with PTSD—implications for the mechanism of the coexistence between PTSD and chronic pain.** *Pain*. 2008; 138(2): 450–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Daly CM, Doyle ME, Radkind M, *et al.*: **Clinical case series: the use of Prazosin for combat-related recurrent nightmares among Operation Iraqi Freedom combat veterans.** *Mil Med*. 2005; 170(6): 513–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
113. Raskind MA, Peskind ER, Hoff DJ, *et al.*: **A parallel group placebo controlled study of prazosin for trauma nightmares and sleep disturbance in combat veterans with post-traumatic stress disorder.** *Biol Psychiatry*. 2007; 61(8): 928–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
114. Taylor FB, Martin P, Thompson C, *et al.*: **Prazosin effects on objective sleep measures and clinical symptoms in civilian trauma posttraumatic stress disorder: a placebo-controlled study.** *Biol Psychiatry*. 2008; 63(6): 629–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
115. Calohan J, Peterson K, Peskind ER, *et al.*: **Prazosin treatment of trauma nightmares and sleep disturbance in soldiers deployed in Iraq.** *J Trauma Stress*. 2010; 23(5): 645–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
116. Raskind MA, Peskind ER, Kanter ED, *et al.*: **Reduction of nightmares and other PTSD symptoms in combat veterans by prazosin: a placebo-controlled study.** *Am J Psychiatry*. 2003; 160(2): 371–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
117. Boynton L, Bentley J, Strachan E, *et al.*: **Preliminary findings concerning the use of prazosin for the treatment of posttraumatic nightmares in a refugee population.** *J Psychiatr Pract*. 2009; 15(6): 454–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
118. **F** Koenigs M, Huey ED, Raymond V, *et al.*: **Focal brain damage protects against post-traumatic stress disorder in combat veterans.** *Nat Neurosci*. 2008; 11(2): 232–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
119. Pitman RK, Rasmusson AM, Koenen KC, *et al.*: **Biological studies of post-traumatic stress disorder.** *Nat Rev Neurosci*. 2012; 13(11): 769–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
120. Sherin JE, Nemeroff CB: **Post-traumatic stress disorder: the neurobiological impact of psychological trauma.** *Dialogues Clin Neurosci*. 2011; 13(3): 263–78.
[PubMed Abstract](#) | [Free Full Text](#)
121. Whalley MG, Rugg MD, Smith AP, *et al.*: **Incidental retrieval of emotional contexts in post-traumatic stress disorder and depression: an fMRI study.** *Brain Cogn*. 2009; 69(1): 98–107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
122. Sailer U, Robinson S, Fischmeister FP, *et al.*: **Altered reward processing in the nucleus accumbens and mesial prefrontal cortex of patients with posttraumatic stress disorder.** *Neuropsychologia*. 2008; 46(11): 2836–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
123. Carrion VG, Weems CF, Watson C, *et al.*: **Converging evidence for abnormalities of the prefrontal cortex and evaluation of midsagittal structures in pediatric posttraumatic stress disorder: an MRI study.** *Psychiatry Res*. 2009; 172(3): 226–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
124. Topal NB, Hakyemez B, Erdogan C, *et al.*: **MR imaging in the detection of diffuse axonal injury with mild traumatic brain injury.** *Neurol Res*. 2008; 30(9): 974–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
125. Lo C, Shifteh K, Gold T, *et al.*: **Diffusion tensor imaging abnormalities in patients with mild traumatic brain injury and neurocognitive impairment.** *J Comput Assist Tomogr*. 2009; 33(2): 293–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
126. Peskind ER, Petrie EC, Cross DJ, *et al.*: **Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms.** *Neuroimage*. 2011; 54(Suppl 1): S76–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
127. **F** MacDonald CL, Johnson AM, Cooper D, *et al.*: **Detection of blast-related traumatic brain injury in U.S. military personnel.** *N Engl J Med*. 2011; 364(22): 2091–100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
128. Baykara B, Aksu I, Buyuk E, *et al.*: **Progesterone treatment decreases traumatic brain injury induced anxiety and is correlated with increased serum IGF-1 levels; prefrontal cortex, amygdala, hippocampus neuron density; and reduced**

- serum corticosterone levels in immature rats. *Biotech Histochem.* 2013; **88**(5): 250–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
129. Baykara B, Cetin F, Baykara B, *et al.*: Anxiety caused by traumatic brain injury correlates to decreased prefrontal cortex VEGF immunoreactivity and neuron density in immature rats. *Turk Neurosurg.* 2012; **22**(5): 604–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
 130. Maxwell WL, MacKinnon MA, Stewart JE, *et al.*: Stereology of cerebral cortex after traumatic brain injury matched to the Glasgow outcome score. *Brain.* 2010; **133**(Pt 1): 139–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 131. Logue OC, Cramer NP, Xu X, *et al.*: Alterations of functional properties of hippocampal networks following repetitive closed-head injury. *Exp Neurol.* 2016; **277**: 227–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 132. Carlson SW, Madathil SK, Sama DM, *et al.*: Conditional overexpression of insulin-like growth factor-1 enhances hippocampal neurogenesis and restores immature neuron dendritic processes after traumatic brain injury. *J Neuropathol Exp Neurol.* 2014; **73**(8): 734–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 133. Casella EM, Thomas TC, Vanino DL, *et al.*: Traumatic brain injury alters long-term hippocampal neuron morphology in juvenile, but not immature, rats. *Childs Nerv Syst.* 2014; **30**(8): 1333–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 134. Zhou H, Chen L, Gao X, *et al.*: Moderate traumatic brain injury triggers rapid necrotic death of immature neurons in the hippocampus. *J Neuropathol Exp Neurol.* 2012; **71**(4): 348–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 135. Smith DH, Chen XH, Pierce JE, *et al.*: Progressive atrophy and neuron death for one year following brain trauma in the rat. *J Neurotrauma.* 1997; **14**(10): 715–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
 136. Saatman KE, Feeko KJ, Pape RL, *et al.*: Differential behavioral and histopathological responses to graded cortical impact injury in mice. *J Neurotrauma.* 2006; **23**(8): 1241–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
 137. Tong W, Igarashi T, Ferriero DM, *et al.*: Traumatic brain injury in the immature mouse brain: characterization of regional vulnerability. *Exp Neurol.* 2002; **176**(1): 105–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
 138. Bremner JD, Innis RB, Southwick SM, *et al.*: Decreased benzodiazepine receptor binding in prefrontal cortex in combat-related posttraumatic stress disorder. *Am J Psychiatry.* 2000; **157**(7): 1120–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 139. Aupperle RL, Allard CB, Grimes EM, *et al.*: Dorsolateral prefrontal cortex activation during emotional anticipation and neuropsychological performance in posttraumatic stress disorder. *Arch Gen Psychiatry.* 2012; **69**(4): 360–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 140. Bazarian JJ, Donnelly K, Peterson DR, *et al.*: The relation between posttraumatic stress disorder and mild traumatic brain injury acquired during Operations Enduring Freedom and Iraqi Freedom. *J Head Trauma Rehabil.* 2013; **28**(1): 1–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
 141. Kern RC, Quinn B, Rosseau G, *et al.*: Post-traumatic olfactory dysfunction. *Laryngoscope.* 2000; **110**(12): 2106–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 142. Doty RL, Yousem DM, Pham LT, *et al.*: Olfactory dysfunction in patients with head trauma. *Arch Neurol.* 1997; **54**(9): 1131–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 143. Vyas A, Mitra R, Shankaranarayana Rao BS, *et al.*: Chronic stress induces contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. *J Neurosci.* 2002; **22**(15): 6810–8.
[PubMed Abstract](#) | [F1000 Recommendation](#)
 144. Mitra R, Jadhav S, McEwen BS, *et al.*: Stress duration modulates the spatiotemporal patterns of spine formation in the basolateral amygdala. *Proc Natl Acad Sci U S A.* 2005; **102**(26): 9371–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 145. Gurvits TV, Metzger LJ, Lasko NB, *et al.*: Subtle neurologic compromise as a vulnerability factor for combat-related posttraumatic stress disorder: results of a twin study. *Arch Gen Psychiatry.* 2006; **63**(5): 571–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 146. Kasai K, Yamasue H, Gilbertson MW, *et al.*: Evidence for acquired pregenual anterior cingulate gray matter loss from a twin study of combat-related posttraumatic stress disorder. *Biol Psychiatry.* 2008; **63**(6): 550–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 147. Carlson K, Kehle S, Meis L, *et al.*: The Assessment and Treatment of Individuals with History of Traumatic Brain Injury and Post-Traumatic Stress Disorder: A Systematic Review of the Evidence [Internet]. Washington DC: Department of Veterans Affairs (US). 2009.
[PubMed Abstract](#)
 148. Evans RW: Post-traumatic headaches. *Neurol Clin.* 2004; **22**(1): 237–49, viii.
[PubMed Abstract](#) | [Publisher Full Text](#)
 149. McCauley SR, Wilde EA, Miller ER, *et al.*: Preinjury resilience and mood as predictors of early outcome following mild traumatic brain injury. *J Neurotrauma.* 2013; **30**(8): 642–52.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Ross Zafonte**

Department of Physical Medicine and Rehabilitation, Harvard Medical School; Spaulding Rehabilitation Hospital; Brigham and Women's Hospital, Boston, MA, USA

Competing Interests: No competing interests were disclosed.

2. **Mary Jo Pugh**

Department of Epidemiology and Biostatistics, University of Texas Health Science Center, San Antonio, TX, USA

Carlos Jaramillo

Physical Medicine and Rehabilitation, South Texas Veterans Health Care System, Texas, USA

Competing Interests: No competing interests were disclosed.

3. **Karen Barlow**

Department of Pediatrics and Clinical Neurosciences, Cumming School of Medicine, University of Calgary, Calgary, Canada

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research