Check for updates

OPINION ARTICLE

Clonal selection versus clonal cooperation: the integrated perception of immune objects [version 1; peer review: 2 approved]

Serge Nataf

Bank of Tissues and Cells, Lyon University Hospital (Hospices Civils de Lyon), CarMeN Laboratory, INSERM 1060, INRA 1397, INSA Lyon, Université Claude Bernard Lyon-1, University Lyon-1, Lyon, France

V1 First published: 05 Sep 2016, 5:2226
<https://doi.org/10.12688/f1000research.9386.1>

Latest published: 05 Sep 2016, 5:2226
<https://doi.org/10.12688/f1000research.9386.1>

Abstract

Analogies between the immune and nervous systems were first envisioned by the immunologist Niels Jerne who introduced the concepts of antigen "recognition" and immune "memory". However, since then, it appears that only the cognitive immunology paradigm proposed by Irun Cohen, attempted to further theorize the immune system functions through the prism of neurosciences. The present paper is aimed at revisiting this analogy-based reasoning. In particular, a parallel is drawn between the brain pathways of visual perception and the processes allowing the global perception of an "immune object". Thus, in the visual system, distinct features of a visual object (shape, color, motion) are perceived separately by distinct neuronal populations during a primary perception task. The output signals generated during this first step instruct then an integrated perception task performed by other neuronal networks. Such a higher order perception step is by essence a cooperative task that is mandatory for the global perception of visual objects. Based on a re-interpretation of recent experimental data, it is suggested that similar general principles drive the integrated perception of immune objects in secondary lymphoid organs (SLOs). In this scheme, the four main categories of signals characterizing an immune object (antigenic, contextual, temporal and localization signals) are first perceived separately by distinct networks of immunocompetent cells. Then, in a multitude of SLO niches, the output signals generated during this primary perception step are integrated by TH-cells at the single cell level. This process eventually generates a multitude of T-cell and B-cell clones that perform, at the scale of SLOs, an integrated perception of immune objects. Overall, this new framework proposes that integrated immune perception and, consequently, integrated immune responses, rely essentially on clonal cooperation rather than clonal selection.

Open Peer Review

Approval Status

	1	2
version 1
05 Sep 2016	view	view

1. **Irun R. Cohen** , Weizmann Institute of Science, Rehovot, Israel

2. **Alfred I. Tauber**, Boston University, Boston, USA

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

theoretical immunology , neuroimmunology , sensory perception , immunity , brain , T-cells

Corresponding author: Serge Nataf (serge.nataf@inserm.fr)

Competing interests: No competing interests were disclosed.

Grant information: This work is supported by the Lyon-1 University and the Lyon University Hospital (Hospices Civils de Lyon), Lyon, France.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Nataf S. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Nataf S. **Clonal selection versus clonal cooperation: the integrated perception of immune objects [version 1; peer review: 2 approved]** F1000Research 2016, 5:2226 <https://doi.org/10.12688/f1000research.9386.1>

First published: 05 Sep 2016, 5:2226 <https://doi.org/10.12688/f1000research.9386.1>

Introduction

Evolution has endowed the human species with the most sophisticated immune and nervous systems. Maintenance of our internal homeostasis and adaptation to our external environment rely essentially on the ability of both systems to sense, memorize and react to a large variety of input signals. These crucial functions are supported by a common organizational grounding base consisting in complex networks of specialized cells that communicate in specific anatomical sites. Similarities between the immune and nervous systems were first highlighted by the immunologist Niels Jerne who introduced the terms “recognition”, “memory” and “learning” in the immunological vocabulary^{1,2}. However, since then, such an analogy-based reasoning was mostly used to demonstrate reminiscent molecular mechanisms between immune and neuronal synapses^{3,4}. Only few works attempted to further theorize the immune system functions through the prism of neurosciences. In this framework, the cognitive immunology paradigm proposed by Irun Cohen^{5,6} is undeniably a key contribution that notably led to the concept of physiological auto-immunity^{7,8}. In particular, Irun Cohen proposed that naturally occurring auto-antibodies provide an indispensable immune system’s representation of our body, the immunological *homunculus*⁵, which resembles its neural counterparts, the somatosensory *homunculus*. Thereafter, other works similarly apprehended immunity as a cognitive process and brought about the emergence of computational immunology^{9,10}. Nevertheless, the line of thought initiated by Jerne appears not to have been nourished by the major conceptual and experimental advances that cognitive neurosciences provided in the last two decades. This context offers a unique opportunity to revisit and explore analogies between the nervous and immune systems in the light of such discoveries. The recently formulated concept of a sensory immune system¹¹ falls into this re-thinking strategy.

Cooperative neuronal networks support the integrated perception of visual objects

Developing further the concept of “perceptive immunity” requires beforehand to provide a basic description of the main mechanisms allowing our brain to perceive sensory inputs. Let us choose the example of visual perception. When considering the perception of a given visual object, different categories of input signals that relate with the shape, color, and motion of this object are captured and integrated independently by distinct neuronal populations. These specialized neuronal networks reside in the so-called primary visual cortex, in the superficial neuronal layers of the brain occipital lobe^{12,13}. Importantly, such a **primary perception** induces the generation of output primary signals (electro-chemical by nature) that converge toward neurons localized in the so-called visual association areas also named higher-order visual areas^{14,15} (Figure 1). There, these specialized neuronal populations capture and integrate varied combinations of output primary signals to perform an **integrated perception** of visual objects. Two major pathways allowing the convergence of output primary signals toward higher-order areas are well characterized: i) the “What” pathway targets associative areas in the temporal cortex and is essential to the recognition and memorization of visual objects¹⁴, ii) the “Where” pathway targets associative areas of the parietal cortex and supports the perception of precise localization¹⁴. Eventually, the interconnections between high-order visual areas allows a fully-integrated perception that takes into account the nature, localization, context and time-related features of a visual object¹⁴.

Thus, visual perception requires not only a specialization of cells depending on their ability to perform primary vs integrated perception tasks but also tight cooperation between neuronal networks. **Primary perception** allows distinct features of a visual object

Figure 1. Brain visual pathways. The signals related to the shape, color and motion of visual objects are integrated by specialized brain neuronal populations residing in the primary visual cortex i.e the superficial neuronal layers of the brain occipital lobe. Output signals are then generated that instruct other cortical areas for higher order integration tasks. The “What” pathway connects the primary visual cortex to areas of the temporal cortex that are essential to the recognition and memorization of objects and forms. The “Where” pathway connects the primary visual cortex to areas of the parietal cortex that support perception of precise localization. The interconnections between higher-order visual areas (dashed arrows) as well as other brain areas not highlighted here, allows a fully-integrated perception that takes into account the “What”, “Where”, “How” and “When” features of a visual object.

(shape, color, motion) to be perceived separately^{12,13}. **Integrated perception** allows a visual object to be perceived as a whole via the integration of distinct categories of primary signals^{14,15}.

Non-neuronal cells are also essential to visual perception

It also important to underscore that the operability of any neuronal network, would it be involved or not in sensory perception, depends on non-neuronal cells that locate in close vicinity to neurons. Astrocytes exert a tight control of interneuronal synaptic transmission^{16,17} and microglia, the resident macrophages of the brain, proceed to a selective trimming of functionally irrelevant or supernumerary synapses^{18,19}. In addition, the blood flow in small arteries and capillaries of the brain is exquisitely tuned by a mechanism of neurovascular coupling that finely adjusts the supply of blood-derived oxygen and glucose to the needs of neuronal networks^{20,21}.

The immune system perceives more than antigens and danger signals: a proposed definition of immune objects

Postulating the existence of analogies between the immune and visual systems implies first that the counterpart of visual objects are immune objects. If so, immune objects cannot be simply reduced to an antigen +/- danger signals. Indeed, in accordance with the

principles of immunogenicity previously enunciated by Rolf Zinkernagel²², one may propose that an immune object (IO) is defined by the association of at least 4 categories of signals: antigenic, contextual, temporal and localization signals. Establishing a parallel between visual and immune perception also implies that the perception of an IO relies first on **primary perception** tasks followed by an **integrated perception** step. It is proposed that distinct IO-related features (antigen, context, localization, time-related signals) are perceived separately by distinct networks of immunocompetent cells in a myriad of SLO niches. Then, output signals generated by such a primary perception step converge toward T-cells which, at the scale of SLOs, perform and orchestrate the integrated perception of IOs.

The primary perception of immune objects is a cooperative task operated by a large range of immunocompetent cells

As previously proposed¹¹, SLOs are likely to be the main anatomical sites where the primary and integrated perception of IOs take place. Below is an attempt to categorize immunocompetent cells according to their functions in the **primary perception of IOs** and the ensuing generation of primary output signals (Figure 2).

- 1) The primary perception of antigenic signals is essentially performed by antigen-presenting cells (APCs) would

Figure 2. Primary perception of antigenic, contextual, localization, and temporal signals. The antigenic, temporo-contextual and localization signals that characterize an immune object are perceived separately by specific populations of cells that perform a primary perception task. The output signals generated by these cells will then instruct an integrated perception step essentially performed by TH-cells. The primary perception of antigenic signals is performed by DCs, macrophages, B-cells and any APC that may reside or migrate in lymph nodes. The primary perception of contextual and temporal signals including notably DAMP, PAMP and cytokines are performed by a large range of immune cells or non-immune cells that reside in SLOs. In addition, immune cells that target SLOs in a context- and time-dependent fashion also perform a primary perception of temporo-contextual signals. These cells include notably Treg and Breg cells irrespective of their target antigens as well as, to some extent, naïve B- or T-cells. Finally, the primary perception of localization signals is essentially performed by APCs that derive from the immune object microenvironment.

they belong or not to the dendritic cells (DC) lineage²³. While tissue-resident DCs are the first line cells exerting such a function, a flurry of APCs that reside or migrate in SLOs also participate to the primary perception of antigens. Depending on the intrinsic properties harbored by APCs with regard to antigen processing and co-stimulation, the primary perception of antigens will result in the presentation of distinct epitopes and the expression of varied combinations of accessory molecules²³.

- 2) Contextual signals are highly diverse in nature and may combine in many different ways under physiological or pathological conditions (development, ageing, trauma, degeneration, infection, cancer...etc.). Danger-associated molecular patterns (DAMP), pathogen-associated molecular patterns (PAMP) and cytokines, which form the great majority of contextual danger signals, bind receptors harboring a large expression pattern in SLOs. The primary perception of contextual signals is thus likely to involve not only immune cells but also SLO-residing endothelial cells and stromal cells²⁴⁻²⁶. Output primary signals consist in a larger array of cell surface and soluble factors that instruct the integrated perception step. Moreover, a variety of immune cells that target SLOs in a context-dependent fashion participate to the primary perception of contextual signals and provide part of the primary response to such signals. These include notably Treg and Breg cells irrespective of their antigen specificity^{27,28}, NK cells²⁹, polymorphonuclear cells³⁰, monocytes^{31,32}, innate lymphoid cells^{33,34} as well as naive T or B lymphocytes^{35,36}. Overall, the primary perception of contextual signals is a cooperative task performed by a large array of cell types that reside in SLOs or migrate toward SLOs. These cells generate of whole of soluble or membranous output signals that instruct the integrated perception of IOs.
- 3) The primary perception of localization signals is essentially performed by dendritic cells and macrophages that are drained from the IO's tissue environment³⁷⁻⁴⁰. The output signals generated by these tissue-derived APCs will imprint the homing properties of T-cells^{38,39,41} and orientate in a tissue-specific manner the polarization of TH cells (notably toward TFH cells)⁴². Interestingly, recent findings indicate that stromal cells also play a major role in the primary perception of localization signals^{43,44}.
- 4) Temporal cues are provided by the duration of the antigenic, contextual and/or localization signals. Sudden vs chronically-installed IOs are not equally seen by the immune system according to the discontinuity theory⁴⁵. However, one may also consider that the distinct patterns of tissue-derived cytokines characterizing acute vs subacute vs chronic inflammation provide crucial

time-related inputs. In any case, as proposed above for contextual signals, the primary perception of temporal signals is a cooperative task performed by a large array of cell types in SLOs.

The integrated perception of immune objects is essentially performed by TH-cells

The highly complex spatial organization of SLOs, essentially determined by stromal cells^{24,46-48}, is currently viewed as a means to tightly control the movements of cells and fluids in SLOs^{49,50}. Such a stromal scaffold formed by endothelial cells and fibroblastic reticular cells also provides a histologic support to a number of niches that exhibit distinct microenvironments. For a given IO, there is indeed a myriad of APCs (subsets of DCs, macrophages, B-cells, other APCs) that interact with naive or central memory T-cells in specific niches localized, for lymph nodes, in the paracortical⁵¹, subcapsular^{52,53} or medullar zone^{54,55}. Such niches are formed by partially overlapping yet distinct compositions of immunocompetent cells that proceed to the primary perception of antigenic, contextual, localization and temporal signals. It can be proposed that in each of these niches, immunocompetent cells having proceeded to the primary perception of an IO generates output signals that converge toward T-cells bearing cognate TCRs. By this mean, T-cells (TH cells or T cytotoxic CD8 T-cells activated via cross-presentation) capture and integrate at the single cell level a whole of signals that relate with the antigenic, contextual, localization and temporal features of an IO.

The integrated perception of immune objects relies on clonal cooperation rather than clonal selection

The multitude of niche-specific integration processes that are performed at the single cell level in SLOs is likely to generate a variety of T-cell subpopulations harboring distinct functional behaviors (TH1, TH2, TH17, TFH, Treg...) and recognizing distinct immunodominant epitopes (Figure 3 and Figure 4). Supporting this view, two recent important studies definitively demonstrated that: i) CD4 T cells primed *in vivo* by pathogens or vaccines are highly heterogeneous with regard to TCRs and TH profiles⁵⁶, ii) germinal center reactions in response to complex antigens generate a highly diverse B-cell population in terms of BCR affinity⁵⁷. Thus, at the scale of a SLO the integrated perception of an IO relies on a multitude of antigen-specific T-cell and B-cell clones that provide a whole of "angles of view" from the same immune object. Whether or not such a diversity is progressively narrowed during the re-occurrence of an immune object (i.e during recall immune perception and recall immune response) would require investigations.

It is now recognized that a relatively high level of functional plasticity is maintained in transcriptionally committed TH cell subsets⁵⁸⁻⁶² as well as in CD8 cytotoxic T-cells⁶³ and B-cells⁶⁴. While SLOs orchestrate an integrated immune perception of IOs, one may postulate that higher order integration steps may take place in the efferent lymphatic system. This process would rely on interclonal

Figure 3. Distinct SLO niches provide distinct combinations of output signals resulting from primary perception tasks. Epitopes presented in the context of MHC molecules are the output signals resulting from the primary perception of antigens. In SLOs, multiple epitopes derived from an immune object are recognized by multiple epitope-specific T cells in distinct niches. These niches are formed by stromal cells (endothelial cells and fibroblastic reticular cells) and by partially overlapping combinations of immune cells performing the primary perception antigens, temporo-contextual signals and localization signals. Three examples of distinct SLO niches are depicted.

Figure 4. Integrated immune perception at the scale of SLOs. The interaction between APCs and epitope-specific T-cells occur in a multitude of niches that provide distinct combinations of output signals from primary perception tasks. In each niche, these output signals are integrated at the single cell level by epitope-specific T-cells. This step results in the generation of a large array of TH or CD8 T-cell subsets that possibly include anergic T-cells. Eventually, the integrated perception of IOs is performed at the scale of SLOs by a large variety of antigen-specific T-cell and B-cell clones. Only TH-cell clones are depicted in the diagram.

communications leading possibly to a functional “cross-imprinting” of TH cells (Figure 5).

The integrated perception of immune objects is performed at the scale of SLOs

At the single cell level, a large number of immune cells may integrate antigenic, temporo-contextual and/or localization signals within or outside SLOs. Besides TH cells and CD8 T-cells, these include B-cells and plasma cells as well as NKT-cells and $\gamma\delta$ T-cells. Moreover, recent findings indicate that innate myeloid or lymphoid cells may also integrate and memorize distinct categories of primary immune signals⁶⁵. However, the immune perception theory sheds a new light on the obvious although frequently neglected statement that SLOs are indispensable to the generation of any integrated immune response and, in the context of perceptive immunity, any integrated immune perception. Indeed SLOs harbor a unique

ability to: i) concentrate a large array of cells involved in primary integration tasks, ii) provide a multitude of niches for single cell integration processes.

Immune perception orientates and adjusts decision making in the immune system

The sensory nervous system allows perceiving as a whole the identity and nature of visual objects, their precise localization, visual context and time-related features (motion, memory traces). Visual perception and other facets of our sensory skills are functionally crucial in the orientation of decision making. Such an orientation may schematically follow three main axes: 1) neglect, 2) engage a neurocognitive activity (memorization, attention, thoughts, emotions...), 3) engage a motor activity (grasp, repel, approach, flee...) (Figure 6). Of note, visual perception is a dynamic process that not only orientates but continuously adjusts decision making. Thus,

Figure 5. Higher order integrated immune perception. Higher order integration tasks may take place in the efferent lymphatic system. This process is proposed to rely on interclonal communications leading to a functional “cross-imprinting” of TH cells.

Figure 6. Functional diagram of the visual system. Distinct primary perception tasks allow the shape, color and motion of a visual object to be perceived separately. Output signals generated from this primary perception step instruct an integrated perception allowing the “What”, “How”, “When” and “Where” of a visual object to be perceived as a whole. Such an integrated perception orientates decision making along three main axes: 1) neglect, 2) engage a neurocognitive activity (memorization, attention, thoughts, emotions...) or 3) engage a motor activity (grasp, repel, approach, flee...).

motor activity and visual perception are finely coupled via a whole of feedforward and feedback mechanisms allowing the execution of motor programs to be adjusted⁶⁶. In a similar manner, the somatosensory perception of movements is essential to the control of motor activity⁶⁷.

Similar to sensory neural perception, it may be suggested that the main function of immune perception is to orientate decision making toward the engagement of proper immune responses or immune programs. When bringing out the danger theory^{68,69}, Polly Matzinger was the first to emphasize the importance of contextual inputs in the initiation of “reject” vs “tolerate” immune responses. Since then, the concept of “protective autoimmunity” enunciated by Michal Schwartz and Jonathan Kipnis⁷⁰ stated that the recognition of tissue-specific auto-antigens allows the immune system to provide a tissue-specific support that is shaped by contextual signals^{71,72}. A semantic adjustment to these major conceptual advances would consist in proposing that contextual inputs orientate decision making along 3 main axes: “Reject”, “Tolerate” or “Support” i.e provide molecular and cellular instructing signals that maintain homeostasis^{73–75} or favor tissue repair^{76,77}. A functional diagram of immune perception and decision making in the immune system could be then aligned with the model of visual perception and decision making in the nervous system (Figure 7). Along this line, it may be proposed that, similar to the visuomotor and sensorimotor feedback processes, effector immune cells that may be drained from tissues to SLOs deliver output signals reflecting the execution of immune programs. The efferent phase of any immune response would be thus constantly adjusted via feedback signals that are captured and integrated in SLOs.

Conclusion

The immune perception theory proposes that immunity is driven by several basic principles that are shared between the immune and nervous system. The first proposed principle is that immune cells are not only recognizing antigens +/- danger signals but are indeed perceiving immune objects that are formed by a whole of antigenic, contextual, temporal and localization signals. The second proposed principle is that immune signals are not only individually captured by immune cells but collectively integrated at the scale of SLOs. Such a cooperative functional organization holds relevance for the communications between innate and adaptive immune cells but also for the interactions between T-cell and B-cell clones that recognize a common immune object. The third basic principle is that immune perception is shaped by a number of parameters that are independent from the perceived immune objects¹¹. These include notably the age, gender, metabolic status and gut microbiota composition of the host.

Over the last decades, the research fields covered by immunology have considerably expanded along with the number of breakthrough discoveries relating with the immune system. As a consequence, capturing an up-to-date global image of the immune system functions has become an increasingly difficult task for education professionals and for students as well. In this regard, the theoretical framework proposed here may be essentially considered as a potentially valuable tool for the teaching of immunology. In addition, while neuroimmunology encompasses, for the most part, the study of neuroimmune interactions, the present work suggests that a larger partnership could be envisioned between neuroscientists and immunologists, on the realm

Figure 7. Functional diagram of the sensory immune system. Distinct primary perception tasks allow the antigenic, temporo-contextual and localization signals characterizing an immune object to be perceived separately. Output signals generated from this primary perception step instruct an integrated perception allowing the “What”, “How”, “When” and “Where” of an immune object to be perceived as a whole. Such an integrated perception orientates decision making along three main axes: 1) tolerate, 2) reject, 3) support. i.e provide molecular and cellular instructing signals that maintain homeostasis or favor tissue repair.

of education. To face the challenge of intimately understanding complex systems such as the immune and nervous systems, a move toward an educational approximation of both disciplines is possibly of major importance to promote future cross-fertilizations of ideas and concepts.

Competing interests

No competing interests were disclosed.

Grant information

This work is supported by the Lyon-1 University and the Lyon University Hospital (Hospices Civils de Lyon), Lyon, France.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

Thanks to Nathalie Davoust-Nataf and Laurent Pays for their scientific and personal support.

References

- Jerne NK: **Towards a network theory of the immune system.** *Ann Immunol (Paris)*. [Accessed October 19, 2015] 1974; **125C**(1–2): 373–89.
[PubMed Abstract](#)
- Tauber A: **The Biological Notion of Self and Non-self.** [Accessed December 9, 2015] 2002.
[Reference Source](#)
- Choudhuri K, Llodrá J, Roth EW, *et al.*: **Polarized release of T-cell-receptor-enriched microvesicles at the immunological synapse.** *Nature*. 2014; **507**(7490): 118–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Grakoui A, Bromley SK, Sumen C, *et al.*: **The immunological synapse: a molecular machine controlling T cell activation.** *Science*. [Accessed November 9, 2015] 1999; **285**(5425): 221–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cohen IR: **The cognitive paradigm and the immunological homunculus.** *Immunol Today*. 1992; **13**(12): 490–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cohen IR: **The cognitive principle challenges clonal selection.** *Immunol Today*. 1992; **13**(11): 441–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Madi A, Bransburg-Zabary S, Maayan-Metzger A, *et al.*: **Tumor-associated and disease-associated autoantibody repertoires in healthy colostrum and maternal and newborn cord sera.** *J Immunol*. 2015; **194**(11): 5272–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Schwartz M, Cohen IR: **Autoimmunity can benefit self-maintenance.** *Immunol Today*. [Accessed December 9, 2015] 2000; **21**(6): 265–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Brownlee J: **Cognition, Immunology, and the Cognitive Paradigm.** 2007; 1–6.
[Reference Source](#)
- Cohen IR: **Real and artificial immune systems: computing the state of the body.** *Nat Rev Immunol*. 2007; **7**(7): 569–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nataf S: **The sensory immune system: a neural twist to the antigenic discontinuity theory.** *Nat Rev Immunol*. 2014; **14**(4): 280.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sincich LC, Horton JC: **The circuitry of V1 and V2: integration of color, form, and motion.** *Annu Rev Neurosci*. 2005; **28**: 303–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Livingstone M, Hubel D: **Segregation of form, color, movement, and depth: anatomy, physiology, and perception.** *Science*. [Accessed November 19, 2015] 1988; **240**(4853): 740–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Haan EH, Cowey A: **On the usefulness of 'what' and 'where' pathways in vision.** *Trends Cogn Sci*. 2011; **15**(10): 460–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Katzner S, Weigelt S: **Visual cortical networks: of mice and men.** *Curr Opin Neurobiol*. 2013; **23**(2): 202–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Araque A, Carmignoto G, Haydon PG, *et al.*: **Glutamate transporters travel in time and space.** *Neuron*. 2014; **81**(4): 728–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Araque A, Parpura V, Sanzgiri RP, *et al.*: **Tripartite synapses: glia, the unacknowledged partner.** *Trends Neurosci*. [Accessed November 29, 2015] 1999; **22**(5): 208–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schafer DP, Lehrman EK, Kautzman AG, *et al.*: **Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner.** *Neuron*. 2012; **74**(4): 691–705.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stephan AH, Barres BA, Stevens B: **The complement system: an unexpected role in synaptic pruning during development and disease.** *Annu Rev Neurosci*. 2012; **35**: 369–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wälchli T, Wacker A, Frei K, *et al.*: **Wiring the Vascular Network with Neural Cues: A CNS Perspective.** *Neuron*. 2015; **87**(2): 271–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Andreone BJ, Lacoste B, Gu C: **Neuronal and vascular interactions.** *Annu Rev Neurosci*. 2015; **38**: 25–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zinkernagel RM: **Localization dose and time of antigens determine immune reactivity.** *Semin Immunol*. 2000; **12**(3): 163–71; discussion 257–344.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Roche PA, Furuta K: **The ins and outs of MHC class II-mediated antigen processing and presentation.** *Nat Rev Immunol*. 2015; **15**(4): 203–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mueller SN, Germain RN: **Stromal cell contributions to the homeostasis and functionality of the immune system.** *Nat Rev Immunol*. 2009; **9**(9): 618–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Card CM, Yu SS, Swartz MA: **Emerging roles of lymphatic endothelium in regulating adaptive immunity.** *J Clin Invest*. 2014; **124**(3): 943–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pegu A, Qin S, Fallert Junecko BA, *et al.*: **Human lymphatic endothelial cells express multiple functional TLRs.** *J Immunol*. [Accessed December 5, 2015] 2008; **180**(5): 3399–405.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rosser EC, Oleinika K, Tonon S, *et al.*: **Regulatory B cells are induced by gut microbiota-driven interleukin-1 β and interleukin-6 production.** *Nat Med*. 2014; **20**(11): 1334–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sage PT, Paterson AM, Lovitch SB, *et al.*: **The coinhibitory receptor CTLA-4 controls B cell responses by modulating T follicular helper, T follicular regulatory, and T regulatory cells.** *Immunity*. 2014; **41**(6): 1026–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Laffont S, Seillet C, Ortaldo J, *et al.*: **Natural killer cells recruited into lymph nodes inhibit alloreactive T-cell activation through perforin-mediated killing of donor allogeneic dendritic cells.** *Blood*. 2008; **112**(3): 661–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hampton HR, Bailey J, Tomura M, *et al.*: **Microbe-dependent lymphatic migration of neutrophils modulates lymphocyte proliferation in lymph nodes.** *Nat Commun*. 2015; **6**: 7139.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Förster R, Braun A, Worbs T: **Lymph node homing of T cells and dendritic cells via afferent lymphatics.** *Trends Immunol*. 2012; **33**(6): 271–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jakubczik C, Gautier EL, Gibbins SL, *et al.*: **Minimal differentiation of classical monocytes as they survey steady-state tissues and transport antigen to lymph nodes.** *Immunity*. 2013; **39**(3): 599–610.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mackley EC, Houston S, Marriott CL, *et al.*: **CCR7-dependent trafficking of ROR γ^+ ILCs creates a unique microenvironment within mucosal draining lymph nodes.** *Nat Commun*. 2015; **6**: 5862.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

34. Yang J, Hu S, Zhao L, *et al.*: **Selective programming of CCR10⁺ innate lymphoid cells in skin-draining lymph nodes for cutaneous homeostatic regulation.** *Nat Immunol.* 2016; **17**(1): 48–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Skisiel GD, Bouchlaka MN, Monjazeb AM, *et al.*: **Out-of-Sequence Signal 3 Paralyzes Primary CD4⁺ T-Cell-Dependent Immunity.** *Immunity.* 2015; **43**(2): 240–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Yang CY, Vogt TK, Favre S, *et al.*: **Trapping of naive lymphocytes triggers rapid growth and remodeling of the fibroblast network in reactive murine lymph nodes.** *Proc Natl Acad Sci U S A.* 2014; **111**(1): E109–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Edele F, Molenaar R, Gütle D, *et al.*: **Cutting edge: instructive role of peripheral tissue cells in the imprinting of T cell homing receptor patterns.** *J Immunol.* [Accessed October 21, 2015] 2008; **181**(6): 3745–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Mora JR, Bono MR, Manjunath N, *et al.*: **Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells.** *Nature.* 2003; **424**(6944): 88–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. McCully ML, Ladell K, Hakobyan S, *et al.*: **Epidermis instructs skin homing receptor expression in human T cells.** *Blood.* 2012; **120**(23): 4591–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Calzascia T, Masson F, Di Bernardino-Besson W, *et al.*: **Homing phenotypes of tumor-specific CD8⁺ T cells are predetermined at the tumor site by crosspresenting APCs.** *Immunity.* 2005; **22**(2): 175–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Mikhak Z, Strassner JP, Luster AD: **Lung dendritic cells imprint T cell lung homing and promote lung immunity through the chemokine receptor CCR4.** *J Exp Med.* 2013; **210**(9): 1855–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Merad M, Sathe P, Helft J, *et al.*: **The dendritic cell lineage: ontogeny and function of dendritic cells and their subsets in the steady state and the inflamed setting.** *Annu Rev Immunol.* 2013; **31**: 563–604.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Molenaar R, Greuter M, van der Marel AP, *et al.*: **Lymph node stromal cells support dendritic cell-induced gut-homing of T cells.** *J Immunol.* 2009; **183**(10): 6395–402.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Cording S, Wahl B, Kulkarni D, *et al.*: **The intestinal micro-environment imprints stromal cells to promote efficient Treg induction in gut-draining lymph nodes.** *Mucosal Immunol.* 2014; **7**(2): 359–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Pradeu T, Jaeger S, Vivier E: **The speed of change: towards a discontinuity theory of immunity?** *Nat Rev Immunol.* 2013; **13**(10): 764–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Fletcher AL, Acton SE, Knoblich K: **Lymph node fibroblastic reticular cells in health and disease.** *Nat Rev Immunol.* 2015; **15**(6): 350–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Qi H, Kastenmüller W, Germain RN: **Spatiotemporal basis of innate and adaptive immunity in secondary lymphoid tissue.** *Annu Rev Cell Dev Biol.* 2014; **30**: 141–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Fasnacht N, Huang HY, Koch U, *et al.*: **Specific fibroblastic niches in secondary lymphoid organs orchestrate distinct Notch-regulated immune responses.** *J Exp Med.* 2014; **211**(11): 2265–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Onder L, Narang P, Scandella E, *et al.*: **IL-7-producing stromal cells are critical for lymph node remodeling.** *Blood.* 2012; **120**(24): 4675–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Malhotra D, Fletcher AL, Turley SJ: **Stromal and hematopoietic cells in secondary lymphoid organs: partners in immunity.** *Immunol Rev.* 2013; **251**(1): 160–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Ingulli E, Mondino A, Khoruts A, *et al.*: **In vivo detection of dendritic cell antigen presentation to CD4⁺ T cells.** *J Exp Med.* [Accessed December 30, 2015] 1997; **185**(12): 2133–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Hickman HD, Takeda K, Skon CN, *et al.*: **Direct priming of antiviral CD8⁺ T cells in the peripheral interfollicular region of lymph nodes.** *Nat Immunol.* 2008; **9**(2): 155–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Barral P, Polzella P, Bruckbauer A, *et al.*: **CD169⁺ macrophages present lipid antigens to mediate early activation of iNKT cells in lymph nodes.** *Nat Immunol.* 2010; **11**(4): 303–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Gonzalez SF, Lukacs-Kornei V, Kulligowski MP, *et al.*: **Capture of influenza by medullary dendritic cells via SIGN-R1 is essential for humoral immunity in draining lymph nodes.** *Nat Immunol.* 2010; **11**(5): 427–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Woodruff MC, Heesters BA, Herndon CN, *et al.*: **Trans-nodal migration of resident dendritic cells into medullary interfollicular regions initiates immunity to influenza vaccine.** *J Exp Med.* 2014; **211**(8): 1611–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Becattini S, Latorre D, Mele F, *et al.*: **T cell immunity. Functional heterogeneity of human memory CD4⁺ T cell clones primed by pathogens or vaccines.** *Science.* 2015; **347**(6220): 400–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Kuraoka M, Schmidt AG, Nojima T, *et al.*: **Complex Antigens Drive Permissive Clonal Selection in Germinal Centers.** *Immunity.* 2016; **44**(3): 542–552.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Gagliani N, Amezcua Vesely MC, Iseppon A, *et al.*: **Th17 cells transdifferentiate into regulatory T cells during resolution of inflammation.** *Nature.* 2015; **523**(7559): 221–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Naldi A, Carneiro J, Chaouiya C, *et al.*: **Diversity and plasticity of Th cell types predicted from regulatory network modelling.** *PLoS Comput Biol.* [Accessed January 6, 2016] 2010; **6**(9): e1000912.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Baumjohann D, Ansel KM: **MicroRNA-mediated regulation of T helper cell differentiation and plasticity.** *Nat Rev Immunol.* [Accessed January 6, 2016] 2013; **13**(9): 666–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Wang C, Collins M, Kuchroo VK: **Effector T cell differentiation: are master regulators of effector T cells still the masters?** *Curr Opin Immunol.* 2015; **37**: 6–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Cannons JL, Lu KT, Schwartzberg PL: **T follicular helper cell diversity and plasticity.** *Trends Immunol.* 2013; **34**(5): 200–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. Baz A, Groves P, Buttigieg K, *et al.*: **Quantitative assessment of the functional plasticity of memory CD8⁺ T cells.** *Eur J Immunol.* 2016; **46**(4): 863–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Seifert M, Przekopowicz M, Taudien S, *et al.*: **Functional capacities of human IgM memory B cells in early inflammatory responses and secondary germinal center reactions.** *Proc Natl Acad Sci U S A.* 2015; **112**(6): E546–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Cerwenka A, Lanier LL: **Natural killer cell memory in infection, inflammation and cancer.** *Nat Rev Immunol.* 2016; **16**(2): 112–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Gilbert CD, Li W: **Top-down influences on visual processing.** *Nat Rev Neurosci.* 2013; **14**(5): 350–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Bensmaia SJ, Miller LE: **Restoring sensorimotor function through intracortical interfaces: progress and looming challenges.** *Nat Rev Neurosci.* 2014; **15**(5): 313–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Matzinger P: **The danger model: a renewed sense of self.** *Science.* 2002; **296**(5566): 301–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Matzinger P: **Tolerance, danger, and the extended family.** *Annu Rev Immunol.* 1994; **12**: 991–1045.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Schwartz M, Kipnis J: **Protective autoimmunity: regulation and prospects for vaccination after brain and spinal cord injuries.** *Trends Mol Med.* [Accessed December 14, 2015] 2001; **7**(6): 252–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Schwartz M, Shaked I, Fisher J, *et al.*: **Protective autoimmunity against the enemy within: fighting glutamate toxicity.** *Trends Neurosci.* 2003; **26**(6): 297–302.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Schwartz M, Baruch K: **Breaking peripheral immune tolerance to CNS antigens in neurodegenerative diseases: Boosting autoimmunity to fight-off chronic neuroinflammation.** *J Autoimmun.* 2014; **54**: 8–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Ziv Y, Ron N, Butovsky O, *et al.*: **Immune cells contribute to the maintenance of neurogenesis and spatial learning abilities in adulthood.** *Nat Neurosci.* 2006; **9**(2): 268–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Lavin Y, Winter D, Blecher-Gonen R, *et al.*: **Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment.** *Cell.* 2014; **159**(6): 1312–1326.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
75. Mortha A, Chudnovskiy A, Hashimoto D, *et al.*: **Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis.** *Science.* 2014; **343**(6178): 1249288.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
76. Sica A, Mantovani A: **Macrophage plasticity and polarization: in vivo veritas.** *J Clin Invest.* 2012; **122**(3): 787–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Schwartz M, Kipnis J, Rivest S, *et al.*: **How do immune cells support and shape the brain in health, disease, and aging?** *J Neurosci.* 2013; **33**(45): 17587–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 12 October 2016

<https://doi.org/10.5256/f1000research.10110.r16681>

© 2016 Tauber A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alfred I. Tauber

Center for Philosophy and History of Science, Boston University, Boston, MA, USA

Serge Nataf has admirably schematized the “cognitive paradigm,” so aptly coined by Irun Cohen 25(!) years ago (see his citations in the accompanying review). The immune system is, indeed, a “mobile” brain (Fridman 1991) and the parallels between the immune and nervous systems is far from metaphorical or idealized modeling: their inter-connections include the origins of their respective developments, that continue with intimate anatomic relationships, as well as shared messengers and receptors. Indeed, *psychoneuroimmunology* comprises a large research field, whose wealth of studies have confirmed the close evolutionary histories of what are, in essence, similar information processing systems (Ader, Felten and Cohen 1991; Ader 2006; Orosz 2001; Forrest and Hofmeyr 2001). (Note, instead of using the visual system as his example, Nataf might have shown the obvious similarities in the molecular-sensing perception of taste or smell to make the same point about perception and, in that example, emphasized the similarities of structural correspondence and subsequent activation employed by both the immune and taste systems.)

Two historical notes:

First, while Nataf credits Jerne with introducing cognitive terminology to immunology (citing the 1974 idiotypic network paper), the origins of immunity as *cognitive* appears earlier in the cybernetics craze of the early 1950s and Burnet’s musings a decade later (Tauber 1994). (Although Burnet first used “cognate” in passing in 1959 [Burnet 1959, p.70], by 1962, he invoked an analogy with language to account for antibody selection [Burnet 1962, p. 94-5].) But historical primacy is not the matter of interest regarding Jerne’s seminal paper, but rather his introduction of how the *context* of antigen presentation determined the immune response. According to his hypothesis, the disruption of the network by antigen, and more to the point, the extent of that interruption in terms of breaking the inter-connections of the lattice-like structure, determined the immunogenicity of the introduced substance. That fundamental idea is the source of Matzinger’s Danger Theory and Pradeu’s more recent continuity/discontinuity theory.

The second historical note is that an important chapter of the cognitive paradigm’s history must include the work of Antonio Coutinho (1991; 1995; 2003), Francisco Varela (Varela *et al.*, 1988;

1991, 1993; Varela and Coutinho 1991), John Stewart (1992; 1994a; 1994b; Stewart and Varela 1989), and Nelson Vaz (2016; Vaz *et al.*, 2006). These publications developed Jerne's cognitive point in its full theoretical array and attempted to draw both theoretical and practical parallels between the nervous and immune systems. Those studies comprise the rich mulch for the emergence of our current thinking about how immunity may be properly regarded as an information processing faculty and as such shares organizational (and ultimately regulatory) characteristics with the brain.

In terms of going beyond Nataf's schema as an educational tool (which it certainly is), a deeper assessment must be made as to how well modelers have applied the integrative principles employed here. In other words, to the extent systems biology has offered models of immune responses or steady-state conditions, has the hierarchical structure employed by Nataf been utilized and if so, what insights does such an approach offer? That assessment would be useful not only in terms of confirming Nataf's schema, but perhaps modelers might develop this approach for their own efforts. A review of this issue, as well as a critical discussion about the cognitive paradigm applied to immunology, is found elsewhere (Tauber 1997; 2013; 2017).

References

1. Ader R.: Psychoneuroimmunology (4th ed.). *Elsevier Academic Press*. 2006. [Reference Source](#)
2. Ader R., Felten D.L., Cohen N.: Psychoneuroimmunology (2nd ed). *Academic Press*. 1991.
3. Coutinho A: Size and connectivity: a bit of the history of immune networks.*J Theor Biol*. 1991; **149** (3): 425-7 [PubMed Abstract](#)
4. Coutinho A: The network theory: 21 years later.*Scand J Immunol*. 1995; **42** (1): 3-8 [PubMed Abstract](#)
5. Coutinho A: A walk with Francisco Varela from first- to second-generation networks: in search of the structure, dynamics and metadynamics of an organism-centered immune system.*Biol Res*. 2003; **36** (1): 17-26 [PubMed Abstract](#)
6. Fridman W.H.: *Le Cerveau Mobile*. Paris: *Hermann*. 1991.
7. Stewart J.: Immunoglobulins did not arise in evolution to fight infection. *Immunol Today*. 1992; **13** (10): 396-399 [Reference Source](#)
8. Stewart J.: The Primordial VRM System and the Evolution of Vertebrate Immunity. *R.G.Landes, Austin, Texas*. 1994a; **124**. [Reference Source](#)
9. Stewart J.: Cognition without neurones: adaptation, learning and memory in the immune system. *Cognitiva*. 1993; **5** (2): 187-202 [Reference Source](#)
10. Stewart J, Varela FJ: Exploring the meaning of connectivity in the immune network.*Immunol Rev* . 1989; **110**: 37-61 [PubMed Abstract](#)
11. Tauber A.I.: The Immune Self. Theory or metaphor?. *Cambridge University Press*. 1994. [Reference Source](#)
12. Tauber AI: Historical and philosophical perspectives concerning immune cognition.*J Hist Biol*. 1997; **30** (3): 419-40 [PubMed Abstract](#)
13. Tauber AI: Immunology's theories of cognition.*Hist Philos Life Sci*. 2013; **35** (2): 239-64 [PubMed Abstract](#)
14. Tauber A.I.: Immunity: The Evolution of an Idea. *Oxford University Press*. 2017.
15. Varela F, Andersson A, Dietrich G, Sundblad A, et al.: Population dynamics of natural antibodies in normal and autoimmune individuals.*Proc Natl Acad Sci U S A*. 1991; **88** (13): 5917-21 [PubMed Abstract](#)
16. Varela FJ, Coutinho A: Second generation immune networks.*Immunol Today*. 1991; **12** (5): 159-66 [PubMed Abstract](#) | [Publisher Full Text](#)
17. Varela F., Coutinho A., Dupire B., Nelson N.: Cognitive networks: Immune, neural, and

otherwise, in A.S. Perelson (ed.). *Theoretical Immunology, Part II. SFI Series on the Science of Complexity*. Addison-Wesley. 359-375

18. Carela F., Coutinho A., Stewart J.: What is the immune network for? in W.D. Stein and F.J. Zerla (eds.), *Thinking About Biology, Studies in the Science of Complexity, Lecture Notes Vol. III*. Redwood City, CA. Addison-Wesley Publishing Co. 1993. 215-230

19. Vaz NM: Self-tolerance revisited. *Stud Hist Philos Biol Biomed Sci*. 2016; **55**: 128-32 [PubMed Abstract](#)

20. Vaz NM, Ramos GC, Pordeus V, Carvalho CR: The conservative physiology of the immune system. A non-metaphoric approach to immunological activity. *Clin Dev Immunol*. **13** (2-4): 133-42 [PubMed Abstract](#) | [Publisher Full Text](#)

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 06 October 2016

<https://doi.org/10.5256/f1000research.10110.r16829>

© 2016 Cohen I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Irun R. Cohen

Department of Immunology, Weizmann Institute of Science, Rehovot, Israel

The paper by Serge Nataf, *Clonal selection versus clonal cooperation: the integrated perception of immune objects*, is aptly named: here Nataf draws an analogy between the immune system (IS) and the central nervous system (CNS); both systems, he points out, are cognitive in the sense that they construct their objects of attention through the integration of discrete features perceived individually. The CNS *constructs* specific visual objects by *integrating* information between networks of neurons that gather discrete and separate information about shape, location, color and motion; similarly, the IS constructs immune objects by integrating discrete information about antigens, cytokines, chemokines, and other signals obtained separately by various cell types such as macrophages, dendritic cells, and B and T lymphocytes. Integration of information between different cells and cell types requires cooperation between the interacting cells; thus the IS, like the CNS, makes functional decisions based on clonal cooperation and not on clonal competition for survival of the fittest. The cognitive analogy is presented in a clear and stimulating manner. However, I think that Serge Nataf could increase the impact of his CNS-IS discussion by sharpening or expanding some additional points:

- *Functional development requires somatic experience*: Both the CNS and the IS combine individual somatic experience with innate programming acquired through evolution of the species. In fact, the CNS and the IS are the only two mammalian systems that require post-natal, individual contact with the world to self-organize their mature structures and functions (see Cohen I.R, (2000)); an IS deprived of immune experience remains undeveloped as does a CNS deprived of formative stimulation.

- *Homunculi direct attention to environmental information that serves adaptive fitness*: Nataf mentions briefly that both the CNS and the IS use homunculi – internal representations of selected features of the self and the outside world; he might add that such homunculi exist to maximize evolutionary fitness. The human brain, for example, is born hard-wired to be attracted to human faces; this primitive homuncular sense of facial recognition is sharpened by individual post-natal experience with real persons, beginning with mother and leading to success in interpersonal bonding, friendship and other important relationships, and to the capacity to communicate by facial expressions.
- The immunological homunculus, which is encoded in innate immune receptors and in shared autoantibody repertoires (Madi A, *et al.* (2015)) and in public T cell receptor repertoires (Madi A, *et al.* (2014)) is much less appreciated. But, like the Neurological Homunculus, these congenital and acquired receptors are likely to enhance fitness, for example, by enabling the IS to better preserve and heal the self and its microbiome while protecting the body from pathogenic invaders from without and from tumor cells from within.
- Important points of difference between the CNS and IS would include hard wiring compared to cell migration, reaction times of milliseconds to hours and days, markedly different cell turnover and differentiation rates, and others; both systems create and integrate their perceived objects in markedly different ways. But, as Nataf points out so convincingly, integrated cooperation is the hallmark of both systems.

References

1. Cohen I.R.: Tending Adam's Garden: evolving the cognitive immune self. *Academic Press*. 2000.
[Reference Source](#)
2. Madi A, Bransburg-Zabary S, Maayan-Metzger A, Dar G, et al.: Tumor-associated and disease-associated autoantibody repertoires in healthy colostrum and maternal and newborn cord sera. *J Immunol*. 2015; **194** (11): 5272-81 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Madi A, Shifrut E, Reich-Zeliger S, Gal H, et al.: T-cell receptor repertoires share a restricted set of public and abundant CDR3 sequences that are associated with self-related immunity. *Genome Res*. 2014; **24** (10): 1603-12 [PubMed Abstract](#) | [Publisher Full Text](#)

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Comments on this article

Version 1

Author Response 19 Jan 2017

Serge Nataf, University Lyon-1, Lyon, France

This a comment from the author:

Based on talks with neuroscientists specialized in the visual system, I would like to add precisions to the description of the visual system and the parallel drawn with the perception of immune objects. It is indeed important to underscore that in the retina, distinct categories of neurons named ganglion cells are able to independently sense the shape, color and motion of a visual object (1, 2). In turn, these neuronal cells generate output signals (action potentials) that are transmitted toward the superficial neuronal layers of the brain occipital lobe. Via synaptic connections, an interaction between retinal ganglion cells and neurons of the primary visual cortex supports thus the primary perception of visual objects. In summary, distinct features harbored by a visual object are sensed separately by specialized neuronal cells at early steps of the visual perception process. Similarly, distinct features harbored by an immune object may be sensed separately by specialized immune cells at early steps of the immune perception process. Please note also that the "Where" and "What" pathways have been inverted in Figure 1 (the legend though is correct as well as the main text describing these pathways of integrated visual perception).

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research