OPINION ARTICLE

REVISED Statin (3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitor)-based therapy for hepatitis C virus (HCV) infection-related diseases in the era of direct-acting antiviral agents [version 3; peer review: 2 approved]

Sara Sobhy Kishta¹, Sobhy Ahmed Kishta², Reem El-Shenawy¹

¹Medical Biotechnology Lab, Microbial Biotechnology Department, National Research Center, Egypt, Cairo, Egypt

²Department of Surgery, Theodor Bilharz Research Institute, Giza, Egypt

V3 First published: 26 Feb 2016, 5:223
<https://doi.org/10.12688/f1000research.7970.1>
 Second version: 07 Jul 2016, 5:223
<https://doi.org/10.12688/f1000research.7970.2>
 Latest published: 25 Jan 2017, 5:223
<https://doi.org/10.12688/f1000research.7970.3>

Abstract

Recent improvements have been made in the treatment of hepatitis C virus (HCV) infection with the introduction of direct-acting antiviral agents (DAAs). However, despite successful viral clearance, many patients continue to have HCV-related disease progression. Therefore, new treatments must be developed to achieve viral clearance and prevent the risk of HCV-related diseases. In particular, the use of pitavastatin together with DAAs may improve the antiviral efficacy as well as decrease the progression of liver fibrosis and the incidence of HCV-related hepatocellular carcinoma. To investigate the management methods for HCV-related diseases using pitavastatin and DAAs, clinical trials should be undertaken. However, concerns have been raised about potential drug interactions between statins and DAAs. Therefore, pre-clinical trials using a replicon system, human hepatocyte-like cells, human neurons and human cardiomyocytes from human-induced pluripotent stem cells should be conducted. Based on these pre-clinical trials, an optimal direct-acting antiviral agent could be selected for combination with pitavastatin and DAAs. Following the pre-clinical trial, the combination of pitavastatin and the optimal direct-acting antiviral agent should be compared to other combinations of DAAs (e.g., sofosbuvir and velpatasvir) according to the antiviral effect on HCV infection, HCV-related diseases and cost-effectiveness.

Keywords

Hepatitis C virus (HCV) infection , HCV infection-related diseases , statins , replicon system , human hepatocyte-like cells from human induced pluripotent stem cells , direct-acting antiviral agents (DAAs)

Open Peer Review

Approval Status

	1	2
version 3 (revision) 25 Jan 2017		
version 2 (revision) 07 Jul 2016		view
version 1 26 Feb 2016	view	view

1. **Susana N. Asin**, Veterans Affairs Medical Center, White River Junction, USA
2. **Joel D. Baines** , Louisiana State University, Baton Rouge, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Sara Sobhy Kishta (kishtassa@yahoo.com)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Kishta SS *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kishta SS, Kishta SA and El-Shenawy R. **Statin (3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitor)-based therapy for hepatitis C virus (HCV) infection-related diseases in the era of direct-acting antiviral agents [version 3; peer review: 2 approved]** F1000Research 2017, 5:223 <https://doi.org/10.12688/f1000research.7970.3>

First published: 26 Feb 2016, 5:223 <https://doi.org/10.12688/f1000research.7970.1>

REVISED Amendments from Version 2

All of the author names were updated to correct them and the order of the authors was also corrected.

See referee reports

Introduction

Hepatitis C virus (HCV) infection is an important public health problem worldwide, as many HCV-infected patients develop liver cirrhosis and/or hepatocellular carcinoma (HCC)¹.

Recent improvements in the treatment of HCV infection have focused on the use of direct-acting antiviral agents (DAAs)¹. However, despite successful viral clearance, many patients with advanced fibrosis continue to have HCV-related disease progression^{2,3}. Therefore, new treatments must be developed to achieve viral clearance and prevent the risk of HCV-related diseases.

Statins (inhibitors of 3-hydroxy-3-methylglutaryl-coenzyme A reductase) have been proposed as a new candidate for the treatment of HCV infection. According to previously conducted clinical studies using statins in HCV-infected patients^{4–8}, the antiviral effect of statins on HCV infection depends on the type of statin used. Among various statins, pitavastatin showed the highest antiviral efficacy against HCV genotype 1b infection *in vitro*⁹. Therefore, pitavastatin represents a novel candidate for the treatment of HCV infection.

Clinical trials for pitavastatin against HCV infection

After performing a search of the PubMed database, we identified three clinical trials using pitavastatin for the treatment of HCV infection^{7,8,10}.

When Shimada *et al.*⁷ considered two reports published in 2010 investigating the antiviral efficacy of pitavastatin against HCV infection *in vitro*^{9,11}, they conducted a randomized controlled trial⁷. The proof-of-concept studies demonstrated the antiviral efficacy and safety of pitavastatin against HCV infection using a replicon system and human hepatocyte-like cells from human induced pluripotent stem cells (hiPSCs)^{9,11}, and these effects of pitavastatin were confirmed in the randomized controlled trial^{7,12}. Indeed, this series of studies^{7,9,11} would be the first to report the clinical applications of hiPSCs¹². After the clinical trial performed by Shimada *et al.*⁷, the antiviral efficacy and safety of pitavastatin against HCV infection were further confirmed in two clinical studies^{8,10}. Thus, investigating the antiviral efficacy and safety of pitavastatin against HCV infection using a replicon system and human hepatocyte-like cells from hiPSCs^{9,11} constitutes a rational approach to discover new drugs and/or new therapeutic methods.

Clinical trials to investigate the management methods of HCV-related diseases in the era of DAAs

Butt *et al.*¹³ showed that statin use was associated with improved antiviral efficacy as well as decreased progression of liver fibrosis

and a reduced incidence of HCC among a large cohort of HCV-positive veterans. Furthermore, the use of statins among patients with HCV and compensated cirrhosis (n=40,512) was associated with a more than 40% lower risk of cirrhosis decomposition and death¹⁴. Moreover, statin users showed a significant reduction in the incidence of HCC¹⁵. In addition, pitavastatin showed anti-cancer effects against human hematoma cell lines^{16–18}.

DAAs in combination with statins have been shown to generate increased antiviral efficacy against HCV infection¹⁹. However, concerns have been raised about the drug interactions between various statins and DAAs²⁰. For instance, simvastatin and lovastatin should be avoided in patients with HCV infection who are using boceprevir or telaprevir as a DAA²¹. Atorvastatin should be avoided in patients with HCV infection who are using telaprevir²¹, and pravastatin plus boceprevir may also pose risks²¹. Although rosvastatin could be considered for use in combination with telaprevir and boceprevir²¹, the drug interactions between pitavastatin and DAAs remain unknown.

Furthermore, according to our search of the PubMed database and UMIN Clinical Trials Registry System (<http://www.umin.ac.jp/icdr/index.html>), no clinical trial has been conducted for the combination of statins and DAAs. Therefore, although there is likely only a minimal additive benefit for viral clearance using statins, as new combinations of DAA (sofosbuvir and velpatasvir) therapy have shown sustained virologic response (SVR) rates above 95%²², conducting a clinical trial for the combination of pitavastatin and DAAs may be meaningful to investigate management methods to prevent fibrosis and cirrhosis or the development of HCC and other HCV-related diseases in the era of DAAs. However, in pre-clinical trials, the antiviral effects of the combination of pitavastatin and DAAs should be evaluated using a replicon system⁹. Furthermore, hepatotoxicities should also be evaluated for the combination of pitavastatin and DAAs using human hepatocyte-like cells from hiPSCs¹¹. Using a replicon system and human hepatocyte-like cells from hiPSCs in a pre-clinical trial^{9,11}, an optimal direct-acting antiviral agent could be selected for use in the combination of pitavastatin and DAAs. After the pre-clinical trial, the combination of pitavastatin and the optimal direct-acting antiviral agent should be compared with other DAA combinations (*e.g.*, sofosbuvir and velpatasvir) according to their antiviral efficacy against HCV infection and prevention of HCV-related diseases. Furthermore, because the cost for DAA combination treatment is very high (\$83,000 to \$153,000 per course of treatment)²², the new and effective hepatitis C treatments seem beyond the reach of low- and middle-income countries²³. However, the cost (\$0.79 to \$2.59 per day) of pitavastatin (2mg)⁷ is low (<http://www.pharmacychecker.com/generic/price-comparison/pitavastatin/2+mg/>). Therefore, the above-mentioned comparison should also be investigated in light of cost-effectiveness.

On the other hand, a replicon system in hepatocellular cells is not likely to duplicate the full gamut of potential side effects seen in patients with HCV infection receiving pitavastatin and DAAs. The risks (potential side effects) that will not be revealed

in a pre-clinical trial using this system should be acknowledged. Therefore, in order to identify the risk (potential side effects) that will not be revealed in a pre-clinical trial using this system, the evaluations using human neurons and human cardiomyocytes from hiPSCs^{24,25} should also be done in pre-clinical trials.

In conclusion, a pre-clinical trial investigating the combination of pitavastatin and DAAs against HCV infection using a replicon system, human hepatocyte-like cells, human neurons and human cardiomyocytes from hiPSCs^{9,11,24,25} represents a rational approach for discovering a new therapeutic method.

Author contributions

All authors (Reem Mohamed Fathy El-Shenawy, Sara Kishta and Sobhy Kishta) equally contributed to the writing of the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Carpentier A, Tesfaye A, Chu V, *et al.*: **Engrafted human stem cell-derived hepatocytes establish an infectious HCV murine model.** *J Clin Invest.* 2014; **124**(11): 4953–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Reddy KR, Zeuzem S, Zoulim F, *et al.*: **Simeprevir versus telaprevir with peginterferon and ribavirin in previous null or partial responders with chronic hepatitis C virus genotype 1 infection (ATTAIN): a randomised, double-blind, non-inferiority phase 3 trial.** *Lancet Infect Dis.* 2015; **15**(1): 27–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Simon TG, King LY, Zheng H, *et al.*: **Statin use is associated with a reduced risk of fibrosis progression in chronic hepatitis C.** *J Hepatol.* 2015; **62**(1): 18–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Patel K, Lim SG, Cheng CW, *et al.*: **Open-label phase 1b pilot study to assess the antiviral efficacy of simvastatin combined with sertraline in chronic hepatitis C patients.** *Antivir Ther.* 2011; **16**(8): 1341–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Patel K, Jhaveri R, George J, *et al.*: **Open-label, ascending dose, prospective cohort study evaluating the antiviral efficacy of rosvastatin therapy in serum and lipid fractions in patients with chronic hepatitis C.** *J Viral Hepat.* 2011; **18**(5): 331–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bader T, Hughes LD, Fazili J, *et al.*: **A randomized controlled trial adding fluvastatin to peginterferon and ribavirin for naïve genotype 1 hepatitis C patients.** *J Viral Hepat.* 2013; **20**(9): 622–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shimada M, Yoshida S, Masuzaki R, *et al.*: **Pitavastatin enhances antiviral efficacy of standard pegylated interferon plus ribavirin in patients with chronic hepatitis C: a prospective randomized pilot study.** *J Hepatol.* 2012; **56**(1): 299–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kohjima M, Enjoji M, Yoshimoto T, *et al.*: **Add-on therapy of pitavastatin and eicosapentaenoic acid improves outcome of peginterferon plus ribavirin treatment for chronic hepatitis C.** *J Med Virol.* 2013; **85**(2): 250–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moriguchi H, Chung RT, Sato C: **New translational research on novel drugs for hepatitis C virus 1b infection by using a replicon system and human induced pluripotent stem cells.** *Hepatology.* 2010; **51**(1): 344–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yokoyama S, Kawakami Y, Chayama K: **Letter: Pitavastatin supplementation of PEG-IFN/ribavirin improves sustained virological response against HCV.** *Aliment Pharmacol Ther.* 2014; **39**(4): 443–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moriguchi H, Chung RT, Sato C: **An identification of the novel combination therapy for hepatitis C virus 1b infection by using a replicon system and human induced pluripotent stem cells.** *Hepatology.* 2010; **51**(1): 351–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moriguchi H: **The development of statin-based therapy for patients with hepatitis C virus (HCV) infection using human induced pluripotent stem (iPS) cell technology.** *Clin Res Hepatol Gastroenterol.* 2015; **39**(5): 541–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Butt AA, Yan P, Bonilla H, *et al.*: **Effect of addition of statins to antiviral therapy in hepatitis C virus-infected persons: Results from ERCHIVES.** *Hepatology.* 2015; **62**(2): 365–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mohanty A, Tate JP, Garcia-Tsao G: **Statins Are Associated With a Decreased Risk of Decompensation and Death in Veterans With Hepatitis C-Related Compensated Cirrhosis.** *Gastroenterology.* 2016; **150**(2): 430–440.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tsan YT, Lee CH, Ho WC, *et al.*: **Statins and the risk of hepatocellular carcinoma in patients with hepatitis C virus infection.** *J Clin Oncol.* 2013; **31**(12): 1514–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arii K, Suehiro T, Ota K, *et al.*: **Pitavastatin induces PON1 expression through p44/42 mitogen-activated protein kinase signaling cascade in Huh7 cells.** *Atherosclerosis.* 2009; **202**(2): 439–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wang J, Xu Z, Zhang M: **Downregulation of survivin expression and elevation of caspase-3 activity involved in pitavastatin-induced HepG 2 cell apoptosis.** *Oncol Rep.* 2007; **18**(2): 383–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wang J, Tokoro T, Higa S, *et al.*: **Anti-inflammatory effect of pitavastatin on NF-kappaB activated by TNF-alpha in hepatocellular carcinoma cells.** *Biol Pharm Bull.* 2006; **29**(4): 634–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Delang L, Paeshuyse J, Vliegen I, *et al.*: **Statins potentiate the *in vitro* anti-hepatitis C virus activity of selective hepatitis C virus inhibitors and delay or prevent resistance development.** *Hepatology.* 2009; **50**(1): 6–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Poordad F, McCone J Jr, Bacon BR, *et al.*: **Boceprevir for untreated chronic HCV genotype 1 infection.** *N Engl J Med.* 2011; **364**(13): 1195–206.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kiser JJ, Burton JR, Anderson PL, *et al.*: **Review and management of drug interactions with boceprevir and telaprevir.** *Hepatology.* 2012; **55**(5): 1620–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ward JW, Mermin JH: **Simple, Effective, but Out of Reach? Public Health Implications of HCV Drugs.** *N Engl J Med.* 2015; **373**(27): 2678–2680.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jayasekera CR, Barry M, Roberts LR, *et al.*: **Treating hepatitis C in lower-income countries.** *N Engl J Med.* 2014; **370**(20): 1869–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Heman-Ackah SM, Bassett AR, Wood MJ: **Precision Modulation of Neurodegenerative Disease-Related Gene Expression in Human iPSC-Derived Neurons.** *Sci Rep.* 2016; **6**: 28420.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sinnecker D, Laugwitz KL, Moretti A: **Induced pluripotent stem cell-derived cardiomyocytes for drug development and toxicity testing.** *Pharmacol Ther.* 2014; **143**(2): 246–252.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 15 August 2016

<https://doi.org/10.5256/f1000research.9843.r15693>

© 2016 Baines J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Joel D. Baines

LSU School of Veterinary Medicine (SVM), Louisiana State University, Baton Rouge, LA, USA

The revisions address my concerns about the preclinical studies.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 07 June 2016

<https://doi.org/10.5256/f1000research.8578.r13945>

© 2016 Baines J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Joel D. Baines

LSU School of Veterinary Medicine (SVM), Louisiana State University, Baton Rouge, LA, USA

This article would be improved with statements quantifying how common progression to disease occurs in patients treated with anti-HCV compounds in the absence of other drugs. The main point of the article, that there should be pre-clinical trials to test for efficacy and toxicity of dual therapy with Pitavastatin are valid. However, a replicon system in hepatocellular cells is not likely to duplicate the full gamut of potential side effects seen in humans receiving pitavastatin and anti-HCV drugs. Although it is an important place to start, the risk that complications will not be

revealed in a preclinical trial using this system should be acknowledged.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 28 Jun 2016

Sara Kishta, National Research Center, Egypt, Cairo, Egypt

First of all, thank you very much for your suggestions. We think that your comments were very helpful. We agree with your suggestions. If you agree with our revisions, we are very happy.

1) According to your suggestions, we added the following sentences to our text (please see, last paragraph and conclusion).

On the other hand, a replicon system in hepatocellular cells is not likely to duplicate the full gamut of potential side effects seen in patients with HCV infection receiving pitavastatin and DAAs. The risks (potential side effects) that will not be revealed in a pre-clinical trial using this system should be acknowledged. Therefore, in order to identify the risk (potential side effects) that will not be revealed in a pre-clinical trial using this system, the evaluations using human neurons and human cardiomyocytes from hiPSCs^{24, 25} should also be done in pre-clinical trials.

In conclusion, a pre-clinical trial investigating the combination of pitavastatin and DAAs against HCV infection using a replicon system, human hepatocyte-like cells, human neurons and human cardiomyocytes from hiPSCs^{9,11,24,25} represents a rational approach for discovering a new therapeutic method.

2) We revised our abstract.

pre-clinical trials using a replicon system, human hepatocyte-like cells, human neurons and human cardiomyocytes from human-induced pluripotent stem cells should be conducted.

3) We added the new references (No. 24 and 25) in the reference section.

Competing Interests: No competing interests were disclosed.

Reviewer Report 11 May 2016

<https://doi.org/10.5256/f1000research.8578.r13680>

© 2016 Asin S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Susana N. Asin

White River Junction, Veterans Affairs Medical Center, White River Junction, VT, USA

Recent improvements in treatment strategies for Hepatitis C virus infected patients have been successful at reaching sustained virological responses yet did not prevent liver fibrosis and progression to hepatocellular carcinoma.

The idea proposed in this paper that new treatments must be developed to achieve viral clearance while preventing disease progression although meritorious is not original and should be complemented with further understanding of mechanisms underlying hepatitis C disease progression in the presence of sustained virological response (i.e. absence/ or low level of viral replication).

The authors stated that pitavastatin represents a novel candidate to treat HCV infection yet the rationale for selecting this statin, as the likely candidate is not clearly justified. The authors based their selection on data from a retrospective analysis demonstrating no significant difference in sustained virological response rate per protocol analysis between patients treated with PEG-IFN/ribavirin in the presence or absence of pitavastatin. Findings from a second prospective trial demonstrates the safety of pitavastatin in combination with Peg IFN plus ribavirin although the decrease in HCV RNA was only significant at 2 (4 and 12 weeks) of 6 evaluated time points of treatment. The third trial used pitavastatin in combination with eicosapentaenoic acid and identified this combination therapy as predictive of sustained virological response in multifactorial analysis only after genetic variation in IL28B was excluded from these factors. Thus the authors could have presented a better rationale for the selection of this statin.

An additional point of discussion that should have been included in this paper would have been the evidence that statin use decreased progression to liver fibrosis and hepatocellular carcinoma yet this effect was independent of having attained a sustained virological response. These findings raised the possibility that the statins-mediated delay in liver fibrosis are related to their immunomodulatory rather than their antiviral effects. This assumption is supported by findings demonstrating little or no effect of statins use on HCV replication.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research