REVIEW

Biology and structure of leukocyte β_2 integrins and their role in inflammation [version 1; peer review: 3 approved]

M. Amin Arnaout

Leukocyte Biology & Inflammation Program, Structural Biology Program, Nephrology, Center for Regenerative Medicine, Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

V1 First published: 04 Oct 2016, 5(F1000 Faculty Rev):2433
<https://doi.org/10.12688/f1000research.9415.1>

Latest published: 04 Oct 2016, 5(F1000 Faculty Rev):2433
<https://doi.org/10.12688/f1000research.9415.1>

Abstract

Integrins comprise a large family of $\alpha\beta$ heterodimeric cell adhesion receptors that are expressed on all cells except red blood cells and that play essential roles in the regulation of cell growth and function. The leukocyte integrins, which include members of the β_1 , β_2 , β_3 , and β_7 integrin family, are critical for innate and adaptive immune responses but also can contribute to many inflammatory and autoimmune diseases when dysregulated. This review focuses on the β_2 integrins, the principal integrins expressed on leukocytes. We review their discovery and role in host defense, the structural basis for their ligand recognition and activation, and their potential as therapeutic targets.

Keywords

inflammation , integrins , leukocytes , integrin structure

Open Peer Review

Approval Status

	1	2	3
version 1 04 Oct 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Klaus Ley**, La Jolla Institute for Allergy and Immunology, La Jolla, USA
2. **Jun Qin**, Lerner Research Institute, Cleveland Clinic, Cleveland, USA
3. **Tobias Ulmer**, University of Southern California, Los Angeles, USA

Any comments on the article can be found at the end of the article.

Corresponding author: M. Amin Arnaout (aarnaout1@mgh.harvard.edu)

Competing interests: The author declares that he has no competing interests.

Grant information: The author's work presented in this review was supported by National Institutes of Health grants DK088327, DK48549, and DK007540 from the National Institutes of Diabetes, Digestive and Kidney Diseases.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Arnaout MA. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Arnaout MA. **Biology and structure of leukocyte β_2 integrins and their role in inflammation [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2433 <https://doi.org/10.12688/f1000research.9415.1>

First published: 04 Oct 2016, 5(F1000 Faculty Rev):2433 <https://doi.org/10.12688/f1000research.9415.1>

Introduction

Leukocytes circulate in the blood in a quiescent state before migrating into tissues to defend against invading pathogens or to participate in other immune functions. Improperly activated leukocytes can also be effectors of pathologic inflammation. Most leukocyte functions are dependent on members of the integrin family (Figure 1). Leukocyte integrins comprise all four β_2 integrins, the two β_7 integrins $\alpha_4\beta_7$ and $\alpha_E\beta_7$, in addition to $\alpha_4\beta_1$, $\alpha_5\beta_1$, $\alpha_9\beta_1$, and $\alpha_V\beta_3$. Leukocyte integrins play key roles in the innate immune response, which include interaction of phagocytic cells with endothelium and the extracellular matrix, ingestion of complement-opsonized pathogens, degranulation, and cytokine production. They are also involved in lymphocyte proliferation, survival, and differentiation in adaptive immunity. Chemokines, cytokines, lipid signaling molecules, and “cross-talk” from other adhesion molecules regulate the functional state, density, and topography of leukocyte integrins. The leukocyte-specific β_2 integrins are the most abundant leukocyte integrins and the first integrins to be studied functionally and structurally in these cells. In this review, we will focus on β_2 integrins and their role in immunity and their structure and mechanism of their inside-out signaling. Many elements of the integrin outside-in signaling networks have been identified and were the subject of excellent reviews^{1–4} but are outside the scope of this concise review.

Discovery of β_2 integrins

The sequential steps leading to an inflammatory response were first documented by Julius Cohnheim in the frog’s tongue⁵. He observed that local mechanical irritation induced first an increase in blood flow and then a slowing, at which time white blood cells began to roll and then halt, lining up the wall of venules, whereas red blood cells sped past them. Then some white blood cells began to creep across the wall into the extravascular space⁵. Elie Metchnikoff discovered the phagocytic function of certain white blood cells by using the transparent avascular starfish larvae⁶. His phagocytosis theory of inflammation complemented Paul Ehrlich’s humoral theory, which attributed bacterial killing to serum-derived “magic bullets”, identified soon after as antibodies and complement proteins. The identity of the molecules involved in leukocyte migration across venules and in phagocytosis remained unknown, however.

In 1979, an experiment of nature led us to the identification of the major surface receptors mediating leukocyte migration and phagocytosis (reviewed in 7). We investigated in a pediatric patient the basis for his life-threatening bacterial infections, impaired wound

healing, persistent marked neutrophilia even during infection-free periods, but a paucity of neutrophils within infected tissues. His neutrophils failed to adhere to substrate, migrate across inflamed endothelium, or ingest serum-opsonized particles. We traced these phagocyte defects to a deficiency of a gp150 surface membrane glycoprotein complex⁸. Monoclonal antibodies (mAbs) raised by us⁹ and by others^{10–15} showed that the gp150 complex comprises four surface glycoproteins now known as CD11a (α_L)^{16–18}, CD11b (α_M)¹⁹, CD11c (α_X)²⁰, and CD11d (α_D)²¹. Each CD11 glycoprotein non-covalently associates with a common 95 kDa glycoprotein (CD18, β_2)^{13–15,18} to form what is now known as the β_2 integrin subfamily. Mutations in the CD18 subunit^{7,22–24} resulted in its partial or complete failure to associate with the synthesized CD11 α -subunits, accounting for the variations in severity of the disease now known as leukocyte adhesion deficiency type I (LAD I)^{18,25}.

Tissue distribution of β_2 integrins

β_2 integrins are expressed only on leukocytes, but their expression varies among the leukocyte subpopulations. CD11a is expressed on all leukocytes but predominates on lymphocytes. CD11b predominates on myeloid cells, being the most abundant integrin on neutrophils, and is also expressed on natural killer (NK) cells, fibrocytes, and some mast cells, B cells, CD8⁺ T cells, and $\gamma\delta$ T cells^{26–33}. CD11c is most abundant on myeloid dendritic cells, predominating on macrophages and dendritic cells of the splenic white pulp and marginal zone and on pulmonary alveolar macrophages, and has a distribution similar to that of CD11b on NK, B, and T cells³⁴. CD11d is basally expressed on the majority of circulating human neutrophils and monocytes, on NK cells, and on a small fraction of circulating T cells^{35,36}. In mice, CD11d expression is restricted to a small percentage of circulating leukocytes under basal conditions but predominates in splenic red pulp macrophages, lymph node medullary cord and sinus macrophages, and hemosiderin-containing bone marrow macrophages and is upregulated on phagocytes at local inflammatory sites^{35–37} and on differentiated macrophages, which may facilitate their retention at sites of inflammation³⁸.

β_2 integrin ligands

CD11a binds intercellular adhesion molecules (ICAMs) 1–5, telencephalin, endothelial cell-specific molecule-1 (ESM-1), and junctional adhesion molecule 1 (JAM1)^{39–41}. CD11b is the most promiscuous β_2 integrin; it has more than 40 reported ligands, including iC3b, ICAM1, 2, 3 and 4, fibrin(ogen), fibronectin, Factor X, Platelet Iba, JAM-3, and some proteases (for example,

Figure 1. Mammalian integrins. This protein family consists of 24 α/β heterodimeric receptors assembled from 18 α -subunits and eight β -subunits. Nine α -subunits (shaded) contain an extra von Willebrand factor type A domain (α_A or α_I). The β_1 integrins are the largest subfamily, with 12 known members.

proteinase 3) CD11c binds ICAM1, 4, iC3b, and vascular cell adhesion protein 1 (VCAM-1)^{42–46}. Like CD11b, CD11c also binds heparin, various polysaccharides, and negative charges in denatured proteins^{26,47–49}. CD11d binds ICAM-3 and VCAM-1⁵⁰ and, like CD11b, also binds several matrix proteins³⁸.

Functional analysis of the individual β_2 integrins

The defects in leukocyte adhesion demonstrated in patients with LAD I and in mice lacking CD18⁵¹ did not allow an assessment of the relative contribution of each of the four β_2 integrins to the phenotypic abnormalities observed. Generation of mice deficient in the individual CD11 subunits revealed that knockout (KO) of CD11a (but not CD11b) in mice caused neutrophilia, which was not as severe as that found in CD18 KO mice, suggesting additional contributions by the other β_2 integrins. No CD11a^{−/−}, CD11b^{−/−}, or CD11d KO mice developed the spontaneous infections observed in CD18 KO mice, suggesting that loss of all CD11/CD18 receptors is necessary to cause spontaneous bacterial infections. Homotypic aggregation and antigen-, mitogen-, and alloantigen-induced lymphoproliferation, which lead to defective host-versus-graft reaction and impaired tumor rejection, were reduced in CD11a^{−/−} but not CD11b^{−/−} or CD11c^{−/−} leukocytes^{52,53}. However, cytotoxic T-cell responses to systemic viral infections were normal in CD11a KO mice^{54,55}, suggesting molecular redundancy or compensatory changes (or both) by other leukocyte integrins such as $\alpha 4\beta 1$ or $\alpha 9\beta 1$ ^{56,57}. This may explain the rarity of viral infections in patients with LAD I. Defective T-cell proliferation in response to the staphylococcal enterotoxin superantigen was more severe in splenocytes from CD18^{−/−}, CD11b^{−/−}, or CD11d KO mice than in CD11a^{−/−} splenocytes but was normal in CD11c^{−/−} splenocytes⁵⁸. The defects in CD11b^{−/−} or CD11d^{−/−} lymphocytes have been traced to transient expression of CD11b and CD11d on thymocytes, which appears to be required for normal T-cell development⁵⁸.

CD11a–d contributed in variable degrees to the adhesion of phagocytes to inflamed endothelium^{21,42,59,60}. Transendothelial neutrophil migration in the tumor necrosis factor-induced air pouch inflammation model was reduced in CD11a KO⁶¹, as in CD18 KO, but was surprisingly increased in CD11b KO mice⁶⁰. Migration within interstitial matrices was integrin independent^{62,63}. Phagocytosis of serum-opsonized particles (with its associated oxygen free radical production, cytokine release, and degranulation) and phagocytosis-induced apoptosis in neutrophils were defective in CD11b^{−/−} null mouse cells⁶⁴, confirming an essential role for CD11b in the programmed elimination of neutrophils that have already phagocytosed their target pathogens. Toll receptor-mediated responses were enhanced in CD11b^{−/−} macrophages, rendering mice more susceptible to sepsis and endotoxin shock⁶⁵. Thus, whereas neutrophil adhesion to endothelium may require all four β_2 integrins, transendothelial migration appears to be mainly CD11a dependent, while phagocytosis is mediated primarily by CD11b⁶⁶. Curiously, CD11b KO mice are obese⁶⁷, a phenotype not seen in patients with LAD I, suggesting a role for CD11b in regulating fat metabolism at least in mice. The number of mast cells in the peritoneal cavity is also reduced in CD11b KO mice²⁷, suggesting an additional role in mast cell development. Mast cells play an important role in the early peritoneal neutrophil response during experimental peritonitis in mice and this may explain the increased mortality of CD11b KO mice after acute septic peritonitis²⁷.

Integrin structure

The αA domain

Structural studies of integrins began with the identification of a novel metal-ion-dependent adhesion site (MIDAS) in an extracellular von Willebrand factor type A (vWFA) domain (αA or αI domain) present in integrin CD11b⁶⁸. The vWFA domain is found in eight additional integrin α -subunits (Figure 1) as well as in several structurally unrelated proteins^{69,70}. αA from CD11b (CD11bA) mediates Mg^{2+} -dependent binding of the receptor to ligands^{68,71}. αA also mediates ligand binding in the other αA -containing integrins. The first crystal structure of recombinant CD11bA showed a compact GTPase-like fold comprising a central, mostly parallel β -sheet surrounded on both sides by seven amphipathic α -helices (Figure 2a). The catalytic site found at the apex in GTPases is replaced with MIDAS, where an Mg^{2+} ion is coordinated by three surface loops (Figure 2b). A solvent-exposed glutamate (E) or aspartate (D) from ligand completes an octahedral coordination sphere around the Mg^{2+} ion⁶⁹. This crystal structure first explained why Mg^{2+} is required for integrin binding to all physiologic ligands and why a solvent-accessible acidic residue from ligand is essential for binding to any integrin. Ligand-binding specificity in αA domains is imparted by the variable surface-exposed side chains surrounding the MIDAS motif.

The αA domain also exists in a second ligand-free “closed” conformation^{72,73}, where the ligand coordinating carboxyl oxygen is replaced with a water molecule (Figure 2c). Superposing the two structures shows the key tertiary changes associated with ligand

Figure 2. Structural comparisons of inactive and active αA domains. (a) Ribbon diagrams showing the superposed structures of inactive (gray) and active (yellow) αA domain from the β_2 integrin CD11b/CD18. Major conformational differences are indicated by arrows. The two phenylalanine residues (F275 and F302) buried in the inactive form are solvent exposed in the active state. A glutamate (E) from ligand is shown in the active (ligand-bound) state, ligating the metal-ion-dependent adhesion site (MIDAS) Mg^{2+} monodentately. (b, c) The MIDAS motif in the active (b) and inactive (c) states. The metal ion at MIDAS is coordinated by residues from three surface loops, and a carboxyl oxygen from ligand completes the octahedral coordinating sphere (b). In the inactive state, an oxygen atom from a water molecule replaces the ligand oxygen, and D242 from the third surface loop moves in to coordinate the metal directly (c). Coordinating oxygen atoms are in red, and hydrogen bonds are shown by dashed red lines. Direct bonds to the metal ion are shown as blue sticks. Water molecules are labeled $\omega 1$ – $\omega 3$.

binding: an inward movement of the N-terminal $\alpha 1$ helix, rearrangements of the metal-coordinating residues at MIDAS, and a 10 Å downward shift of the C-terminal $\alpha 7$ helix at the opposite pole to MIDAS^{72,74} (Figure 2a). The key residues that stabilize the closed conformation have been identified, and mutations of some of these residues converted the closed into the open conformation^{75–79}. Locking the open conformation with a pair of disulfides allowed crystallization of this form in the absence of ligand^{80,81}. Crystal structures of αA domains from other integrins (for example, $\alpha_2\beta_1$ ⁸²), complement factors (for example, factors B and C2^{83,84}), certain matrix proteins⁸⁵, and microorganisms (for example, anthrax⁸⁶) were subsequently determined. These structures displayed the same basic conformational changes observed in CD11bA, underscoring their functional importance. In solution, recombinant wild-type CD11bA exists in an equilibrium where the proportion of the closed to the open state is nearly 9:1^{75,79}; the presence of ligand shifts this equilibrium in favor of the open state.

The integrin ectodomain

The modular nature of an integrin was first revealed with the determination of the crystal structure of the ectodomain of the αA -lacking integrin $\alpha_v\beta_3$ in its unliganded state⁸⁷ and when occupied by a cyclic peptide ligand containing the prototypical Arg-Gly-Asp motif⁸⁸. The α_v subunit is composed of a seven-bladed

propeller domain, followed by a thigh domain and two large Ig-like Calf domains. The β_3 subunit comprises an N-terminal plexin-semaphorin-integrin (PSI) domain, an Ig-like “hybrid” domain in which an αA -like domain (βA) is inserted, four successive epidermal growth factor (EGF)-like domains (IE1–4), and a novel membrane-proximal β -tail domain (βTD) (Figure 3a, b). In the full-length integrin, Calf2 and βTD each is attached to a transmembrane (TM) domain and a short cytoplasmic tail. An unexpected feature of the $\alpha_v\beta_3$ ectodomain is a sharp bending in the structure at the α -genu (between the thigh and calf1 domains) and the β -genu (within IE2) (Figure 3a). Extension at the knees is expected to produce an extended integrin (Figure 3b), which resembles the shape seen previously using rotary shadowing electron microscopy⁸⁹.

In αA -lacking integrins, the integrin head is formed of the βA and propeller domains (Figure 3a, b), which associate non-covalently in a manner that resembles the association of the $G\alpha$ and $G\beta$ subunits of heterotrimeric G proteins⁸⁷. In αA -containing integrins, the head also contains the αA domain, which projects from a surface loop in the propeller (Figure 3c). The heterodimer-disruptive point mutations found in the β_2 (CD18) and β_3 subunits in patients with LAD I and Glanzmann’s thrombasthenia (a bleeding disorder), respectively, map to the βA domain and commonly involve residues at the βA -propeller interface⁸⁷. As in αA domains, an acidic residue

Figure 3. Structure of the integrin ectodomain. (a) Ribbon drawing of the bent ectodomain from integrin $\alpha_v\beta_3$. α_v is in light blue, and the 12 domains of β_3 are shown in different colors for better visualization. The two tails would extend into the plasma membrane in the native integrin. (b) Model of $\alpha_v\beta_3$ ectodomain linearized by introducing breaks at the α and β genu (arrows). The modular nature of the ectodomain can be readily appreciated. H, hybrid domain; PSI, plexin-semaphorin-integrin. (c) The integrin head from the αA -containing β_2 integrin CD11c/CD18. (d) Interactions between arginine-glycine-aspartate (RGD)-containing ligand peptide (yellow) and the $\alpha_v\beta_3$ head. The peptide aspartate (D) completes the metal ion coordination sphere at metal-ion-dependent adhesion site (MIDAS), and ligand arginine forms salt bridges in the propeller pocket. αV and β_3 residues are labeled light blue and orange, respectively. The three metal ions in β_3 at MIDAS, adjacent to MIDAS (ADMIDAS), and ligand-associated metal binding site (LIMBS) are shown in cyan, magenta, and gray, respectively, and their coordinating residues displayed. The upper portion of the $\alpha 1$ helix and the loop between strand-F and $\alpha 7$ helix (F- $\alpha 7$) are also shown. Oxygen and nitrogen atoms are in red and blue, respectively. Hydrogen bonds and salt bridges (distance cutoff, 3.5 Å) are represented with dotted lines. (e) Superposed structures of αA and βA domains. Shown are the two inserted loops in βA : the specificity determining loop (SDL) and heterodimer-association loop. The hydrophobic phenylalanine residue at the top of $\alpha 7$ helix that contacts $\alpha 1$ helix in αA is replaced in βA with an ionic interaction mediated by ADMIDAS ion.

from ligand completes the octahedral coordination of Mg^{2+} at MIDAS, an interaction stabilized by the arginine residue in the prototypical arginine-glycine-aspartate (RGD) motif, which inserts into a pocket in the propeller domain, making contacts with acidic residues in the pocket (Figure 3d). Five metal ions (Ca^{2+} or Mn^{2+}) occupy the bases of blades 4–7 of the α_v propeller and the α -genu (Figure 3a, b); these may help rigidify the interfaces the thigh domain makes with the propeller base proximally and the top of Calfl distally.

The structure of inactive βA is largely superimposable onto that of αA , except for two loop insertions: one forming the core of the interface with the α -subunit's propeller and the second—the specificity determining loop, SDL—contributing to ligand binding as well as to the βA /propeller interface in some integrins (for example, $\alpha_{IIB}\beta_3$) (Figure 3e). In addition, a Ca^{2+} ion at a site adjacent to MIDAS (ADMIDAS) in βA links the two activation-sensitive $\alpha 1$ and $\alpha 7$ helices, stabilizing this domain in the closed state; in αA , this ionic interaction is replaced by a hydrophobic one (Figure 3e). In addition to the ADMIDAS ion, ligand-bound βA contains a ligand-associated metal binding site (LIMBS), which is occupied by Ca^{2+} in ligand- or pseudoligand-bound integrins^{88,90}. The structure of LIMBS in ligand-free integrins is regulated by the α -subunit's propeller domain⁹¹ and this may explain the variable metal ion occupancy of this site (sometimes also called synergy metal binding site).

In αA -containing integrins, the ligand-associated downward shift of the C-terminal $\alpha 7$ helix enables an invariant glutamate at the bottom of $\alpha 7$ to ligate the βA MIDAS ion (Figure 4); mutation of this residue to alanine blocked integrin function⁹². This led us to propose that αA serves as an intrinsic ligand for βA in αA -containing integrins. Blocking this coordination by the synthetic molecule XVA143 severs the αA link to βA and blocks integrin signaling⁹³. Support for this “ligand-relay” model came from the recent crystal structure of the CD11c/CD18 ectodomain⁹⁴. Thus, the βA domain transduces outside-in signals that are triggered by either extrinsic (in αA -lacking integrins) or intrinsic (in αA -containing integrins) ligands.

Integrin transmembrane and cytoplasmic tails

The structure of the lipid-embedded α_{IIB} and β_3 single-pass TM helices was determined by using solution nuclear magnetic resonance (NMR) spectroscopy⁹⁵. The structure revealed two dominant integrin TM association motifs or clasps: an outer (membrane-proximal) and an inner (membrane-distal) one that extends to include the adjacent cytoplasmic salt bridge between α_{IIB} and β_3 ⁹⁶. The two clasps maintain the integrin in the inactive state⁹⁷. Another structure in hydrophobic organic solvent invokes several differences in the membrane-proximal clasp regions, especially the helical conformation of α_{IIB} in the latter versus a reverse turn in the former structure⁹⁸. It is unclear at present whether this difference in the membrane proximal regions in the NMR structures reflects the nature of the lipid-like TM environment in which the TM domains were incorporated or reflects potential changes in response to binding of cytosolic regulators such as filamin^{99,100}.

Figure 4. The ligand-relay model. The downward movement of the c-terminal $\alpha 7$ helix (dark blue) triggered by ligand binding to αA allows an invariant glutamate (E) at the bottom of the $\alpha 7$ helix to reach and ligate the βA metal-ion-dependent adhesion site (MIDAS) ion (cyan), thus relaying the ligand occupancy state of αA to βA .

Binding of the N-terminal talin head to the membrane proximal NPxY/F motif in the β cytoplasmic tail destabilizes the α - β TM association^{101,102}. Recruitment of talin to the plasma membrane requires ras-related protein 1 (Rap1) and its effector Rap1-GTP-interacting adaptor molecule (RIAM), and the latter is critical *in vivo* for inside-out signaling of β_2 but not β_1 or β_3 integrins^{103,104}. Kindlins have been reported to modulate receptor affinity¹⁰⁵ or avidity¹⁰⁶ or both. Kindlins bind the distal NPxY/F motif and a preceding threonine-containing region of the β cytoplasmic tail¹⁰⁷ but do not appear to destabilize α - β TM association¹⁰⁸. The structural basis for regulation of integrins by kindlins remains to be elucidated. Loss of kindlin 3 causes LAD III, a disease characterized by bleeding diathesis (defective $\alpha_{IIB}\beta_3$ function) and defective leukocyte recruitment to sites of infection (defective β_2 integrin function)¹⁰⁵.

Integrin activation

Integrins are normally expressed in an inactive state on the cell surface. This is critical, as it allows leukocytes and platelets, for example, to freely circulate in blood with minimal aggregation or interaction with blood vessel walls. Binding of an agonist such as a chemokine or a cytokine (for example, granulocyte-macrophage colony-stimulating factor¹⁰⁹) to their respective receptors initiates inside-out signals that rapidly switch the integrin into the active

state. Integrins stored in intracellular pools (for example, CD11b/CD18^{18,110,111} and $\alpha_{\text{IIb}}\beta_3$ ¹¹²) are also recruited to the cell surface in response to agonists, but this process appears to follow the switch of the integrin to the active state^{113,114}.

The structural basis for integrin inside-out signaling is debated. Following publication of the bent ectodomain structure⁸⁷, a “switchblade” model envisioned that in the bent state, the ligand-binding site in βA (and αA in αA -containing integrin) is inaccessible to soluble ligand because of its proposed proximity to the plasma membrane. It is suggested, therefore, that the integrin linearizes to expose the ligand-binding site¹¹⁵, which also allows an approximately 80° swingout of the hybrid domain and a switch of βA into high affinity⁹⁰ (Figure 5). An alternate β TD-centric deadbolt model¹¹⁶ proposed that the ligand-binding site in βA is already accessible to soluble macromolecular ligand in the native integrin¹¹⁷ and can assume high affinity in the compact structure¹¹⁸ and that genuextension occurs following binding of ligands or

ligand-mimetic drugs to the cellular integrin¹¹⁹. Movements of the membrane proximal β TD resulting from unpacking of the immediately distal TM segments disrupt β TD contacts with βA and hybrid domains, allowing the central switch of βA into the active state with minimal hybrid domain swingout¹¹⁸.

Both models are supported by experimental data. Two-dimensional imaging using negative-stain electron microscopy (EM) showed a greater proportion of extended integrin ectodomains in the presence of the metal ion Mn^{2+} (used as a mimic of inside-out signaling), and hydrodynamic studies showed an increase in the stokes radius of the $\alpha_v\beta_3$ ectodomain in Mn^{2+} ¹¹⁵. However, cryoelectron tomography showed that $\alpha_{\text{IIb}}\beta_3$ maintained the compact (bent) conformation after Mn^{2+} activation in a membrane environment¹²⁰. Differences in sample preparation, sampling bias in EM, and differences in ectodomain constructs may explain these discrepancies. A recent EM study of full-length integrin $\alpha_{\text{IIb}}\beta_3$ in lipid-embedded nanodiscs showed a small increase in the extended conformation when the integrin was activated by talin¹²¹. More recently, negative-stain EM of membrane-embedded full-length $\alpha_{\text{IIb}}\beta_3$ showed that the active ligand-free $\alpha_{\text{IIb}}\beta_3$ is mainly bent but that the ligand-bound receptor is predominantly extended¹²². High-resolution quantitative dynamic footprinting microscopy combined with homogenous conformation-reporter binding assays showed that a substantial fraction of β_2 integrins on the surface of human neutrophils assumed a high-affinity bent conformation¹²³. Because of the profound influence of the TM domains on integrin activation by inside-out signaling, settling the ongoing debate regarding the structural basis of integrin activation will likely require a three-dimensional crystal structure determination of a full-length native integrin in its native inactive and high-affinity states.

Ligand-bound integrins cluster, especially when occupied by multivalent ligands, and transduce outside-in signals leading to cell adhesion via new connections established between the integrin cytoplasmic tails and filamentous actin¹²⁴. In migrating cells, inward movement of the actin cytoskeleton from the site of assembly at the leading edge toward the cell center generates a pulling force across the nascent-integrin-matrix linkages and this unbends the liganded integrin and strengthens adhesion at these sites by accelerating recruitment of additional cytoskeletal and signaling proteins to the clustered integrins¹²⁵. As this pulling force increases in the moving cell, integrin-ligand bonds eventually break and integrins are endocytosed and this allows rear detachment and directional cell movement at the leading edge. Known adaptor proteins involved in integrin uptake and recycling have been recently reviewed¹²⁶.

β_2 integrins as therapeutic targets

Although β_2 integrins are critical for innate and adaptive immunity, they can also induce serious pathology if improperly activated. Hyperadherent leukocytes may, for example, bind and injure the blood vessel wall, leukoaggregate intravascularly resulting in blocked capillaries or emboli, or compromise immune surveillance, thus contributing to inflammatory and autoimmune diseases. The finding that CD18 deficiency impaired the inflammatory response suggested that knockout of CD18 or CD11 or

Figure 5. Structural changes in the βA domain following ligand binding. The superposed structures of βA domain of the β_3 subunit in its unliganded (pdb 3ije) state and bound to cacodylate (acting as a pseudoligand, L) (pdb 1ty3) are shown in magenta and green, respectively. The main movements involve the $\alpha 1$ and $\alpha 7$ helices, loop F- $\alpha 7$, and the hybrid domain. In the unliganded state, helix $\alpha 1$ and F- $\alpha 7$ loop are connected via the adjacent to MIDAS (ADMIDAS) ion (magenta), and no metal-ion-dependent adhesion site (MIDAS) or ligand-associated metal binding site (LIMBS) atoms are detected. In the liganded state, a ligand oxygen coordinates MIDAS, and the $\alpha 1$ helix moves inwards (reported by tyrosine 122, Y122), bringing the ADMIDAS ion closer to the MIDAS ion and breaking the ionic contact with the F- $\alpha 7$ loop. These changes are coupled with a one-turn descent of the $\alpha 7$ helix and a 135° swingout of the hybrid domain in structures lacking the integrin leg domains.

inhibiting their functions in leukocytes using antibodies may be beneficial in treating inflammatory or autoimmune diseases⁷. A similar logic has been successful in targeting platelet $\alpha_{IIb}\beta_3$ to inhibit pathologic thrombosis and this resulted in two orthosteric inhibitors, eptifibatide and tirofiban, and an allosteric inhibitor Abciximab, all three in clinical use¹²⁷.

Genetic deficiency of CD18, CD11a, or CD11b or targeting β_2 integrins with various inhibitory antibodies in rodents ameliorated ischemia-reperfusion injury (IRI) in heart attacks, cerebral stroke, burns, and traumatic shock as well as autoimmune injury of the brain (multiple sclerosis), lung (asthma), and skin (psoriasis) and in native or transplanted kidneys (reviewed in 128). However, humanized forms of these mAbs failed when tested in patients with myocardial infarction, stroke, traumatic shock, multiple sclerosis, asthma, or acute rejection (reviewed in 128). An anti-CD11a mAb that showed promise in treating psoriasis was withdrawn because of fatal brain infections resulting from reactivation of JC virus¹²⁹. Inadequate design of some of the trials¹²⁸, important differences in immune responses between rodents and humans¹³⁰, and the relatively short follow-up period in the preclinical studies may have contributed to these failures. In addition, most clinical studies evaluating IRI syndromes used anti-CD18 antibodies, which might have acted allosterically to switch the integrin into the active proadhesive state. This scenario has precedence in β_3 integrin-targeted mAb or small-molecule drugs, which act as partial agonists, unbending the integrin, thus exposing neoepitopes recognized by natural antibodies and leading to immune thrombocytopenia and bleeding,

or inducing proadhesive outside-in signaling leading to paradoxical thrombosis^{131,132}. Therefore, recent attempts have been made to solve the problem of partial agonism, making use of the advances made in structural biology of integrins. The central role of the A-domain in integrin activation and signaling made it a main focus of drug development efforts. The non-RGD-containing small molecules RUC-1, RUC-2, and UR-2922 were identified and act by inserting into the arginine-binding pocket in the propeller domain^{133,134}, thus interfering with the stable binding of RGD-containing ligands. RUC-2 also binds to the β_3 MIDAS residue E220 thus displacing the Mg^{2+} at MIDAS¹³³. *In vivo* studies of RUC-1 administered intraperitoneally demonstrated anti-thrombotic effects in microvascular injury models in mice¹³⁵.

We have approached the problem of partial agonism by identifying orthosteric inhibitors of integrin β_2 (mAb107,¹¹⁷) and β_3 (a mutant high-affinity form of fibronectin-10, hFN10¹³⁶) that do not induce the activating proadhesive changes in the αA or βA domains, respectively. mAb107 stabilized the inhibitory Ca^{2+} in place of the proadhesive Mg^{2+} at the CD11bA MIDAS, freezing the β_2 integrin CD11b/CD18 in the inactive conformation¹¹⁷ (Figure 6a). hFN10 bound the βA MIDAS of integrin $\alpha_v\beta_3$ and blocked the activating inward movement of the $\alpha 1$ helix (Figure 6b), which is critical for integrin unbending and outside-in signaling¹³⁶. *In vivo* studies in monkeys showed that mAb107 ameliorated leukocyte-mediated inflammation in a severe IRI kidney model, salvaging kidney function from otherwise irreversible failure several months after a single injection of the mAb at the onset of IRI¹³⁷.

Figure 6. Structural basis of integrin inhibition by “pure” orthosteric inhibitors. (a) Ribbon drawing showing structure of αA from the β_2 integrin CD11b/CD18 bound to the pure ligand-mimetic antagonist mAb107 (in magenta). For clarity, only the ligand-Asp of mAb107 is shown. The unusual symmetric bidentate ligation of the antibody-derived ligand Asp to a hepta-coordinated metal-ion-dependent adhesion site (MIDAS) Ca^{2+} (blue sticks) prevents the tertiary changes associated with Mg^{2+} -dependent ligand binding. The superposed structure in gray is that of unliganded αA from CD11b/CD18. (b) Structure of unliganded βA from $\alpha_v\beta_3$ (pdb 3ije) (gray), superposed on the structure of βA in complex with a fibronectin-10-derived “pure antagonist” (magenta). Only the RGDW residues (in ball and stick) from ligand are shown (pdb 4mmz). Ligand-associated inward movement of the $\alpha 1$ helix and the resulting activating tertiary changes are prevented by a π - π interaction involving the ligand tryptophan (W) and βA 's tyrosine 122 (Y122). The ionic bridge (dashed red lines) between $\alpha 1$ and $\alpha 7$ helices is unaffected by binding of the pure orthosteric inhibitor. The metal ions at ADMIDAS, MIDAS, and LIMBS are in magenta (or gray), cyan, and dark gray, respectively.

Conclusions

Much has been learned since Cohnheim's and Metchnikoff's respective descriptions of leukocyte transendothelial migration and phagocytosis. The receptors involved have been identified, their critical role in innate and adaptive immunity defined, and their structures elucidated, revealing the atomic basis for their Mg^{2+} dependency, ligand binding, and activation. Although putting the myriad interactions mediated by integrins into structural and biologic contexts remains a major challenge, the recent advances already made form a basis for structure-based discovery of effective and safer anti-inflammatory and anti-thrombosis therapeutics targeting these dynamic receptors.

Competing interests

The author declares that he has no competing interests.

Grant information

The author's work presented in this review was supported by National Institutes of Health grants DK088327, DK48549, and DK007540 from the National Institutes of Diabetes, Digestive and Kidney Diseases.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Harburger DS, Calderwood DA: **Integrin signalling at a glance.** *J Cell Sci.* 2009; **122**(Pt 2): 159–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gahmberg CG, Fagerholm SC, Nurmi SM, *et al.*: **Regulation of integrin activity and signalling.** *Biochim Biophys Acta.* 2009; **1790**(6): 431–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Horton ER, Astudillo P, Humphries MJ, *et al.*: **Mechanosensitivity of integrin adhesion complexes: role of the consensus adhesome.** *Exp Cell Res.* 2016; **343**(1): 7–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mócsai A, Walzog B, Lowell CA: **Intracellular signalling during neutrophil recruitment.** *Cardiovasc Res.* 2015; **107**(3): 373–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cohnheim J: **Lectures On General Pathology: A Handbook for Practitioners and Students.** 2nd edition, (translated from the second German edition) London, The New Sydenham Society. 1889; 1.
[Reference Source](#)
- Metchnikoff E: **Sur la lutte des cellules de l'organisme contre l'invasion des microbes.** *Ann Inst Pasteur I.* 1887; 321.
[Reference Source](#)
- Arnaout MA: **Leukocyte adhesion molecules deficiency: its structural basis, pathophysiology and implications for modulating the inflammatory response.** *Immunol Rev.* 1990; **114**(1): 145–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arnaout MA, Pitt J, Cohen HJ, *et al.*: **Deficiency of a granulocyte-membrane glycoprotein (gp150) in a boy with recurrent bacterial infections.** *N Engl J Med.* 1982; **306**(12): 693–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dana N, Pitt J, Todd RF, *et al.*: **Deficiency of a monocyte-granulocyte surface glycoprotein (Mo1) in man.** [abstract]. *Clin Res.* 1983; **31**: 489.
- Springer T, Galfré G, Secher DS, *et al.*: **Mac-1: a macrophage differentiation antigen identified by monoclonal antibody.** *Eur J Immunol.* 1979; **9**(4): 301–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Remold-O'Donnell E: **Macrophage component gp160, a major trypsin-sensitive surface glycoprotein.** *J Exp Med.* 1980; **152**(6): 1699–708.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Todd RF 3rd, Nadler LM, Schlossman SF: **Antigens on human monocytes identified by monoclonal antibodies.** *J Immunol.* 1981; **126**(4): 1435–42.
[PubMed Abstract](#)
- Trowbridge IS, Omary MB: **Molecular complexity of leukocyte surface glycoproteins related to the macrophage differentiation antigen Mac-1.** *J Exp Med.* 1981; **154**(5): 1517–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sanchez-Madrid F, Nagy JA, Robbins E, *et al.*: **A human leukocyte differentiation antigen family with distinct alpha-subunits and a common beta-subunit: the lymphocyte function-associated antigen (LFA-1), the C3bi complement receptor (OKM1/Mac-1), and the p150,95 molecule.** *J Exp Med.* 1983; **158**(6): 1785–803.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beatty PG, Ochs HD, Harlan JM, *et al.*: **Absence of monoclonal-antibody-defined protein complex in boy with abnormal leukocyte function.** *Lancet.* 1984; **1**(8376): 535–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- LeBien TW, Kersey JH: **A monoclonal antibody (TA-1) reactive with human T lymphocytes and monocytes.** *J Immunol.* 1980; **125**(5): 2208–14.
[PubMed Abstract](#)
- Davignon D, Martz E, Reynolds T, *et al.*: **Lymphocyte function-associated antigen 1 (LFA-1): a surface antigen distinct from Lys-2,3 that participates in T lymphocyte-mediated killing.** *Proc Natl Acad Sci U S A.* 1981; **78**(7): 4535–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Arnaout MA, Spits H, Terhorst C, *et al.*: **Deficiency of a leukocyte surface glycoprotein (LFA-1) in two patients with Mo1 deficiency. Effects of cell activation on Mo1/LFA-1 surface expression in normal and deficient leukocytes.** *J Clin Invest.* 1984; **74**(4): 1291–300.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dana N, Todd RF 3rd, Pitt J, *et al.*: **Deficiency of a surface membrane glycoprotein (Mo1) in man.** *J Clin Invest.* 1984; **73**(1): 153–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lanier LL, Arnaout MA, Schwarting R, *et al.*: **p150/95, Third member of the LFA-1/CR3 polypeptide family identified by anti-Leu M5 monoclonal antibody.** *Eur J Immunol.* 1985; **15**(7): 713–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Danilenko DM, Rossitto PV, van der Vieren M, *et al.*: **A novel canine leukointegrin, alpha d beta 2, is expressed by specific macrophage subpopulations in tissue and a minor CD8+ lymphocyte subpopulation in peripheral blood.** *J Immunol.* 1995; **155**(1): 35–44.
[PubMed Abstract](#)
- Springer TA, Thompson WS, Miller LJ, *et al.*: **Inherited deficiency of the Mac-1, LFA-1, p150,95 glycoprotein family and its molecular basis.** *J Exp Med.* 1984; **160**(6): 1901–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dana N, Clayton LK, Tennen DG, *et al.*: **Leukocytes from four patients with complete or partial Leu-CAM deficiency contain the common beta-subunit precursor and beta-subunit messenger RNA.** *J Clin Invest.* 1987; **79**(3): 1010–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nelson C, Rabb H, Arnaout MA: **Genetic cause of leukocyte adhesion molecule deficiency. Abnormal splicing and a missense mutation in a conserved region of CD18 impair cell surface expression of beta 2 integrins.** *J Biol Chem.* 1992; **267**(5): 3351–7.
[PubMed Abstract](#)
- Anderson DC, Schmalstieg FC, Finegold MJ, *et al.*: **The severe and moderate phenotypes of heritable Mac-1, LFA-1 deficiency: their quantitative definition and relation to leukocyte dysfunction and clinical features.** *J Infect Dis.* 1985; **152**(4): 668–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arnaout MA: **Structure and function of the leukocyte adhesion molecules CD11/CD18.** *Blood.* 1990; **75**(5): 1037–50.
[PubMed Abstract](#)
- Rosenkranz AR, Coxon A, Maurer M, *et al.*: **Impaired mast cell development and innate immunity in Mac-1 (CD11b/CD18, CR3)-deficient mice.** *J Immunol.* 1998; **161**(12): 6463–7.
[PubMed Abstract](#)
- Wagner C, Hansch GM, Stegmaier S, *et al.*: **The complement receptor 3, CR3 (CD11b/CD18), on T lymphocytes: activation-dependent up-regulation and regulatory function.** *Eur J Immunol.* 2001; **31**(4): 1173–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lahmers KK, Hedges JF, Jutila MA, *et al.*: **Comparative gene expression by WC1+ gamma delta and CD4+ alpha beta T lymphocytes, which respond to Anaplasma**

- marginale**, demonstrates higher expression of chemokines and other myeloid cell-associated genes by WC1⁺ gamma delta T cells. *J Leukoc Biol*. 2006; 80(4): 939–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Pilling D, Fan T, Huang D, *et al.*: Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. *PLoS One*. 2009; 4(10): e7475.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Rubtsova K, Rubtsov AV, van Dyk LF, *et al.*: T-box transcription factor T-bet, a key player in a unique type of B-cell activation essential for effective viral clearance. *Proc Natl Acad Sci U S A*. 2013; 110(34): E3216–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 32. Clements M, Gershenovich M, Chaber C, *et al.*: Differential Ly6C Expression after Renal Ischemia-Reperfusion Identifies Unique Macrophage Populations. *J Am Soc Nephrol*. 2016; 27(1): 159–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 33. Ghosh EE, Yang Y, Tung J, *et al.*: CD11b expression distinguishes sequential stages of peritoneal B-1 development. *Proc Natl Acad Sci U S A*. 2008; 105(13): 5195–200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Keizer GD, Borst J, Visser W, *et al.*: Membrane glycoprotein p150,95 of human cytotoxic T cell clone is involved in conjugate formation with target cells. *J Immunol*. 1987; 138(10): 3130–6.
[PubMed Abstract](#)
 35. van der Vieren M, Le Trong H, Wood CL, *et al.*: A novel leukointegrin, alpha d beta 2, binds preferentially to ICAM-3. *Immunity*. 1995; 3(6): 683–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Miyazaki Y, Vieira-de-Abreu A, Harris ES, *et al.*: Integrin $\alpha_p\beta_2$ (CD11d/CD18) is expressed by human circulating and tissue myeloid leukocytes and mediates inflammatory signaling. *PLoS One*. 2014; 9(11): e112770.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 37. Noti JD: Expression of the myeloid-specific leukocyte integrin gene CD11d during macrophage foam cell differentiation and exposure to lipoproteins. *Int J Mol Med*. 2002; 10(6): 721–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Yakubenko VP, Belevych N, Mishchuk D, *et al.*: The role of integrin $\alpha_p\beta_2$ (CD11d/CD18) in monocyte/macrophage migration. *Exp Cell Res*. 2008; 314(14): 2569–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Tian L, Yoshihara Y, Mizuno T, *et al.*: The neuronal glycoprotein telencephalin is a cellular ligand for the CD11a/CD18 leukocyte integrin. *J Immunol*. 1997; 158(2): 928–36.
[PubMed Abstract](#)
 40. Béchard D, Scherpereel A, Hammad H, *et al.*: Human endothelial-cell specific molecule-1 binds directly to the integrin CD11a/CD18 (LFA-1) and blocks binding to intercellular adhesion molecule-1. *J Immunol*. 2001; 167(6): 3099–106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 41. Ostermann G, Weber KS, Zernecke A, *et al.*: JAM-1 is a ligand of the β_2 integrin LFA-1 involved in transendothelial migration of leukocytes. *Nat Immunol*. 2002; 3(2): 151–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Sadhu C, Ting HJ, Lipsky B, *et al.*: CD11c/CD18: novel ligands and a role in delayed-type hypersensitivity. *J Leukoc Biol*. 2007; 81(6): 1395–403.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Bilsland CA, Diamond MS, Springer TA: The leukocyte integrin p150,95 (CD11c/CD18) as a receptor for IC3b. Activation by a heterologous beta subunit and localization of a ligand recognition site to the I domain. *J Immunol*. 1994; 152(9): 4582–9.
[PubMed Abstract](#)
 44. Malhotra V, Hogg N, Sim RB: Ligand binding by the p150,95 antigen of U937 monocytic cells: properties in common with complement receptor type 3 (CR3). *Eur J Immunol*. 1986; 16(9): 1117–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Blackford J, Reid HW, Pappin DJ, *et al.*: A monoclonal antibody, 3/22, to rabbit CD11c which induces homotypic T cell aggregation: evidence that ICAM-1 is a ligand for CD11c/CD18. *Eur J Immunol*. 1996; 26(3): 525–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Ihanus E, Uotila LM, Toivanen A, *et al.*: Red-cell ICAM-4 is a ligand for the monocyte/macrophage integrin CD11c/CD18: characterization of the binding sites on ICAM-4. *Blood*. 2007; 109(2): 802–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Xia Y, Velticka V, Yan J, *et al.*: The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function in generating a primed state of the receptor that mediates cytotoxic activation in response to IC3b-opsonized target cells. *J Immunol*. 1999; 162(4): 2281–90.
[PubMed Abstract](#)
 48. Yakubenko VP, Solovjov DA, Zhang L, *et al.*: Identification of the binding site for fibrinogen recognition peptide γ 383-395 within the $\alpha_m\beta_2$ -domain of integrin $\alpha_m\beta_2$. *J Biol Chem*. 2001; 276(17): 13995–4003.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Humphries JD, Byron A, Humphries MJ: Integrin ligands at a glance. *J Cell Sci*. 2006; 119(pt 19): 3901–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. van der Vieren M, Crowe DT, Hoekstra D, *et al.*: The leukocyte integrin alpha D beta 2 binds VCAM-1: evidence for a binding interface between I domain and VCAM-1. *J Immunol*. 1999; 163(4): 1984–90.
[PubMed Abstract](#)
 51. Scharffetter-Kochanek K, Lu H, Norman K, *et al.*: Spontaneous skin ulceration and defective T cell function in CD18 null mice. *J Exp Med*. 1998; 188(1): 119–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Shier P, Ngo K, Fung-Leung WP: Defective CD8+ T cell activation and cytolytic function in the absence of LFA-1 cannot be restored by increased TCR signaling. *J Immunol*. 1999; 163(9): 4826–32.
[PubMed Abstract](#)
 53. Shier P, Otulakowski G, Ngo K, *et al.*: Impaired immune responses toward alloantigens and tumor cells but normal thymic selection in mice deficient in the beta2 integrin leukocyte function-associated antigen-1. *J Immunol*. 1996; 157(12): 5375–86.
[PubMed Abstract](#)
 54. Bose TO, Pham QM, Jellison ER, *et al.*: CD11a regulates effector CD8 T cell differentiation and central memory development in response to infection with *Listeria monocytogenes*. *Infect Immun*. 2013; 81(4): 1140–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 55. Schmits R, Kündig TM, Baker DM, *et al.*: LFA-1-deficient mice show normal CTL responses to virus but fail to reject immunogenic tumor. *J Exp Med*. 1996; 183(4): 1415–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. Johnston B, Kubes P: The alpha4-integrin: an alternative pathway for neutrophil recruitment? *Immunol Today*. 1999; 20(12): 545–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 57. Taooka Y, Chen J, Yednock T, *et al.*: The integrin alpha9beta1 mediates adhesion to activated endothelial cells and transendothelial neutrophil migration through interaction with vascular cell adhesion molecule-1. *J Cell Biol*. 1999; 145(2): 413–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Wu H, Rodgers JR, Perrard XY, *et al.*: Deficiency of CD11b or CD11d results in reduced staphylococcal enterotoxin-induced T cell response and T cell phenotypic changes. *J Immunol*. 2004; 173(1): 297–306.
[PubMed Abstract](#) | [Publisher Full Text](#)
 59. Amaout MA, Lanier LL, Faller DV: Relative contribution of the leukocyte molecules Mo1, LFA-1, and p150,95 (LeuM5) in adhesion of granulocytes and monocytes to vascular endothelium is tissue- and stimulus-specific. *J Cell Physiol*. 1988; 137(2): 305–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Ding ZM, Babensee JE, Simon SI, *et al.*: Relative contribution of LFA-1 and Mac-1 to neutrophil adhesion and migration. *J Immunol*. 1999; 163(9): 5029–38.
[PubMed Abstract](#)
 61. Lu H, Smith CW, Perrard J, *et al.*: LFA-1 is sufficient in mediating neutrophil emigration in Mac-1-deficient mice. *J Clin Invest*. 1997; 99(6): 1340–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Luo D, McGettrick HM, Stone PC, *et al.*: The roles of integrins in function of human neutrophils after their migration through endothelium into interstitial matrix. *PLoS One*. 2015; 10(2): e0118593.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 63. Lämmermann T, Bader BL, Monkley SJ, *et al.*: Rapid leukocyte migration by integrin-independent flowing and squeezing. *Nature*. 2008; 453(7191): 51–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 64. Coxon A, Rieu P, Barkalow FJ, *et al.*: A novel role for the beta 2 integrin CD11b/CD18 in neutrophil apoptosis: a homeostatic mechanism in inflammation. *Immunity*. 1996; 5(6): 653–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. Han C, Jin J, Xu S, *et al.*: Integrin CD11b negatively regulates TLR-triggered inflammatory responses by activating Syk and promoting degradation of MyD88 and TRIF via Cbl-b. *Nat Immunol*. 2010; 11(8): 734–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Milde R, Ritter J, Tennent GA, *et al.*: Multinucleated Giant Cells Are Specialized for Complement-Mediated Phagocytosis and Large Target Destruction. *Cell Rep*. 2015; 13(9): 1937–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 67. Dong ZM, Gutierrez-Ramos JC, Coxon A, *et al.*: A new class of obesity genes encodes leukocyte adhesion receptors. *Proc Natl Acad Sci U S A*. 1997; 94(14): 7526–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 68. Michishita M, Videm V, Amaout MA: A novel divalent cation-binding site in the A domain of the beta 2 integrin CR3 (CD11b/CD18) is essential for ligand binding. *Cell*. 1993; 72(6): 857–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 69. Lee JO, Rieu P, Amaout MA, *et al.*: Crystal structure of the A domain from the alpha subunit of integrin CR3 (CD11b/CD18). *Cell*. 1995; 80(4): 631–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. Whittaker CA, Hynes RO: Distribution and evolution of von Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and elsewhere. *Mol Biol Cell*. 2002; 13(10): 3369–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Diamond MS, Garcia-Aguilar J, Bickford JK, *et al.*: The I domain is a major recognition site on the leukocyte integrin Mac-1 (CD11b/CD18) for four distinct

- adhesion ligands. *J Cell Biol.* 1993; **120**(4): 1031–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Lee JO, Bankston LA, Arnaout MA, *et al.*: Two conformations of the integrin A-domain (I-domain): a pathway for activation? *Structure.* 1995; **3**(12): 1333–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 73. Qu A, Leahy DJ: Crystal structure of the I-domain from the CD11a/CD18 (LFA-1, alpha L beta 2) integrin. *Proc Natl Acad Sci U S A.* 1995; **92**(22): 10277–81.
[PubMed Abstract](#) | [Free Full Text](#)
 74. **F** Bajic G, Yattine L, Sim RB, *et al.*: Structural insight on the recognition of surface-bound opsonins by the integrin I domain of complement receptor 3. *Proc Natl Acad Sci U S A.* 2013; **110**(41): 16426–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 75. Li R, Rieu P, Griffith DL, *et al.*: Two functional states of the CD11b A-domain: correlations with key features of two Mn²⁺-complexed crystal structures. *J Cell Biol.* 1998; **143**(6): 1523–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 76. Xiong JP, Li R, Essafi M, *et al.*: An isoleucine-based allosteric switch controls affinity and shape shifting in integrin CD11b A-domain. *J Biol Chem.* 2000; **275**(49): 38762–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 77. Shimaoka M, Shifman JM, Jing H, *et al.*: Computational design of an integrin I domain stabilized in the open high affinity conformation. *Nat Struct Biol.* 2000; **7**(8): 674–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Aquilina A, Korda M, Bergelson JM, *et al.*: A novel gain-of-function mutation of the integrin alpha2 VVFA domain. *Eur J Biochem.* 2002; **269**(4): 1136–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. McCleverty CJ, Liddington RC: Engineered allosteric mutants of the integrin alphaMbeta2 I domain: structural and functional studies. *Biochem J.* 2003; **372**(Pt 1): 121–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Ponnuraj K, Xu Y, Macon K, *et al.*: Structural analysis of engineered Bb fragment of complement factor B: insights into the activation mechanism of the alternative pathway C3-convertase. *Mol Cell.* 2004; **14**(1): 17–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. **F** Shimaoka M, Xiao T, Liu JH, *et al.*: Structures of the alpha L I domain and its complex with ICAM-1 reveal a shape-shifting pathway for integrin regulation. *Cell.* 2003; **112**(1): 99–111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 82. Emsley J, King SL, Bergelson JM, *et al.*: Crystal structure of the I domain from integrin alpha2beta1. *J Biol Chem.* 1997; **272**(45): 28512–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Bhattacharya AA, Luper ML Jr, Staunton DE, *et al.*: Crystal structure of the A domain from complement factor B reveals an integrin-like open conformation. *Structure.* 2004; **12**(3): 371–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Krishnan V, Xu Y, Macon K, *et al.*: The crystal structure of C2a, the catalytic fragment of classical pathway C3 and C5 convertase of human complement. *J Mol Biol.* 2007; **367**(1): 224–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. Emsley J, Cruz M, Handin R, *et al.*: Crystal structure of the von Willebrand Factor A1 domain and implications for the binding of platelet glycoprotein Ib. *J Biol Chem.* 1998; **273**(17): 10396–401.
[PubMed Abstract](#) | [Publisher Full Text](#)
 86. Santelli E, Bankston LA, Leppla SH, *et al.*: Crystal structure of a complex between anthrax toxin and its host cell receptor. *Nature.* 2004; **430**(7002): 905–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. **F** Xiong JP, Stehle T, Diefenbach B, *et al.*: Crystal structure of the extracellular segment of integrin alpha Vbeta3. *Science.* 2001; **294**(5541): 339–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 88. **F** Xiong JP, Stehle T, Zhang R, *et al.*: Crystal structure of the extracellular segment of integrin alpha Vbeta3 in complex with an Arg-Gly-Asp ligand. *Science.* 2002; **296**(5565): 151–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 89. Nermut MV, Green NM, Eason P, *et al.*: Electron microscopy and structural model of human fibronectin receptor. *EMBO J.* 1988; **7**(13): 4093–9.
[PubMed Abstract](#) | [Free Full Text](#)
 90. **F** Xiao T, Takagi J, Collier BS, *et al.*: Structural basis for allostery in integrins and binding to fibrinogen-mimetic therapeutics. *Nature.* 2004; **432**(7013): 59–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 91. Rui X, Mehrbod M, van Aghoven JF, *et al.*: The α -subunit regulates stability of the metal ion at the ligand-associated metal ion-binding site in β_3 integrins. *J Biol Chem.* 2014; **289**(33): 23256–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. **F** Alonso JL, Essafi M, Xiong JP, *et al.*: Does the integrin alphaA domain act as a ligand for its betaA domain? *Curr Biol.* 2002; **12**(10): R340–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 93. Shimaoka M, Salas A, Yang W, *et al.*: Small molecule integrin antagonists that bind to the β_2 subunit I-like domain and activate signals in one direction and block them in the other. *Immunity.* 2003; **19**(3): 391–402.
[PubMed Abstract](#) | [Publisher Full Text](#)
 94. **F** Sen M, Yuki K, Springer TA: An internal ligand-bound, metastable state of a leukocyte integrin, $\alpha_5\beta_2$. *J Cell Biol.* 2013; **203**(4): 629–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. Lau T, Kim C, Ginsberg MH, *et al.*: The structure of the integrin α IIb β 3 transmembrane complex explains integrin transmembrane signalling. *EMBO J.* 2009; **28**(9): 1351–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 96. Hughes PE, Diaz-Gonzalez F, Leong L, *et al.*: Breaking the integrin hinge. A defined structural constraint regulates integrin signaling. *J Biol Chem.* 1996; **271**(12): 6571–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 97. Ginsberg MH: Integrin activation. *BMB Rep.* 2014; **47**(12): 655–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 98. **F** Yang J, Ma YQ, Page RC, *et al.*: Structure of an integrin α IIb β 3 transmembrane-cytoplasmic heterocomplex provides insight into integrin activation. *Proc Natl Acad Sci U S A.* 2009; **106**(42): 17729–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 99. Sharma CP, Ezzell RM, Arnaout MA: Direct interaction of filamin (ABP-280) with the beta 2-integrin subunit CD18. *J Immunol.* 1995; **154**(7): 3461–70.
[PubMed Abstract](#)
 100. **F** Liu J, Das M, Yang J, *et al.*: Structural mechanism of integrin inactivation by filamin. *Nat Struct Mol Biol.* 2015; **22**(7): 383–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 101. Calderwood DA, Campbell ID, Critchley DR: Talins and kindlins: partners in integrin-mediated adhesion. *Nat Rev Mol Cell Biol.* 2013; **14**(8): 503–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 102. Anthis NJ, Wegener KL, Ye F, *et al.*: The structure of an integrin/talin complex reveals the basis of inside-out signal transduction. *EMBO J.* 2009; **28**(22): 3623–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 103. **F** Klaproth S, Sperandio M, Pinheiro EM, *et al.*: Loss of the Rap1 effector RIAM results in leukocyte adhesion deficiency due to impaired β_2 integrin function in mice. *Blood.* 2015; **126**(25): 2704–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 104. **F** Su W, Wynne J, Pinheiro EM, *et al.*: Rap1 and its effector RIAM are required for lymphocyte trafficking. *Blood.* 2015; **126**(25): 2695–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 105. **F** Moser M, Nieswandt B, Ussar S, *et al.*: Kindlin-3 is essential for integrin activation and platelet aggregation. *Nat Med.* 2008; **14**(3): 325–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 106. **F** Ye F, Petrich BG, Anekal P, *et al.*: The mechanism of kindlin-mediated activation of integrin α IIb β 3. *Curr Biol.* 2013; **23**(22): 2288–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 107. Harburger DS, Bouaouina M, Calderwood DA: Kindlin-1 and -2 directly bind the C-terminal region of β integrin cytoplasmic tails and exert integrin-specific activation effects. *J Biol Chem.* 2009; **284**(17): 11485–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 108. **F** Bledzka K, Liu J, Xu Z, *et al.*: Spatial coordination of kindlin-2 with talin head domain in interaction with integrin β cytoplasmic tails. *J Biol Chem.* 2012; **287**(29): 24585–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 109. Metelitsa LS, Gillies SD, Super M, *et al.*: Antidialoganglioside/granulocyte macrophage-colony-stimulating factor fusion protein facilitates neutrophil antibody-dependent cellular cytotoxicity and depends on Fc γ RII (CD32) and Mac-1 (CD11b/CD18) for enhanced effector cell adhesion and azurophilic granule exocytosis. *Blood.* 2002; **99**: 4166–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
 110. Todd RF 3rd, Arnaout MA, Rosin RE, *et al.*: Subcellular localization of the large subunit of Mo1 (Mo1 alpha; formerly gp 110), a surface glycoprotein associated with neutrophil adhesion. *J Clin Invest.* 1984; **74**(4): 1280–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. Arnaout MA, Wang EA, Clark SC, *et al.*: Human recombinant granulocyte-macrophage colony-stimulating factor increases cell-to-cell adhesion and surface expression of adhesion-promoting surface glycoproteins on mature granulocytes. *J Clin Invest.* 1986; **78**(2): 597–601.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 112. Shattil SJ, Kashiwagi H, Pampori N: Integrin signaling: the platelet paradigm. *Blood.* 1998; **91**(8): 2645–57.
[PubMed Abstract](#)
 113. Vedder NB, Harlan JM: Increased surface expression of CD11b/CD18 (Mac-1) is not required for stimulated neutrophil adherence to cultured endothelium. *J Clin Invest.* 1988; **81**(3): 676–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 114. **F** Kim M, Carman CV, Yang W, *et al.*: The primacy of affinity over clustering in regulation of adhesiveness of the integrin $\{\alpha\}_n\{\beta\}_2$. *J Cell Biol.* 2004; **167**(6): 1241–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

115. **F** Takagi J, Petre BM, Walz T, *et al.*: **Global conformational rearrangements in integrin extracellular domains in outside-in and inside-out signaling.** *Cell*. 2002; **110**(5): 599–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
116. Xiong JP, Stehle T, Goodman SL, *et al.*: **New insights into the structural basis of integrin activation.** *Blood*. 2003; **102**(4): 1155–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
117. Mahalingam B, Ajroud K, Alonso JL, *et al.*: **Stable coordination of the inhibitory Ca²⁺ ion at the metal ion-dependent adhesion site in integrin CD11b/CD18 by an antibody-derived ligand aspartate: implications for integrin regulation and structure-based drug design.** *J Immunol*. 2011; **187**(12): 6393–401.
[PubMed Abstract](#) | [Publisher Full Text](#)
118. **F** Adair BD, Xiong JP, Maddock C, *et al.*: **Three-dimensional EM structure of the ectodomain of integrin {alpha}V{beta}3 in a complex with fibronectin.** *J Cell Biol*. 2005; **168**(7): 1109–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
119. Cox D, Smith R, Quinn M, *et al.*: **Evidence of platelet activation during treatment with a GPIIb/IIIa antagonist in patients presenting with acute coronary syndromes.** *J Am Coll Cardiol*. 2000; **36**(5): 1514–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
120. Ye F, Liu J, Winkler H, *et al.*: **Integrin $\alpha_{IIb}\beta_3$ in a Membrane Environment Remains the Same Height after Mn²⁺ Activation when Observed by Cryoelectron Tomography.** *J Mol Biol*. 2008; **378**(5): 976–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
121. **F** Ye F, Hu G, Taylor D, *et al.*: **Recreation of the terminal events in physiological integrin activation.** *J Cell Biol*. 2010; **188**(1): 157–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
122. **F** Dai A, Ye F, Taylor DW, *et al.*: **The Structure of a Full-length Membrane-embedded Integrin Bound to a Physiological Ligand.** *J Biol Chem*. 2015; **290**(45): 27168–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
123. **F** Fan Z, McArdle S, Marki A, *et al.*: **Neutrophil recruitment limited by high-affinity bent β_2 integrin binding ligand in cis.** *Nat Commun*. 2016; **7**: 12658.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
124. Friedland JC, Lee MH, Boettiger D: **Mechanically activated integrin switch controls $\alpha_5\beta_1$ function.** *Science*. 2009; **323**(5914): 642–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
125. Puklin-Faucher E, Sheetz MP: **The mechanical integrin cycle.** *J Cell Sci*. 2009; **122**(Pt 2): 179–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
126. Morse EM, Brahme NN, Calderwood DA: **Integrin cytoplasmic tail interactions.** *Biochemistry*. 2014; **53**(5): 810–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
127. Collier BS: **Translating from the rivers of Babylon to the coronary bloodstream.** *J Clin Invest*. 2012; **122**(11): 4293–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
128. Yonekawa K, Harlan JM: **Targeting leukocyte integrins in human diseases.** *J Leukoc Biol*. 2005; **77**(2): 129–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
129. Major EO: **Progressive multifocal leukoencephalopathy in patients on immunomodulatory therapies.** *Annu Rev Med*. 2010; **61**: 35–47.
[PubMed Abstract](#) | [Publisher Full Text](#)
130. Mestas J, Hughes CC: **Of mice and not men: differences between mouse and human immunology.** *J Immunol*. 2004; **172**(5): 2731–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
131. Aster RH, Curtis BR, McFarland JG, *et al.*: **Drug-induced immune thrombocytopenia: pathogenesis, diagnosis, and management.** *J Thromb Haemost*. 2009; **7**(6): 911–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
132. Cox D: **Oral GPIIb/IIIa antagonists: what went wrong?** *Curr Pharm Des*. 2004; **10**(14): 1587–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
133. **F** Zhu J, Choi WS, McCoy JG, *et al.*: **Structure-guided design of a high-affinity platelet integrin $\alpha_{IIb}\beta_3$ receptor antagonist that disrupts Mg²⁺ binding to the MIDAS.** *Sci Transl Med*. 2012; **4**(125): 125ra32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
134. Cox D, Brennan M, Moran N: **Integrins as therapeutic targets: lessons and opportunities.** *Nat Rev Drug Discov*. 2010; **9**(10): 804–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
135. Blue R, Kowalska MA, Hirsch J, *et al.*: **Structural and therapeutic insights from the species specificity and *in vivo* antithrombotic activity of a novel $\alpha_{IIb}\beta_3$ -specific $\alpha_{IIb}\beta_3$ antagonist.** *Blood*. 2009; **114**(1): 195–201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
136. van Aghoven JF, Xiong JP, Alonso JL, *et al.*: **Structural basis for pure antagonism of integrin $\alpha_V\beta_3$ by a high-affinity form of fibronectin.** *Nat Struct Mol Biol*. 2014; **21**(4): 383–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
137. Cosimi A, Dehnadi A, Smith R, *et al.*: **Targeting Leukocyte Integrin CD11b/CD18 With a Novel mAb Salvages Renal Function Following an Otherwise Irreversible Ischemic Insult in Cynomolgus Monkeys.** [abstract]. *Am J Transplant*. 2015; **15**(suppl 3).
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Tobias Ulmer

Department of Biochemistry & Molecular Biology and Zilkha Neurogenetic Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

Competing Interests: No competing interests were disclosed.

2. Jun Qin

Department of Molecular Cardiology, Lerner Research Institute, Cleveland Clinic, Cleveland, OH, USA

Competing Interests: No competing interests were disclosed.

3. Klaus Ley

Division of Inflammation Biology; Department of Bioengineering, La Jolla Institute for Allergy and Immunology, La Jolla, CA, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research