REVIEW

Perioperative regional anaesthesia and postoperative longer-term outcomes [version 1; peer review: 3 approved]

Jan G. Jakobsson ¹, Mark Z. Johnson ^{2,3}

¹Department of Anaesthesia & Intensive Care, Institution for Clinical Science, Karolinska Institutet, Danderyds University Hospital, Stockholm, Sweden

²Department of Anaesthesia & Critical Care, Mater Misericordiae University Hospital, Dublin, Ireland

³School of Medicine, University College Dublin, Dublin, Ireland

V1 First published: 11 Oct 2016, 5(F1000 Faculty Rev):2501
<https://doi.org/10.12688/f1000research.9100.1>

Latest published: 11 Oct 2016, 5(F1000 Faculty Rev):2501
<https://doi.org/10.12688/f1000research.9100.1>

Abstract

Regional anaesthesia provides effective anaesthesia and analgesia in the perioperative setting. Central neuraxial blocks—that is, spinal and epidural blocks—are well established as an alternative or adjunct to general anaesthesia. Peripheral blocks may be used as part of multimodal anaesthesia/analgesia in perioperative practice, reducing the need for opioid analgesics and enhancing early recovery. Furthermore, regional anaesthesia has increased in popularity and may be done with improved ease and safety with the introduction of ultrasound-guided techniques. The effects of local anaesthetics and regional anaesthesia on long-term outcomes such as morbidity, mortality, the quality of recovery beyond the duration of analgesia, and whether it can expedite the resumption of activities of daily living are less clear. It has also been suggested that regional anaesthesia may impact the risk of metastasis after cancer surgery. This article provides an overview of current evidence around quality of recovery, risk for delirium, long-term effects, and possible impact on cancer disease progression associated with the clinical use of local and regional anaesthetic techniques. In summary, there is still a lack of robust data that regional anaesthesia has a clinical impact beyond its well-acknowledged beneficial effects of reducing pain, reduced opioid consumption, and improved quality of early recovery. Further high-quality prospective studies on long-term outcomes are warranted.

Keywords

local anaesthesia , neuraxial block , inflammation , immune function , cancer re-occurrence , metastasis

Open Peer Review

Approval Status

	1	2	3
version 1 11 Oct 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Markus W. Hollmann**, University of Amsterdam, Amsterdam, The Netherlands
2. **Donal Buggy**, University College Dublin, Dublin, Ireland
3. **Juan P. Cata**, University of Texas MD Anderson Cancer Center, Houston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Jan G. Jakobsson (jan.jakobsson@ki.se)

Competing interests: JJ declares that he has no competing interests in relation to this article. MJ declares that he has no competing interests.

Grant information: JJ has received research grants from and taken part in advisory boards for Abbott, Baxter, AstraZeneca, Pfizer, MSD, and Marquette and acts as safety physician for Linde Healthcare.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Jakobsson JG and Johnson MJ. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Jakobsson JG and Johnson MJ. **Perioperative regional anaesthesia and postoperative longer-term outcomes [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2501
<https://doi.org/10.12688/f1000research.9100.1>

First published: 11 Oct 2016, 5(F1000 Faculty Rev):2501 <https://doi.org/10.12688/f1000research.9100.1>

Introduction

The effect of local anaesthesia on nerve conductivity has a long history. Modern amide local anaesthetics are reassuringly safe, providing dose-dependent, reversible nerve conduction blockade. The beneficial effects of regional anaesthesia on postoperative pain are well known. Local anaesthesia may be administered by local infiltration, peripheral nerve block, or central neuraxial block depending on what effects are sought. Local infiltration anaesthesia is simple and safe. Peripheral nerve blocks are increasingly popular because of ultrasound techniques, which provide enhanced visualisation of its performance. The positive effects on postoperative pain and reduced opioid consumption are well acknowledged. There are recent reviews of upper-extremity blocks¹, lower-extremity blocks², and abdominal-wall blocks³ that demonstrate these effects. There may not be a major difference between local infiltration and peripheral nerve blocks for the management of postoperative pain in the first 24 hours following some surgical procedures (for example, total hip replacement⁴) in contrast to total knee arthroplasty (TKA), where wide variability in postoperative pain scores is documented. In a Cochrane meta-analysis, Chan *et al.* supported the use of regional anaesthesia for postoperative analgesia following TKA⁵. In May 2016, Hu *et al.* published a meta-analysis showing that the two techniques were similarly effective for early pain management following TKA⁶.

There is also increasing interest in the systemic effects of local anaesthetics. The potential beneficial effect of local anaesthesia on inflammation and immune function has long been suggested. In 2000, Hollmann and Durieux⁷ had already addressed the potential anti-inflammatory effects of local anaesthetics, reviewing the literature on local anaesthetic effects on the inflammatory response and especially on inflammatory cells (mainly polymorphonuclear granulocytes but also macrophages and monocytes)⁸. The exact mechanisms of action are not clear but seem to involve a reversible interaction with membrane proteins and lipids, thus regulating cell metabolic activity, migration, exocytosis, and phagocytosis^{9,10}. The clinical anti-inflammatory effect of intravenous lidocaine is today reasonably well documented. The beneficial effects of intravenous lidocaine on recovery have been described; this technique has been shown to reduce the time to return of bowel function and shorter hospital stay after abdominal surgery¹¹. The most recent meta-analysis, published in 2016¹², supports the beneficial effects on pain but calls for further studies, as the quality of evidence was limited as a result of trial heterogeneity and small sample sizes. There are also several studies looking at the potential effect of local anaesthetics on immune function^{13–15}. The findings of these studies have not necessarily been shown to have clinical significance and have yet to be translated into routine clinical practice.

Central blocks, epidural and spinal anaesthesia, and morbidity and mortality in general surgery

Neuraxial blocks are widely used in addition or as an alternative to general anaesthesia for general surgery as well as for postoperative pain management. The beneficial effects during the early phase of recovery are well documented. A recent meta-analysis comparing general anaesthetic with epidural and general anaesthetic with opioid analgesia showed improved pain control and bowel function in the epidural and general anaesthesia group¹⁶. The impact

of regional anaesthesia on longer-term outcomes has become of increasing interest¹⁷. An overview of Cochrane systematic reviews of neuraxial blockade for the prevention of postoperative mortality and major morbidity found weak evidence for reduced mortality and pneumonia in high-risk patients who received neuraxial anaesthesia¹⁸. There is still a lack of evidence to show a difference in morbidity and mortality between regional anaesthesia and general anaesthesia for urological and vascular surgery^{19,20}. In 2014, Popping *et al.* published a meta-analysis supporting the use of combined general and epidural anaesthesia, since epidural analgesia reduced mortality and improved a multitude of cardiovascular, respiratory, and gastrointestinal morbidity endpoints compared with patients receiving systemic analgesia²¹. However, this article had numerous limitations, including certain methodological flaws and the inclusion of unpublished data.

Regional anaesthesia and orthopaedic surgery

In a meta-analysis published in 2013, Barbosa *et al.* found no significant long-term benefit associated with regional anaesthesia as compared with general anaesthesia for orthopaedic surgery²². Memtsoudis *et al.* found that regional anaesthesia reduced the risk of major complications in a subset of patients with obstructive sleep apnoea undergoing joint arthroplasty in comparison with patients who received combined neuraxial and general or general anaesthesia alone (16.0%, 17.2%, and 18.1%, respectively; $P = 0.0177$)²³.

However, the benefits associated with epidural anaesthesia have been questioned. The quality of studies included in meta-analyses is low and some of the positive effects are merely dependent on single studies in high-risk patients²⁴. The potential benefit on morbidity and mortality from regional anaesthesia for acute femoral fracture repair surgery has also recently been challenged. Neuman *et al.* showed shorter hospital stay but no difference in mortality²⁵. Seitz *et al.* showed regional anaesthesia to be associated with similar mortality but a lower need for intensive care unit admission compared with general anaesthesia in older adults with dementia who underwent surgery for hip fracture repair²⁶. Brox *et al.* found one-year mortality to be only marginally lower in patients with neuraxial anaesthesia compared with general anaesthesia (odds ratio (OR) = 0.84, confidence interval (CI) 0.70–1.0) but to be similar in patients with combined regional and general anaesthesia (OR = 1.3, CI 0.70–2.3)²⁷. White *et al.* likewise found no difference in mortality associated with neuraxial block or general anaesthesia but emphasised the importance of maintaining adequate blood pressure²⁸. In February 2016, Johnson *et al.* published a meta-analysis of neuraxial anaesthesia compared with general anaesthesia for hip or knee arthroplasty (or both) showing no significant differences in any outcomes studied: mortality, surgical duration, surgical site or chest infections, nerve palsies, postoperative nausea and vomiting, or thromboembolic disease when anti-thrombotic prophylaxis was used²⁹.

Thus, regional anaesthesia provides effective early postoperative analgesia and is a valid alternative to general anaesthesia. The effects on early postoperative pain are well documented, but its impact beyond the early period and effects on major morbidity and mortality are not obvious.

Regional anaesthesia and delirium

Postoperative delirium (PD) is commonly seen in the elderly. The risk in the elderly having orthopaedic surgery has been reported to be around 17% with huge variability³⁰. The impact of anaesthetic technique on early PD and more long-lasting cognitive deficits such as postoperative cognitive dysfunction (POCD) has been under scrutiny for decades. In 1995, Williams-Russo *et al.*³¹ compared epidural and general anaesthesia in 262 patients undergoing elective primary total knee replacement with a median age of 69 years. The type of anaesthesia—general or neuraxial—did not affect the magnitude or pattern of POCD. The International Study of Postoperative Cognitive Dysfunction (ISPOCD) group³² published the results of a multi-centre study assessing POCD and the impact of anaesthetic technique, general anaesthesia versus regional anaesthesia. Early testing (one week postoperatively) showed a higher incidence among patients having general anaesthesia, but no difference was noted at 3 months. In 2010, Mason *et al.* conducted a meta-analysis and found no impact of anaesthetic technique on the risk for PD, but general anaesthesia was found to have a small, non-significant association with an increased rate of POCD (OR = 1.34, CI 0.93–1.95)³³. There was no evidence of publication bias. Davis *et al.* came to a similar conclusion in a meta-analysis published in 2014; of 16 included studies, 13 showed no difference and three showed some minor effect³⁴. There is a need for further high-quality studies, and there are ongoing studies on this topic.

Regional anaesthesia and quality of recovery

The impact of regional anaesthesia on the quality of recovery, as assessed by objective multi-dimensional tools (for example, Quality of Recovery 15 or Postoperative Quality of Recovery score), beyond the early postoperative phase is sparsely studied. Catro-Alves *et al.* compared spinal anaesthesia and general anaesthesia for elective hysterectomy and assessed recovery by using the global quality of recovery-40 questionnaire (QoR-40) up to 48 hours after surgery but not beyond³⁵. The beneficial effects on pain and quality of early recovery were confirmed. Liu *et al.* conducted a study amongst elderly patients undergoing knee arthroplasty in general anaesthesia or peripheral block and followed intraoperative and postoperative course up to day 7 with the Postoperative Quality of Recovery Scale (PostopQRS) tool³⁶. The intraoperative and early postoperative course favoured the block technique, but at day 7 no difference was found. Resumption of activities of daily living and patient satisfaction were similar between the two groups.

Regional anaesthesia and cancer

There is considerable interest in whether local anaesthetics and regional anaesthesia may impact cancer progression and recurrence. The potential mechanisms by which cancer progression may be influenced include attenuation of the surgical stress response, reduced pain and opioid requirements, and also the direct action that some local anaesthetics may have on cancer cells. It is also suggested that local anaesthetics and regional anaesthesia may

have an effect on immune function and inflammation. There are studies assessing immune function in the experimental setting³⁷. Despite mounting *in vitro* data suggesting positive effects^{38,39}, *in vivo* studies and well-designed human trials are lacking⁴⁰. The clinical relevance of these results must be interpreted with great caution and cannot necessarily be directly translated into human clinical practice. Several recent articles have raised the question of whether anaesthetic technique and drug choice could influence long-term tumour progression^{41,42}. However, these articles are retrospective analyses of studies designed to address a different clinical question. In 2006, Exadaktylos *et al.* published a retrospective study suggesting that paravertebral block in combination with general anaesthesia for primary breast cancer surgery could reduce the risk for tumour progression as compared with standard anaesthesia and postoperative opioids for pain management⁴³. In 2014, however, a Cochrane systematic review concluded that the evidence is insufficient to make any firm recommendations⁴⁴.

Although regional anaesthesia provides effective early postoperative analgesia and promotes enhanced recovery, there is insufficient data to promote a change in clinical practice in order to affect cancer progression^{45–48}. We await results from ongoing studies to better define the role of anaesthesia for patients with cancer⁴⁹.

Conclusions

Regional anaesthesia is effective at reducing pain and opioid consumption during the early postoperative phase. The impact of regional anaesthesia on the risk for cognitive side effects of surgery, PD, and POCD seems minor. Data on the effects of local anaesthesia, local infiltration, and regional anaesthesia on quality of recovery assessed by multi-domain tools are sparse. The documentation around long-term outcomes and the potential beneficial effects on morbidity and mortality is also weak, and further high-quality studies are warranted. The potential effect on cancer progression is not proven. There are several ongoing prospective randomised studies that may help define whether regional anaesthesia techniques could not only provide effective postoperative pain but also impact cancer progression. However, it should be acknowledged that the number of patients needed to have sufficient power to show any potential significant effect is large, such studies are expensive, and long follow-up times are required.

Competing interests

JJ declares that he has no competing interests in relation to this article. MJZ declares that he has no competing interests.

Grant information

JJ has received research grants from and taken part in advisory boards for Abbott, Baxter, AstraZeneca, Pfizer, MSD, and Marquette and acts as safety physician for Linde Healthcare.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Brattwall M, Jildensstål P, Warren Stombérg M, *et al.*: **Upper extremity nerve block: how can benefit, duration, and safety be improved? An update [version 1; referees: 3 approved].** *F1000Res.* 2016; 5: pii: F1000 Faculty Rev-907.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Kehlet H, Aasvang EK: **Regional or general anesthesia for fast-track hip and knee replacement - what is the evidence? [version 1; referees: 2 approved].** *F1000Res.* 2015; 4: pii: F1000 Faculty Rev-1449.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Jakobsson J, Wickerts L, Forsberg S, *et al.*: **Transversus abdominal plane (TAP) block for postoperative pain management: a review [version 1; referees: 2 approved].** *F1000Res.* 2015; 4: pii: F1000 Faculty Rev-1359.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Jimenez-Almonte JH, Wyles CC, Wyles SP, *et al.*: **Is Local Infiltration Analgesia Superior to Peripheral Nerve Blockade for Pain Management After THA: A Network Meta-analysis.** *Clin Orthop Relat Res.* 2016; 474(2): 495–516.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
5. Chan EY, Fransen M, Parker DA, *et al.*: **Femoral nerve blocks for acute postoperative pain after knee replacement surgery.** *Cochrane Database Syst Rev.* 2014; (5): CD009941.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
6. Hu B, Lin T, Yan SG, *et al.*: **Local Infiltration Analgesia Versus Regional Blockade for Postoperative Analgesia in Total Knee Arthroplasty: A Meta-analysis of Randomized Controlled Trials.** *Pain Physician.* 2016; 19(4): 205–14.
[PubMed Abstract](#) | [F1000 Recommendation](#)
7. Hollmann MW, Durieux ME: **Local anesthetics and the inflammatory response: a new therapeutic indication?** *Anesthesiology.* 2000; 93(3): 858–75.
[PubMed Abstract](#)
8. Picardi S, Cartellieri S, Groves D, *et al.*: **Local anesthetic-induced inhibition of human neutrophil priming: the influence of structure, lipophilicity, and charge.** *Reg Anesth Pain Med.* 2013; 38(1): 9–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
9. Cassuto J, Sinclair R, Bonderovic M: **Anti-inflammatory properties of local anesthetics and their present and potential clinical implications.** *Acta Anaesthesiol Scand.* 2006; 50(3): 265–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. van der Wal SEI, van den Heuvel SA, Radema SA, *et al.*: **The *in vitro* mechanisms and *in vivo* efficacy of intravenous lidocaine on the neuroinflammatory response in acute and chronic pain.** *Eur J Pain.* 2016; 20(5): 655–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. Herroeder S, Pecher S, Schönherr ME, *et al.*: **Systemic lidocaine shortens length of hospital stay after colorectal surgery: a double-blinded, randomized, placebo-controlled trial.** *Ann Surg.* 2007; 246(2): 192–200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
12. Weibel S, Jokinen J, Pace NL, *et al.*: **Efficacy and safety of intravenous lidocaine for postoperative analgesia and recovery after surgery: a systematic review with trial sequential analysis.** *Br J Anaesth.* 2016; 116(6): 770–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
13. Le Cras AE, Galley HF, Webster NR: **Spinal but not general anesthesia increases the ratio of T helper 1 to T helper 2 cell subsets in patients undergoing transurethral resection of the prostate.** *Anesth Analg.* 1998; 87(6): 1421–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Wada H, Seki S, Takahashi T, *et al.*: **Combined spinal and general anesthesia attenuates liver metastasis by preserving TH1/TH2 cytokine balance.** *Anesthesiology.* 2007; 106(3): 499–506.
[PubMed Abstract](#)
15. Werdehausen R, Braun S, Fazeli S, *et al.*: **Lipophilicity but not stereospecificity is a major determinant of local anaesthetic-induced cytotoxicity in human T-lymphoma cells.** *Eur J Anaesthesiol.* 2012; 29(1): 35–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
16. Guay J, Nishimori M, Kopp S: **Epidural local anaesthetics versus opioid-based analgesic regimens for postoperative gastrointestinal paralysis, vomiting and pain after abdominal surgery.** *Cochrane Database Syst Rev.* 2016; (7): CD001893.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
17. Xu J, Chen XM, Ma CK, *et al.*: **Peripheral nerve blocks for postoperative pain after major knee surgery.** *Cochrane Database Syst Rev.* 2014; (12): CD010937.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Guay J, Choi P, Suresh S, *et al.*: **Neuraxial blockade for the prevention of postoperative mortality and major morbidity: an overview of Cochrane systematic reviews.** *Cochrane Database Syst Rev.* 2014; (1): CD010108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
19. Barbosa FT, Castro AA: **Neuraxial anesthesia versus general anesthesia for urological surgery: systematic review.** *Sao Paulo Med J.* 2013; 131(3): 179–86.
[PubMed Abstract](#) | [F1000 Recommendation](#)
20. Barbosa FT, Jucá MJ, Castro AA, *et al.*: **Neuraxial anaesthesia for lower-limb revascularization.** *Cochrane Database Syst Rev.* 2013; (7): CD007083.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
21. Popping DM, Elia N, van Aken HK, *et al.*: **Impact of epidural analgesia on mortality and morbidity after surgery: systematic review and meta-analysis of randomized controlled trials.** *Ann Surg.* 2014; 259(6): 1056–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
22. Barbosa FT, Castro AA, Sousa-Rodrigues CF: **Neuraxial anesthesia for orthopedic surgery: systematic review and meta-analysis of randomized clinical trials.** *Sao Paulo Med J.* 2013; 131(6): 411–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
23. Memtsoudis SG, Stundner O, Rasul R, *et al.*: **Sleep apnea and total joint arthroplasty under various types of anesthesia: a population-based study of perioperative outcomes.** *Reg Anesth Pain Med.* 2013; 38(4): 274–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
24. Kooij FO, Schlack WS, Preckel B, *et al.*: **Does regional analgesia for major surgery improve outcome? Focus on epidural analgesia.** *Anesth Analg.* 2014; 119(3): 740–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. Neuman MD, Rosenbaum PR, Ludwig JM, *et al.*: **Anesthesia technique, mortality, and length of stay after hip fracture surgery.** *JAMA.* 2014; 311(24): 2508–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
26. Seitz DP, Gill SS, Bell CM, *et al.*: **Postoperative medical complications associated with anesthesia in older adults with dementia.** *J Am Geriatr Soc.* 2014; 62(11): 2102–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. Brox WT, Chan PH, Cafri G, *et al.*: **Similar mortality with general or regional anesthesia in elderly hip fracture patients.** *Acta Orthop.* 2016; 87(2): 152–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. White SM, Moppett IK, Griffiths R, *et al.*: **Secondary analysis of outcomes after 11,085 hip fracture operations from the prospective UK Anaesthesia Sprint Audit of Practice (ASAP-2).** *Anaesthesia.* 2016; 71(5): 506–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. Johnson RL, Kopp SL, Burkle CM, *et al.*: **Neuraxial vs general anaesthesia for total hip and total knee arthroplasty: a systematic review of comparative-effectiveness research.** *Br J Anaesth.* 2016; 116(2): 163–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. Scott JE, Mathias JL, Kneebone AC: **Incidence of delirium following total joint replacement in older adults: a meta-analysis.** *Gen Hosp Psychiatry.* 2015; 37(3): 223–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. Williams-Russo P, Sharrock NE, Mattis S, *et al.*: **Cognitive effects after epidural vs general anesthesia in older adults. A randomized trial.** *JAMA.* 1995; 274(1): 44–50.
[PubMed Abstract](#)
32. Rasmussen LS, Johnson T, Kuipers HM, *et al.*: **Does anaesthesia cause postoperative cognitive dysfunction? A randomised study of regional versus general anaesthesia in 438 elderly patients.** *Acta Anaesthesiol Scand.* 2003; 47(3): 260–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Mason SE, Noel-Storr A, Ritchie CW: **The impact of general and regional anesthesia on the incidence of post-operative cognitive dysfunction and post-operative delirium: a systematic review with meta-analysis.** *J Alzheimers Dis.* 2010; 22(Suppl 3): 67–79.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Davis N, Lee M, Lin AY, *et al.*: **Postoperative cognitive function following general versus regional anesthesia: a systematic review.** *J Neurosurg Anesthesiol.* 2014; 26(4): 369–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Catro-Alves LJ, De Azevedo VL, De Freitas Braga TF, *et al.*: **The effect of neuraxial versus general anesthesia techniques on postoperative quality of recovery and analgesia after abdominal hysterectomy: a prospective, randomized, controlled trial.** *Anesth Analg.* 2011; 113(6): 1480–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Liu J, Yuan W, Wang X, *et al.*: **Peripheral nerve blocks versus general anesthesia for total knee replacement in elderly patients on the postoperative quality of recovery.** *Clin Interv Aging.* 2014; 9: 341–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. Mikawa K, Akamarsu H, Nishina K, *et al.*: **Effects of ropivacaine on human neutrophil function: comparison with bupivacaine and lidocaine.** *Eur J Anaesthesiol.* 2003; 20(2): 104–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Chang YC, Hsu YC, Liu CL, *et al.*: **Local anesthetics induce apoptosis in human thyroid cancer cells through the mitogen-activated protein kinase pathway.** *PLoS One.* 2014; 9(2): e89563.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

39. **F** Xuan W, Zhao H, Hankin J, *et al.*: **Local anesthetic bupivacaine induced ovarian and prostate cancer apoptotic cell death and underlying mechanisms in vitro.** *Sci Rep.* 2016; **6**: 26277.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. Sessler DI, Ben-Eliyahu S, Mascha EJ, *et al.*: **Can regional analgesia reduce the risk of recurrence after breast cancer? Methodology of a multicenter randomized trial.** *Contemp Clin Trials.* 2008; **29**(4): 517–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Niwa H, Rowbotham DJ, Lambert DG, *et al.*: **Can anesthetic techniques or drugs affect cancer recurrence in patients undergoing cancer surgery?** *J Anesth.* 2013; **27**(5): 731–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Fodale V, D'Arrigo MG, Triolo S, *et al.*: **Anesthetic techniques and cancer recurrence after surgery.** *ScientificWorldJournal.* 2014; **2014**: 328513.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. **F** Exadaktylos AK, Buggy DJ, Moriarty DC, *et al.*: **Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis?** *Anesthesiology.* 2006; **105**(4): 660–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. **F** Cakmakaya OS, Kolodzie K, Apfel CC, *et al.*: **Anaesthetic techniques for risk of malignant tumour recurrence.** *Cochrane Database Syst Rev.* 2014; (11): CD008877.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
45. **F** Cassinello F, Prieto I, del Olmo M, *et al.*: **Cancer surgery: how may anesthesia influence outcome?** *J Clin Anesth.* 2015; **27**(3): 262–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
46. **F** Byrne K, Levins KJ, Buggy DJ: **Can anesthetic-analgesic technique during primary cancer surgery affect recurrence or metastasis?** *Can J Anaesth.* 2016; **63**(2): 184–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Le-Wendling L, Nin O, Capdevila X: **Cancer Recurrence and Regional Anesthesia: The Theories, the Data, and the Future in Outcomes.** *Pain Med.* 2016; **17**(4): 756–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. **F** Cata JP, Hernandez M, Lewis VO, *et al.*: **Can regional anesthesia and analgesia prolong cancer survival after orthopaedic oncologic surgery?** *Clin Orthop Relat Res.* 2014; **472**(5): 1434–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Piegeler T, Beck-Schimmer B: **Anesthesia and colorectal cancer -The perioperative period as a window of opportunity?** *Eur J Surg Oncol.* 2016; **42**(9): 1286–95.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Juan P. Cata**

Department of Anesthesiology and Perioperative Medicine, University of Texas MD Anderson Cancer Center, Houston, TX, USA

Competing Interests: No competing interests were disclosed.

2. **Donal Buggy**

Mater University Hospital, School of Medicine, University College Dublin, Dublin, Ireland

Competing Interests: No competing interests were disclosed.

3. **Markus W. Hollmann**

Department of Anesthesiology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research