REVIEW

Recent advances in the management of migraine [version 1; peer review: 3 approved]

Mark Obermann¹, Dagny Holle²

¹Center for Neurology, Asklepios Hospitals Schildautal, Seesen, Germany

²Department of Neurology and West-German Headache Center, University of Duisburg-Essen, Essen, Germany

V1 First published: 21 Nov 2016, 5(F1000 Faculty Rev):2726
<https://doi.org/10.12688/f1000research.9764.1>

Latest published: 21 Nov 2016, 5(F1000 Faculty Rev):2726
<https://doi.org/10.12688/f1000research.9764.1>

Abstract

Migraine remains one of the most disabling disorders worldwide. The high prevalence in the general population and the often-delicate treatment of patients account for that. Therapeutic management of migraine relies mainly on non-specific medical treatment and is affected by low patient adherence to the treatment regimens applied. The introduction of specific anti-migraine treatment occurred over 20 years ago when the first triptan was approved by regulatory authorities (sumatriptan, 28 December 1992). Triptan use is limited by side effects, time- and frequency-restricted application, and the risk of developing medication overuse headache. Within the past few years, new and promising drugs such as more specific 5-HT 1F receptor agonists (that is, lasmiditan) and monoclonal calcitonin gene-related peptide (CGRP) receptor antibodies entered advanced development phases while non-invasive neuromodulatory approaches were suggested to be potentially effective as non-pharmaceutical interventions for migraine.

Keywords

migraine management, recent advances, CGRP, lasmiditan, vagus nerve stimulation

Open Peer Review

Approval Status

	1	2	3
version 1 21 Nov 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Joost Haan**, Alrijne Ziekenhuis, Leiden, The Netherlands
2. **Kevin C. Brennan**, University of Utah, Salt Lake City, USA
3. **Hanns Ulrich Zeilhofer**, University of Zurich, Zurich, Switzerland
ETH Zurich, Zurich, Switzerland

Any comments on the article can be found at the end of the article.

Corresponding author: Mark Obermann (mark.obermann@uni-due.de)

Competing interests: MO has received scientific support, travel support, and/or honoraria from Biogen Idec, Novartis, Sanofi-Aventis, Genzyme, Pfizer, Teva, and Heel. He received research grants from Allergan, Electrocore, Heel, and the German Ministry for Education and Research (BMBF). DH has received scientific support and honoraria from Allergan, Grünenthal, Desitin, Hormosan, and NeuroDepeche. She received research grants from Allergan, Electrocore, and the BMBF.

Grant information: MO received research grants from Allergan, Electrocore, Heel, and the German Ministry for Education and Research (BMBF). DH received research grants from Allergan, Electrocore, and the BMBF.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Obermann M and Holle D. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Obermann M and Holle D. **Recent advances in the management of migraine [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2726 <https://doi.org/10.12688/f1000research.9764.1>

First published: 21 Nov 2016, 5(F1000 Faculty Rev):2726 <https://doi.org/10.12688/f1000research.9764.1>

Introduction

The Global Burden of Disease Survey 2010 ranked migraine the second most common disease worldwide¹, with a prevalence of 14.7%. Migraine is associated with a substantially reduced quality of life for the individual and a high level of economic burden for society. Migraine was recognized among the seven highest specific causes of disability worldwide. The costs of migraine in the European Union were estimated to exceed 100 billion euros per year². Although most of these costs are related to economic deficits such as inability to work and absence from work, one major factor is medical treatment. Within the past decade, different substances were introduced to treat migraine but with limited success in some patients. Only triptans have been developed for the treatment of migraine specifically and although their efficacy is generally good in most patients, their usage is limited by side effects and the number of times they can be used safely before exposing the patient to the risk of cardiac side effects or the development of medication overuse headache. After several setbacks in the development of novel promising drug classes that led to the discontinuation of the respective development program (that is, the calcitonin gene-related peptide [CGRP] receptor antagonists olcegepant and telcagepant and the glutamate receptor antagonist BGG492), there are now several quite promising therapeutic agents emerging from phase II and early phase III studies that may well fulfill these aims.

Even though final confirmation of efficacy for most of the drugs discussed below will have to follow within the next few years, they will likely change the way we treat migraine in the future and considerably add therapeutic power to the specific treatment of migraine.

Non-pharmaceutical interventions such as behavioral therapy and non-invasive neuromodulatory approaches such as vagus nerve stimulation (VNS) and transcutaneous supraorbital nerve stimulation were successfully tested to treat migraine and represent a treatment alternative for those patients unwilling or unable to manage their disorder pharmacologically.

5-hydroxytryptamine 1F receptor agonists

Lasmiditan is a 5-hydroxytryptamine (5-HT) 1F receptor agonist that works similarly to triptans (5-HT 1B/1D agonists) but without the vasoconstrictive side effects well known to this

substance class³. It was given the generic stem name “ditan” to distinguish it from the “triptans” and other drug classes. The results of the SAMURAI (A Study of two doses of lasmiditan compared to placebo in the acute treatment of migraine) trial were presented recently⁴. The SAMURAI trial was a randomized, double-blinded, placebo-controlled parallel group phase III clinical trial that compared 100 and 200 mg oral administration of lasmiditan with placebo for 2-hour pain freedom from migraine headache as the primary endpoint. The trial randomly assigned 2,231 migraine patients in the United States, including those patients with cardiovascular risk factors that would not be appropriate for treatment with triptans. The mean age of randomly assigned patients was 41.6 years, 83% were women, and the mean disease duration was 19 years. They had an average of five migraine attacks per month and an average MIDAS (migraine disability assessment) score of 31. One-quarter of patients used prophylactic migraine medication, and 82% of patients had cardiovascular risk factors. The most common ones were obesity, family history of coronary artery disease, smoking, hypertension, hyperlipidemia, and diabetes mellitus type 2. Patients also had cardiovascular conditions such as arrhythmias, mitral valve disease, angina, atrial fibrillation, congestive heart failure, prior myocardial infarction, or ischemic stroke. A single migraine attack was treated and 28.2% (odds ratio 2.2, 95% confidence interval [CI] 1.6 to 3.0, $P < 0.0001$) of patients were pain-free at 2 hours with 100 mg lasmiditan, 32.2% (odds ratio 2.6, 95% CI 2.0 to 3.6, $P < 0.001$) were pain-free with 200 mg, and only 15.3% with placebo, thus reaching the primary endpoint. These results are in a comparable efficacy range of key clinical trials done with triptans without having access to direct comparison trials. Lasmiditan was generally well tolerated with mild or moderate side effects. The most common side effects were dizziness, paresthesia, somnolence, nausea, fatigue, and lethargy (Table 1). In conclusion, lasmiditan showed a good treatment effect in highly affected migraine patients that is comparable to triptans but with a much better cardiovascular side effect profile. It may become a valuable treatment option for patients at risk for cardiovascular events due to vasoconstriction.

These promising results will have to be confirmed by the ongoing SPARTAN (Three Doses of Lasmiditan [50, 100, and 200 mg] Compared to Placebo in the Acute Treatment of Migraine) trial that uses the same endpoints and same statistical power. It also tests the 50 mg dose of lasmiditan in order to establish the lowest effective dose for an even better adverse event profile⁴.

Table 1. Most common side effects of lasmiditan in the SAMURAI trial⁴.

Side effect	100 mg, n = 630	200 mg, n = 609	Placebo, n = 617
Dizziness	75 (11.9%)	94 (15.4%)	19 (3.1%)
Paresthesia	36 (5.7%)	46 (7.6%)	13 (2.1%)
Somnolence	33 (5.2%)	32 (5.3%)	14 (2.3%)
Nausea	16 (2.5%)	29 (4.8%)	9 (1.5%)
Fatigue	24 (3.8%)	18 (3.0%)	1 (0.2%)
Lethargy	12 (1.9%)	14 (2.3%)	1 (0.2%)

Monoclonal calcitonin gene-related peptide antibodies

Four monoclonal antibodies targeting the CGRP pathways were introduced within the past few years and all have successfully completed their phase II study and are entering phase III clinical trials at the moment (Table 2;^{5–10}). The main difference between the different substances is the frequency and route of administration (subcutaneously or intravenously). AMG 334 targets the CGRP receptor, whereas the other three antibodies bind the CGRP peptide. The CGRP receptor is a complex transmembrane receptor (CLR/RAMP 1 complex) that is 5,000 times more selective for CGRP than other related receptors in this class. The 70 mg dose of AMG 334 was effective with a mean monthly migraine day reduction of 3.4 days compared with placebo (–2.28 days; $P = 0.021$), whereas the lower doses with 7 and 21 mg, respectively, were not significant⁶. Patients received subcutaneous injections every 4 weeks. The double-blind treatment phase was completed by 448 (93%) patients. Adverse events were mild to moderate in severity and did not differ between the active drug and placebo. The most common complaints were nasopharyngitis and fatigue. Importantly, given the experience with small-molecule antagonists, no hepatotoxicity was observed in these patients⁶.

ALD403 is one of the antibodies binding CGRP⁷. One gram was administered intravenously every 4 weeks and led to a reduction of 5.6 migraine days compared with 4.6 days in the placebo group ($P = 0.0306$). Eleven percent of 67 treated patients showed a 100% response (that is, no more migraine attacks) after 12 weeks⁷. Similar efficacy was observed with the subcutaneous administration of 150 mg LY2951742 every 2 weeks compared with placebo⁸. The number of migraine headache days was decreased by 4.2 days after 12 weeks compared with baseline, whereas placebo reduced the headache days by only 3.0 days ($P = 0.003$). The 100% responder rate was 15% in a *post hoc* analysis, and no serious adverse events occurred on treatment⁸.

TEV-48125 was tested in high-frequency episodic migraine as well as chronic migraine in two separate controlled clinical trials^{9,10}. Episodic migraine patients with 8 to 14 headache days per month received either 225 or 600 mg TEV-48125 every 4 weeks subcutaneously and showed reductions in headache days of 4.86 days in the 225 mg group and 4.80 days in the 600 mg group, whereas placebo reduced the headache days by 3.10 days

(both $P < 0.0001$). The most common treatment-related adverse events were mild injection site pain or erythema without differences between the groups⁹.

TEV-48125 was equally effective in chronic migraine patients who had more than 15 headache days per month. The number of headache days after 12 weeks was reduced by 6.04 days in the group receiving up to 675 mg ($P = 0.0386$), 6.2 days in the 900 mg group ($P = 0.0057$), and only 4.2 days in the placebo group. Four patients reported serious adverse events, one in the placebo group. One patient had acute kidney stone pain, one patient developed pneumonia after influenza infection, one had irritable bowel syndrome, and one had an episode of depression with suicidal ideation. None of these events was related to the study drug by the investigators¹⁰.

Non-invasive neuromodulatory approaches: transcutaneous vagus nerve stimulation

The EVENT (Chronic migraine headache prevention with noninvasive VNS) study investigated the efficacy and safety of non-invasive, transcutaneous VNS for the treatment of chronic migraine¹¹. The stimulation device delivers a low voltage (peak of 24 V) and maximum output current of 60 mA to the right side of the neck close to the vagus nerve. Three 2-minute treatment sessions were applied per day over 2 months followed by a 6-month open-label extension phase. In total, 59 patients (30 VNS and 29 sham) were evaluated in an intention-to-treat analysis. Although the mean change in headache days from baseline after 2 months was not different between groups ($P = 0.56$), the number of headache days continued to decrease in the open-label phase with a mean of 17.2 (95% CI 13.8 to 20.5) headache days and a mean change from baseline of –3.6 (95% CI –6.3 to –0.87; $P = 0.02$) after 8 months of treatment. Adverse events that were deemed possibly related to the device included eye twitch, facial pain/numbness, paresthesia, treatment site skin reaction, worsening migraine, and gastrointestinal symptoms. All adverse events were distributed equally across groups. Adherence was over 90% in both groups throughout the study. Transcutaneous VNS may become an interesting adjunct non-pharmacological treatment option for the treatment of chronic migraine. Further studies with more patients and a longer blinded study period will have to determine the value of this treatment¹¹.

Table 2. Monoclonal antibodies targeting the calcitonin gene-related peptide pathway.

LY2951742	Superior to placebo (number of migraine headache days –4.2 days after 12 weeks compared with baseline, placebo –3.0 days; $P = 0.003$) ⁸
ALD403	Superior to placebo (comparing baseline migraine days to weeks 5–8 reduction of 5.6 migraine days compared with 4.6 days in the placebo group; $P = 0.0306$) ⁷
AMG 334	Superior to placebo (mean monthly migraine days –3.4 days compared with placebo; –2.28 days; $P = 0.021$) ⁶
TEV-48125 (LBR-101)	Superior to placebo (high-frequent episodic migraine reduction in headache days of –4.86 days for the 225 mg group and –4.80 days for the 600 mg group; both $P < 0.0001$) ⁹
TEV-48125 (LBR-101)	Superior to placebo (chronic migraine –6.04 days in the 675 mg group, $P = 0.0386$, –6.16 days in the 900 mg group, $P = 0.0057$) ¹⁰

Conclusions

The next couple of years will be an exciting time for patients with migraine and their physicians. The range of different treatment modalities considerably widened within the past few years and may continue to do so in the years to come. Medications with new mechanisms of action and fewer side effects may prove to be safe and effective for the treatment of episodic and chronic migraine. Additionally, non-invasive neuromodulation may prove to be a genuine therapeutic asset in the management of migraine.

Competing interests

MO has received scientific support, travel support, and/or honoraria from Biogen Idec, Novartis, Sanofi-Aventis, Genzyme, Pfizer,

Teva, and Heel. He received research grants from Allergan, Electrocore, Heel, and the German Ministry for Education and Research (BMBF). DH has received scientific support and honoraria from Allergan, Grünenthal, Desitin, Hormosan, and NeuroDepeche. She received research grants from Allergan, Electrocore, and the BMBF.

Grant information

MO received research grants from Allergan, Electrocore, Heel, and the German Ministry for Education and Research (BMBF). DH received research grants from Allergan, Electrocore, and the BMBF.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Vos T, Flaxman AD, Naghavi M, *et al.*: Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; **380**(9859): 2163–96. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
2. Linde M, Gustavsson A, Stovner LJ, *et al.*: The cost of headache disorders in Europe: the Eurolight project. *Eur J Neurol*. 2012; **19**(5): 703–11. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
3. Reuter U, Israel H, Neeb L: The pharmacological profile and clinical prospects of the oral 5-HT_{1F} receptor agonist lasmiditan in the acute treatment of migraine. *Ther Adv Neurol Disord*. 2015; **8**(1): 46–54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
4. CoLucid: CoLucid Pharmaceuticals Announces Achievement of Secondary Endpoints in SAMURAI. 2016. [Reference Source](#)
5. Shi L, Rao S, King CH, *et al.*: AMG 334, the first potent and selective human monoclonal antibody antagonist against the CGRP receptor. *Headache*. 2014; **54**(8): 1417–18. [Reference Source](#)
6. Lenz R, Silberstein S, Dodick D, *et al.*: Results of a randomized, double-blind, placebo controlled, phase 2 study to evaluate the efficacy and safety of AMG 334 for the prevention of episodic migraine. *Cephalalgia*. 2015; **35**: 5. [Reference Source](#)
7. Dodick DW, Goadsby PJ, Silberstein SD, *et al.*: Safety and efficacy of ALD403, an antibody to calcitonin gene-related peptide, for the prevention of frequent episodic migraine: a randomised, double-blind, placebo-controlled, exploratory phase 2 trial. *Lancet Neurol*. 2014; **13**(11): 1100–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
8. Dodick DW, Goadsby PJ, Spierings EL, *et al.*: Safety and efficacy of LY2951742, a monoclonal antibody to calcitonin gene-related peptide, for the prevention of migraine: a phase 2, randomised, double-blind, placebo-controlled study. *Lancet Neurol*. 2014; **13**(9): 885–92. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
9. Bigal ME, Dodick DW, Rapoport AM, *et al.*: Safety, tolerability, and efficacy of TEV-48125 for preventive treatment of high-frequency episodic migraine: a multicentre, randomised, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol*. 2015; **14**(11): 1081–90. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
10. Bigal ME, Edvinsson L, Rapoport AM, *et al.*: Safety, tolerability, and efficacy of TEV-48125 for preventive treatment of chronic migraine: a multicentre, randomised, double-blind, placebo-controlled, phase 2b study. *Lancet Neurol*. 2015; **14**(11): 1091–100. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. Silberstein SD, Calhoun AH, Lipton RB, *et al.*: Chronic migraine headache prevention with noninvasive vagus nerve stimulation: The EVENT study. *Neurology*. 2016; **87**(5): 529–38. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Hanns Ulrich Zeilhofer**

¹ Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland

² Institute of Pharmaceutical Sciences, ETH Zurich, Zurich, Switzerland

Competing Interests: No competing interests were disclosed.

2. **Kevin C. Brennan**

Department of Neurology, University of Utah, Salt Lake City, UT, USA

Competing Interests: No competing interests were disclosed.

3. **Joost Haan**

Alrijne Ziekenhuis, Leiden, The Netherlands

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research