Check for updates

CORRESPONDENCE

Matching target dose to target organ [version 1; peer review: 2 approved]

Desmond I. Bannon, Marc A. Williams

Toxicology Directorate, Health Effects Division, U.S. Army Public Health Center, Edgewood, MD, USA

V1 First published: 29 Nov 2016, 5:2785
<https://doi.org/10.12688/f1000research.10055.1>
Latest published: 30 Mar 2017, 5:2785
<https://doi.org/10.12688/f1000research.10055.2>

Abstract

In vitro assays have become a mainstay of modern approaches to toxicology with the promise of replacing or reducing the number of *in vivo* tests required to establish benchmark doses, as well as increasing mechanistic understanding. However, matching target dose to target organ is an often overlooked aspect of *in vitro* assays, and the calibration of *in vitro* exposure against *in vivo* benchmark doses is often ignored, inadvertently or otherwise. An example of this was recently published in *Environmental Health Perspectives* by Wagner *et al.*, where neural stem cells were used to model the molecular toxicity of lead. On closer examination of the *in vitro* work, the doses used in media reflected *in vivo* lead doses that would be at the highest end of lead toxicity, perhaps even lethal. Here we discuss the doses used and suggest more realistic doses for future work with stem cells or other neuronal cell lines.

Keywords

In vitro, lead, in vivo, dose, stem cells.

Open Peer Review

Approval Status

1

2

version 2

(revision)

30 Mar 2017

version 1

29 Nov 2016

view

view

1. **Mir Ahamed Hossain**, Department of Neurology, Baltimore, USA
2. **Donald Smith**, University of California Santa Cruz, Santa Cruz, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Desmond I. Bannon (desmond.i.bannon.civ@mail.mil)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Bannon DI and Williams MA. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bannon DI and Williams MA. Matching target dose to target organ [version 1; peer review: 2 approved] F1000Research 2016, 5:2785 <https://doi.org/10.12688/f1000research.10055.1>

First published: 29 Nov 2016, 5:2785 <https://doi.org/10.12688/f1000research.10055.1>

A recent article by Wagner *et al.* reported the involvement of the anti-oxidant Nrf2 transcription factor signaling pathway in the toxicity of lead using neural stem cells in an *in vitro* model of neuronal differentiation¹. While this work was completed in a similar way to other studies involving *in vitro* lead exposure, the work avoids a critical, often neglected issue of what constitutes a relevant physiological dose *in vitro*. The assumption that the selected dose of 1 μM (or 20.7 $\mu\text{g/dL}$) for neuronal stem cell exposure was “4 times the CDC levels of concern (LOC) for blood lead (5 $\mu\text{g/dL}$) and is within the range of exposed populations” requires further examination. Since the *in vitro* exposure was completed in media (the equivalent of plasma or serum) and not in whole blood, the assumption that the *in vitro* lead level would be equivalent to that found in blood of lead-exposed humans is somewhat inaccurate. Lead in serum (or plasma) represents only a fraction (~1%) of the level found in whole blood^{2,3}, with the major fraction of lead bound inside erythrocytes⁴. For arguments sake, if the proportion of lead used in this study was 1% of that in whole blood, the equivalent blood lead value would be 2073 $\mu\text{g/dL}$, a level over 400 times the CDC LOC, and one that would be acutely toxic and perhaps lethal.

Another study, which was cited by Wagner *et al.*¹, showed that measurable effects in stem cells *in vitro* could occur at doses as low as 0.4 μM ; this dose would represent a blood lead level of 800 $\mu\text{g/dL}$ by our calculations⁵. In a study by Chan *et al.*, the lowest dose of 1 μM lead used in a study of newborn rat neuronal stem cells would represent 1000 $\mu\text{g/L}$ in serum and a massive systemic blood lead level of about 10,000 $\mu\text{g/dL}$ ⁶. Other studies examining the toxicity of lead in cell cultures have also failed to adequately match the *in vitro* doses^{7–10} with those found *in vivo*, by taking account of the well documented difference between plasma and whole blood lead values. More importantly, with measurable effects on differentiation only beginning at 10 μM for Chan *et al.*⁶, could these data suggest the alternative interpretation that neuronal stem cells *in vivo* are more resistant to toxic insult by lead than our current understanding would have us believe – at least in the short term?

What is clear is that at current blood lead levels in the US population, serum or plasma levels will represent a very low fraction of those values and *in vitro* work could more realistically model

chronic neurological effects in humans if target doses were better matched to the doses found at target sites. Thus, the model proposed in this and other work, while presenting novel effects, may be more appropriate for high acute exposures. To ensure that doses used in *in vitro* assays are complimentary to a target *in vivo* blood lead level of 20 $\mu\text{g/dL}$, exposure to cells *in vitro* should more accurately correspond to 1% of the blood lead value, or a dose of 0.2 $\mu\text{g/dL}$ (0.01 μM). At the current CDC 5 $\mu\text{g/dL}$ LOC for children, the *in vitro* dose would become 0.05 $\mu\text{g/dL}$ (0.002 μM), a dose that would present difficulties to laboratories that cannot eliminate background levels from residual lead on glassware and other sources of possible contamination or confounding of the reported data.

In the study by Wagner *et al.*¹, much of this may have been considered by the authors, and key assumptions may have been made; however, the question still remains whether the upregulation of genes in the Nrf2-mediated anti-oxidative stress pathway would have been observed if a more physiologically relevant dose of 0.2 $\mu\text{g/dL}$ (0.1 μM) in the media (i.e., representing a blood lead level of 20 $\mu\text{g/dL}$) had been used.

Disclaimer

The views expressed in this article are those of the author(s) and do not necessarily reflect the official policy of the Department of Defense, Department of the Army, U.S. Army Medical Department or the U.S.

Author contributions

DB conceptualized the article and analyzed the original critiqued article reported herein. MW provided technical writing support and analysis of the original critiqued article reported herein.

Competing interests

No competing interests were disclosed.

Grant information

The authors confirm that no grant(s) were involved in supporting this work.

References

- Wagner PJ, Park HR, Wang Z, *et al.*: **In Vitro Effects of Lead on Gene Expression in Neural Stem Cells and Associations between Upregulated Genes and Cognitive Scores in Children.** *Environ Health Perspect.* 2016. [Epub Ahead of Print]. [PubMed Abstract](#) | [Publisher Full Text](#)
- Manton WL, Cook JD: **High accuracy (stable isotope dilution) measurements of lead in serum and cerebrospinal fluid.** *Br J Ind Med.* 1984; 41(3): 313–19. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Smith D, Hernandez-Avila M, Tellez-Rojo MM, *et al.*: **The relationship between lead in plasma and whole blood in women.** *Environ Health Perspect.* 2002; 110(3): 263–68. [PubMed Abstract](#) | [Free Full Text](#)
- Simons TJ: **Passive transport and binding of lead by human red blood cells.** *J Physiol.* 1986; 378(1): 267–86. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Senut MC, Sen A, Cingolani P, *et al.*: **Lead exposure disrupts global DNA methylation in human embryonic stem cells and alters their neuronal differentiation.** *Toxicol Sci.* 2014; 139(1): 142–61. [PubMed Abstract](#) | [Free Full Text](#)
- Chan YH, Gao M, Wu W: **Are newborn rat-derived neural stem cells more sensitive to lead neurotoxicity?** *Neural Regen Res.* 2013; 8(7): 581–92. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Engstrom A, Wang H, Xia Z: **Lead decreases cell survival, proliferation, and neuronal differentiation of primary cultured adult neural precursor cells through activation of the JNK and p38 MAP kinases.** *Toxicol In Vitro.* 2015; 29(5): 1146–155. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Qasimian Lemraski M, Soodi M, Taha Fakhr M, *et al.*: **Study of lead-induced neurotoxicity in neural cells differentiated from adipose tissue-derived stem cells.** *Toxicol Mech Methods.* 2015; 25(2): 128–35. [PubMed Abstract](#) | [Publisher Full Text](#)
- Jia Q, Ha X, Yang Z, *et al.*: **Oxidative stress: a possible mechanism for lead-induced apoptosis and nephrotoxicity.** *Toxicol Mech Methods.* 2012; 22(9): 705–10. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kermani S, Karbalaie K, Madani SH, *et al.*: **Effect of lead on proliferation and neural differentiation of mouse bone marrow-mesenchymal stem cells.** *Toxicol In Vitro.* 2008; 22(4): 995–1001. [PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 26 January 2017

<https://doi.org/10.5256/f1000research.10834.r19115>

© 2017 Smith D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Donald Smith

Microbiology and Environmental Toxicology, University of California Santa Cruz, Santa Cruz, CA, USA

This commentary is well written, very well justified, and timely. While there are countless published papers on the myriad effects of lead in biological systems, the consideration of dose extrapolation from *in vitro* to *in vivo* studies and their relationships to the human condition often goes unappreciated. Indeed, since toxicology is driven by the dose of the poison, establishing environmental or occupational relevance of the dose is absolutely key to the relevance of the findings. This commentary points this out in a concise and evidence-driven fashion, and is worthy of publication.

Below are a few minor comments to consider.

1. Pg. 2, 1st para: *For arguments sake, ...*

Comment: A caveat here might be that is known that the proportion of whole blood lead in plasma increases with increasing blood lead, so it is likely that the blood lead level that would produce a 1 uM plasma lead would be lower than 2,073 ug/dL, but this does not detract from the point the authors are making, which is a good and important one.

2. Pg. 2, 3rd para: *Thus, the model proposed in this and other work...*

Comment: It is not clear whose work 'this work' is referring to - Chan *et al*?

3. Pg. 2, 3rd para: *To ensure that doses used in in vitro assays are complimentary to a target in vivo blood lead level of 20 µg/dL...*

Comment: This suggestion by the authors is reasonable, assuming that plasma lead reflects extracellular fluid lead, though it might also be worth looking at the relationship between blood lead and CSF lead levels (in the literature) to see if it follows an appx 1% relationship as does plasma to further substantiate this suggestion.

4. Pg. 2, 3rd para: *...eliminate background levels from residual lead on glassware and other sources of possible contamination or confounding of the reported data...*

Comment: This too raises an important point in that the vast majority of studies do not make sufficient effort to reduce background lead levels in control cultures, so it is quite possible that here and in those other studies the control cultures, even with modestly elevated background lead levels will also be affected, requiring higher exposure doses to demonstrate a difference or 'effect' in the lead-exposed treatments. It is good that the authors pointed this out.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 02 Mar 2017

Mark A Williams, Army Public Health Center, USA

Reviewer 2. We thank reviewer #2 for knowledgeable and helpful comments on our article. Here are our responses to specific comments.

Comment 1. This point is well made – we agree that the proportion of lead in plasma would increase as blood lead increases, so that equivalent plasma lead at blood lead values greater than 100 µg/dL could be upwards of 2%. As it was we selected 1% plasma/blood ratio as the blood lead under question was 20 µg/dL but of course there is some inbuilt error in our calculations at high doses. Nonetheless, our extrapolated exposure scenario is meant to demonstrate that the assumptions under which many *in vitro* studies lie with respect to their relationship to *in vivo* blood lead values are often violated; the reviewer also acknowledges our efforts to point this out. We have added more text to acknowledge this non-linear relationship at increasing doses between whole blood lead and plasma lead.

Comment 2. This sentence has been restructured to indicate that we referring to the Wagner *et al* study, as well as other studies that have made similar assumption.

Comment 3. We agree that cerebrospinal fluid measures would further corroborate our assumptions. The work by Manton *et al* (cited in our article) showed that cerebrospinal fluid levels were about 50% of serum levels, though it should be pointed out that this work was carried out in only one subject. We have added more text to acknowledge this fact.

Comment 4. We agree with the further elaboration of this sentence and have added additional text to incorporate the details of the comment.

Competing Interests: None.

Reviewer Report 18 January 2017

<https://doi.org/10.5256/f1000research.10834.r19455>

© 2017 Hossain M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mir Ahamed Hossain

Johns Hopkins University School of Medicine, Department of Neurology, Baltimore, MD, USA

In vitro assays have become a mainstay of modern approaches to toxicology with a high promise of understanding the underlying mechanisms of toxicity. The results reported by Wagner *et al.*, (2016) in the August 26 issue of the Environmental Health Perspectives, where neural stem cells were used to model the toxicity of lead. The results support the notion that lead treatment of cells leads to upregulation of vascular gene expression (JBC 275:27874-27882, 2000). While this work presents interesting effects, this reviewer's opinion is in agreement with the correspondence (critiqued article) authors Bannon and Williams that it may be more appropriate for high acute exposures particularly in case of neural stem/progenitor cells, which lack many of the characteristic features of mature neurons.

It is also likely that neural stem cells (NSCs) could be more resistance to toxic insult by lead - at least in the short term. Thus the *in vitro* work could more realistically model chronic neurological effects if doses are better matched with the doses at the target site, as supported by the fact that serum or plasma levels represent a very low fraction of the total blood lead levels. Thus the concentrations of lead used in this study, which elicits upregulation of genes in the Nrf2-mediated anti-oxidative stress pathway, appear to be in the low micromolar range, which is much higher than the *in vitro* dose equivalent of the current CDC levels of concentrations (5 ug/dL) for children. Thus the concentrations used in the study does not reflect the likely exposure of lead in the environment, that is to say, concentrations which are likely to be cytotoxic particularly in case of NSCs. This is clearly a near impossible issue to address empirically, but if some information available along these lines using a more physiologically relevant dose in the media of *in vitro* NSC cultures to show gene expression in the Nrf2-mediated anti-oxidative stress pathway would be helpful for the reader as suggested by Bannon and Williams in the critiqued article. It will also be interesting to see how the differentiated neurons from lead exposed NSCs express neurons specific features or exhibit mature neuronal function.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 02 Mar 2017

Mark A Williams, Army Public Health Center, USA

Reviewer 1. We thank reviewer #1 for helpful comments on our article. We address some

specific aspects below.

The reviewer agreed with our principle argument, but goes on to state that the Wagner *et al* “results support the notion that lead treatment of cells leads to upregulation of vascular gene expression”, citing an *in vitro* microarray study using astrocytes, (Hossain *et al*, 2002, ref 9 above) when in fact two of the three VEGF transcripts listed in Supplemental Table 1 of Wagner *et al* were downregulated by lead, with only one – VEGFA downregulated by 0.8-fold – being statistically significant. Therefore the cited publication by Hossain *et al* is contradicted by the Wagner *et al* data for the VEGF gene.

The fact that the Hossain *et al* study used 10 μ M lead acetate to dose astrocytes *in vitro* further supports our main point – that most lead concentrations *in vitro* would reflect highly lethal lead concentrations *in vivo* if the difference between lead in whole blood (red blood cells) and plasma were taken into account. Hossain *et al* did cite Audersirk (Audersirk G, *et al*. In Vitro Cell Dev Biol. 1989 Dec;25(12):1121-8) as supporting evidence for the use of 10 μ M lead as a dosing solution for astrocytes, where Audersirk measured free lead (Pb^{2+}) in the nanomolar range in the presence of full experimental media dosed with micromolar lead acetate using an ion selective electrode. However, Audersirk’s work in snail and chick neurons did not examine the potential lethality of the *in vitro* working doses to the whole organism, taking account of plasma/whole blood differences.

Competing Interests: None.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research