REVIEW

Recent advances in understanding *Streptomyces* [version 1; peer review: 4 approved]

Keith F. Chater

Department of Molecular Microbiology, John Innes Centre, Norwich, UK

V1 First published: 30 Nov 2016, 5(F1000 Faculty Rev):2795
<https://doi.org/10.12688/f1000research.9534.1>

Latest published: 30 Nov 2016, 5(F1000 Faculty Rev):2795
<https://doi.org/10.12688/f1000research.9534.1>

Abstract

About 2,500 papers dated 2014–2016 were recovered by searching the PubMed database for *Streptomyces*, which are the richest known source of antibiotics. This review integrates around 100 of these papers in sections dealing with evolution, ecology, pathogenicity, growth and development, stress responses and secondary metabolism, gene expression, and technical advances. Genomic approaches have greatly accelerated progress. For example, it has been definitively shown that interspecies recombination of conserved genes has occurred during evolution, in addition to exchanges of some of the tens of thousands of non-conserved accessory genes. The closeness of the association of *Streptomyces* with plants, fungi, and insects has become clear and is reflected in the importance of regulators of cellulose and chitin utilisation in overall *Streptomyces* biology. Interestingly, endogenous cellulose-like glycans are also proving important in hyphal growth and in the clumping that affects industrial fermentations. Nucleotide secondary messengers, including cyclic di-GMP, have been shown to provide key input into developmental processes such as germination and reproductive growth, while late morphological changes during sporulation involve control by phosphorylation. The discovery that nitric oxide is produced endogenously puts a new face on speculative models in which regulatory Wbl proteins (peculiar to actinobacteria) respond to nitric oxide produced in stressful physiological transitions. Some dramatic insights have come from a new model system for *Streptomyces* developmental biology, *Streptomyces venezuelae*, including molecular evidence of very close interplay in each of two pairs of regulatory proteins. An extra dimension has been added to the many complexities of the regulation of secondary metabolism by findings of regulatory crosstalk within and between pathways, and even between species, mediated by end products. Among many outcomes from the application of chromosome immunoprecipitation sequencing (ChIP-seq) analysis and other methods based on “next-generation sequencing” has been the finding that 21% of *Streptomyces* mRNA species lack leader sequences and conventional ribosome binding sites. Further technical advances now emerging should lead to

Open Peer Review

Approval Status

	1	2	3	4
version 1				
30 Nov 2016				

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Rosemary Loria**, University of Florida, Gainesville, USA
2. **Klas Flardh**, University of Lund, Lund, Sweden
3. **Paul J Dyson**, Swansea University Medical School, Swansea, UK
4. **Wolfgang Wohlleben**, University of Tuebingen, Tuebingen, Germany

Any comments on the article can be found at the end of the article.

continued acceleration of knowledge, and more effective exploitation, of these astonishing and critically important organisms.

Keywords

streptomyces, genomics, streptomyces ecology, streptomyces evolution, pathogenic streptomyces

Corresponding author: Keith F. Chater (keith.chater@nbi.ac.uk)

Competing interests: The author declares that he has no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Chater KF. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Chater KF. **Recent advances in understanding *Streptomyces* [version 1; peer review: 4 approved]**
F1000Research 2016, 5(F1000 Faculty Rev):2795 <https://doi.org/10.12688/f1000research.9534.1>

First published: 30 Nov 2016, 5(F1000 Faculty Rev):2795 <https://doi.org/10.12688/f1000research.9534.1>

Introduction

The majority of antibiotics used in medicine, veterinary practice, and agriculture originate from *Streptomyces* bacteria. Genomic analysis has shown that any one strain has the potential to make tens of such secondary metabolites, and metagenomic analysis has revealed vast numbers of relevant biosynthetic gene sets^{1,2}. These organisms are therefore being studied ever more intensively in the expectation that they will contribute significantly to the provision of new therapeutic agents to combat the global emergence of antibiotic resistance among pathogenic bacteria, as well as providing other bioactive agents with medical applications. However, this is only one aspect of the interest and importance of streptomycetes. Ecologically, streptomycetes have key roles in the natural recycling of the globally abundant cell walls of fungi and plants. Some of them have evolved intimate partnerships with insects or plants, and a few have acquired pathogenic attributes. The interplay of these associations with *Streptomyces* molecular physiology is a strong theme in this review and has been underpinned by continuing in-depth analysis of the model organism *Streptomyces coelicolor* A3(2), which has been studied genetically for about 60 years³.

Unusually for bacteria, streptomycetes grow as branching hyphal filaments to form a mat of fungus-like mycelium, from which emerge aerial branches that bear chains of spores (Figure 1). Very significant progress has been made recently in the analysis of this complex growth and development in both *S. coelicolor* and the emerging model species *Streptomyces venezuelae*, which sporulates very readily, even in liquid culture.

These two model organisms are providing new insights into how antibiotic production is integrated into overall physiology, taking advantage of the fact that, among the more than 20 secondary

metabolites of *S. coelicolor*, two are pigmented antibiotics (the polyketide actinorhodin [ACT] and the red tripyrrole prodiginines [RED]), while *S. venezuelae* makes chloramphenicol and the polyketide jadomycin. These studies have pointed to an unexpected role for antibiotics themselves as regulatory ligands mediating autoregulation, cross-regulation, and interspecies interactions.

This wide-ranging topical snapshot of the genus has been compiled from a selection of the approximately 2,500 papers about *Streptomyces* published in 2014–2016. Where possible, I have cited review articles that provide context to the recent advances. In concentrating on whole organism physiology and evolution, I have excluded certain major *Streptomyces* research areas, notably taxonomy, systematics, and natural product chemistry.

Ecology and evolution

More than just soil organisms...

Streptomycetes, which are abundant in soil, are believed to have originated around 400 million years ago, when the land was being colonised by green plants. Their major role in the solubilisation of cell wall or surface components of plants, fungi, and insects⁴ suggests that they played a part in the composting of early land plants and hence in the formation of primeval soil. This conserved ecology over evolutionary time is reflected at the genome level: for example, *Streptomyces reticuli*, which is particularly active in cellulose degradation, has an estimated 456 genes for proteins involved in the binding, degradation, and utilisation of cellulose and other complex and simple carbohydrates⁵. *S. coelicolor* also has eight genes for cellulase-like proteins, at least one of which is a true cellulase, even though *S. coelicolor* cannot grow with cellulose as its sole carbon source⁶. This situation is widespread among streptomycetes, prompting speculation that most isolates have

Figure 1. Colonies of *Streptomyces coelicolor* A3(2). The fuzzy surface of these mould-like colonies is made up of aerial hyphae carrying chains of spores (photo, K.F. Chater).

evolved to live in mixed, cooperating communities, within which only a few species are competent to release amenable products from native cellulose⁷. In addition, streptomycetes have had a long evolutionary association with fungi, with fungal cell walls being a key source of nutrition; thus, *S. coelicolor* has (at least) 13 chitinases regulated by the important global regulator DasR, which coordinates many aspects of *Streptomyces* biology in response to the chitin-derived ligands N-acetylglucosamine or N-acetylglucosamine-phosphate^{4,8–10}, as well as two chitosanases and a dedicated oligoglucosamine uptake system that are regulated independently of DasR¹¹.

Reinforcing their close association with plants (reviewed in 12), streptomycetes are abundantly represented in metagenomic analysis of the rhizosphere¹³, and they are among the most frequent endophytes: a search of PubMed for “*Streptomyces* endophyte” yielded 74 papers published in the last 15 years, with 38 dated 2013–2016. Many studies have focused on endophytes from traditional medicinal plants, often in the hope that the endophytes may be responsible for the medicinal properties (e.g. 14–19). Indeed, *Streptomyces* endophytes produce various antagonists of plant disease agents and pests such as bacteria, fungi, and insects²⁰, consistent with potentially ancient adaptive benefits that might be turned to agricultural and medical use in the modern world. Streptomycetes may also stimulate plant growth^{21,22}, sometimes by producing phytohormones (e.g. 23,24). Endophytic streptomycetes are also found in diverse non-medicinal dicot plants^{25,26} and in monocots including rice²², suggesting (but not proving) that such interactions may date back hundreds of millions of years. Metagenomic evidence of particularly exuberant diversity of soil streptomycetes at low latitudes^{27,28} might reflect greater plant and insect diversity in the tropics (along with some degree of specificity in interactions between *Streptomyces* and plants or insects).

Streptomycetes are partners in other mutualistic interactions involving plants. Inoculation of the rhizosphere of rooted oak cuttings with *Streptomyces* sp. AcH 505 not only stimulated mycorrhizal development but also elicited plant defence mechanisms, leading to increased resistance to powdery mildew infection²⁹. Streptomycetes are also carried by ants that cultivate black food yeasts inside cavities (“domatia”) in host plants. In this mutualism, the plants secrete sugary exudates to the ants and yeasts, while the ants ward off predators and generally clean up their host. The streptomycetes produce inhibitors of other, unwanted, fungi that might invade the domatia³⁰. In a broad survey, the streptomycetes involved in these plant–ant interactions fell into just four distinct taxa, while many other taxa were unrepresented, implying specific adaptations³⁰. Other cases of insect-borne streptomycetes preventing unwanted fungal invasion have been extensively reported in the last 20 years, such as in the cases of leaf-cutter ants and beewolf wasps¹²: in the latter case, some degree of co-evolution of hosts and microbes has been taking place for perhaps 68 million years^{12,31}. Streptomycetes isolated from associations with diverse arthropods that feed on plant biomass are also highly enriched for two taxa having the ability to degrade cellulose completely, compared to random soil isolates, suggesting that this ability is valuable in the symbiosis⁷.

The production of antibiotics by streptomycetes has long suggested that they may be formidable competitors in natural environments,

but it was a surprise that, when growing on agar spread with other bacteria, the growing hyphal tips of most strains caused local lysis of what can reasonably be interpreted as their prey³². Such predatory activity may add a new perspective to the benefits of streptomycetes to plants and insect mutualistic partners.

In a further illustration of the ecological diversity of streptomycetes, strains adapted to marine conditions are prominent in the microbiota of marine sponges³³.

Streptomycetes as pathogens: evolution and regulation of potato scab

Not all interactions of streptomycetes with complex eukaryotes are beneficial. Scab disease of potato and other tuber/root crops is caused by several different streptomycetes and (as in many cases of bacterial pathogenesis) involves a laterally acquired pathogenicity island that presumably evolved with the emergence of scab hosts, which are all flowering plants that evolved long after simpler plants had colonised the land. In *Streptomyces turgidiscabies*, this entire DNA island can excise from the host genome and transfer into a new host, where it can again integrate via a site-specific recombinase at the palindromic sequence 5'-TTCATGAA-3'³⁴. The presence of internal copies of this sequence can lead to modular excision of the island. Fixation of the pathogenicity genes in some other potato scab agents (*Streptomyces scabies* and *Streptomyces acidiscabies*) is correlated with degradation of the recombinase gene and/or its target sites³⁴ or with loss of a module that encodes all the functions needed for mobilisation³⁵.

Remarkably, the expression of some of the pathogenicity genes, including those for the phytotoxin thaxtomin, is determined in part by the cellobiose sensor CebR, the universal regulator of *Streptomyces* cellulolytic genes, integrating pathogenicity-linked cellulolytic activity with more ancient overall responses to cellulose-related metabolites³⁶. The connection of pathogenicity with cellulose is further emphasised by the fact that thaxtomin is an inhibitor of plant cellulose synthase. The biosynthetic clusters for thaxtomin and a second phytotoxin, coronafacoyl phytotoxin, are under further control from pathway-specific regulatory genes that appear to respond to plant metabolites³⁷ and from global regulatory genes (*bld* genes that regulate colony differentiation and antibiotic production, see below)³⁸. In an extension of this regulatory scenario, the induction of a large swathe of glycohydrolitic enzymes of *S. scabies* is enhanced when cellulose in the medium is supplemented with suberin, a complex lipid- and polyaromatic-containing major component of the potato tuber periderm³⁹.

Although scab research has focused on the pathogenicity islands, comparative genomics has revealed 64 other genes that are common to four distinct scab agents but absent from non-pathogens, as well as an exceptionally high proportion of enzymes such as pectinases and cutinases that would aid or take advantage of pathogenicity^{35,40}.

Genome and pan-genome evolution, the role of recombination, and its molecular basis

Comparative genomics has been dramatically accelerated by the advent of cheap and rapid sequencing (but note that the quality of the deposited *Streptomyces* genome sequences is variable⁴¹).

Early deductions about the core and accessory genes of streptomycetes have been refined⁴²: the genomes of 17 diverse strains (ranging from 6.7 to 12.3 Mb) contained from 5,382 to 10,022 CDSs, of which 2,018 were universally present (the core genome, which occupies most of the central region of *Streptomyces* linear chromosomes), with most of the remaining genes being present in only one or a few genomes. This accessory genome totalled 32,574 genes, so the *Streptomyces* pan-genome comprises at least 34,592 genes. The accessory genes are mostly clustered at the chromosome ends, which are well known to be unstable and subject to replacement by recombination with incoming linear plasmids or conjugally transferred segments of chromosomes from other streptomycetes, but recent analyses have revealed an unexpectedly high level of historical recombination within representative core genes too, resulting in phylogenetic incongruence among these genes^{40,43,44}. The recombination levels detected are high enough to raise the question of whether the evolutionary radiation of the genus has taken place over a significantly shorter time than has been widely accepted⁴⁴. During divergence from common ancestors, such genetic exchange continued at a rate that decreased as divergence increased, generating a reticulate origin of extant streptomycetes, largely defined by a relatively small number of ancestral recombination events. Recombination associated with nascent divergence continues into the near-present, since recombination events detected by genome analysis of independent isolates of the same species are about 100-fold more frequent than those between species⁴³. The fixation of particular recombination events is thought to reflect periods of rapid demographic expansion^{27,28}.

The high frequency of genetic exchange implied by these studies may be because conjugal DNA transfer in streptomycetes is much more efficient than it is in most other bacteria, at least in the laboratory: instead of a complex type IV secretion apparatus transferring single-stranded DNA, streptomycetes employ the products of *traB* genes encoded by autonomous or chromosomally integrated plasmids to transfer double-stranded DNA⁴⁵. TraB proteins belong to the FtsK/SpoIIIE family needed in many bacteria to complete the segregation of chromosomes into newly formed daughter cells, but, while FtsK/SpoIIIE proteins assemble in the nascent septum, TraB proteins assemble at growing hyphal tips. Transfer, which is ATP dependent, involves the interaction of TraB with short recognition sequences (consensus 5'-GACCCGGA-3' in plasmid pSVH1): 25 clusters of four such sequences are also spaced along the chromosome of *S. coelicolor*, accounting for chromosome mobilisation⁴⁵. Initial plasmid transfer is followed by the remarkable spread of the plasmid, sometimes throughout most of the recipient mycelium, in a process in which specialised Spd proteins assemble with TraB in the vegetative septa that occur roughly once every 10–20 μm ^{45,46} (Figure 2). The importance of TraB for plasmid spread (presumably providing the ATPase-driven motor function lacking from Spd proteins) was nicely demonstrated by showing that the presence of TraB in a recipient permitted efficient spreading following the occasional transfer of a *traB*-deleted plasmid⁴⁶.

It was a surprise when a recent cytological study showed that plasmid invasion events were associated with contact between the lateral walls of donor and recipient hyphae, and events involving

donor tips were not seen⁴⁶. This unexpected result may still be compatible with tip involvement: TraB may also assemble at nascent branch tips, some distance away from leading tips. In any case, the cytological analysis could presumably detect only transfer events that had occurred a significant time earlier, after subsequent plasmid multiplication and spread had already occurred, so the state of hyphae at the time of primary transfer could not be assessed.

Within-species comparative genomics

Generally, comparative genomics of closely related streptomycetes has had little attention, though marine isolates from two distinct sponges were interestingly closely related to the terrestrial *Streptomyces albus* G derivative J1074 (which is often used for heterologous expression work) in a study identifying candidate marine-adaptation genes present only in the two marine isolates and in some other marine actinomycetes^{33,47}. Four further J1074-related organisms were subsequently found in 140 *Streptomyces* genomes available at the time, making this species group the most readily isolated to date^{33,47}. This provided an opportunity to examine the dynamics of acquisition and loss of secondary metabolite biosynthetic gene sets in relatively recent evolution (i.e. within a species group). Ten biosynthetic gene clusters were specific to, and conserved among, the J1074 group, while a roughly equal number showed some degree of isolate specificity^{33,47}.

New perspectives in growth and development of *Streptomyces*

Germination is controlled by several second messengers

Spore germination involves the action of “resuscitation-promoting factors” (Rpfs, specialised peptidoglycan hydrolases), probably as part of a peptidoglycan remodelling system⁴⁸. At least one of five partially functionally redundant Rpfs of *S. coelicolor* is subject to multilevel regulation by different nucleotide second messengers: transcription initiation, controlled by the cAMP-binding protein previously associated with germination; transcriptional attenuation via a riboswitch responsive to cyclic di-AMP; and changing rates of proteolysis in response to levels of ppGpp⁴⁹ (Figure 2).

Importance of cellulose-like glycan synthesis for hyphal growth

Streptomyces growth is unusual, but not unique, among bacteria in taking place at tips through the activities of a protein complex (“polarisome”). The key polarisome component, DivIVA, coordinates aspects of intracellular and cell surface growth in ways that may differ during vegetative and reproductive growth (for reviews, see 50,51). It is emerging that normal growth and development require deposition at hyphal tips of an uncharacterised glycan through the combined action of a cellulose synthase-like protein (ClsA) and GlxA, the product of the gene downstream of *clsA*, which is a copper oxidase that may oxidise the glycan as it is secreted⁵² (Figure 2). Like ClsA, GlxA is tip-located at least at some growth stages⁵³, though the two proteins have not been analysed together. The synthesis of extracellular polysaccharides soon after germination, under the control of *clsA*, *glxA*, and the newly identified locus *matAB*, is responsible for the aggregation of germlings, leading to the formation of mycelial clumps in submerged culture, a well-known problem in industrial fermentations^{54,55}.

Figure 2. Recent discoveries about germination and hyphal growth. Three different nucleotide signalling molecules are involved in stimulating the production of peptidoglycan hydrolase(s) (Rpf, for resuscitation-promoting factors), leading to remodelling of the spore wall and the emergence of a germ tube⁴⁹. Continued hyphal elongation is co-ordinated by the DivIVA-based polarisome⁵⁰ and (in addition to cell wall growth) involves the extracellular production of cellulose-like glycan catalysed by polarisome-linked CslA (cellulose synthase-like) associated with the copper oxidase GlxA^{52,53}. MatAB proteins have an uncharacterised role in glycan production⁵⁴. Vesicular membrane structures form in apparently irregular locations within hyphae, and some of them extend across the hyphal compartment, separating nucleoids^{57,58}.

Membranous structures appear to subdivide the cytoplasm of vegetative hyphae into compartments

The recent application of advanced microscopic methods has suggested that vesicular, DNA-impermeable membranous structures may delimit compartments in vegetative hyphae in the absence of conventional septa, may account for the viability of some hyphal fragments of *ftsZ* mutants, and may be implicated in the heterogeneous staining of vegetative hyphae with vital stains (interpreted as programmed death^{56–58}) (Figure 2). These intriguing observations raise questions of how the structures are placed in time and space, how they affect the spread of plasmids through the mycelium (see above), and whether they might limit the spread of initially tip-localised phage infections. Addressing such questions may require inventive genetic approaches.

Complex and subtle influences on the initiation of aerial development, involving transcription, proteolysis, and hormone-like molecules

The onset of reproductive aerial growth is sensitive to diverse stresses and signals. In *S. coelicolor*, most of the nine σ^B -like

sigma factors (compared to four in *Bacillus subtilis*) are involved in stress responses and/or differentiation. There is growing evidence of complex crosstalk interactions among the even larger number of anti-sigma and anti-anti-sigma factors for this class of σ factor, and it has been suggested that the signal input to regulate these interactions may involve some of the variable, but always multiple, collection of *whiJ*-like gene sets present in *Streptomyces* genomes: the clusters include anti-sigma genes of this type and have been implicated in development and antibiotic production^{51,59}. In one such cluster, which is known to influence antibiotic production and differentiation, a small RNA antisense to the Sco4676–4677 intergenic region has been found to have complicated regulatory effects on Sco4676–4677 expression⁶⁰.

Furthermore, particularly since the discovery that actinomycetes have a eukaryotic-like proteasome system, with ubiquitylation of targets being replaced by pupylation⁶¹, regulation at the level of protein degradation is becoming increasingly apparent. This has been reinforced by the cataloguing of pupylation/proteasome targets and the phenotypes of mutants in the system, which show

changes in differentiation and antibiotic production levels^{62–64}. Presumably, controlled proteolysis is involved in the death of some hyphal compartments that accompanies stages of growth and development⁵⁶.

It has been known for decades that in *Streptomyces griseus* the repression of a central developmental regulatory gene, *adpA*, by the TetR-like protein ArpA is relieved by the accumulation of A-factor, an extracellular hormone-like gamma-butyrolactone (GBL). It now seems that *adpA* in most, perhaps all, other streptomycetes may be subject to regulation by similar ArpA-like repressors, but with species-specific GBL ligands, permitting coordinated sporulation of one species without triggering sporulation of nearby different species⁶⁵. In an opposite scenario, *S. coelicolor* and *S. venezuelae* produce an identical GBL called SCB1 and SVB1, respectively, making it possible for them to respond to each other^{66,67}.

Spectacular insights using a new model organism, *Streptomyces venezuelae*: time-lapse cell biology, heterodimeric regulatory proteins, global regulation by a cyclic di-GMP receptor, and developmental differences between species

Streptomyces colony development has usually been studied on solid agar medium, since *S. coelicolor* (like many other streptomycetes) does not undergo a full developmental cycle in liquid culture. Studies of surface-grown cultures are bedevilled by developmental asynchrony and heterogeneity, which are particularly problematic for high-resolution biochemical analysis, including “omic” approaches to the definition of the regulons controlled by developmental regulatory genes. These problems have been circumvented by the adoption of *S. venezuelae*, which grows very rapidly and undergoes comprehensive and near-synchronous differentiation in liquid culture, including under coverslips⁶⁸. In a first illustration of the value of this system, a detailed time-lapse study has shown that the ParA and ParB chromosome segregation proteins play an important coordinating role in the transition from ongoing aerial growth to growth cessation and the onset of sporulation septation, and even have opposing influences on the rate of tip extension and eventual length of pre-sporulation hyphae⁶⁹. Further advances are promised by the development of a microfluidics system that allows continuous direct observation of fluorescently marked key proteins through the entire developmental cycle⁶⁸.

Submerged sporulation has allowed the highly successful combined application of microarray-based transcriptomics and chromosome immunoprecipitation sequencing (ChIP-seq) analysis to reveal two exciting and novel cases of very close interactions between different developmental regulators. In one case⁷⁰, an atypical response regulator, BldM, has two rounds of activity: first, it forms a homodimer to activate other developmental regulatory genes and morphogenetic functions needed for aerial hyphae to grow and, second, it forms a heterodimer with another atypical response regulator, WhiI, to activate genes needed for the aerial hyphae to turn into spore chains. In another case⁷¹, with wide implications for all actinomycetes including pathogenic mycobacteria, two enigmatic proteins conserved across all actinomycetes have been found to bind to the same target promoters, activating some and repressing others. These proteins are WhiA, which is also present in Gram-positive Firmicutes and has a structure closely similar to that of eukaryotic homing

endonucleases (but lacking key catalytic residues), and WhiB, the archetype of the large “Wbl” family of iron-sulphur proteins represented throughout actinomycetes but found nowhere else (other Wbl proteins conserved among streptomycetes include two other developmental regulators, WblA and WhiD)⁵¹. The ChIP-seq/transcriptome profiles of targets of Bld and Whi regulators in *S. venezuelae* have provided many clues about the nature and coordination of the components contributing actively to development^{70–72}.

It is likely that BldM, WhiI, WhiA, and WhiB respond to different signal inputs, and the elucidation of these signals is a major challenge (but see the next section). These four developmental activators, and most of the well-known developmental regulatory *bld* and *whi* genes and many of their targets, are among the large number of genes directly controlled by BldD⁷², a master repressor of development whose regulatory ligand has been revealed as cyclic di-GMP⁷³. This important discovery begins to provide an explanation for the presence of eight actual or predicted diguanylate cyclase genes in *S. coelicolor*, some with associated cyclic di-GMP phosphodiesterase domains⁷⁴. Notably, the [BldD]₂:[cyclic di-GMP]₄ complex is structurally new to science^{73,75}.

Although it is highly probable that the developmental programmes of all streptomycetes are broadly the same, some of the details may differ between species. This seems to be the case with *whiG*, which encodes a sigma factor important in initiating sporulation septation of aerial hyphae. Although *whiG* transcription depends on WhiA and WhiB in *S. venezuelae*⁷¹, in *S. coelicolor* its transcription was nearly the same in *whiA* or *whiB* mutants as in the wild-type⁷⁶. In another example, the promoter of the WhiI-dependent *inoA* gene of *S. coelicolor* (for inositol phosphate synthetase) could bind truncated WhiI *in vitro*⁷⁷, indicating that WhiI could bind DNA in the absence of BldM; yet in ChIP-seq analysis of an *S. venezuelae* *bldM* mutant, WhiI did not bind to any targets⁷⁰. Other inter-species differences were previously noted between *S. coelicolor* and *S. griseus*⁷⁸. A different kind of surprise concerning the WhiG sigma was the discovery in *Streptomyces chatanoogensis* that it can bind to target promoters in the absence of RNA polymerase core enzyme, including some in the gene cluster for natamycin production: binding was either to typical WhiG-binding sites or to CGTCA repeat elements, and was needed for gene activation⁷⁹.

Could endogenous nitric oxide have an important developmental regulatory role?

Speculation about a possible role of nitric oxide as a regulatory ligand for WhiB and other Wbl proteins was stimulated by the extremely high affinity for nitric oxide of some Wbl proteins of streptomycetes and mycobacteria, which has been studied in detail at the molecular level^{80–83}. It has generally been implied that the source of nitric oxide would be exogenous (e.g. produced by eukaryotic hosts as a defence mechanism). However, comparative genomics revealed the general presence in Wbl-containing actinomycetes of other nitric oxide-interacting proteins (or biochemically unstudied homologues) that might be involved in cycling nitric oxide, raising the possibility that *endogenous* nitric oxide might regulate development by interacting with Wbl proteins⁵¹.

This model was compromised by the absence of evidence for the conserved endogenous production of nitric oxide across actinomycetes, most of which do not have a conventional nitric oxide synthase. However, endogenous nitric oxide production involving conserved actinobacterial genes has now been convincingly demonstrated in *S. coelicolor*⁸⁴ and mycobacteria^{85,86} (Figure 3). Thus, in *S. coelicolor*, organic nitrogen is converted by an unknown route to nitrate, which is reduced by nitrate reductase to nitrite⁸⁴. This mainly involves NarG2, one of three nitrate reductases in *S. coelicolor*: NarG2 is predominant in growing hyphae and is induced by moderate hypoxic downshift, while NarG3 has a subordinate role and NarG1 is mainly active during germination⁸. Nitrite was shown in turn to be the source of nitric oxide (albeit again by an unknown route)⁸⁴. This source of nitric oxide mirrors a pathway important in eukaryotes⁸⁸, where endogenous nitric oxide is a major physiological signalling molecule, even influencing epigenetic changes⁸⁹.

The recycling of endogenous nitric oxide was also studied by 84 and is illustrated in Figure 3, which also shows how recycling might be connected to the regulatory functions of Wbl proteins. Nitric oxide re-enters the nitrogen oxide cycle via flavohaemoglobin, which oxidises nitric oxide to nitrate. In a feedforward mechanism, nitric oxide is the ligand of NsrR, which represses the *hmpA1* and *hmpA2* genes for flavohaemoglobin and its own promoter^{84,90}. Remarkably, the affinity of NsrR for the *hmpA* promoters is more sensitive to nitric oxide than is its affinity for the *nsrR* promoter⁹¹, permitting different strengths of induction of flavohaemoglobin as NO concentration increases. Another dimension of the cycle is provided by the interaction of endogenous nitric oxide with DevS, a protein kinase that activates the cognate response regulator DevR, which in turn activates the *narG2* gene for the major nitrate reductase needed for nitric oxide production⁸⁴. Interestingly, an earlier report⁹² found that DevS could also phosphorylate the SCO3818 response regulator, which is very similar to DevR, and might therefore

Figure 3. Model integrating the nitrogen oxide cycle (ochre) with mycothiol cycling (yellow) and with morphological development mediated by Wbl proteins (red) in *Streptomyces coelicolor*. In the nitrogen oxide cycle⁸⁴, nitrate generated intracellularly by an unknown pathway is reduced to nitrite mainly by the NarG2 enzyme⁸⁷. Nitrite is the source of nitric oxide (NO) (mechanism unknown)⁸⁴. Nitric oxide is oxidised to nitrate by flavohaemoglobins specified by SCO7428 (incorrectly given as SCO7472 in 84) and SCO7094, which are induced when nitric oxide binds to NsrR, the repressor of both genes^{84,90}. Internal nitric oxide also binds the haem-containing sensor kinase DevS, which then phosphorylates the cognate response regulator DevR. DevR-P induces Nar2 in an autoinducing feedforward loop⁸⁴. Nitric oxide binds strongly to Wbl proteins such as WblA, WhiB, and WhiD, modulating their regulatory activity⁸¹ and hence coordinating differentiation. Other regulatory components may take part in the action of Wbl proteins: for example, WhiB target genes are all also dependent on WhiA protein⁷¹. It is speculated that the Wbl proteins are denitrosylated by the SCO4179 (nitrobindin?)-mediated transfer of nitric oxide to mycothiol (MSH) to generate MSNO⁵¹. MSNO is denitrosylated by MSNO reductase, with the resulting oxidised mycothiol then being reduced to MSH by an undetermined mechanism.

also be involved in regulating the nitrogen oxide cycle: orthologues of SCO3818 are very widespread across actinobacteria, whereas DevS orthologues are more sporadic across the phylum (<http://streptomyces.org.uk/actinoblast/>).

Artificially engineered production of endogenous nitric oxide triggers the onset of production of at least one secondary metabolite (undecylprodiginines [RED]) by increasing the expression of the major RED pathway activator gene *redD*, but it inhibits the onset of aerial growth⁸⁴. It remains to be determined whether this is a straightforward reflection of the normal signalling activity of nitric oxide or whether the experimental procedures disrupt the normal balance of the nitrogen oxide cycle. In a further twist, it appears that endogenous nitrite can also serve as an extracellular signal to nearby mycelium⁸⁴.

Regulation of septum positioning and cell wall changes during the formation of spores

Many of the processes involved in sporulation septation and spore maturation require high levels of proteins that might interfere with growth if expressed inappropriately, so mechanisms to prevent untimely expression can be expected. Thus, production of some of these proteins, such as FtsZ, is repressed by the BldD:cyclic di-GMP complex⁷². Timely increased expression of such division genes, presumably triggered by cyclic di-GMP hydrolysis, must

also be accompanied by a mechanism for the regular positioning of sporulation septa. New evidence suggests that SepG, a small membrane protein encoded by a gene next to *divIVA* in many Gram-positive bacteria, provides the most basic anchor yet described for the building of sporulation septa, recruiting protein SsgB, which in turn recruits FtsZ⁹³ (Figure 4). After septation, SepG relocates to the prespore periphery, where it may interact with the cell wall synthesising complex during spore wall remodelling. Five protein kinases, encoded by a cluster of adjacent genes, phosphorylate key proteins of this complex, keeping them inactive until sporulation septation begins: it is assumed that activation involves one or more of the more than 50 phosphatases encoded by the *S. coelicolor* genome⁹⁴.

The variable size, thinner walls, diffuse nucleoids, and heat sensitivity of *sepG* mutant spores are intriguingly reminiscent of the phenotype of mutants in the conserved *whiD* gene⁹⁵, which encodes a Wbl protein whose nitric oxide interactions have been particularly closely studied (e.g. 83 [see above]). Conceivably, *sepF* could be an important WhiD target (Figure 4).

Other aspects of cellular biochemistry that change during growth and development: lipids and glycogen

In a new perspective on development, the lipid composition of *S. coelicolor* differed strikingly when comparing cultures grown

Figure 4. Schematic diagram of recent discoveries about late stages in sporulation. As aerial hyphae approach sporulation, c-di-GMP phosphodiesterase releases *ftsZ* from repression by BldD:c-di-GMP^{72,74,75}. FtsZ attaches to SsgB, held at regularly spaced positions by membrane-bound SepG⁹³. Another SsgB-like protein, SsgA, also plays a role in SsgB location, which is less well defined¹³⁹. FtsZ then condenses into Z-rings, which guide septation, and SepG relocates to the periphery of the prespore compartments⁹³. The spore wall remodelling complex, held in an inactive phosphorylated state, is activated by one or more of many phosphorylases⁹⁴ and, apparently with the involvement of SepG, causes prespores to become rounder and thicker walled. I tentatively suggest that the increased amounts of SepG at this stage may depend on the Wbl-type regulator WhiD (possibly influenced by nitric oxide [NO] – see Figure 3), since *sepG* and *whiD* null mutants have very similar phenotypes^{93,95}.

on solid or in liquid media, and mutants defective in ornithine lipid biosynthesis showed precocious development and antibiotic production⁹⁶. It would be of interest to apply such an analysis to *S. venezuelae*.

Studies of carbon storage metabolism in *Streptomyces* and mycobacteria had revealed a new and widespread pathway for glycogen biosynthesis^{97,98}. This GlgE pathway is duplicated in most streptomycetes, with the two sets being expressed differently during development, and the pathway generally exists alongside the classical GlgAC pathway. In an exception to this generalisation, *S. venezuelae* has only one *glgE* gene set and lacks *glgAC*. This made it possible to construct glycogen-free mutants, leading to the finding that spores could be generated without the developmental accumulation of glycogen. Sporulation appeared to be normal, unless the strain/medium combination resulted in the accumulation of the toxic GlgE substrate maltose-1-phosphate⁹⁸.

Stress responses and secondary metabolism Connecting redox homeostasis to primary metabolism

Global stress responses often involve the action of alternative sigma factors, whose target regulons may be informative in understanding physiological and cell biological adjustments that have evolved to accommodate stresses. Thus, it was recently found that one SigR target, the *ndgR* gene, encodes a regulator of sulphur assimilation and branched-chain amino acid synthesis. NdgR is presumed to affect redox homeostasis at the levels of precursor and cofactor supply to redox buffering agents such as mycothiol and supply of CoA derivatives of branched-chain fatty acids for membrane biosynthesis⁹⁹.

From global physiology to the expression of biosynthetic genes for secondary metabolites: new global regulators and expanding roles for cluster-situated regulatory genes

As a broad generalisation, antibiotic biosynthetic gene clusters depend on regulatory genes located within the cluster, which are themselves directly regulated by many higher level regulatory proteins, encoded by genes such as the developmental *bld* genes scattered around the chromosome¹⁰⁰. New regulators continue to be identified: a TetR-like protein specified by Sco3201 has been found to bind to the promoters of several known developmental and antibiotic biosynthetic genes¹⁰¹, and the *abrC* cluster (Sco4596–8) encoding a response regulator (AbrC3) and two histidine protein kinases has been shown to influence secondary metabolism^{102,103}. An accumulation of papers has shown that the promoter of the *actII-orf4* gene that activates the ACT biosynthetic genes of *S. coelicolor* binds at least eight, and possibly as many as 12, regulatory proteins¹⁰⁰, a number to which AbrC3 has been added¹⁰³. A different scenario was observed in the case of the novobiocin biosynthetic genes when heterologously expressed in *S. coelicolor*: a promoter of these biosynthetic genes (instead of a regulatory gene) was a target for as many as 11 host-specified proteins and the cluster-situated NovG regulatory protein¹⁰⁴. Only one of these 11 proteins, AbsC, is also known to bind to *actII-orf4*.

Regulatory genes such as *actII-orf4* and *novG* that are situated within biosynthetic clusters were previously called “pathway specific”, but numerous observations have shown that they may

have targets outside the cluster, so their renaming as “cluster-situated regulators” (CSRs) by 105 has proved prescient. In a striking recent example involving *Streptomyces avermitilis*, a CSR belonging to a class with PAS-LuxR-like domains was shown to interact with genes for different pathways and with many other promoters for housekeeping genes¹⁰⁶.

Small-molecule ligands for proteins that regulate antibiotic production: feedback and feedforward regulation and crosstalk within and between organisms

Some of the proteins regulating antibiotic production interact with known small-molecule ligands associated with different aspects of primary metabolism¹⁰⁰. Another widespread scenario is the involvement of diffusible GBL and related signal molecules as the ligands of a subfamily of TetR-like regulators, like the A-factor/ArpA pair of *S. griseus* (see above). This may operate differently in different species and different pathways. In *S. venezuelae*, the SVB1 GBL receptor JadR3 has complex interactions with the promoters of both its own biosynthetic genes and jadomycin biosynthetic genes. This results in an integrated feedback and feedforward regulatory scheme that accounts for both the onset of the major period of jadomycin production and its subsequent down-regulation⁶⁶. In *S. coelicolor*, genome-wide ChIP-seq analysis has revealed an extensive network of regulatory interactions centred on ScbR (GBL receptor) and ScbR2 (pseudo-GBL receptor that binds antibiotics)^{107,108}.

A major development in the regulation of antibiotic production concerns the activities of secondary metabolites themselves as regulatory ligands^{65,100}. In one recent example from *Streptomyces antibioticus*, the spiroketone chlorothricin and some of its biosynthetic intermediates act as regulatory ligands for the CSR ChlF1¹⁰⁹. In another case involving the aminocoumarin caci-biocin, a repressor of the biosynthetic genes, CabR, loses its DNA-binding activity in the presence of the end product or other aminocoumarins¹¹⁰.

Internal cross-pathway regulation mediated by end products has been subjected to closer analysis following the discovery of cross-talk between the chloramphenicol and jadomycin biosynthetic pathways in *S. venezuelae*. JadR1, the main activator of the jadomycin gene cluster, was implicated in repressing chloramphenicol biosynthesis¹¹¹, but a recent further analysis of this situation revealed that cross-regulation, though still observed, was markedly different when the experiments were done on a different medium¹¹². It is not yet known whether these differences involve *jadX*, another newly identified regulatory gene in the jadomycin cluster¹¹³. Knocking out *jadX* prevented the well-known induction of jadomycin production by stress conditions while at the same time putting chloramphenicol biosynthesis under the control of these stresses, and it was shown that both jadomycins and chloramphenicol could interact with JadX¹¹³. Interestingly, genes for JadX-like proteins are present in some other antibiotic biosynthetic clusters¹¹³.

In *S. coelicolor*, ScbR2, a protein closely similar to GBL-binding proteins, has been found to interact with jadomycins from *S. venezuelae* (and other heterologously produced angucyclines) to relieve the ScbR2-mediated repression both of the global

regulatory gene *adpA* and of *redD*, encoding an activator of the RED pathway genes¹¹⁴. Thus, *S. coelicolor* can respond to the presence of another species by undergoing developmental changes and switching on antibiotic production. Another example of such a response has been documented for lidamycin biosynthesis in *Streptomyces globisporus*, involving a widely conserved TetR-like global activator of antibiotic production, AtrA¹⁰⁷. AtrA directly activates lidamycin biosynthetic genes, but its activity is prevented by binding either a lidamycin biosynthetic intermediate, heptaene, or heterologously produced ACT¹⁰⁷. Inter-species interactions affecting antibiotic production are proving to be widespread among streptomycetes: a large-scale study of inter-organism interactions revealed many cases of production of secondary metabolites induced by growth close to another species^{115,116}, and another study found that low levels of lincomycin induced ACT production by *S. coelicolor*¹¹⁷. It will be interesting to see whether ScbR2-, JadR1-, and JadX-like proteins are often involved in such crosstalk, or even in some of the more divergent interactions of streptomycetes with other microbes that continue to emerge from co-culture studies (for recent examples, see 118–121).

A high-resolution global survey of gene expression in *S. coelicolor* at different growth stages and the importance of translation-level regulation

Through the use of RNA sequencing to analyse the global transcription profile, and ribosome profiling (ribo-seq) to analyse translation, a near-comprehensive gene expression picture of *S. coelicolor* at different stages of growth in liquid culture has been obtained¹²². This survey identified most of the transcription start sites in the entire genome, including those for 230 small RNAs. Amazingly, about 21% of protein-coding RNAs are leaderless (and lack conventional ribosome-binding sites). Moreover, clear changes were seen in translation rates between different growth stages, including increased efficiency of translation of mRNAs for major CSR genes at the end of the main growth phase but decreased efficiency for the biosynthetic genes that they control. It is not clear whether these changes reflect different codon usage, a relevant question given that several of the CSR genes for antibiotic production contain the very rare codon TTA^{65,100}. The protein-level expression of TTA-containing genes is usually dependent on the *bldA* gene, which encodes the cognate tRNA; abundance of this tRNA increases on the transition into stationary phase^{123,124}.

In an interesting counterpoint to the *bldA* story, the TTA codon in *ccrA*, the CSR gene controlling cephamycin C and clavulanic acid biosynthesis in *S. clavuligerus*, does not make the production of CcrA *bldA*-dependent¹²⁵. It has now turned out that many, but not all, TTA-containing genes in *S. clavuligerus* display the same anomaly¹²⁶. It would perhaps be worth examining the *bldA* dependence of these genes when introduced into *S. coelicolor* and of TTA-containing genes from *S. coelicolor* upon introduction into *S. clavuligerus*.

Technological advances

The last few years have seen steps forward in the development and application of diverse technologies for the study and exploitation of streptomycetes. The first systems for applying genome editing to streptomycetes will soon accelerate the construction of diverse

kinds of mutant^{127–130}, while the discovery of a CRISPR-Cas system in *S. avermitilis* may open up new approaches¹³¹. Along with new strategies for the cloning and deletion of very large segments of *Streptomyces* DNA, sufficient to encompass almost all antibiotic biosynthetic gene clusters^{132,133}, procedures are in place to exploit the ever-growing numbers of uncharacterised clusters emanating from genome sequencing, along the lines summarised by 134 and 135. Bioinformatic tools have been developed to predict the nature of the products of such clusters¹³⁶. One component of systems-based exploitation will be a panel of strong promoters, such as 20 promoters significantly stronger than the widely used *ermE** that have been characterised in *S. albus* G¹³⁷. Another promoter survey has identified promoters with very stable expression levels at different growth stages in two strains of *S. coelicolor* A3(2) and in other species¹³⁸. These are likely to be valuable standards in transcription studies.

Concluding remarks

The rapid recent progress summarised in this article seems set to accelerate further, particularly in light of the imminent application of powerful additions to the techniques available for *Streptomyces* research. Research targets will include further elucidation of the ligands that determine the activity of regulatory proteins, the ways in which developmentally important proteolysis is controlled, the search for the molecular factors involved in symbiotic associations, and the structural and functional characterisation of the DivIVA-centred polarisome that determines tip growth. In the course of such studies, an integrated picture of these diverse aspects of *Streptomyces* biology will begin to emerge. It is likely that this will require increasing input from computational biologists. The accumulating understanding of the biology of these key antibiotic-producing organisms can be expected to facilitate the development of new therapeutic agents in the global struggle against antibiotic-resistant pathogens.

Abbreviations

ACT, actinorhodin; ChIP-seq, chromosome immunoprecipitation sequencing; CSR, cluster-situated regulator; GBL, gamma-butyrolactone; RED, undecylprodiginine.

Competing interests

The author declares that he has no competing interests.

Grant information

The authors declared that no grants were involved in supporting this work.

Acknowledgments

I thank the John Innes Centre for an Emeritus Fellowship. The writing of this (my final) review article has not directly involved anyone else, but it has been made possible by countless stimulating interactions with many colleagues over many years. I have also benefited greatly from the privilege of being a member of F1000, with organized access to the richness of *Streptomyces* literature. Finally, I thank Jean Chater for allowing me 50 years of indulgence in my fascination with bacterial genetics.

References

1. Charlop-Powers Z, Owen JG, Reddy BV, *et al.*: **Chemical-biogeographic survey of secondary metabolism in soil.** *Proc Natl Acad Sci U S A.* 2014; 111(10): 3757–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Charlop-Powers Z, Owen JG, Reddy BV, *et al.*: **Global biogeographic sampling of bacterial secondary metabolism.** *eLife.* 2015; 4: e05048.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
3. Hopwood DA: ***Streptomyces* in Nature and Medicine: The Antibiotic Makers.** Oxford University Press, New York, NY. 2007.
[Publisher Full Text](#)
4. Chater KF, Biró S, Lee KJ, *et al.*: **The complex extracellular biology of *Streptomyces*.** *FEMS Microbiol Rev.* 2010; 34(2): 171–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Wibberg D, Al-Dilaimi A, Busche T, *et al.*: **Complete genome sequence of *Streptomyces reticuli*, an efficient degrader of crystalline cellulose.** *J Biotechnol.* 2016; 222: 13–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Lim JH, Lee CR, Dhakshnamoorthy V, *et al.*: **Molecular characterization of *Streptomyces coelicolor* A(3) SCO6548 as a cellulose 1,4- β -cellobiosidase.** *FEMS Microbiol Lett.* 2016; 363(3): pii: fnv245.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Book AJ, Lewin GR, McDonald BR, *et al.*: **Evolution of High Cellulolytic Activity in Symbiotic *Streptomyces* through Selection of Expanded Gene Content and Coordinated Gene Expression.** *PLoS Biol.* 2016; 14(6): e1002475.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
8. Barka EA, Vatsa P, Sanchez L, *et al.*: **Taxonomy, Physiology, and Natural Products of *Actinobacteria*.** *Microbiol Mol Biol Rev.* 2015; 80(1): 1–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Tenconi E, Urem M, Świątek-Polatyńska MA, *et al.*: **Multiple allosteric effectors control the affinity of DasR for its target sites.** *Biochem Biophys Res Commun.* 2015; 464(1): 324–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
10. Świątek-Polatyńska MA, Bucca G, Laing E, *et al.*: **Genome-wide analysis of *in vivo* binding of the master regulator DasR in *Streptomyces coelicolor* identifies novel non-canonical targets.** *PLoS One.* 2015; 10(4): e0122479.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Viens P, Dubeau MP, Kimura A, *et al.*: **Uptake of chitosan-derived D-glucosamine oligosaccharides in *Streptomyces coelicolor* A3(2).** *FEMS Microbiol Lett.* 2015; 362(9): pii: fnv048.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Seipke RF, Kaltenpoth M, Hutchings MI: ***Streptomyces* as symbionts: an emerging and widespread theme?** *FEMS Microbiol Rev.* 2012; 36(4): 862–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Viaene T, Langendries S, Beirincx S, *et al.*: ***Streptomyces* as a plant's best friend?** *FEMS Microbiol Ecol.* 2016; 92(8): pii: fiw119.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Liu M, Abdel-Mageed WM, Ren B, *et al.*: **Endophytic *Streptomyces* sp. Y3111 from traditional Chinese medicine produced antitubercular pluramycins.** *Appl Microbiol Biotechnol.* 2014; 98(3): 1077–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Yang X, Yang Y, Peng T, *et al.*: **A new cyclopeptide from endophytic *Streptomyces* sp. YIM 64018.** *Nat Prod Commun.* 2013; 8(12): 1753–4.
[PubMed Abstract](#)
16. Conti R, Chagas FO, Caraballo-Rodríguez AM, *et al.*: **Endophytic Actinobacteria from the Brazilian Medicinal Plant *Lychnophora ericoides* Mart. and the Biological Potential of Their Secondary Metabolites.** *Chem Biodivers.* 2016; 13(6): 727–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Chen X, Pizzatti C, Bonaldi M, *et al.*: **Biological Control of Lettuce Drop and Host Plant Colonization by Rhizospheric and Endophytic Streptomycetes.** *Front Microbiol.* 2016; 7: 714.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Zloch M, Thiem D, Gadzala-Kopciuch R, *et al.*: **Synthesis of siderophores by plant-associated metallo-tolerant bacteria under exposure to Cd²⁺.** *Chemosphere.* 2016; 156: 312–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Miao GP, Zhu CS, Feng JT, *et al.*: **Effects of plant stress signal molecules on the production of wilforfingine in an endophytic actinomycete isolated from *Tripterium wilfordii* Hook.f.** *Curr Microbiol.* 2015; 70(4): 571–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Shi Y, Zhang X, Lou K: **Isolation, characterization, and insecticidal activity of an endophyte of drunken horse grass, *Achnatherum inebrians*.** *J Insect Sci.* 2013; 13(1): 151.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Singh SP, Gaur R: **Evaluation of antagonistic and plant growth promoting activities of chitinolytic endophytic actinomycetes associated with medicinal plants against *Sclerotium rolfsii* in chickpea.** *J Appl Microbiol.* 2016; 121(2): 506–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Wang W, Zhai Y, Cao L, *et al.*: **Illumina-based analysis of core actinobacteriome in roots, stems, and grains of rice.** *Microbiol Res.* 2016; 190: 12–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Passari AK, Chandra P, Zothanpuia, *et al.*: **Detection of biosynthetic gene and phytohormone production by endophytic actinobacteria associated with *Solanum lycopersicum* and their plant-growth-promoting effect.** *Res Microbiol.* 2016; 167(8): 692–705.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Mukasheva T, Berzhanova R, Ignatova L, *et al.*: **Bacterial endophytes of Trans-Ili Alatau region's plants as promising components of a microbial preparation for agricultural use.** *Acta Biochim Pol.* 2016; 63(2): 321–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Tchinda RA, Boudjeko T, Simao-Beaunoir AM, *et al.*: **Morphological, Physiological, and Taxonomic Characterization of Actinobacterial Isolates Living as Endophytes of Cacao Pods and Cacao Seeds.** *Microbes Environ.* 2016; 31(1): 56–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Peng SM, Wang BL, Xu JZ, *et al.*: **[Effect of different treatment on endophytic bacterial communities in continuous cropping of *Chrysanthemum morifolium*].** *Zhongguo Zhong Yao Za Zhi.* 2014; 39(24): 4763–8.
[PubMed Abstract](#)
27. Andam CP, Doroghazi JR, Campbell AN, *et al.*: **A Latitudinal Diversity Gradient in Terrestrial Bacteria of the Genus *Streptomyces*.** *MBio.* 2016; 7(2): e02200–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Martiny JB: **History Leaves Its Mark on Soil Bacterial Diversity.** *MBio.* 2016; 7(3): pii: e00784-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Kurth F, Mailänder S, Bönn M, *et al.*: ***Streptomyces*-induced resistance against oak powdery mildew involves host plant responses in defense, photosynthesis, and secondary metabolism pathways.** *Mol Plant Microbe Interact.* 2014; 27(9): 891–900.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. Hanshew AS, McDonald BR, Díaz Díaz C, *et al.*: **Characterization of actinobacteria associated with three ant-plant mutualisms.** *Microb Ecol.* 2015; 69(1): 192–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. Kaltenpoth M, Roeser-Mueller K, Koehler S, *et al.*: **Partner choice and fidelity stabilize coevolution in a Cretaceous-age defensive symbiosis.** *Proc Natl Acad Sci U S A.* 2014; 111(17): 6359–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. Kumbhar C, Mudliar P, Bhatia L, *et al.*: **Widespread predatory abilities in the genus *Streptomyces*.** *Arch Microbiol.* 2014; 196(4): 235–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Ian E, Malko DB, Sekurova ON, *et al.*: **Genomics of sponge-associated *Streptomyces* spp. closely related to *Streptomyces albus* J1074: insights into marine adaptation and secondary metabolite biosynthesis potential.** *PLoS One.* 2014; 9(5): e96719.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Huguet-Tapia JC, Bignell DR, Loria R: **Characterization of the integration and modular excision of the integrative conjugative element PAIS1 in *Streptomyces turgidiscabies* Car8.** *PLoS One.* 2014; 9(6): e99345.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Chapeau M, Guertin JF, Farrokhi A, *et al.*: **Identification of genetic and environmental factors stimulating excision from *Streptomyces scabies* chromosome of the toxicogenic region responsible for pathogenicity.** *Mol Plant Pathol.* 2016; 17(4): 501–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Francis IM, Jourdan S, Fanara S, *et al.*: **The cellobiose sensor CebR is the gatekeeper of *Streptomyces scabies* pathogenicity.** *MBio.* 2015; 6(2): e02018.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. Cheng Z, Bown L, Tahlan K, *et al.*: **Regulation of coronafacoyl phytotoxin production by the PAS-LuxR family regulator CfaR in the common scab pathogen *Streptomyces scabies*.** *PLoS One.* 2015; 10(3): e0122450.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Bignell DR, Francis IM, Fyans JK, *et al.*: **Thaxtomin A production and virulence are controlled by several *bid* gene global regulators in *Streptomyces scabies*.** *Mol Plant Microbe Interact.* 2014; 27(8): 875–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
39. Padilla-Reynaud R, Simao-Beaunoir AM, Lerat S, *et al.*: **Suberin Regulates the Production of Cellulolytic Enzymes in *Streptomyces scabiei*, the Causal Agent of Potato Common Scab.** *Microbes Environ.* 2015; 30(3): 245–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Huguet-Tapia JC, Lefebvre T, Badger JH, *et al.*: **Genome Content and Phylogenomics Reveal both Ancestral and Lateral Evolutionary Pathways in Plant-Pathogenic *Streptomyces* Species.** *Appl Environ Microbiol.* 2016; 82(7): 2146–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Studholme DJ: **Genome Update. Let the consumer beware: *Streptomyces***

- genome sequence quality. *Microb Biotechnol.* 2016; 9(1): 3–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Kim J, Kim Y, Jeong Y, *et al.*: Comparative Genomics Reveals the Core and Accessory Genomes of *Streptomyces* Species. *J Microbiol Biotechnol.* 2015; 25(10): 1599–605.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Doroghazi JR, Buckley DH: Widespread homologous recombination within and between *Streptomyces* species. *ISME J.* 2010; 4(9): 1136–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. **F** Cheng K, Rong X, Huang Y: Widespread interspecies homologous recombination reveals reticulate evolution within the genus *Streptomyces*. *Mol Phylogenet Evol.* 2016; 102: 246–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 45. Thoma L, Muth G: The conjugative DNA-transfer apparatus of *Streptomyces*. *Int J Med Microbiol.* 2015; 305(2): 224–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. **F** Thoma L, Vollmer B, Muth G: Fluorescence microscopy of *Streptomyces* conjugation suggests DNA-transfer at the lateral walls and reveals the spreading of the plasmid in the recipient mycelium. *Environ Microbiol.* 2016; 18(2): 598–608.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 47. **F** Seipke RF: Strain-level diversity of secondary metabolism in *Streptomyces albus*. *PLoS One.* 2015; 10(1): e0116457.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 48. Sexton DL, St-Onge RJ, Haiser HJ, *et al.*: Resuscitation-promoting factors are cell wall-lytic enzymes with important roles in the germination and growth of *Streptomyces coelicolor*. *J Bacteriol.* 2015; 197(5): 848–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. **F** St-Onge RJ, Haiser HJ, Yousef MR, *et al.*: Nucleotide second messenger-mediated regulation of a muralytic enzyme in *Streptomyces*. *Mol Microbiol.* 2015; 96(4): 779–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 50. Flårdh K, Richards DM, Hempel AM, *et al.*: Regulation of apical growth and hyphal branching in *Streptomyces*. *Curr Opin Microbiol.* 2012; 15(6): 737–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Chandra G, Chater KF: Developmental biology of *Streptomyces* from the perspective of 100 actinobacterial genome sequences. *FEMS Microbiol Rev.* 2014; 38(3): 345–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. **F** Chaplin AK, Petrus ML, Mangiameli G, *et al.*: GlxA is a new structural member of the radical copper oxidase family and is required for glycan deposition at hyphal tips and morphogenesis of *Streptomyces lividans*. *Biochem J.* 2015; 469(3): 433–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 53. Liman R, Facey PD, van Keulen G, *et al.*: A laterally acquired galactose oxidase-like gene is required for aerial development during osmotic stress in *Streptomyces coelicolor*. *PLoS One.* 2013; 8(1): e54112.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. van Dissel D, Claessen D, Roth M, *et al.*: A novel locus for mycelial aggregation forms a gateway to improved *Streptomyces* cell factories. *Microb Cell Fact.* 2015; 14: 44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. Zacchetti B, Willemsse J, Recter B, *et al.*: Aggregation of germlings is a major contributing factor towards mycelial heterogeneity of *Streptomyces*. *Sci Rep.* 2016; 6: 27045.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. Yagüe P, Lopez-Garcia MT, Rioseras B, *et al.*: New insights on the development of *Streptomyces* and their relationships with secondary metabolite production. *Curr Trends Microbiol.* 2012; 8: 65–73.
[PubMed Abstract](#) | [Free Full Text](#)
 57. **F** Celler K, Koning RI, Willemsse J, *et al.*: Cross-membranes orchestrate compartmentalization and morphogenesis in *Streptomyces*. *Nat Commun.* 2016; 7: ncomms11836.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 58. **F** Yagüe P, Willemsse J, Koning RI, *et al.*: Subcompartmentalization by cross-membranes during early growth of *Streptomyces* hyphae. *Nat Commun.* 2016; 7: 12467.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 59. **F** Mingyar E, Sevcikova B, Rezuchova B, *et al.*: The α -specific anti-sigma factor RsfA is one of the protein kinases that phosphorylates the pleiotropic anti-anti-sigma factor BldG in *Streptomyces coelicolor* A3(2). *Gene.* 2014; 538(2): 280–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 60. Hindra, Moody MJ, Jones SE, *et al.*: Complex intra-operonic dynamics mediated by a small RNA in *Streptomyces coelicolor*. *PLoS One.* 2014; 9(1): e85856.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Maupin-Furlow JA: Prokaryotic ubiquitin-like protein modification. *Annu Rev Microbiol.* 2014; 68: 155–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Boubakri H, Seghezzi N, Duchateau M, *et al.*: The Absence of Pupylation (Prokaryotic Ubiquitin-Like Protein Modification) Affects Morphological and Physiological Differentiation in *Streptomyces coelicolor*. *J Bacteriol.* 2015; 197(21): 3388–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. Compton CL, Fernandopulle MS, Nagari RT, *et al.*: Genetic and Proteomic Analyses of Pupylation in *Streptomyces coelicolor*. *J Bacteriol.* 2015; 197(17): 2747–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Mao XM, Ren NN, Sun N, *et al.*: Proteasome involvement in a complex cascade mediating SigT degradation during differentiation of *Streptomyces coelicolor*. *FEBS Lett.* 2014; 588(4): 608–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. Niu G, Chater KF, Tian Y, *et al.*: Specialised metabolites regulating antibiotic biosynthesis in *Streptomyces* spp. *FEMS Microbiol Rev.* 2016; 40(4): 554–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. **F** Zou Z, Du D, Zhang Y, *et al.*: A γ -butyrolactone-sensing activator/repressor, JadR3, controls a regulatory mini-network for jadomycin biosynthesis. *Mol Microbiol.* 2014; 94(3): 490–505.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 67. Nodwell JR: Are you talking to me? A possible role for γ -butyrolactones in interspecies signalling. *Mol Microbiol.* 2014; 94(3): 483–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. Schlimpert S, Flårdh K, Buttner M: Fluorescence Time-lapse Imaging of the Complete *S. venezuelae* Life Cycle Using a Microfluidic Device. *J Vis Exp.* 2016; (108): 53863.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. **F** Donczew M, Mackiewicz P, Wróbel A, *et al.*: ParA and ParB coordinate chromosome segregation with cell elongation and division during *Streptomyces* sporulation. *Open Biol.* 2016; 6(4): 150263.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 70. **F** Al-Bassam MM, Bibb MJ, Bush MJ, *et al.*: Response regulator heterodimer formation controls a key stage in *Streptomyces* development. *PLoS Genet.* 2014; 10(8): e1004554.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 71. **F** Bush MJ, Chandra G, Bibb MJ, *et al.*: Genome-Wide Chromatin Immunoprecipitation Sequencing Analysis Shows that WhiB Is a Transcription Factor That Cocontrols Its Regulon with WhiA To Initiate Developmental Cell Division in *Streptomyces*. *MBio.* 2016; 7(2): e00523–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 72. Bush MJ, Tschowri N, Schlimpert S, *et al.*: c-di-GMP signalling and the regulation of developmental transitions in streptomycetes. *Nat Rev Microbiol.* 2015; 13(12): 749–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 73. Tschowri N, Schumacher MA, Schlimpert S, *et al.*: Tetrameric c-di-GMP mediates effective transcription factor dimerization to control *Streptomyces* development. *Cell.* 2014; 158(5): 1136–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Tran NT, Den Hengst CD, Gomez-Escribano JP, *et al.*: Identification and characterization of CdgB, a diguanylate cyclase involved in developmental processes in *Streptomyces coelicolor*. *J Bacteriol.* 2011; 193(12): 3100–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 75. Tschowri N: Cyclic Dinucleotide-Controlled Regulatory Pathways in *Streptomyces* Species. *J Bacteriol.* 2016; 198(1): 47–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 76. Kelemen GH, Brown GL, Kormanec J, *et al.*: The positions of the sigma-factor genes, *whiG* and *sigF*, in the hierarchy controlling the development of spore chains in the aerial hyphae of *Streptomyces coelicolor* A3(2). *Mol Microbiol.* 1996; 21(3): 593–603.
[PubMed Abstract](#) | [Publisher Full Text](#)
 77. Zhang G, Tian Y, Hu K, *et al.*: Importance and regulation of inositol biosynthesis during growth and differentiation of *Streptomyces*. *Mol Microbiol.* 2012; 83(6): 1178–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Chater KF, Horinouchi S: Signalling early developmental events in two highly diverged *Streptomyces* species. *Mol Microbiol.* 2003; 48(1): 9–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. **F** Liu S, Yu P, Yuan P, *et al.*: Sigma factor WhiG_{ch} positively regulates natamycin production in *Streptomyces chattanoogensis* L10. *Appl Microbiol Biotechnol.* 2015; 99(6): 2715–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 80. Green J, Rolfe MD, Smith LJ: Transcriptional regulation of bacterial virulence gene expression by molecular oxygen and nitric oxide. *Virulence.* 2014; 5(8): 794–809.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 81. Crack JC, Green J, Thomson AJ, *et al.*: Iron-sulfur clusters as biological sensors: the chemistry of reactions with molecular oxygen and nitric oxide. *Acc Chem Res.* 2014; 47(10): 3196–205.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Chen Z, Hu Y, Cumming BM, *et al.*: Mycobacterial WhiB6 Differentially Regulates ESX-1 and the Dos Regulon to Modulate Granuloma Formation and Virulence in Zebrafish. *Cell Rep.* 2016; 16(9): 2512–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Serrano PN, Wang H, Crack JC, *et al.*: Nitrosylation of Nitric-Oxide-Sensing

- Regulatory Proteins Containing [4Fe-4S] Clusters Gives Rise to Multiple Iron-Nitrosyl Complexes.** *Angew Chem Int Ed Engl.* 2016; 55(47): 14575–14579.
PubMed Abstract | Publisher Full Text
84. **F** Sasaki Y, Oguchi H, Kobayashi T, *et al.*: Nitrogen oxide cycle regulates nitric oxide levels and bacterial cell signaling. *Sci Rep.* 2016; 6: 22038.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 85. **F** Cunningham-Bussell A, Zhang T, Nathan CF: Nitrite produced by *Mycobacterium tuberculosis* in human macrophages in physiologic oxygen impacts bacterial ATP consumption and gene expression. *Proc Natl Acad Sci U S A.* 2013; 110(45): E4256–65.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 86. Cunningham-Bussell A, Bange FC, Nathan CF: Nitrite impacts the survival of *Mycobacterium tuberculosis* in response to isoniazid and hydrogen peroxide. *Microbiologyopen.* 2013; 2(6): 901–11.
PubMed Abstract | Publisher Full Text | Free Full Text
 87. Fischer M, Falke D, Pawlik T, *et al.*: Oxygen-dependent control of respiratory nitrate reduction in the mycelium of *Streptomyces coelicolor* A3(2). *J Bacteriol.* 2014; 196(23): 4152–62.
PubMed Abstract | Publisher Full Text | Free Full Text
 88. Lundberg JO, Weitzberg E, Gladwin MT: The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov.* 2008; 7(2): 156–67.
PubMed Abstract | Publisher Full Text
 89. Vasudevan D, Bovee RC, Thomas DD: Nitric oxide, the new architect of epigenetic landscapes. *Nitric Oxide.* 2016; 59: 54–62.
PubMed Abstract | Publisher Full Text
 90. Crack JC, Munnoch J, Dodd EL, *et al.*: NsrR from *Streptomyces coelicolor* is a nitric oxide-sensing [4Fe-4S] cluster protein with a specialized regulatory function. *J Biol Chem.* 2015; 290(20): 12689–704.
PubMed Abstract | Publisher Full Text | Free Full Text
 91. **F** Crack JC, Svistunenko DA, Munnoch J, *et al.*: Differentiated, Promoter-specific Response of [4Fe-4S] NsrR DNA Binding to Reaction with Nitric Oxide. *J Biol Chem.* 2016; 291(16): 8663–72.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 92. Wang W, Shu D, Chen L, *et al.*: Cross-talk between an orphan response regulator and a noncognate histidine kinase in *Streptomyces coelicolor*. *FEMS Microbiol Lett.* 2009; 294(2): 150–6.
PubMed Abstract | Publisher Full Text
 93. **F** Zhang L, Willemse J, Claessen D, *et al.*: SepG coordinates sporulation-specific cell division and nucleoid organization in *Streptomyces coelicolor*. *Open Biol.* 2016; 6(4): 150164.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 94. **F** Ladwig N, Franz-Wachtel M, Hezel F, *et al.*: Control of Morphological Differentiation of *Streptomyces coelicolor* A3(2) by Phosphorylation of MreC and PBP2. *PLoS One.* 2015; 10(4): e0125425.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 95. Molle V, Palframan WJ, Findlay KC, *et al.*: WhiD and WhiB, homologous proteins required for different stages of sporulation in *Streptomyces coelicolor* A3(2). *J Bacteriol.* 2000; 182(5): 1286–95.
PubMed Abstract | Publisher Full Text | Free Full Text
 96. **F** Sandoval-Calderón M, Nguyen DD, Kapono CA, *et al.*: Plasticity of *Streptomyces coelicolor* Membrane Composition Under Different Growth Conditions and During Development. *Front Microbiol.* 2015; 6: 1465.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 97. Chandra G, Chater KF, Bornemann S: Unexpected and widespread connections between bacterial glycogen and trehalose metabolism. *Microbiology.* 2011; 157(Pt 6): 1565–72.
PubMed Abstract | Publisher Full Text
 98. Miah F, Bibb MJ, Barclay JE, *et al.*: Developmental delay in a *Streptomyces venezuelae* *glgE* null mutant is associated with the accumulation of α -maltose 1-phosphate. *Microbiology.* 2016; 162(7): 1208–19.
PubMed Abstract | Publisher Full Text | Free Full Text
 99. **F** Kim JN, Jeong Y, Yoo JS, *et al.*: Genome-scale analysis reveals a role for NdgR in the thiol oxidative stress response in *Streptomyces coelicolor*. *BMC Genomics.* 2015; 16(1): 116.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 100. Liu G, Chater KF, Chandra G, *et al.*: Molecular regulation of antibiotic biosynthesis in *Streptomyces*. *Microbiol Mol Biol Rev.* 2013; 77(1): 112–43.
PubMed Abstract | Publisher Full Text | Free Full Text
 101. Xu D, Waack P, Zhang Q, *et al.*: Structure and regulatory targets of SCO3201, a highly promiscuous TetR-like regulator of *Streptomyces coelicolor* M145. *Biochem Biophys Res Commun.* 2014; 450(1): 513–8.
PubMed Abstract | Publisher Full Text
 102. Rodríguez H, Rico S, Yepes A, *et al.*: The two kinases, AbrC1 and AbrC2, of the atypical two-component system AbrC are needed to regulate antibiotic production and differentiation in *Streptomyces coelicolor*. *Front Microbiol.* 2015; 6: 450.
PubMed Abstract | Publisher Full Text | Free Full Text
 103. Rico S, Santamaría RI, Yepes A, *et al.*: Deciphering the regulon of *Streptomyces coelicolor* AbrC3, a positive response regulator of antibiotic production. *Appl Environ Microbiol.* 2014; 80(8): 2417–28.
PubMed Abstract | Publisher Full Text | Free Full Text
 104. **F** Bekiesch P, Franz-Wachtel M, Kulik A, *et al.*: DNA affinity capturing identifies new regulators of the heterologously expressed novobiocin gene cluster in *Streptomyces coelicolor* M512. *Appl Microbiol Biotechnol.* 2016; 100(10): 4495–509.
PubMed Abstract | Publisher Full Text | F1000 Recommendation
 105. Huang J, Shi J, Molle V, *et al.*: Cross-regulation among disparate antibiotic biosynthetic pathways of *Streptomyces coelicolor*. *Mol Microbiol.* 2005; 58(5): 1276–87.
PubMed Abstract | Publisher Full Text
 106. **F** Vicente CM, Payero TD, Santos-Aberturas J, *et al.*: Pathway-specific regulation revisited: cross-regulation of multiple disparate gene clusters by PAS-LuxR transcriptional regulators. *Appl Microbiol Biotechnol.* 2015; 99(12): 5123–35.
PubMed Abstract | Publisher Full Text | F1000 Recommendation
 107. **F** Li X, Wang J, Li S, *et al.*: ScbR- and ScbR2-mediated signal transduction networks coordinate complex physiological responses in *Streptomyces coelicolor*. *Sci Rep.* 2015; 5: 14831.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 108. **F** Li X, Yu T, He Q, *et al.*: Binding of a biosynthetic intermediate to AtrA modulates the production of lidamycin by *Streptomyces globisporus*. *Mol Microbiol.* 2015; 96(6): 1257–71.
PubMed Abstract | Publisher Full Text | F1000 Recommendation
 109. **F** Li Y, Li J, Tian Z, *et al.*: Coordinative Modulation of Chlorothricin Biosynthesis by Binding of the Glycosylated Intermediates and End Product to a Responsive Regulator ChlF1. *J Biol Chem.* 2016; 291(10): 5406–17.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 110. **F** Wolański M, Lebowski T, Kois-Ostrowska A, *et al.*: Two transcription factors, CabA and CabR, are independently involved in multilevel regulation of the biosynthetic gene cluster encoding the novel aminocoumarin, cacibiocin. *Appl Microbiol Biotechnol.* 2016; 100(7): 3147–64.
PubMed Abstract | Publisher Full Text | F1000 Recommendation
 111. **F** Xu G, Wang J, Wang L, *et al.*: “Pseudo” gamma-butyrolactone receptors respond to antibiotic signals to coordinate antibiotic biosynthesis. *J Biol Chem.* 2010; 285(35): 27440–8.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 112. Sekurova ON, Zhang J, Kristiansen KA, *et al.*: Activation of chloramphenicol biosynthesis in *Streptomyces venezuelae* ATCC 10712 by ethanol shock: insights from the promoter fusion studies. *Microb Cell Fact.* 2016; 15: 85.
PubMed Abstract | Publisher Full Text | Free Full Text
 113. **F** Robertson AW, Forget SM, Martinez-Farina CF, *et al.*: Jadx is a Disparate Natural Product Binding Protein. *J Am Chem Soc.* 2016; 138(7): 2200–8.
PubMed Abstract | Publisher Full Text | F1000 Recommendation
 114. **F** Wang W, Ji J, Li X, *et al.*: Angucyclines as signals modulate the behaviors of *Streptomyces coelicolor*. *Proc Natl Acad Sci U S A.* 2014; 111(15): 5688–93.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 115. **F** Abrudan MI, Smakman F, Grimbergen AJ, *et al.*: Socially mediated induction and suppression of antibiosis during bacterial coexistence. *Proc Natl Acad Sci U S A.* 2015; 112(35): 11054–9.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 116. Cornforth DM, Foster KR: Antibiotics and the art of bacterial war. *Proc Natl Acad Sci U S A.* 2015; 112(35): 10827–8.
PubMed Abstract | Publisher Full Text | Free Full Text
 117. **F** Imai Y, Sato S, Tanaka Y, *et al.*: Lincomycin at Subinhibitory Concentrations Potentiates Secondary Metabolite Production by *Streptomyces* spp. *Appl Environ Microbiol.* 2015; 81(11): 3869–79.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 118. Bertrand S, Bohni N, Schnee S, *et al.*: Metabolite induction via microorganism co-culture: a potential way to enhance chemical diversity for drug discovery. *Biotechnol Adv.* 2014; 32(6): 1180–204.
PubMed Abstract | Publisher Full Text
 119. Schäberle TF, Orland A, König GM: Enhanced production of undecylprodigiosin in *Streptomyces coelicolor* by co-cultivation with the coralalpyronin A-producing myxobacterium, *Corallocozoc coralloides*. *Biotechnol Lett.* 2014; 36(3): 641–8.
PubMed Abstract | Publisher Full Text
 120. **F** Vargas-Bautista C, Rahlwes K, Straight P: Bacterial competition reveals differential regulation of the *pkS* genes by *Bacillus subtilis*. *J Bacteriol.* 2014; 196(4): 717–28.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation
 121. Traxler MF, Kolter R: Natural products in soil microbe interactions and evolution. *Nat Prod Rep.* 2015; 32(7): 956–70.
PubMed Abstract | Publisher Full Text
 122. **F** Jeong Y, Kim JN, Kim MW, *et al.*: The dynamic transcriptional and translational landscape of the model antibiotic producer *Streptomyces coelicolor* A3(2). *Nat Commun.* 2016; 7: 11605.
PubMed Abstract | Publisher Full Text | Free Full Text | F1000 Recommendation

123. Chater KF, Chandra G: **The use of the rare UUA codon to define “expression space” for genes involved in secondary metabolism, development and environmental adaptation in *Streptomyces*.** *J Microbiol.* 2008; **46**(1): 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
124. Chandra G, Chater KF: **Evolutionary flux of potentially *bldA*-dependent *Streptomyces* genes containing the rare leucine codon TTA.** *Antonie Van Leeuwenhoek.* 2008; **94**(1): 111–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
125. Jensen SE: **Biosynthesis of clavam metabolites.** *J Ind Microbiol Biotechnol.* 2012; **39**(10): 1407–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
126. Ferguson NL, Peña-Castillo L, Moore MA, *et al.*: **Proteomics analysis of global regulatory cascades involved in clavulanic acid production and morphological development in *Streptomyces clavuligerus*.** *J Ind Microbiol Biotechnol.* 2016; **43**(4): 537–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
127. Zeng H, Wen S, Xu W, *et al.*: **Highly efficient editing of the actinorhodin polyketide chain length factor gene in *Streptomyces coelicolor* M145 using CRISPR/Cas9-CodA(sm) combined system.** *Appl Microbiol Biotechnol.* 2015; **99**(24): 10575–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
128. Liu Y, Tao W, Wen S, *et al.*: ***In Vitro* CRISPR/Cas9 System for Efficient Targeted DNA Editing.** *MBio.* 2015; **6**(6): e01714–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
129. Tong Y, Charusanti P, Zhang L, *et al.*: **CRISPR-Cas9 Based Engineering of Actinomycetal Genomes.** *ACS Synth Biol.* 2015; **4**(9): 1020–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
130. Huang H, Zheng G, Jiang W, *et al.*: **One-step high-efficiency CRISPR/Cas9-mediated genome editing in *Streptomyces*.** *Acta Biochim Biophys Sin (Shanghai).* 2015; **47**(4): 231–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
131. Qiu Y, Wang S, Chen Z, *et al.*: **An Active Type I-E CRISPR-Cas System Identified in *Streptomyces avermitilis*.** *PLoS One.* 2016; **11**(2): e0149533.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
132. Du D, Wang L, Tian Y, *et al.*: **Genome engineering and direct cloning of antibiotic gene clusters via phage ϕ BT1 integrase-mediated site-specific recombination in *Streptomyces*.** *Sci Rep.* 2015; **5**: 8740.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
133. Antoraz S, Santamaria RI, Díaz M, *et al.*: **Toward a new focus in antibiotic and drug discovery from the *Streptomyces arsenal*.** *Front Microbiol.* 2015; **6**: 461.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
134. Baltz RH: **Genetic manipulation of secondary metabolite biosynthesis for improved production in *Streptomyces* and other actinomycetes.** *J Ind Microbiol Biotechnol.* 2016; **43**(2–3): 343–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
135. Breitling R, Takano E: **Synthetic Biology of Natural Products.** *Cold Spring Harb Perspect Biol.* 2016; **8**(10): pii: a023994.
[PubMed Abstract](#) | [Publisher Full Text](#)
136. Weber T, Blin K, Duddela S, *et al.*: **antiSMASH 3.0-a comprehensive resource for the genome mining of biosynthetic gene clusters.** *Nucleic Acids Res.* 2015; **43**(W1): W237–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
137. Luo Y, Zhang L, Barton KW, *et al.*: **Systematic Identification of a Panel of Strong Constitutive Promoters from *Streptomyces albus*.** *ACS Synth Biol.* 2015; **4**(9): 1001–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
138. Li S, Wang W, Li X, *et al.*: **Genome-wide identification and characterization of reference genes with different transcript abundances for *Streptomyces coelicolor*.** *Sci Rep.* 2015; **5**: 15840.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
139. Willemse J, Borst JW, de Waal E, *et al.*: **Positive control of cell division: FtsZ is recruited by SsgB during sporulation of *Streptomyces*.** *Genes Dev.* 2011; **25**(1): 89–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Wolfgang Wohlleben**
Interfaculty Institute of Microbiology and Infection Medicine, University of Tuebingen, Tuebingen, Germany
Competing Interests: No competing interests were disclosed.
2. **Paul J Dyson**
Institute of Life Science, Swansea University Medical School, Swansea, UK
Competing Interests: No competing interests were disclosed.
3. **Klas Flardh**
Department of Biology, University of Lund, Lund, Sweden
Competing Interests: No competing interests were disclosed.
4. **Rosemary Loria**
Department of Plant Pathology, University of Florida, Gainesville, FL, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research