Check for updates

REVIEW

Ahead of the Curve: New Insights into Microtubule Dynamics

[version 1; peer review: 2 approved]Ryoma Ohi¹, Marija Zanic^{1,2}¹Department of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville, Tennessee, USA²Department of Chemical and Biomolecular Engineering, Vanderbilt University, Nashville, Tennessee, USA**V1** First published: 10 Mar 2016, 5(F1000 Faculty Rev):314
<https://doi.org/10.12688/f1000research.7439.1>Latest published: 10 Mar 2016, 5(F1000 Faculty Rev):314
<https://doi.org/10.12688/f1000research.7439.1>**Abstract**

Microtubule dynamics are fundamental for many aspects of cell physiology, but their mechanistic underpinnings remain unclear despite 40 years of intense research. In recent years, the continued union of reconstitution biochemistry, structural biology, and modeling has yielded important discoveries that deepen our understanding of microtubule dynamics. These studies, which we review here, underscore the importance of GTP hydrolysis-induced changes in tubulin structure as microtubules assemble, and highlight the fact that each aspect of microtubule behavior is the output of complex, multi-step processes. Although this body of work moves us closer to appreciating the key features of microtubule biochemistry that drive dynamic instability, the divide between our understanding of microtubules in isolation versus within the cellular milieu remains vast. Bridging this gap will serve as fertile grounds of cytoskeleton-focused research for many years to come.

Keywords

Microtubule dynamics, microtubule biochemistry, cytoskeleton

Open Peer Review**Approval Status**

	1	2
version 1
10 Mar 2016		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **William Hancock**, Pennsylvania State University, University Park, USA
2. **Chad Pearson**, University of Colorado School of Medicine, Aurora, USA
Domenico Gelati, University of Colorado School of Medicine, Aurora, USA
Marisa Ruehle, University of Colorado School of Medicine, Aurora, USA

Any comments on the article can be found at the end of the article.

Corresponding authors: Ryoma Ohi (ryoma.ohi@vanderbilt.edu), Marija Zanic (marija.zanic@vanderbilt.edu)

Competing interests: The authors declare that they have no competing interests.

Grant information: Work in the laboratory of RO is supported by a grant from the National Institutes of Health (GM086610) and a Scholar Career Development Award from the Leukemia and Lymphoma Society. MZ is supported by a Career Development Award from the Human Frontier Science Program.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Ohi R and Zanic M. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ohi R and Zanic M. **Ahead of the Curve: New Insights into Microtubule Dynamics [version 1; peer review: 2 approved]** F1000Research 2016, 5(F1000 Faculty Rev):314 <https://doi.org/10.12688/f1000research.7439.1>

First published: 10 Mar 2016, 5(F1000 Faculty Rev):314 <https://doi.org/10.12688/f1000research.7439.1>

Introduction

Microtubules are hollow, cylindrical polymers of $\alpha\beta$ -tubulin that are vital for many biological processes, including cell division, polarization, and migration. In cells, microtubules are composed of 13 laterally associated protofilaments, strands of $\alpha\beta$ -tubulin subunits lined up in a head-to-tail fashion. Key to their functional versatility is that microtubules are capable of assembling and disassembling over many cycles, as first revealed by Shinya Inoue's seminal polarization microscopy studies of mitosis¹. The dynamic nature of microtubules is intrinsic to tubulin, as microtubules formed *in vitro* with tubulin alone coexist as growing and shortening polymers that switch between these states². This non-equilibrium behavior, termed dynamic instability, is in turn dependent on the ability of β -tubulin to bind and hydrolyze GTP: an $\alpha\beta$ -tubulin heterodimer that contains GTP- β -tubulin can add on to the growing end of a microtubule, but polymerization induces GTP hydrolysis. GDP- β -tubulin is restricted to the lattice, as ends rich in GDP- β -tubulin are unstable and prone to depolymerization. These observations laid the foundation for the "GTP cap" model, which postulates that microtubules can grow only when non-hydrolyzed GTP- β -tubulin subunits crown the end of a microtubule^{3,4}. Although GTP hydrolysis plays an indisputable role in fueling dynamic instability, structural aspects of microtubule subunit interactions—both longitudinally and laterally—ultimately lie at the heart of microtubule dynamics.

Whereas the general principles of dynamic instability are well established, individual aspects of a microtubule's life—nucleation, growth, and the growth-to-shortening transition (catastrophe)—are complex and represent the output of poorly understood multi-step processes. In this commentary, we review recent progress in the field, focusing on nucleation and microtubule assembly, where significant advances have been made. This progress reflects our improved understanding of key microtubule-associated proteins (MAPs), development of *in vitro* assays that probe novel aspects of microtubule assembly and disassembly, and technological breakthroughs that have increased the resolving power of cryo-electron microscopy (cryo-EM)-based structural approaches.

Nucleation

For a microtubule to form, it must first be *nucleated*. What does this mean in the context of microtubules? A nucleus is a multimer that forms through sequential subunit addition and allows growth to be thermodynamically favorable⁵. Nuclei can assemble in the absence of cellular factors and have been observed by EM to be two-dimensional sheets^{6,7} or short oligomers⁸, which presumably grow until tube closure is possible. Such spontaneous nucleation is a slow and energetically unfavorable process, involving a considerable lag phase. Importantly, cells bypass the kinetic barrier to nucleation by using factors that accelerate microtubule formation.

The best-understood microtubule nucleation factor is a protein complex that contains γ -tubulin, a protein closely related to α - and β -tubulin. The γ -tubulin ring complex (γ -TuRC) specifically templates 13-protofilament microtubules⁹ and participates in microtubule nucleation at microtubule-organizing centers such as the centrosome, as well as in the chromosome-mediated^{10–12}, Golgi-mediated^{13,14}, and microtubule-dependent^{15–17} microtubule

nucleation pathways. The γ -TuRC fulfills the expected function of a microtubule nucleating factor; that is, microtubules form more rapidly in its presence¹⁸, and it is easy to imagine how it does so: the γ -TuRC mimics an early assembly intermediate of $\alpha\beta$ -tubulin¹⁹. However, studies of the chromosome-mediated microtubule nucleation pathway have demonstrated that two additional MAPs—TPX2²⁰ and XMAP215²¹—also play key roles in facilitating microtubule nucleation^{22,23}. In fact, these proteins can nucleate microtubules independently of the γ -TuRC^{23,24} when added to concentrations of tubulin at which polymer does not form. As MAPs, the γ -TuRC, TPX2, and XMAP215 have distinct biochemical properties. XMAP215, through the concerted activities of an array of tubulin-binding TOG domains, processively catalyzes microtubule assembly²⁵, increasing microtubule growth rates up to 10-fold²⁶. This activity depends on the ability of XMAP215 to bind tubulin subunits while retaining a grip on the microtubule plus end²⁷. TPX2, on the other hand, has not been reported to bind tubulin subunits and therefore seems to be a conventional MAP that stabilizes and crosslinks microtubules²⁴. On a molecular level, these findings suggest that multiple non-redundant activities can be integrated during microtubule nucleation, providing support that nucleation is a multi-step process. Interestingly, TPX2 and XMAP215 interact with proteins that co-locate their activities with the γ -TuRC at the centrosome. TPX2 binds RHAMM²⁸, which is present in a microtubule nucleation complex that contains γ -TuRC, and NEDD1, a centrosome-targeting factor²⁹. XMAP215 family proteins are targeted to the centrosome through direct interaction with a group of coiled-coil proteins called TACCs³⁰. Therefore, it is likely that TPX2 and XMAP215 family proteins synergize with γ -TuRC in the context of the cell, but how they do so remains unclear.

At a molecular level, recent studies have shed light on how TPX2 and XMAP215 promote nucleation^{31,32}. Previous *in vitro* work showed that TPX2 induces the formation of disordered tubulin aggregates, which were speculated to be small oligomers capable of elongation²⁴. This finding has recently been recapitulated by the Surrey laboratory by using total internal reflection microscopy-based assays; TPX2 induces the formation of granular tubulin foci ("stubs"), which elongate in solution³¹. In addition, Roostalu *et al.* show that chTOG, the human ortholog of XMAP215, only mildly promotes microtubule nucleation³¹. Interestingly, this result differs from conclusions reached through previous work, where XMAP215 alone was sufficient for robust microtubule formation²³. However, it is worth noting that the ability of XMAP215 to nucleate microtubules in the previous study was dependent on its conjugation to beads, an experimental setup that locally concentrates the protein. Strikingly, Roostalu *et al.* find that a combination of chTOG and TPX2 produces more microtubule polymer, leading to the idea that chTOG and TPX2 perform distinct functions during the nucleation process. The authors speculate that TPX2 promotes nucleation by stabilizing early oligomeric intermediates, whereas chTOG acts by accelerating subunit addition to nuclei³¹. chTOG activity may be crucial for oligomers to form a sheet large enough to fold into a tube. In this view, the γ -TuRC may simply act to ensure that microtubules are built using 13 protofilaments, as microtubules nucleated by TPX2 and chTOG are likely of mixed protofilament composition.

Whereas nucleation *de novo* can easily be imagined to be a multi-step process, microtubule assembly on pre-existing templates such as axonemes is also complex, as initially demonstrated by Walker *et al.*³³. This problem was revisited in a recent article from the Brouhard laboratory³². Wicczorek *et al.* found that a lag phase always precedes microtubule assembly, regardless of whether the nucleating source is a centrosome, axoneme, or a pre-formed microtubule end³². This lag phase is attenuated by TPX2 and XMAP215, or extended by catastrophe factors such as MCAK and EB1. Although the Roostalu *et al.* and Wicczorek *et al.* studies demonstrate a role for TPX2 and XMAP215 in microtubule nucleation, it is important to note that the two proteins may act differently during templated nucleation versus microtubule formation *de novo*. Both reports show that TPX2 is a strong anti-catastrophe factor that slows depolymerization^{31,32}. Therefore, it is possible that TPX2 acts as a traditional MAP during templated nucleation, simply increasing a filament's probability to extend. Mechanism aside, the picture we are left with is that the birth of a microtubule is complex, requiring the formation of a plus-end structure that is compatible with subunit addition.

Microtubule assembly and tubulin structure

Once formed from a nucleus, the growing microtubule will continue to *elongate*, sometimes for minutes at a time. Our understanding of the structure of the elongating microtubule end is shaped by early cryo-EM studies^{34,35}. These images show that growing microtubule ends display “sheets” of interconnected protofilaments, which are thought to dynamically close into a tube as the microtubule grows. After more than 20 years, direct visualization of the structure of growing microtubule ends in real time remains an open challenge.

The highest-resolution studies of microtubule growth dynamics to date employed optical trapping methods to observe changes in microtubule length with up to 3.5 nm resolution^{36,37}. In these experiments, microtubules were grown against barriers, such that the observed length fluctuations represent the length changes of the longest individual protofilament or group of protofilaments, and do not uncover the structure of the end. Nevertheless, these studies showed that microtubule growth is irregular with frequent shortening excursions that can retract the longest protofilaments more than 40 nm (corresponding to the length of five tubulin dimer subunits) while a microtubule remains in the overall growth phase.

Large fluctuations in polymer length observed during microtubule growth can be understood as a consequence of a very unproductive growth process³⁸. Indeed, further studies by Gardner *et al.*³⁹ found that the vast majority of tubulin subunits that associate with the growing microtubule end rapidly dissociate. The underlying cause of this high tubulin off-rate is unclear; efficient subunit incorporation into the microtubule lattice might require an additional step (for example, a structural alteration that would promote formation of stabilizing lateral bonds). The exact structure of tubulin subunits when bound to different nucleotides, both in solution as well as within the microtubule polymer, has been somewhat controversial. Whereas earlier studies proposed that GTP-tubulin is straight^{40–42}, allowing it to readily incorporate into the microtubule lattice, more recent studies suggest that GTP-tubulin dimers are curved, similar to their hydrolyzed, GDP-bound counterparts^{43–45}. Additionally,

detailed structural changes that accompany GTP hydrolysis once a subunit is incorporated in the microtubule polymer have, until recently, been unknown.

A recent study by Alushin *et al.*⁴⁶ used high-resolution cryo-EM, combined with computational modeling, to investigate the effect of GTP-hydrolysis on the structure of tubulin dimers within the microtubule lattice. With a 5 Å resolution, the authors report that GDP-bound tubulin dimers undergo longitudinal compaction close to the exchangeable nucleotide site with tubulin dimers within microtubules grown with a slowly hydrolyzable GTP-analog GMPCPP. In contrast to a previous lower-resolution EM study, performed with a different nucleotide analog (GTPyS)⁴⁷, Alushin *et al.* found no evidence for changes in lateral interactions between the tubulin dimers. Rather, the authors hypothesize that the observed structural rearrangements in the intermediate domain and the H7 helix of α -tubulin increase lattice strain, which ultimately results in microtubule lattice destabilization. A new study by Geyer *et al.*⁴⁸ supports the idea that structural changes associated with GTP-hydrolysis underlie microtubule instability. Here, the authors studied the polymerization dynamics of purified yeast tubulin with a mutation in helix H7 of β -tubulin (T238A), which is expected to block H7 movement upon nucleotide hydrolysis. Although the GTPase activity of these microtubules was unaffected, the mutation indeed appeared to prevent structural changes that accompany hydrolysis. Interestingly, microtubules assembled from T238A tubulin are hyperstable, suggesting that allosteric effects of GTP hydrolysis, rather than hydrolysis itself, drive microtubule instability. Future studies with additional tubulin mutants are likely to provide a more detailed link between tubulin structure and microtubule dynamics.

Microtubule-associated proteins recognize and modulate microtubule structure

Structural features at the microtubule plus end also govern the action of MAPs. XMAP215, for example, is thought to promote microtubule assembly by tethering a weakly bound tubulin dimer to the microtubule end until it becomes stably incorporated into the microtubule lattice²⁵. In the absence of soluble tubulin, XMAP215 is thought to convert a tightly bound subunit at the microtubule end into one that is only loosely associated. This can explain why XMAP215 can promote microtubule depolymerization⁴⁹ in addition to assembly. Indeed, recent structural studies with TOG domains of Stu2, the yeast homolog of XMAP215, report that TOG domains preferably bind the curved GTP-like conformation of tubulin⁵⁰. The authors propose that the TOG domain dissociates from the tubulin dimer once it straightens, a structural transition that presumably accompanies its stable incorporation into the microtubule lattice. This “hand-off” mechanism in turn allows the TOG domain to move forward and processively add the next tubulin dimer. Thus, XMAP215 is thought to bind a specific curved conformation of tubulin dimers expected to be found only at the very end of the growing microtubule.

EB proteins comprise another major family of proteins known for their ability to bind growing microtubule ends. *In vitro* studies with nucleotide analogs established that EBs recognize the nucleotide state of tubulin dimers in the microtubule, preferentially

binding to GTP-like tubulin over GDP-tubulin^{51,52}. Interestingly, EBs discriminate between microtubules formed from two GTP mimics—GMPCPP and GTP γ S—favoring the latter⁵¹. In this context, it is noteworthy that EBs form comets that lag behind the distal microtubule tip occupied by XMAP215, both *in vitro* and in cells^{53,54}. It is thus speculated that GTP γ S mimics GDP-Pi-tubulin, a post-GTP hydrolysis state wherein phosphate has not yet dissociated.

Given that XMAP215 and EB proteins bind different features of the growing microtubule end, it is interesting that the two proteins synergize *in vitro* to promote fast microtubule growth, with rates matching those previously observed only inside of cells⁵⁵. This synergy is not realized through direct interaction between XMAP215 and EB1. Rather, it is due to an allosteric interaction involving the microtubule end structure. The authors hypothesized that EBs induce structural changes, such as protofilament straightening, that could in turn promote lateral protofilament interactions and sheet closure. Such EB-induced structural changes at the plus end could increase the polymerase activity of XMAP215 by accelerating subunit “hand-off”⁵⁰.

Previous studies have reported that EB family proteins can affect the structure of the microtubule lattice⁵⁶ as well as modulate the number of microtubule protofilaments^{47,57}. Binding of EBs at the interface of four tubulin dimers⁴⁷ could facilitate such structural effects. The latest evidence that EBs modulate the structure of the tubulin dimers in the microtubule lattice comes from a new study by the Nogales lab⁵⁸. Here, the authors determined the structures of GMPCPP-, GTP γ S-, and GDP-bound microtubules copolymerized with EB3 at an unprecedented resolution of 3.5 Å. The authors found all three structures grown with EB3 to be compacted, similar to GDP lattice in the absence of EB3, suggesting that EB binding induces compaction of the microtubule lattice. Unfortunately, the authors were unable to obtain the structure of the GTP γ S lattice in the absence of EBs, leaving open the question of whether compaction occurs prior to, or after, phosphate release. In either case, induction of lattice compaction via EBs is consistent with the view that EBs promote GTP hydrolysis in the microtubule lattice.

Allosteric interactions between MAPs are not limited to XMAP215 and EB1⁵⁹. A recent study found similar interactions between EB1 and another TOG-domain protein, CLASP⁶⁰. The authors reported that EBs have a lower binding affinity for microtubules that are grown in the presence of CLASP. The exact features encoded in the microtubule by CLASP that are recognized by EBs remain unknown. Interestingly, the CLASP TOG2 domain exhibits a strongly bent conformation, raising the possibility that CLASP is binding highly curved protofilaments at the microtubule end⁶¹. The theme of MAPs recognizing aspects of microtubule curvature is highlighted by other recent studies. Doublecortin, which preferentially binds 13-protofilament microtubules^{62,63}, enriches on curved microtubule segments⁶⁴. TPX2 was found to strongly bind curved microtubule ends³¹. CENP-F associates more strongly with vinblastine-generated tubulin curls compared with straight protofilaments that are found within the microtubule lattice⁶⁵. Kinesin-5 has been found to associate with curved tubulin protofilaments at growing microtubule ends,

where it stimulates microtubule assembly⁶⁶. Lastly, kinesin-13s, the most potent catastrophe factors known, are well appreciated to recognize and stabilize a bent tubulin protofilament conformation^{67,68} observed on depolymerizing microtubule ends^{34,35}.

Catastrophe

Even though the link between GTP hydrolysis, structure, and dynamics might be established, we still do not know what features define the stabilizing cap. The observations that EB end binding is largely lost well before the onset of catastrophe^{47,53,55} might imply that GDP-Pi subunits also confer stability to the growing end, if GTP γ S tubulin dimers are indeed to be viewed as a model of GDP-Pi state. In that context, and given that EBs bind very strongly to GTP γ S microtubule lattice, and much more weakly to the GDP lattice, it is interesting that the only structural difference observed between these two is a small relative rotation of tubulin dimers along a protofilament, resulting in a different lattice twist⁵⁸. Whether it is this twist that ultimately leads to the high off-rate of GDP tubulin remains to be understood.

Whatever the stabilizing cap looks like, its loss is a complex process intimately linked to structural features of the microtubule end. Recently, Gardner *et al.* reported that the probability of undergoing catastrophe grows with microtubule age⁶⁹, a finding that confirms older work performed by Odde *et al.*⁷⁰. Thus, catastrophe cannot be caused by a single-step mechanism unless catalyzed by protein factors such as kinesin-13⁶⁹. The process by which microtubule aging causes catastrophes could involve changes in protofilament numbers and/or structural evolution of the growing microtubule end such as tapering and protofilament curling^{71–74}. In any case, the fate of the microtubule is likely to be encoded in the structure of its end.

Closing statements

The work reviewed here has significantly advanced our understanding of microtubule assembly and disassembly, but many questions remain. The complexity of the microtubule cytoskeleton in cells, difficult to capture in reconstitution-based approaches, can and should provide a useful framework for posing further questions. An interesting discrepancy concerning the relationship between microtubule end structure and dynamics, for example, is highlighted by the observation that all protofilaments are curved at the plus ends of microtubules during mitosis⁷⁵. The implication of this finding is that microtubule growth in cells may be governed by different constraints that permit assembly to occur without a sheet-like intermediate that has been observed *in vitro*. Cell cycle-dependent variations involved in the regulation of microtubule biology are also likely to exist. Recent studies on microtubule nucleation have focused on factors that are principally active during cell division. TPX2, for example, is sequestered in the nucleus during interphase⁷⁶ and requires Ran-GTP to become active during mitosis²⁴. The factors and mechanisms that regulate microtubule nucleation during interphase remain to be elucidated. Lastly, given that most, if not all, aspects of microtubule dynamics involve multi-step processes, an important challenge will be to understand the emergent properties of the network of MAPs that synergistically modulate the kinetics of microtubule assembly and disassembly in ways relevant for cell physiology.

Competing interests

The authors declare that they have no competing interests.

Grant information

Work in the laboratory of RO is supported by a grant from the National Institutes of Health (GM086610) and a Scholar Career Development Award from the Leukemia and Lymphoma Society.

MZ is supported by a Career Development Award from the Human Frontier Science Program.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank William Hancock, Chad Pearson, Gary Brouhard, Elizabeth Lawrence, and Marija Podolski for comments on the manuscript.

References

- Inoué S, Sato H: **Cell motility by labile association of molecules. The nature of mitotic spindle fibers and their role in chromosome movement.** *J Gen Physiol.* 1967; **50**(6): Suppl: 259–292.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mitchison T, Kirschner M: **Dynamic instability of microtubule growth.** *Nature.* 1984; **312**(5991): 237–242.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Carlier MF, Hill TL, Chen Y: **Interference of GTP hydrolysis in the mechanism of microtubule assembly: an experimental study.** *Proc Natl Acad Sci U S A.* 1984; **81**(3): 771–775.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Carlier MF, Pantaloni D: **Kinetic analysis of guanosine 5'-triphosphate hydrolysis associated with tubulin polymerization.** *Biochemistry.* 1981; **20**(7): 1918–1924.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Erickson HP, Pantaloni D: **The role of subunit entropy in cooperative assembly. Nucleation of microtubules and other two-dimensional polymers.** *Biophys J.* 1981; **34**(2): 293–309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Erickson HP: **Microtubule surface lattice and subunit structure and observations on reassembly.** *J Cell Biol.* 1974; **60**(1): 153–167.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kirschner MW, Williams RC, Weingarten M, et al.: **Microtubules from mammalian brain: some properties of their depolymerization products and a proposed mechanism of assembly and disassembly.** *Proc Natl Acad Sci U S A.* 1974; **71**(4): 1159–1163.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mozziconacci J, Sandblad L, Wachsmuth M, et al.: **Tubulin dimers oligomerize before their incorporation into microtubules.** *PLoS One.* 2008; **3**(11): e3821.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kollman JM, Polka JK, Zelter A, et al.: **Microtubule nucleating gamma-TuSC assembles structures with 13-fold microtubule-like symmetry.** *Nature.* 2010; **466**(7308): 879–882.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Carazo-Salas RE, Gruss OJ, Mattaj JW, et al.: **Ran-GTP coordinates regulation of microtubule nucleation and dynamics during mitotic-spindle assembly.** *Nat Cell Biol.* 2001; **3**(3): 228–234.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kalab P, Pu RT, Dasso M: **The ran GTPase regulates mitotic spindle assembly.** *Curr Biol.* 1999; **9**(9): 481–484.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wiese C, Wilde A, Moore MS, et al.: **Role of importin-beta in coupling Ran to downstream targets in microtubule assembly.** *Science.* 2001; **291**(5504): 653–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chabin-Brion K, Marceiller J, Perez F, et al.: **The Golgi complex is a microtubule-organizing organelle.** *Mol Biol Cell.* 2001; **12**(7): 2047–2060.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Efimov A, Kharitonov A, Efimova N, et al.: **Asymmetric CLASP-dependent nucleation of noncentrosomal microtubules at the trans-Golgi network.** *Dev Cell.* 2007; **12**(6): 917–930.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Janson ME, Setty TG, Paoletti A, et al.: **Efficient formation of bipolar microtubule bundles requires microtubule-bound gamma-tubulin complexes.** *J Cell Biol.* 2005; **169**(2): 297–308.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Murata T, Sonobe S, Baskin TI, et al.: **Microtubule-dependent microtubule nucleation based on recruitment of gamma-tubulin in higher plants.** *Nat Cell Biol.* 2005; **7**(10): 961–968.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Petry S, Groen AC, Ishihara K, et al.: **Branching microtubule nucleation in *Xenopus* egg extracts mediated by augmin and TPX2.** *Cell.* 2013; **152**(4): 768–777.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Zheng Y, Wong ML, Alberts B, et al.: **Nucleation of microtubule assembly by a gamma-tubulin-containing ring complex.** *Nature.* 1995; **378**(6557): 578–583.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kollman JM, Merdes A, Mourey L, et al.: **Microtubule nucleation by γ -tubulin complexes.** *Nat Rev Mol Cell Biol.* 2011; **12**(11): 709–721.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wittmann T, Wilm M, Karsenti E, et al.: **TPX2, A novel *Xenopus* MAP involved in spindle pole organization.** *J Cell Biol.* 2000; **149**(7): 1405–1418.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tournebise R, Popov A, Kinoshita K, et al.: **Control of microtubule dynamics by the antagonistic activities of XMAP215 and XKCM1 in *Xenopus* egg extracts.** *Nat Cell Biol.* 2000; **2**(1): 13–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gruss OJ, Carazo-Salas RE, Schatz CA, et al.: **Ran induces spindle assembly by reversing the inhibitory effect of importin alpha on TPX2 activity.** *Cell.* 2001; **104**(1): 83–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Popov AV, Severin F, Karsenti E: **XMAP215 is required for the microtubule-nucleating activity of centrosomes.** *Curr Biol.* 2002; **12**(15): 1326–1330.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Schatz CA, Santarella R, Hoenger A, et al.: **Importin alpha-regulated nucleation of microtubules by TPX2.** *EMBO J.* 2003; **22**(9): 2060–2070.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brouhard GJ, Stear JH, Noetzel TL, et al.: **XMAP215 is a processive microtubule polymerase.** *Cell.* 2008; **132**(1): 79–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Gard DL, Kirschner MW: **Microtubule assembly in cytoplasmic extracts of *Xenopus* oocytes and eggs.** *J Cell Biol.* 1987; **105**(5): 2191–2201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Widlund PO, Stear JH, Pozniakovskiy A, et al.: **XMAP215 polymerase activity is built by combining multiple tubulin-binding TOG domains and a basic lattice-binding region.** *Proc Natl Acad Sci U S A.* 2011; **108**(7): 2741–2746.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Scrofani J, Sardon T, Meunier S, et al.: **Microtubule nucleation in mitosis by a RanGTP-dependent protein complex.** *Curr Biol.* 2015; **25**(2): 131–140.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Teixido-Travesa N, Roig J, Lüders J: **The where, when and how of microtubule nucleation - one ring to rule them all.** *J Cell Sci.* 2012; **125**(Pt 19): 4445–4456.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee MJ, Gergely F, Jeffers K, et al.: **Mps/XMAP215 interacts with the centrosomal protein D-TACC to regulate microtubule behaviour.** *Nat Cell Biol.* 2001; **3**(7): 643–649.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Roostalu J, Cade NI, Surrey T: **Complementary activities of TPX2 and chTOG constitute an efficient importin-regulated microtubule nucleation module.** *Nat Cell Biol.* 2015; **17**(11): 1422–1434.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

32. **F** Wieczorek M, Bechstedt S, Chaaban S, *et al.*: **Microtubule-associated proteins control the kinetics of microtubule nucleation.** *Nat Cell Biol.* 2015; **17**(7): 907–916.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Walker RA, O'Brien ET, Pryer NK, *et al.*: **Dynamic instability of individual microtubules analyzed by video light microscopy: rate constants and transition frequencies.** *J Cell Biol.* 1988; **107**(4): 1437–1448.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Chrétien D, Fuller SD, Karsenti E: **Structure of growing microtubule ends: two-dimensional sheets close into tubes at variable rates.** *J Cell Biol.* 1995; **129**(5): 1311–1328.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Mandelkow EM, Mandelkow E, Milligan RA: **Microtubule dynamics and microtubule caps: a time-resolved cryo-electron microscopy study.** *J Cell Biol.* 1991; **114**(5): 977–991.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. **F** Kerssemakers JW, Munteanu EL, Laan L, *et al.*: **Assembly dynamics of microtubules at molecular resolution.** *Nature.* 2006; **442**(7103): 709–712.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
37. **F** Schek HT 3rd, Gardner MK, Cheng J, *et al.*: **Microtubule assembly dynamics at the nanoscale.** *Curr Biol.* 2007; **17**(17): 1445–1455.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
38. Howard J, Hyman AA: **Growth, fluctuation and switching at microtubule plus ends.** *Nat Rev Mol Cell Biol.* 2009; **10**(8): 569–574.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. **F** Gardner MK, Charlebois BD, Jánosi IM, *et al.*: **Rapid microtubule self-assembly kinetics.** *Cell.* 2011; **146**(4): 582–592.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. Melki R, Carlier MF, Pantaloni D, *et al.*: **Cold depolymerization of microtubules to double rings: geometric stabilization of assemblies.** *Biochemistry.* 1989; **28**(23): 9143–9152.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Nogales E, Wolf SG, Downing KH: **Structure of the alpha beta tubulin dimer by electron crystallography.** *Nature.* 1998; **391**(6663): 199–203.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. **F** Wang HW, Nogales E: **Nucleotide-dependent bending flexibility of tubulin regulates microtubule assembly.** *Nature.* 2005; **435**(7044): 911–915.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. Buey RM, Diaz JF, Andreu JM: **The nucleotide switch of tubulin and microtubule assembly: a polymerization-driven structural change.** *Biochemistry.* 2006; **45**(19): 5933–5938.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. **F** Nawrotek A, Knossow M, Gigant B: **The determinants that govern microtubule assembly from the atomic structure of GTP-tubulin.** *J Mol Biol.* 2011; **412**(1): 35–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
45. **F** Rice LM, Montabana EA, Agard DA: **The lattice as allosteric effector: structural studies of alphabeta- and gamma-tubulin clarify the role of GTP in microtubule assembly.** *Proc Natl Acad Sci U S A.* 2008; **105**(14): 5378–5383.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. **F** Alushin GM, Lander GC, Kellogg EH, *et al.*: **High-resolution microtubule structures reveal the structural transitions in $\alpha\beta$ -tubulin upon GTP hydrolysis.** *Cell.* 2014; **157**(5): 1117–1129.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
47. **F** Maurer SP, Fourniol FJ, Bohner G, *et al.*: **EBs recognize a nucleotide-dependent structural cap at growing microtubule ends.** *Cell.* 2012; **149**(2): 371–382.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. **F** Geyer EA, Burns A, Lalonde BA, *et al.*: **A mutation uncouples the tubulin conformational and GTPase cycles, revealing allosteric control of microtubule dynamics.** *eLife.* 2015; **4**: e10113.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. **F** Shirasu-Hiza M, Coughlin P, Mitchison T: **Identification of XMAP215 as a microtubule-destabilizing factor in *Xenopus* egg extract by biochemical purification.** *J Cell Biol.* 2003; **161**(2): 349–358.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
50. **F** Ayaz P, Ye X, Huddleston P, *et al.*: **A TOG: $\alpha\beta$ -tubulin complex structure reveals conformation-based mechanisms for a microtubule polymerase.** *Science.* 2012; **337**(6096): 857–860.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. **F** Maurer SP, Bieling P, Cope J, *et al.*: **GTPgammaS microtubules mimic the growing microtubule end structure recognized by end-binding proteins (EBs).** *Proc Natl Acad Sci U S A.* 2011; **108**(10): 3988–3993.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. **F** Zaníc M, Stear JH, Hyman AA, *et al.*: **EB1 recognizes the nucleotide state of tubulin in the microtubule lattice.** *PLoS One.* 2009; **4**(10): e7585.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
53. **F** Maurer SP, Cade NI, Bohner G, *et al.*: **EB1 accelerates two conformational transitions important for microtubule maturation and dynamics.** *Curr Biol.* 2014; **24**(4): 372–384.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
54. Nakamura S, Grigoriev I, Nogi T, *et al.*: **Dissecting the nanoscale distributions and functions of microtubule-end-binding proteins EB1 and ch-TOG in interphase HeLa cells.** *PLoS One.* 2012; **7**(12): e51442.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. **F** Zaníc M, Widlund PO, Hyman AA, *et al.*: **Synergy between XMAP215 and EB1 increases microtubule growth rates to physiological levels.** *Nat Cell Biol.* 2013; **15**(6): 688–693.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. **F** des Georges A, Katsuki M, Drummond DR, *et al.*: **Mal3, the *Schizosaccharomyces pombe* homolog of EB1, changes the microtubule lattice.** *Nat Struct Mol Biol.* 2008; **15**(10): 1102–1108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
57. **F** Vitre B, Coquelle FM, Heichette C, *et al.*: **EB1 regulates microtubule dynamics and tubulin sheet closure *in vitro*.** *Nat Cell Biol.* 2008; **10**(4): 415–421.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. **F** Zhang R, Alushin GM, Brown A, *et al.*: **Mechanistic Origin of Microtubule Dynamic Instability and Its Modulation by EB Proteins.** *Cell.* 2015; **162**(4): 849–859.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
59. Grimaldi AD, Zaníc M, Kaverina I: **Encoding the microtubule structure: Allosteric interactions between the microtubule +TIP complex master regulators and TOG-domain proteins.** *Cell Cycle.* 2015; **14**(9): 1375–1378.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Grimaldi AD, Maki T, Fitton BP, *et al.*: **CLASPs are required for proper microtubule localization of end-binding proteins.** *Dev Cell.* 2014; **30**(3): 343–352.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Leano JB, Rogers SL, Slep KC: **A cryptic TOG domain with a distinct architecture underlies CLASP-dependent bipolar spindle formation.** *Structure.* 2013; **21**(6): 939–950.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. **F** Bechstedt S, Brouhard GJ: **Doublecortin recognizes the 13-prot filament microtubule cooperatively and tracks microtubule ends.** *Dev Cell.* 2012; **23**(1): 181–192.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
63. **F** Moores CA, Perderiset M, Francis F, *et al.*: **Mechanism of microtubule stabilization by doublecortin.** *Mol Cell.* 2004; **14**(6): 833–839.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
64. Bechstedt S, Lu K, Brouhard GJ: **Doublecortin recognizes the longitudinal curvature of the microtubule end and lattice.** *Curr Biol.* 2014; **24**(20): 2366–2375.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. **F** Volkov VA, Grissom PM, Arzhanik VK, *et al.*: **Centromere protein F includes two sites that couple efficiently to depolymerizing microtubules.** *J Cell Biol.* 2015; **209**(6): 813–828.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
66. Chen Y, Hancock WO: **Kinesin-5 is a microtubule polymerase.** *Nat Commun.* 2015; **6**: 8160.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Desai A, Verma S, Mitchison TJ, *et al.*: **Kin I kinesins are microtubule-destabilizing enzymes.** *Cell.* 1999; **96**(1): 69–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Moores CA, Yu M, Guo J, *et al.*: **A mechanism for microtubule depolymerization by KinI kinesins.** *Mol Cell.* 2002; **9**(4): 903–909.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. **F** Gardner MK, Zaníc M, Gell C, *et al.*: **Depolymerizing kinesins Kip3 and MCAK shape cellular microtubule architecture by differential control of catastrophe.** *Cell.* 2011; **147**(5): 1092–1103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
70. Odde DJ, Cassimeris L, Buettner HM: **Kinetics of microtubule catastrophe assessed by probabilistic analysis.** *Biophys J.* 1995; **69**(3): 796–802.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. Bowne-Anderson H, Zaníc M, Kauer M, *et al.*: **Microtubule dynamic instability: a new model with coupled GTP hydrolysis and multistep catastrophe.** *Bioessays.* 2013; **35**(5): 452–461.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Castle BT, Odde DJ: **Brownian dynamics of subunit addition-loss kinetics and thermodynamics in linear polymer self-assembly.** *Biophys J.* 2013; **105**(11): 2528–2540.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
73. **F** Coombes CE, Yamamoto A, Kenzie MR, *et al.*: **Evolving tip structures can explain age-dependent microtubule catastrophe.** *Curr Biol.* 2013; **23**(14): 1342–1348.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. Zakharov P, Gudimchuk N, Voevodin V, *et al.*: **Molecular and Mechanical Causes of Microtubule Catastrophe and Aging.** *Biophys J.* 2015; **109**(12): 2574–2591.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
75. **F** McIntosh JR, O'Toole E, Zhudenko K, *et al.*: **Conserved and divergent features of kinetochores and spindle microtubule ends from five species.** *J Cell Biol.* 2013; **200**(4): 459–474.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
76. Wittmann T, Wilm M, Karsenti E, *et al.*: **TPX2, A novel *Xenopus* MAP involved in spindle pole organization.** *J Cell Biol.* 2000; **149**(7): 1405–1418.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Chad Pearson

Department of Cell and Developmental Biology, University of Colorado School of Medicine, Aurora, USA

Domenico Gelati

Department of Cell and Developmental Biology, University of Colorado School of Medicine, Aurora, Colorado, USA

Marisa Ruehle

Department of Biochemistry and Molecular Genetics, University of Colorado School of Medicine, Aurora, Colorado, USA

Competing Interests: No competing interests were disclosed.

2. William Hancock

Pennsylvania State University, University Park, Pennsylvania, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research