REVIEW

Overexpression of angiotensin-converting enzyme in myelomonocytic cells enhances the immune response

[version 1; peer review: 3 approved]

Kenneth E. Bernstein, Zakir Khan, Jorge F. Giani, Tuantuan Zhao, Masahiro Eriguchi, Ellen A. Bernstein, Romer A. Gonzalez-Villalobos, Xiao Z. Shen

Department of Biomedical Sciences and the Department of Pathology, Cedars-Sinai Medical Center, Los Angeles, CA, USA

V1 First published: 23 Mar 2016, 5(F1000 Faculty Rev):393
<https://doi.org/10.12688/f1000research.7508.1>

Latest published: 23 Mar 2016, 5(F1000 Faculty Rev):393
<https://doi.org/10.12688/f1000research.7508.1>

Abstract

Angiotensin-converting enzyme (ACE) converts angiotensin I to the vasoconstrictor angiotensin II and thereby plays an important role in blood pressure control. However, ACE is relatively non-specific in its substrate specificity and cleaves many other peptides. Recent analysis of mice overexpressing ACE in monocytes, macrophages, and other myelomonocytic cells shows that these animals have a marked increase in resistance to experimental melanoma and to infection by *Listeria monocytogenes* or methicillin-resistant *Staphylococcus aureus* (MRSA). Several other measures of immune responsiveness, including antibody production, are enhanced in these animals. These studies complement a variety of studies indicating an important role of ACE in the immune response.

Keywords

Angiotensin-converting enzyme , ACE , angiotensin , immune response , renin-angiotensin system ,

Open Peer Review

Approval Status

	1	2	3
version 1 23 Mar 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Paul Overbeek**, Baylor College of Medicine, Houston, USA
2. **Markus Kalkum**, Beckman Research Institute of City of Hope, Duarte, USA
3. **Ifor Williams**, Emory University School of Medicine, Atlanta, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Kenneth E. Bernstein (kenneth.bernstein@cshs.org)

Competing interests: Romer A. Gonzalez-Villalobos is a visiting scientist at Cedars-Sinai Medical Center. He is employed by the Pfizer Inc Drug Safety Research and Development (DSRD) Immunotoxicology Center of Emphasis (CoE) (Groton, CT, USA). The other authors declare that they have no competing interests.

Grant information: This work was supported by National Institutes of Health grants R01 HL110353, R21 AI114965, and R03DK101592. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2016 Bernstein KE *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bernstein KE, Khan Z, Giani JF *et al.* **Overexpression of angiotensin-converting enzyme in myelomonocytic cells enhances the immune response [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):393 <https://doi.org/10.12688/f1000research.7508.1>

First published: 23 Mar 2016, 5(F1000 Faculty Rev):393 <https://doi.org/10.12688/f1000research.7508.1>

Introduction

The renin-angiotensin system (RAS) is composed of the substrate angiotensinogen, a protein produced by the liver, which is sequentially degraded by the enzymes renin and angiotensin-converting enzyme (ACE) into the vasoconstrictor angiotensin II. The workings of this system and its role as a central regulator of blood pressure control have been the subject of thousands of scientific publications¹. Although there are voluminous biochemical data on each of the components of the RAS, an interesting question is the origins of this system. Such a question recognizes that blood pressure is a function of higher organisms, whereas the origins of the proteins composing the RAS go back to a time when simpler organisms had neither blood nor blood pressure. Angiotensinogen is a liver acute-phase protein; it is an α_2 globulin related to haptoglobin and α_2 -microglobulin. Renin, made in the kidney, is an aspartyl protease related to pepsin, whereas ACE, made by endothelium and many other tissues, is a zinc-dependent peptidase. Whereas angiotensinogen and renin have evolved to the point where they participate only in blood pressure regulation, ACE plays additional roles and has been implicated in normal male reproduction, several aspects of hematopoiesis, the degradation of β -amyloid peptides, and several other physiologic and pathologic processes²⁻⁷.

One of the most interesting areas of ACE biology is its role in the immune response. Early evidence for this came from the analysis of ACE expression by the macrophages and giant cells found in granuloma. The formation of a granuloma is the immune system's response to organisms or inorganic materials that are difficult to phagocytize⁸. The classic granuloma is associated with tuberculosis, but granuloma formation is also seen in sarcoid, leprosy, and several other diseases. It has been known for many years that the epithelial macrophages and giant cells that are a prominent feature of granulomas express ACE⁹. Thus, there is an association between the immune activation of the monocytic-derived cells in a granuloma and ACE expression. What remained unknown was whether ACE expression was also a contributor to this process.

ACE 10/10 mice

Our experience with ACE and the immune response first came from the study of mice with genetic mutations in the ACE gene. The wild-type (WT) ACE protein contains two independent catalytic domains that resulted from an ancient gene duplication, estimated to have occurred over 300 million years ago¹. Each catalytic domain independently binds zinc, which is required for catalytic activity. To understand the function of each catalytic domain, mice were made in which point mutations were introduced into the mouse genome that specifically inactivated either the ACE N- or C-domain zinc binding site^{10,11}. Thus, the mutations specifically inactivated the N- or C-domain catalytic sites. These mice are called ACE N-KO and ACE C-KO, respectively. In these animals, the non-targeted domain remained fully catalytic and the tissue expression levels of the mutated ACE proteins were identical to those of WT mice. When peritoneal macrophages from these animals were isolated and exposed *in vitro* overnight to lipopolysaccharide (LPS), the macrophages were activated and secreted tumor necrosis factor alpha (TNF α) into the media (Figure 1)¹². While mice lacking ACE C-domain catalytic activity produced roughly equivalent levels of

Figure 1. Tumor necrosis factor-alpha (TNF α) expression in macrophages lacking the angiotensin-converting enzyme N- or C-catalytic domain. Thioglycolate-elicited peritoneal macrophages were collected and purified by adhesion. After an overnight incubation with 1 μ g/mL lipopolysaccharide, the concentration of TNF α was determined by enzyme-linked immunosorbent assay. The figure shows data for individual mice and the group mean $\pm$ standard error of the mean. C-KO, **macrophages lacking the angiotensin-converting enzyme C-catalytic domain**; N-KO, **macrophages lacking the angiotensin-converting enzyme N-catalytic domain**.

TNF α as WT mice, macrophages from mice lacking N-domain activity (ACE N-KO) produced approximately fourfold higher levels. Thus, ACE activity can have profound effects on the ability of macrophages to produce the important pro-inflammatory cytokine TNF α .

A second model giving insight into the role of ACE in the immune function is a genetically engineered mouse line termed ACE 10/10. Here, gene targeting was used to insert a neomycin resistance cassette and strong transcriptional stop signal 3' to the ACE promoter¹³. This acts as a barrier to endogenous ACE promoter activity. 3' to the neomycin cassette, we inserted a *c-fms* promoter cassette that now controls tissue expression of the ACE gene. *c-fms* encodes the receptor for macrophage colony-stimulating factor and is normally expressed at high levels by myelomonocytic lineage cells^{14,15}. Mice that are heterozygous or homozygous for the ACE 10/10 allele overexpress active ACE in monocytic cells such as monocytes and macrophages. Other myelomonocytic cells, such as neutrophils and dendritic cells, also overexpress ACE but at only 4% and 17% of macrophage levels, respectively. ACE expression by T or B cells is very low, similar to WT mice. Homozygous ACE 10/10 mice lack ACE expression by endothelial cells and renal epithelial cells. Despite this, ACE 10/10 mice have normal body weights, serum ACE levels, renal function, and blood pressure. They live normal life spans and have no evidence of auto-immune disease.

The first inkling that ACE 10/10 mice were unusual came from challenging the mice with B16-F10 (H-2^b) melanoma tumor cells¹³. Tumor volumes were assessed 2 weeks after the intradermal injection of tumor cells. A typical response is shown in Figure 2. Whereas tumors in WT mice averaged 540 mm³, tumors in ACE 10/10 mice averaged only 90 mm³. This difference was seen using different B16 sublines and in both inbred and outbred ACE 10/10 mice. The decrease in tumor growth was dependent on CD8⁺ T cells; in ACE 10/10 mice, depletion of CD8⁺ T cells, but not CD4⁺ T cells, led to rapid tumor growth. When B16 tumor cells constitutively expressing ovalbumin were used, tetramer analysis of T-cell receptors showed that ACE 10/10 mice have more circulating tumor-specific CD8⁺ T cells with specificity for the ovalbumin epitope SIINFEKL and the B16 TRP-2 epitope (SVYDFFVWL) than WT mice. Tumor resistance was dependent on ACE catalytic activity, as demonstrated by WT levels of tumor growth in ACE 10/10 mice treated with an ACE inhibitor¹³. In contrast, treatment of the mice with an angiotensin II receptor antagonist had no effect on tumor growth.

An insight into the mode of tumor resistance in ACE 10/10 mice was provided by histological examination of the small tumors present in these mice. This revealed far larger numbers of inflammatory cells, including monocytes, macrophages, and some lymphocytes within the tumor blood vessels and the tumor itself than in tumors from WT mice. Furthermore, tumor resistance was transferable by bone marrow transplantation from ACE 10/10 into WT mice. This transplant experiment was important since WT mice chimeric for ACE 10/10 bone marrow have normal ACE expression in all tissues except bone marrow-derived cells. Thus, these data and several other lines of investigation indicated that it was the presence of ACE activity in bone marrow-derived cells, and not the lack of ACE expression in endothelium, that was important in the enhanced immune response of the ACE 10/10 mice.

The ACE 10/10 mice were also tested in a tumor metastasis model where the mice were injected intravenously with B16 tumor cells¹⁶. After 14 days, melanotic modules in the lungs were quantitated. In this protocol, ACE 10/10 mice averaged only one third the number of visible lung metastases as in WT mice.

A detailed analysis of macrophage function in ACE 10/10 mice indicated that these cells have a pronounced pro-inflammatory “M1” phenotype, as compared with equivalent WT cells. Specifically, macrophages from ACE 10/10 mice produced more interleukin-12 (IL-12) p40, TNF α , and nitric oxide synthase II (inducible nitric oxide synthase, or iNOS) in response to tumor cells or LPS. On the other hand, the ACE 10/10 cells made less of the immunosuppressive cytokine IL-10^{13,17,18}.

Innate immunity in ACE 10/10 mice

Resistance to B16 melanoma is typically thought to be mediated by the adaptive immune response. In order to assess innate immunity, the resistance of the ACE 10/10 mice to infection with either *L. monocytogenes* or methicillin-resistant *S. aureus* (MRSA) was evaluated¹⁷. Resistance to *L. monocytogenes* was assessed following intravenous injection of the strain EGD. Mice were sacrificed 3 or 5 days after infection, and bacterial counts were determined in the spleen and liver. This showed consistently fewer bacteria in ACE 10/10 mice as compared with WT. Pre-treatment of the ACE 10/10 mice for several days with the ACE inhibitor ramipril eliminated any significant differences between these mice and WT. In contrast, the angiotensin II receptor antagonist losartan had no significant effect.

Peritoneal macrophages were also tested *in vitro* for their ability to kill *L. monocytogenes*. In the absence of interferon-gamma (IFN γ) priming, there was no significant difference between

Figure 2. Growth of the B16 melanoma in angiotensin-converting enzyme (ACE) 10/10 mice. Both ACE 10/10 and wild-type (WT) mice were injected intradermally with one million B16-F10 melanoma cells. After 14 days, there was a very significant difference in tumor growth, with WT mice having much larger tumors than ACE 10/10 mice. These photos show typical results. The fur of the ACE 10/10 was shaved to demonstrate the reduction in tumor size in these animals.

ACE 10/10 and WT macrophages in the killing of *L. monocytogenes*. As ACE 10/10 macrophages express abundant surface ACE, these data demonstrate that ACE has no direct bactericidal effect. However, after IFN γ priming, ACE 10/10 macrophages were much more effective in killing bacteria than WT macrophages were¹⁷.

ACE 10/10 mice were also challenged by intradermal infection with MRSA (USA300, strain SAF8300). After 4 days, bacterial counts in the skin lesions were determined. ACE 10/10 mice averaged 50-fold less bacteria within the MRSA lesions than WT mice. Again, this difference was eliminated by the ACE inhibitor lisinopril, and enhanced resistance to MRSA could be conferred by bone marrow transplant of ACE 10/10 bone marrow cells into WT mice¹⁷.

Immunological memory and antibody production

To investigate immunological memory, female ACE 10/10 and WT mice were infected with polyoma virus. By 28 days, the mice clear serum polyoma and were re-challenged with either WT vaccinia virus (V-WT) or vaccinia virus modified to express the polyoma large T epitope LT359–368 (V-PLT)¹⁸. Four days after V-WT infection, there was no difference in ovarian vaccinia viral titers as measured by a plaque assay (Figure 3). However, when mice were challenged with V-PLT, there was a marked difference in viral titers. Thus, in a viral recall assay, these data indicate an increased immune response in the ACE 10/10 mice.

Finally, antibody production in ACE 10/10 mice was assessed after immunization with ovalbumin in complete Freund's adjuvant (Figure 4). Ten days after immunization, plasma was collected and titers of antibody subtypes specific to ovalbumin were determined by enzyme-linked immunosorbent assay. The major anti-ovalbumin antibody by enzyme-linked immunosorbent assay. The major anti-ovalbumin

antibody was IgG1, which showed a 22-fold increase in the ACE 10/10 mice versus WT. A similar pattern of antibody increase was seen for IgG2b, IgG2c, and IgG3 antibodies, though at substantially lower levels of antibody expression.

Figure 3. Polyoma virus recall. Wild-type (WT) and angiotensin-converting enzyme (ACE) 10/10 mice were infected with mouse polyoma virus. After 1 month, the mice were challenged with either WT vaccinia virus (V-WT) or a modified vaccinia virus which expressed a portion of the polyoma large T protein (V-PLT). Viral titers in the ovary were measured using a plaque assay. Whereas there is no difference in response to V-WT, the ACE 10/10 mice have a much better immune recall response to V-PLT.

Figure 4. Anti-ovalbumin antibodies. Wild-type (WT) and angiotensin-converting enzyme (ACE) 10/10 mice were immunized subcutaneously with 100 µg of ovalbumin in complete Freund's adjuvant. Ten days later, plasma was collected and titers of antibody subtypes specific to ovalbumin were determined by enzyme-linked immunosorbent assay. The major anti-ovalbumin antibody was IgG1, which showed a marked increase in the ACE 10/10 mice versus WT (990 versus 45 µg/mL). A similar pattern of increased antibody expression was seen for IgG2b, IgG2c, and IgG3 antibodies, though at lower levels of antibody expression. These data are for illustrative purposes only. The results are currently unpublished and have not been peer-reviewed but will be peer-reviewed and published elsewhere.

The data summarized here support the idea that ACE plays an important role in the immune response in addition to its role in regulating blood pressure. These data complement other studies implicating ACE in immunity to disease. These include a role of ACE and angiotensin II in the immune response to experimental auto-immune encephalomyelitis^{19–22}, and rheumatoid and other types of arthritis^{23–26}, in the production of the immuno-dominant epitope of the HIV protein gp160^{27,28}, and in generating the peptide repertoire diversity displayed by major histocompatibility complex (MHC) class I proteins^{29,30}.

Conclusions

ACE 10/10 mice are a model in which ACE is overexpressed in myelomonocytic cells, particularly monocytes and macrophages. It may be that this model recapitulates the expression of ACE by monocytic cells found in granuloma but that now supranormal expression levels generate a more pronounced immune phenotype than under natural conditions. We think that the most likely mechanistic explanation for the ACE 10/10 phenotype is that ACE overexpression fundamentally modifies the differentiation program of monocytes and macrophages. As indicated by the elimination of the phenotype after the administration of ACE inhibitors, the ACE 10/10 phenotype requires the catalytic activity of ACE and thus must be hypothesized to be a downstream effect of some ACE product(s). The precise peptides that have such a dramatic effect on the immune response are not known at present. The discovery that ACE overexpression enhances both innate and acquired monocytic function holds promise for an entirely new approach for improving

the immune response to a variety of stimuli, including infections and tumors.

Abbreviations

ACE, angiotensin-converting enzyme; IFN γ , interferon gamma; IL, interleukin; LPS, lipopolysaccharide; MRSA, methicillin-resistant *Staphylococcus aureus*; RAS, renin-angiotensin system; TNF α , tumor necrosis factor alpha; V-PLT, polyoma large T protein; V-WT, wild-type vaccinia virus; WT, wild-type.

Competing interests

Romer A. Gonzalez-Villalobos is a visiting scientist at Cedars-Sinai Medical Center. He is employed by the Pfizer Inc Drug Safety Research and Development (DSRD) Immunotoxicology Center of Emphasis (CoE) (Groton, CT, USA). The other authors declare that they have no competing interests.

Grant information

This work was supported by National Institutes of Health grants R01 HL110353, R21 AI114965, and R03DK101592.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

The authors thank Brian Taylor for administrative assistance.

References

- Bernstein KE, Ong FS, Blackwell WB, *et al.*: **A modern understanding of the traditional and nontraditional biological functions of angiotensin-converting enzyme.** *Pharmacol Rev.* 2013; **65**(1): 1–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Krege JH, John SW, Langenbach LL, *et al.*: **Male-female differences in fertility and blood pressure in ACE-deficient mice.** *Nature.* 1995; **375**(6527): 146–148.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Esther CR, Howard TE, Marino EM, *et al.*: **Mice lacking angiotensin-converting enzyme have low blood pressure, renal pathology, and reduced male fertility.** *Lab Invest.* 1996; **74**(5): 953–965.
[PubMed Abstract](#)
- Fuchs S, Frenzel K, Hubert C, *et al.*: **Male fertility is dependent on dipeptidase activity of testis ACE.** *Nat Med.* 2005; **11**(11): 1140–2; author reply 1142–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hubert C, Savary K, Gasc J, *et al.*: **The hematopoietic system: a new niche for the renin-angiotensin system.** *Nat Clin Pract Cardiovasc Med.* 2006; **3**(2): 80–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lin C, Datta V, Okwan-Duodu D, *et al.*: **Angiotensin-converting enzyme is required for normal myelopoiesis.** *FASEB J.* 2011; **25**(4): 1145–1155.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bernstein KE, Koronyo Y, Salumbides BC, *et al.*: **Angiotensin-converting enzyme overexpression in myelomonocytes prevents Alzheimer's-like cognitive decline.** *J Clin Invest.* 2014; **124**(3): 1000–1012.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ramakrishnan L: **Revisiting the role of the granuloma in tuberculosis.** *Nat Rev Immunol.* 2012; **12**(5): 352–366.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stanton L, Fenhalls G, Lucas A, *et al.*: **Immunophenotyping of macrophages in human pulmonary tuberculosis and sarcoidosis.** *Int J Exp Pathol.* 2003; **84**(6): 289–304.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Fuchs S, Xiao HD, Cole JM, *et al.*: **Role of the N-terminal catalytic domain of angiotensin-converting enzyme investigated by targeted inactivation in mice.** *J Biol Chem.* 2004; **279**(16): 15946–15953.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fuchs S, Xiao HD, Hubert C, *et al.*: **Angiotensin-converting enzyme C-terminal catalytic domain is the main site of angiotensin I cleavage in vivo.** *Hypertension.* 2008; **51**(2): 267–274.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ong FS, Lin CX, Campbell DJ, *et al.*: **Increased angiotensin II-induced hypertension and inflammatory cytokines in mice lacking angiotensin-converting enzyme N domain activity.** *Hypertension.* 2012; **59**(2): 283–290.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shen XZ, Li P, Weiss D, *et al.*: **Mice with enhanced macrophage angiotensin-converting enzyme are resistant to melanoma.** *Am J Pathol.* 2007; **170**(6): 2122–2134.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Himes SR, Tagoh H, Goonetilleke N, *et al.*: **A highly conserved c-fms gene intronic element controls macrophage-specific and regulated expression.** *J Leukoc Biol.* 2001; **70**(5): 812–820.
[PubMed Abstract](#) | **F1000 Recommendation**
- Sasmono RT, Oceandy D, Pollard JW, *et al.*: **A macrophage colony-stimulating factor receptor-green fluorescent protein transgene is expressed throughout**

- the mononuclear phagocyte system of the mouse. *Blood*. 2003; **101**(3): 1155–1163.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
16. Shen XZ, Okwan-Duodu D, Blackwell WL, *et al.*: Myeloid expression of angiotensin-converting enzyme facilitates myeloid maturation and inhibits the development of myeloid-derived suppressor cells. *Lab Invest*. 2014; **94**(5): 536–544.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Okwan-Duodu D, Datta V, Shen XZ, *et al.*: Angiotensin-converting enzyme overexpression in mouse myelomonocytic cells augments resistance to *Listeria* and methicillin-resistant *Staphylococcus aureus*. *J Biol Chem*. 2010; **285**(50): 39051–39060.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Gonzalez-Villalobos RA, Shen XZ, Bernstein EA, *et al.*: Rediscovering ACE: novel insights into the many roles of the angiotensin-converting enzyme. *J Mol Med (Berl)*. 2013; **91**(10): 1143–1154.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. **F** Constantinescu CS, Ventura E, Hilliard B, *et al.*: Effects of the angiotensin converting enzyme inhibitor captopril on experimental autoimmune encephalomyelitis. *Immunopharmacol Immunotoxicol*. 1995; **17**(3): 471–491.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 20. **F** Platten M, Youssef S, Hur EM, *et al.*: Blocking angiotensin-converting enzyme induces potent regulatory T cells and modulates TH1- and TH17-mediated autoimmunity. *Proc Natl Acad Sci U S A*. 2009; **106**(35): 14948–14953.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 21. Stegbauer J, Lee DH, Seubert S, *et al.*: Role of the renin-angiotensin system in autoimmune inflammation of the central nervous system. *Proc Natl Acad Sci U S A*. 2009; **106**(35): 14942–14947.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Steinman L: Development of therapies for autoimmune disease at Stanford: a tale of multiple shots and one goal. *Immunol Res*. 2014; **58**(2–3): 307–314.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. Lühder F, Lee DH, Gold R, *et al.*: Small but powerful: short peptide hormones and their role in autoimmune inflammation. *J Neuroimmunol*. 2009; **217**(1–2): 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. **F** Dalbeth N, Edwards J, Fairchild S, *et al.*: The non-thiol angiotensin-converting enzyme inhibitor quinapril suppresses inflammatory arthritis. *Rheumatology (Oxford, England)*. 2005; **44**(1): 24–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 25. **F** Fahmy Wahba MG, Shehata Messiha BA, Abo-Saif AA: Ramipril and haloperidol as promising approaches in managing rheumatoid arthritis in rats. *Eur J Pharmacol*. 2015; **765**: 307–315.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 26. **F** Sagawa K, Nagatani K, Komagata Y, *et al.*: Angiotensin receptor blockers suppress antigen-specific T cell responses and ameliorate collagen-induced arthritis in mice. *Arthritis Rheum*. 2005; **52**(6): 1920–1928.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 27. **F** Kozlowski S, Corr M, Takeshita T, *et al.*: Serum angiotensin-1 converting enzyme activity processes a human immunodeficiency virus 1 gp160 peptide for presentation by major histocompatibility complex class I molecules. *J Exp Med*. 1992; **175**(6): 1417–1422.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 28. **F** Nakagawa Y, Takeshita T, Berzofsky JA, *et al.*: Analysis of the mechanism for extracellular processing in the presentation of human immunodeficiency virus-1 envelope protein-derived peptide to epitope-specific cytotoxic T lymphocytes. *Immunology*. 2000; **101**(1): 76–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 29. Eisenlohr LC, Bacik I, Bennink JR, *et al.*: Expression of a membrane protease enhances presentation of endogenous antigens to MHC class I-restricted T lymphocytes. *Cell*. 1992; **71**(6): 963–972.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. **F** Shen XZ, Billet S, Lin C, *et al.*: The carboxypeptidase ACE shapes the MHC class I peptide repertoire. *Nat Immunol*. 2011; **12**(11): 1078–1085.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Ifor Williams**

Department of Pathology, Emory University School of Medicine, Atlanta, GA, USA

Competing Interests: No competing interests were disclosed.

2. **Markus Kalkum**

Department of Immunology, Beckman Research Institute of City of Hope, Duarte, CA, USA

Competing Interests: No competing interests were disclosed.

3. **Paul Overbeek**

Department of Molecular & Cellular Biology, Baylor College of Medicine, Houston, TX, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research