REVIEW

Molecular biology of breast tumors and prognosis [version 1; peer review: 3 approved]

Gustavo Baldassarre, Barbara Belletti

Department of Experimental Oncology, Department of Translational Research, CRO Aviano IRCCS, National Cancer Institute, Italy, Aviano, Italy

V1 First published: 21 Apr 2016, 5(F1000 Faculty Rev):711
<https://doi.org/10.12688/f1000research.8158.1>
 Latest published: 21 Apr 2016, 5(F1000 Faculty Rev):711
<https://doi.org/10.12688/f1000research.8158.1>

Abstract

Breast cancer is the most common cancer among women worldwide. Great scientific, economical, and organizational efforts are in place to understand the causes of onset, identify the critical molecular players of progression, and define new lines of intervention providing more benefits and less toxicity. These efforts have certainly not been vain, since overall survival, especially in specific subsets of breast cancer, has greatly improved during the last decades. At present, breast cancer patients' treatment and care have reached a high standard of quality, and currently one of the most urgent needs resides in the necessity to better distinguish the tumors that need to be more aggressively treated and identify the best therapeutic option tailored to each patient. This objective will be achievable only if the information clarifying the biology of breast cancer can be successfully transferred to the clinic. A common effort by scientists and clinicians toward this integration and toward the use of multidisciplinary approaches will be necessary to reach this important goal.

Keywords

breast cancer, breast cancer treatment and care, breast cancer prognosis

Open Peer Review

Approval Status

	1	2	3
version 1 21 Apr 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Andrea Morrione**, Thomas Jefferson University, Philadelphia, USA
2. **Michael Retsky**, Harvard TH Chan School of Public Health, Boston, USA
University College London, London, UK
3. **Alfredo Budillon**, Istituto Nazionale Tumori G. Pascale Napoli, Naples, Italy

Any comments on the article can be found at the end of the article.

Corresponding author: Gustavo Baldassarre (gbaldassarre@cro.it)

Competing interests: The authors declare that they have no competing interests.

Grant information: The authors' work is supported by AIRC (Associazione Italiana Ricerca sul Cancro) IG 15902 (Barbara Belletti) and IG 16865 (Gustavo Baldassarre).

Copyright: © 2016 Baldassarre G and Belletti B. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Baldassarre G and Belletti B. **Molecular biology of breast tumors and prognosis [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):711 <https://doi.org/10.12688/f1000research.8158.1>

First published: 21 Apr 2016, 5(F1000 Faculty Rev):711 <https://doi.org/10.12688/f1000research.8158.1>

Introduction

Breast cancer (BC) is the most common cancer among women worldwide, accounting for approximately one-quarter of all cancers in females worldwide and 27% of cancers in developed countries with a Western lifestyle, thus representing a real health emergency¹. Here, we briefly focus on some specific biological and clinical aspects of BC that are still matter of controversy and in which there is urgent need for competent integration to implement diagnostic and therapeutic options.

Gene expression profiles define different breast cancer subtypes

Large-scale gene expression profile (GEP) studies demonstrated that BCs are not a single entity but can be divided into at least four major subtypes: luminal A (LBC-A), luminal B (LBC-B), HER2-positive, and triple negative/basal-like². This classification has been recently confirmed and integrated with genomic data demonstrating that the diverse subtypes are indeed associated with different recurrent genomic alterations³. LBC in the Western world represents the most common subtype, accounting for more than 60% of all diagnosed BC. In clinical practice, a few characteristics, such as estrogen receptor (ER), progesterone receptor (PR), HER2, and Ki67 expression, are currently used to distinguish LBC-A (ER+ and/or PR+, HER2-, low Ki67) from LBC-B (ER+ and/or PR+, HER2- or HER2+, high Ki67)⁴. This classification has therapeutic implications, since based on the relative expression of the above markers, LBC patients will or will not receive hormone-, chemo-, or targeted-therapies⁴. When compared with LBC-A, LBC-B displays a higher rate of early recurrence and worse prognosis, representing a subgroup for which the choice of the optimal therapy still represents a difficult task for the clinician⁴. In fact, clear biomarkers to select the most appropriate (hormone- with or without chemo-) therapy in LBC-B still have to be validated and introduced into the clinic⁴.

The above-mentioned molecular classification also has prognostic relevance, with triple-negative and HER2+ BC having a more aggressive progression. Moreover, since the same molecular characteristics are already present in the first stages of BC and in *in situ* lesions, it is expected that early diagnosis could help in the identification (and removal) of potentially malignant tumors that need to be aggressively treated.

Significance of identifying *in situ* breast cancer and the risk of overtreatment

Based on this idea, several BC screening programs have been introduced worldwide in women aged >50 years. The introduction of screening programs has substantially increased the number of early stage *in situ* lesions, mainly ductal carcinoma *in situ* (DCIS). It is debated whether low-grade DCIS would result in an invasive stage⁵. The general view was that the increase in DCIS diagnoses would result in decreased incidence of invasive BC and, eventually, decreased mortality for BC. However, this has not always been the case, as demonstrated by epidemiological analyses, raising some concern over the possibility of “over-diagnosis” linked to the wide spread of BC screening programs⁶.

The importance of early diagnosis in BC was recently addressed by a large observational study estimating the mortality for BC following a diagnosis of DCIS⁷. This important study, enrolling more than 100,000 women, provided the interesting observation that women diagnosed with DCIS display a BC-related death risk at 20 years comparable to that estimated for the general population and that using aggressive treatments for all DCIS does not reduce the mortality for BC⁷. Yet a diagnosis of DCIS in black women or in women under the age of 40 and the presence of high risk factors such as HER2 expression are associated with an increased risk of BC-related death⁷. This clinical evidence reinforces the concept that, even when *in situ*, different molecular alterations in BC strongly impact on the outcome of the disease and on patients’ survival.

Breast cancer: age matters

This concept introduces the relevance of molecular studies to precisely identify the cancer that needs to be aggressively treated and to discover the most appropriate treatment for each patient. Addressing these two unmet clinical needs is the only way to further improve BC cure while limiting the risk of overtreatment.

The recent study by Narod and colleagues suggests that the group of BC patients that receives a diagnosis of BC before the age of 40 is the one that more urgently needs to be accurately classified at molecular level⁷. Young age at diagnosis has emerged as an independent factor associated with higher risk of relapse and death in several large studies on BC^{8,9}. Several factors have been linked to this poor prognosis, including large tumor size at diagnosis, higher tumor grade, mitotic index, lymphovascular invasion, increased expression of HER2, and lower ER and PR expression¹⁰. More recently, it has been proposed that for these patients it is also worth testing the presence of mutation in the BC susceptibility genes *BRCA1* and *BRCA2*, independently of their family history¹¹. A general consensus in the scientific community has been reached to define BC in Young Women (BCYW), although fairly rare (~7% of all diagnosed BC), as a distinct entity that merits being studied and treated using specific guidelines and following specific research priorities¹².

More aggressive subtypes are more common in BCYW. In particular, when compared with older patients, BCYW more frequently displays the triple-negative and HER2+ BC subtypes. However, in the Western world, LBC-B still accounts for more than 60% of all BCYW, remaining the most common histotype and displaying a particularly bad prognosis^{8,9}. These observations raised the question of whether BCYW has a unique biology or whether this just represents a surrogate of the higher incidence of aggressive molecular subtypes. But, even after correction for stage and tumor characteristics, young age at diagnosis remains an independent risk factor for relapse and BC-related death¹³.

Unraveling the biological uniqueness of BCYW is fundamental, as it not only increases our understanding of the disease process but also underlies the decision of whether or not to offer the same therapeutic options reserved to the high-risk older patients or to choose therapeutic approaches based on a specific biology¹⁴.

Accumulating evidence suggests that differences in the mammary stroma composition and changes that occur with pregnancy and breastfeeding likely contribute to the different biology of BCYW. Moreover, these tumors are enriched with processes related to immune-related gene signatures and immature mammary cell populations (RANKL, c-kit, BRCA1-mutated phenotype, mammary stem cells, and luminal progenitors)¹⁵.

The comprehensive analysis of BC with respect to age on a large compendium of publicly available gene expression datasets (more than 3500 BCYW)¹⁵ has demonstrated that specific pathways are altered in BCYW with respect to older patients. However, the identification of altered signaling pathways is not sufficient to identify the driver alterations responsible for tumor onset and/or progression.

A recent conference centered on the management of BCYW confirmed that, although some progress has been made in the understanding of the clinical and biological behaviors of BC in patients younger than 40, we still need to clarify many aspects to properly treat this particular subgroup of patients¹⁶. In particular, there is a consensus on the fact that biology appears to be different in BCYW and that this is particularly true for the endocrine-sensitive tumors. Based on this consideration, it is necessary to study the tumor genetic profile in larger cohorts to further understand if a common and unique pattern of gene expression exists in BCYW. Then, it becomes clear that the study of primary tumors will not be sufficient and that characterizing the recurrent tumors will be mandatory. Finally, it could be of special relevance to understand the interaction of the endocrine and immune systems in these young patients and to cleverly look at response patterns to the current treatments. Only in this way will we have the possibility to improve the treatment, quality of life, and survival of BCYW patients¹⁶.

Is it the time, in the post-genomic era, to return to functional studies?

It is now clear that sequencing and gene expression profile studies must be associated with high-throughput functional analyses to precisely identify the genes and/or the pathway each tumor type is addicted to. A recent resource has proposed that it is possible to identify the vulnerability of each type of BC using functional assays, even using a panel of cell lines, if these well recapitulate the original disease¹⁷.

We are convinced, however, that these types of assays should be integrated by the generation of more reliable models of validation. We recently explored this possibility focusing on the onset of local recurrences in BC by setting up a mouse model closely resembling

the course of the human pathology¹⁸. Our work highlighted that targeting specific signaling pathways at the time of surgery has great potential to prevent the re-appearance of BC in mice^{18–21}. In particular, we observed that the specific inhibition of p70S6K1 had little effect on blocking the growth of established breast tumors but prevented the onset of BC recurrences when therapy was administered with a perisurgical treatment schedule. p70S6K1 activity was necessary for the survival of isolated BC cells residually present in the post-surgery setting, making it an ideal target to improve the efficacy of surgery^{18,19}. Similar results were also shown for the PAR-4 protein using different models of recurrence formation²². In these models, PAR-4 acted as an inhibitor of recurrence formation by inducing multinucleation in oncogene-addicted cells²².

In accord with the importance of timely delivery of the therapy, it has been recently demonstrated that the application of intraoperative radiotherapy (IORT) in BC patients has different effects if used immediately after tumor removal or as a second procedure after pathological examination²³. At the molecular level, this clinical observation could be explained by the recently demonstrated direct effect of IORT in the tumor microenvironment, where it modulates the EGF-EGFR-p70S6K1 signaling axis, via the induction of miR-223 expression in the local peri-tumoral microenvironment²¹.

It would be interesting to evaluate whether in high-risk BC patients, such as BCYW, similar mechanisms of cell survival exist and are at least partially responsible for their aggressive phenotype and whether targeting these specific pathways at the right time could significantly impact on the patient's disease-free and overall survival. To this aim, better models recapitulating the biology of BCYW and more integration between clinicians and preclinical researchers are primarily and urgently needed.

Competing interests

The authors declare that they have no competing interests.

Grant information

The authors' work is supported by AIRC (Associazione Italiana Ricerca sul Cancro) IG 15902 (Barbara Belletti) and IG 16865 (Gustavo Baldassarre).

Acknowledgements

We thank all the members of the Breast Unit and of the SCICC lab at CRO and all our external collaborators for their support, dedicated work, and interesting and stimulating discussions.

References

1. Ferlay J, Shin HR, Bray F, *et al.*: **Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008.** *Int J Cancer*. 2010; **127**(12): 2893–917.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
2. Perou CM, Sorlie T, Eisen MB, *et al.*: **Molecular portraits of human breast**

tumours. *Nature*. 2000; **406**(6797): 747–52.

[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

3. Cancer Genome Atlas Network: **Comprehensive molecular portraits of human breast tumours.** *Nature*. 2012; **490**(7418): 61–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

4. **F** Ades F, Zardavas D, Bozovic-Spasojevic I, *et al.*: **Luminal B breast cancer: molecular characterization, clinical management, and future perspectives.** *J Clin Oncol.* 2014; **32**(25): 2794–803.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
5. **F** Ernster VL, Ballard-Barbash R, Barlow WE, *et al.*: **Detection of ductal carcinoma in situ in women undergoing screening mammography.** *J Natl Cancer Inst.* 2002; **94**(20): 1546–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
6. **F** Bleyer A, Welch HG: **Effect of three decades of screening mammography on breast-cancer incidence.** *N Engl J Med.* 2012; **367**(21): 1998–2005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
7. **F** Narod SA, Iqbal J, Giannakeas V, *et al.*: **Breast Cancer Mortality After a Diagnosis of Ductal Carcinoma In Situ.** *JAMA Oncol.* 2015; **1**(7): 888–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
8. **F** El Saghir NS, Seoud M, Khalil MK, *et al.*: **Effects of young age at presentation on survival in breast cancer.** *BMC Cancer.* 2006; **6**: 194.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
9. **F** Canello G, Maisonneuve P, Mazza M, *et al.*: **Pathological features and survival outcomes of very young patients with early breast cancer: how much is “very young”?** *Breast.* 2013; **22**(6): 1046–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
10. **F** Copson E, Eccles B, Maishman T, *et al.*: **Prospective observational study of breast cancer treatment outcomes for UK women aged 18–40 years at diagnosis: the POSH study.** *J Natl Cancer Inst.* 2013; **105**(13): 978–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. **F** Rosenberg SM, Ruddy KJ, Tamimi RM, *et al.*: **BRCA1 and BRCA2 Mutation Testing in Young Women With Breast Cancer.** *JAMA Oncol.* 2016.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
12. **F** Partridge AH, Pagani O, Abulkhair O, *et al.*: **First international consensus guidelines for breast cancer in young women (BCY1).** *Breast.* 2014; **23**(3): 209–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
13. **F** Anders CK, Fan C, Parker JS, *et al.*: **Breast carcinomas arising at a young age: unique biology or a surrogate for aggressive intrinsic subtypes?** *J Clin Oncol.* 2011; **29**(1): e18–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
14. **F** Fredholm H, Eaker S, Frisell J, *et al.*: **Breast cancer in young women: poor survival despite intensive treatment.** *PLoS One.* 2009; **4**(11): e7695.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
15. **F** Azim HA Jr, Michiels S, Bedard PL, *et al.*: **Elucidating prognosis and biology of breast cancer arising in young women using gene expression profiling.** *Clin Cancer Res.* 2012; **18**(5): 1341–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
16. **F** BCY 2 - Second Breast Cancer in Young Women Conference 4th–5th November 2014 Dublin, Ireland. *Breast Care (Basel).* 2015; **10**(1): 55–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
17. **F** Marcotte R, Sayad A, Brown KR, *et al.*: **Functional Genomic Landscape of Human Breast Cancer Drivers, Vulnerabilities, and Resistance.** *Cell.* 2016; **164**(1–2): 293–309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
18. Segatto I, Berton S, Sonogo M, *et al.*: **Inhibition of breast cancer local relapse by targeting p70S6 kinase activity.** *J Mol Cell Biol.* 2013; **5**(6): 428–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Segatto I, Berton S, Sonogo M, *et al.*: **p70S6 kinase mediates breast cancer cell survival in response to surgical wound fluid stimulation.** *Mol Oncol.* 2014; **8**(3): 766–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Segatto I, Berton S, Sonogo M, *et al.*: **Surgery-induced wound response promotes stem-like and tumor-initiating features of breast cancer cells, via STAT3 signaling.** *Oncotarget.* 2014; **5**(15): 6267–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Fabris L, Berton S, Citron F, *et al.*: **Radiotherapy-induced miR-223 prevents relapse of breast cancer by targeting the EGF pathway.** *Oncogene.* 2016.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. **F** Alvarez JV, Pan TC, Ruth J, *et al.*: **Par-4 downregulation promotes breast cancer recurrence by preventing multinucleation following targeted therapy.** *Cancer Cell.* 2013; **24**(1): 30–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. **F** Vaidya JS, Wenz F, Bulsara M, *et al.*: **Risk-adapted targeted intraoperative radiotherapy versus whole-breast radiotherapy for breast cancer: 5-year results for local control and overall survival from the TARGIT-A randomised trial.** *Lancet.* 2014; **383**(9917): 603–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Alfredo Budillon**

Experimental Pharmacology Unit, Istituto Nazionale Tumori G. Pascale Napoli, Naples, Italy

Competing Interests: No competing interests were disclosed.

2. **Michael Retsky**

¹ Molecular and Integrative Physical Science, Harvard TH Chan School of Public Health, Boston, MA, USA

² University College London, London, UK

Competing Interests: No competing interests were disclosed.

3. **Andrea Morrione**

Department of Urology and Biology of Prostate Cancer Program, Thomas Jefferson University, Philadelphia, PA, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research